

Decade Long Upsurge In Target Site Pyrethroid Resistance In Bed Bug Populations In The United States

Cari D. Lewis

The University of Tulsa <https://orcid.org/0000-0002-1596-0317>

Brenna A. Levine

The University of Tulsa <https://orcid.org/0000-0002-4326-591X>

Coby Schal

North Carolina State University <https://orcid.org/0000-0001-7195-6358>

Edward L. Vargo

Texas A&M University <https://orcid.org/0000-0002-8712-1248>

Warren Booth (✉ warren-booth@utulsa.edu)

The University of Tulsa <https://orcid.org/0000-0003-2355-0702>

Research Article

Keywords: Cimex lectularius, knockdown resistance, voltage-gated sodium channel, urban evolution

Posted Date: October 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-887494/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Over the past three decades, the bed bug *Cimex lectularius* has resurged as a prominent indoor pest on a global scale. Knockdown-associated insecticide resistance (*kdr*) involving the voltage-gated sodium channel, targeted by organochlorine and pyrethroid insecticides, was first reported in *C. lectularius* within a few years of the widespread use of Dichlorodiphenyltrichloroethane

(DDT) and has been implicated as a significant factor contributing to the species recent resurgence. Since then, selection with pyrethroid insecticides has intensified, yet little is known regarding its short-term impacts on the frequency of *kdr*-associated mutations. Here, we report temporal changes in the frequencies of three *kdr*-associated mutations in *C. lectularius* populations collected across the United States from two time periods, sampled approximately a decade apart. Results reveal a significant increase in the frequencies of *kdr*-associated mutations over this period, and absence of the insecticide-susceptible genotype in recent collections. Furthermore, a significant transition towards infestations possessing multiple *kdr*-associated mutations was observed. These results suggest that the persistent use of pyrethroid insecticides over the past decade continues to impose strong selection pressure on *C. lectularius* populations, driving the proliferation of *kdr*-associated mutations. They demonstrate that, if unabated, strong anthropogenic selection can drive the rapid evolution of adaptive traits.

Introduction

Globally, urban areas are undergoing rapid expansion and are increasing in density (Homer et al. 2020), promoting the establishment and spread of urban-adapted invasive species (Francis and Chadwick 2015; Reed et al. 2020). Within these anthropogenic environments, a myriad of selection pressures are exerted on resident populations, often resulting in the rapid evolution of phenotypes and genotypes (Johnson and Munchi-South 2017; Beans 2019). Ranging from the evolution of unique personality traits and morphologies, to genotypes that confer resistance to pesticides, many such adaptations have been shown to be beneficial, promoting population persistence and spread (Wada-Katsumate et al. 2013; Donihue and Lambert 2015; Winchell et al. 2016; Johnson and Munshi-South 2017; Zalucki and Furlong 2017).

Some pest species share a long history of anthropogenic association (Aplin et al. 2011; Puckett et al. 2016; Potter 2018; Tang et al. 2019). They are readily transported, inadvertently by humans, adapt well to the human-built environment, and are of public health, veterinary, and economic significance (Brenner 1995; Corrigan 2011; Doggett et al. 2018; Hwang et al. 2018; Perron et al. 2018). As such, these pests have been the target of pest control using various pesticide formulations (Rose 2001; Corrigan 2011; Potter 2011; Lee et al. 2018). The persistent use of pesticides exerts selection pressures that can drive the evolution of a suite of resistance-conferring adaptive mechanisms, including mutations in receptor target sites that reduce sensitivity to certain pesticides [e.g., knockdown resistance (*kdr*)] (Davies et al. 2008; Dong et al. 2014).

One group of insecticides, the sodium channel modulators, which includes the organochlorides (e.g., dichloro-diphenyl trichloroethane [DDT]), pyrethrins, and pyrethroids, alter the activity of *para*-type voltage-gated sodium channels (VGSCs), which are necessary for neuronal function. These insecticides maintain the sodium channels in an activated state, preventing cellular repolarization, resulting in paralysis and death (Soderlund and Knipple 2003). Nonsynonymous mutations within the VGSC gene have been identified that reduce *kdr*-associated target-site sensitivity to these insecticides across a variety of arthropod pests (Milani 1956; Martinez-Torres et al. 1998; Soderlund and Knipple 2003; Matowo et al. 2017). Despite being well-documented, temporal data on the distribution and frequency of *kdr*-associated mutations are surprisingly lacking for urban pest species in general, with the exception of *Anopheles* and *Aedes* mosquitoes (Mathias et al. 2011; Abdalla et al. 2014; Linss et al. 2014; Aguirre-Obando et al. 2016; Foster et al. 2016; Ismail et al. 2018; Amoudji et al. 2019; Biduda et al. 2019; Mandeng et al. 2019; Tancredi et al. 2020; Baltzegar et al. 2021) and house flies (Freeman et al. 2019). The lack of understanding of the evolutionary response of populations to the prolonged use of these insecticides may impede the continued development of effective control strategies over time.

The common bed bug, *C. lectularius*, is an obligate hematophagous ectoparasite, primarily of humans, that has re-emerged in recent decades to become a prominent indoor pest on a near global scale. This resurgence has been aided by an increase in travel (Potter et al. 2010), exchange of secondhand furniture, ineffective pest control, and the evolution of mechanisms conferring insecticide resistance (Boase 2001; Doggett et al. 2004; Romero et al. 2007; Reinhardt et al. 2008; Zhu et al. 2010; Gordon et al. 2014). Within *C. lectularius* populations, various insecticide resistance mechanisms have evolved (Dang et al. 2017), with *kdr* mutations particularly well documented. First reported shortly after the widespread implementation of DDT (Busvine 1958), *kdr* mutations have been reported in populations around the globe (Romero et al. 2007; Yoon et al. 2008; Seong et al. 2010; Tawatsin et al. 2011; Wang and Wen 2011; Durand et al. 2012; Tomita et al. 2012; Booth et al. 2015; Dang et al. 2015; Lilly et al. 2015, 2017; Palenchar et al. 2015; Raab et al. 2016; Balvin and Booth 2018; Holleman et al. 2019; Cho et al. 2020; Vander Pan et al. 2020; Akhouni et al. 2021). Nevertheless, organochlorides, pyrethroids, and pyrethrins, remain commonly used insecticides on a global scale, and thus persist as selection pressures promoting *kdr*-associated mutations that confer resistance to these insecticides.

In *C. lectularius*, three nonsynonymous mutations in the VGSC-gene (V419L, L925I, and I936F) have been found to be associated with *kdr* resistance (Yoon et al. 2008; Dang et al. 2015). The population frequency and geographic distribution of these mutations have been reported from a variety of locations, including the U.S. (Yoon et al. 2008; Zhu et al. 2010; Raab et al. 2016; Holleman et al. 2018), Europe (Durand et al. 2012; Booth et al. 2015; Balvin and Booth 2016; Vander Pan et al. 2020; Akhouni et al. 2021), the Middle East (Palenchar et al. 2015), Asia (Seong et al. 2010; Tomita et al. 2012; Cho et al. 2020), and Australia (Dang et al. 2015). However, we are unaware of any studies that assessed temporal changes in the frequencies of *kdr*-associated mutations at any landscape level. Although it would be most instructive to track the frequency of *kdr* mutations over time within the same populations, this design is particularly challenging with residential infestations due to their ephemerality and turnover of residents. Here, we report temporal changes in the frequency of the three *kdr*-associated mutations from populations of *C. lectularius* collected across the United States at two time periods approximately 10-years apart. We hypothesized that both the number of infestations exhibiting *kdr*-associated mutations, and the number of adaptive mutations per individual bed bug would increase over time, due to selection imposed by the frequent and widespread use of pyrethrin and pyrethroid insecticides.

Materials And Methods

Bed bugs were collected during two separate time periods: 30 August 2005 to 20 December 2009 (electronic supplementary material, Table S1), and 27 June 2018 to 27 December 2019 (electronic supplementary material, Table S2), representing 394 unique populations (161 in 2005–2009 and 233 in 2018–2019). Collections spanned 38 U.S. states (Table 1, Fig. 1a) and were made by pest management companies (electronic supplementary material, Table S3). Upon collection, specimens were preserved in 100% ethanol and stored at -20°C until DNA extraction. Genomic DNA was extracted using the Qiagen DNeasy Blood and Tissue Kit (QIAGEN, Germantown, MD).

Table 1
Distribution of *Cimex lectularius* populations by state for each sampling period.

State	2005–2009	2018–2019
Alabama	1	-
Alaska	-	3
Arizona	1	-
Arkansas	2	-
California	7	27
Colorado	3	-
Florida	13	10
Georgia	2	2
Hawaii	1	-
Idaho	-	1
Illinois	3	79
Indiana	5	4
Iowa	5	-
Kansas	1	5
Kentucky	3	-
Louisiana	1	-
Maine	1	-
Maryland	4	-
Massachusetts	2	-
Michigan	3	-
Minnesota	1	-
Missouri	1	4
New Jersey	12	20
New York	3	1
Nebraska	3	-
Nevada	5	-
North Carolina	16	-
Ohio	20	33
Oklahoma	3	9
Oregon	4	15
Pennsylvania	17	5
South Carolina	1	-
Tennessee	-	1
Texas	4	13
Utah	3	-
Virginia	5	-
Washington	3	-
Wisconsin	2	1
Total	161	233

Due to the highly inbred nature of *C. lectularius* populations (Booth et al. 2012, 2018; Saenz et al. 2012), which results from population foundation by a single gravid female or small group of highly related individuals, the genetic diversity and hence *kdr*-associated mutation profile of a given population can be

determined through the sequencing of a single individual. Therefore, each population is represented here by a single specimen. Two genomic fragments that contain the three *kdr*-associated mutations were PCR-amplified and sequenced using primer combinations BBParaF1/BBParaR1 (V419L) and BBParaF3/BBParaR3 (L925I, I936F), following the methods of Holleman et al. (2018). The resulting chromatograms were visualized and scored using CLC Genomics Workbench (QIAGEN). Knockdown resistance genotype designations are reported as follows: wild type (pyrethroid-susceptible in the absence of other resistance mechanisms; L925, V419, I936), L925I (mutant L925I [leucine (L) replaced by isoleucine (I)], wild type V419&I936), L925I&V419L (mutations in regions L925I and V419L [valine (V) replaced by leucine (L)], wild type I936), L925I&I936F (mutations in both L925I and I936F [isoleucine (I) replaced by phenylalanine (F)], wild type V419), and I936F (mutant I936F, wild type L925&V419). The V419L (wild type L925, I936, mutant V419L) genotype was not detected in any population screened. Where the mutations are heterozygous, these are reported with a superscript “*het*”.

Genotype frequencies were compared using Chi-square tests. Additionally, the overall homozygous genotype frequencies of populations present in the same U.S. states across both sample periods were compared.

Results

Only samples with unambiguous sequences for both PCR fragments were analyzed in this study. Of the 161 populations sampled between 2005 and 2009, genotypes L925I and L925I&V419L accounted for 36% and 50% of the populations, respectively (Table 2 and Fig. 1a). The wild type (‘susceptible’) genotype was found in 2.5% of the populations. Approximately 10% of samples screened were found to be L925I&V419L^{het}, while the I936F, L925I^{het}&I936F^{het}, and I936F^{het} genotypes each comprised < 0.1% of the remaining infestations (Table 2 and Fig. 1a). In contrast, genotype L925I&V419L predominated in samples collected between 2018 and 2019, accounting for 84% of the populations, while genotypes L925I (~ 5%), L925I^{het} (~ 1%), L925I&V419L^{het} (~ 9%), and L925I^{het}&V419L (~ 1%) comprised the remaining populations (Table 2 and Fig. 1a). The wild type and I936F genotypes were not detected in the 2018–2019 populations.

Table 2
Frequency of *kdr*-associated genotypes reported in *Cimex lectularius* populations sampled in the United States sampled between 2005–2009 and 2018–

Sample periods	No. of States	No. of pops.	No. of individuals	<i>kdr</i> genotypes							
				Wild type	L925I	L925I ^{het}	L925I&V419L	L925I&V419L ^{het}	L925I ^{het} &V419L	I936F	I936F ^{het}
2005–2009	35	161	161	4 (0.02)	57 (0.35)	1 (0.006)	81 (0.50)	16 (0.10)	0 (0.00)	1 (0.006)	1 (0.006)
2018–2019	18	233	233	0 (0.00) [5.85]*	12 (0.05) [60.32]**	2 (0.01) [0.07]	194 (0.84) [49.04]**	22 (0.09) [0.03]	3 (0.01) [2.09]	0 (0.00) [1.45]	0 (0.00) [1.45]

For each genotype, the number of observations is reported. Parentheses indicate the frequency of each genotype and the brackets report the χ^2 statistic when = 394)=[χ^2 stat], $p > 0.05$ or * $p < 0.05$, ** $p < 0.001$.

Temporally, there were significant differences in the frequencies of all homozygous [wild type, $p < 0.05$; L925I, $p < 0.001$; L925I&V419L, $p < 0.001$], but not the heterozygous genotypes and I936F, likely due to their low frequency across datasets (Table 2). Comparisons conducted using only populations sampled in the same U.S. states in both timeframes again revealed significant differences in the frequencies of all homozygous genotypes (electronic supplementary material, Table S4).

Discussion

By surveying 394 distinct populations of *C. lectularius* across the U.S., collected from two time-periods separated by approximately a decade, this study represents the most comprehensive screening to date of *kdr*-associated mutation frequencies for the species. To our knowledge, this is also the first temporal comparison of *kdr* mutation frequencies in any urban pest insect in the U.S., outside of house flies collected from animal production facilities (Freeman et al. 2019) and mosquitoes (Mathias et al. 2011; Abdalla et al. 2014; Linss et al. 2014; Aguirre-Obando et al. 2016; Foster et al. 2016; Ismail et al. 2018; Amoudji et al. 2019; Biduda et al. 2019; Mandeng et al. 2019; Tancredi et al. 2020; Baltzegar et al. 2021). Comparing our homozygous populations from 2005–2009 (N = 143) to those of Zhu et al. (2010) [N = 93] revealed significant differences in *kdr* genotype frequencies (wild type: $\chi^2(1, n = 236) = 9.11, p < 0.05$; L925I: $\chi^2(1, n = 236) = 0.45, p > 0.05$; L925I&V419L: $\chi^2(1, n = 236) = 7.25, p < 0.05$; V419L: $\chi^2(1, n = 236) = 4.67, p < 0.05$), supporting the lower baseline frequency of *kdr* mutations present in the U.S. in 2006–2007 (Zhu et al. 2010), and a reduction in wild type and increase in L925I&V419L by 2008–2009 when the majority of our data were collected. Prior to our study, the most recent study conducted in the U.S. reported *kdr* genotype frequencies for populations collected across Midwestern states between 2014 and 2015 (Holleman et al. 2018), with the frequencies of L925I (~ 21%) and L925I&V419L (~ 76%) intermediate between the data sets presented here (Fig. 1b), as expected for a shift towards increased resistance. Thus, our results, when viewed together with those of Zhu et al. (2010) and Holleman et al (2018), reveal a significant increase in the frequency of *kdr* mutations over approximately a decade, with a shift towards multiple *kdr* mutations per population (Fig. 1b). This is likely in response to persistent selection driven by the long-term use of pyrethroid insecticides.

Prior bioassay-based research in *C. lectularius* has linked genotypes L925I and L925I&V419L to increased levels of pyrethroid resistance (Zhu et al. 2010), including > 12,765-fold deltamethrin resistance (Romero et al. 2007), when compared to a susceptible wild type population. Furthermore, the presence of multiple *kdr* mutations has been associated with elevated levels of pyrethroid resistance in other pest insects (Soderlund 2008; Abdalla et al. 2014). Thus, we presume that a higher proportion of the bed bugs collected in 2018–2019 would exhibit greater resistance to pyrethroids than those collected a decade earlier.

Comparable findings of temporal change in *kdr* mutations have been reported in mosquito populations exposed to the prolonged use of insecticides. Congruent with the results presented here, several long-term studies (> 10 years) showed an increase in *kdr* mutation frequencies in populations of *Anopheles gambiae* (Mathias et al. 2011) and *Aedes aegypti* (Linns et al. 2014; Baltzegar et al. 2021), from different geographic regions. Recently, Baltzegar et al (2021), documented the complete replacement of the wild type pyrethroid-susceptible genotype, with either single or double *kdr*-mutants in *Ae. Aegypti* mosquitos sampled in Peru, with the replacement occurring over a period of 13 years following the introduction of a city-wide pyrethroid insecticide control strategy. Similarly, the malaria vector, *An. arabiensis*, was shown to respond to strong selection for deltamethrin resistance over four years, which was correlated with the L1014F *kdr* mutation (Ismail et al. 2018). Verhaeghen et al. (2010) documented a five-year increase in *kdr* mutations in *An. gambiae* from multiple geographic locations, and similar region-specific fluctuations in *kdr* mutation frequencies have been shown in *Ae. albopictus* at multiple time intervals (Tancredi et al. 2020). A significant increase was also reported for the 1b resistance allele in house flies collected over a 10-year period from animal production facilities across five U.S. States (Freeman et al. 2019). While these studies span different regional pest management practices, the trend of increasing *kdr* mutation frequencies is consistent. However, a unique characteristic of *C. lectularius* is the high level of intra-population inbreeding, and an apparent lack of significant levels of gene flow (Booth et al. 2012, 2015, 2018; Saenz et al. 2012; Fountain et al. 2014). As a result, resistance alleles rapidly propagate and are then spread in a fixed-state through propagule transfer. While there is still uncertainty surrounding the fitness costs of resistance in *C. lectularius* (Polanco et al. 2011; Gordon et al. 2015), the frequency and distribution of *kdr* mutations and the prevalence of the species suggests this may be minimal.

The zygosity of *kdr* mutations and its impact on insecticide resistance adds additional complexity. For example, in the house fly, *Musca domestica*, mutations exhibit a recessive pattern of inheritance (Rinkevich et al. 2012). In contrast, polygenic modes of inheritance have been documented in the codling moth, *Cydia pomonella* (Bouvier et al. 2001), with both recessive and incomplete dominance described. Unlike Zhu et al. (2010), this study reports the zygosity of the *kdr*-associated mutations detected. This information may be relevant for control, for while the level of resistance exhibited by *C. lectularius* heterozygous for a given mutation is unclear, it has been suggested that heterozygotes may show incomplete dominance (Romero et al. 2007). Heterozygotes may therefore exhibit intermediate levels of resistance between wild type and homozygous mutants, although this has yet to be confirmed. If *kdr* exhibits incomplete dominance in *C. lectularius*, heterozygotes would be expected in areas undergoing genotype replacement from wild type to resistant, either through population spread or continued pyrethroid applications. However, bed bugs heterozygous for *kdr* mutations would also be expected to succumb to control more rapidly than homozygous mutant individuals. In both time periods sampled, L925I&V419F^{het} was the most common heterozygous genotype observed, representing 89% (2005–2009) and 78% (2018–2019) of the heterozygous individuals. Given the short generation time of *C. lectularius*, high fecundity, and lack of obvious inbreeding depression (Fountain et al. 2014), the transition within a population to the double homozygous mutant (L925I&V419L) would be expected to occur rapidly following selection by insecticide treatment.

In summary, the significant temporal shift towards increased frequencies of *kdr* mutations reveals how anthropogenic selection can influence the evolution of a species of public health, veterinary, and economic significance, and suggests that indoor pest insects may represent ideal model systems to study the effects of anthropogenic influences on the evolution of urban species. Furthermore, these findings are practically significant in the development of management strategies for the control of *C. lectularius* populations and those of other urban and indoor pest species commonly controlled with pesticides.

Declarations

Acknowledgements

We thank all pest control companies for providing samples (electronic supplementary material, Table 3) and Rick Santangelo for the management of the 2005-2009 samples.

Funding

Funding was provided by competitive grants from USDA-NIFA (AFRI 2008-35302- 04565), NSF (DEB-1754394), Oklahoma Center for the Advancement of Science and Technology (HR13-211 and HR20-142), the National Pest Management Foundation, the Blanton J. Whitmire endowment at North Carolina State University, the Urban Entomology Endowment at Texas A&M University, and student research grants awarded by The University of Tulsa.

Conflicts of Interest

The authors declare no conflicts of interest

Availability of Data and Material

Data supporting the results can be found in the Online Supplementary Material.

Code Availability

Not applicable

Authors' contributions

CDL, WB, BAL, ELV, and CS designed the experiment. CDL and WB collected and analyzed the data. All authors contributed to the writing of the manuscript and gave final approval for publication.

References

1. Abdalla H, Wilding CS, Nardini L, Pignatelli P, Koekemoer LL, Ranson H, Coetzee M (2014) Insecticide resistance in *Anopheles arabiensis* in Sudan: temporal trends and underlying mechanisms. *Parasit Vectors* 7:123
2. Aguirre-Obando OA, Pietrobon AJ, Dalla Bona AC, Navarro-Silva MA (2016) Contrasting patterns of insecticide resistance and knockdown resistance (*knr*) in *Aedes aegypti* populations from Jacarezinho (Brazil) after a Dengue outbreak. *Med Vet Entomol* 60:94–100
3. Akhouni M, Chebbah D, Sereno D, Marteau A, Jan J, Bruel C, Elissa N, Izri A (2021) Widespread mutations in voltage-gated sodium channel gene of *Cimex lectularius* (Hemiptera: Cimicidae) population in Paris. *Int J Environ Res Public Health* 18:407
4. Amoudji AD, Ahadji-Dabla KM, Hien AS, Apétogbo YG, Yaméogo B, Soma DD, Bamogo R, Atcha-Oubou RT, Dabiré RK, Ketoh GK (2019) Insecticide resistance profiles of *Anopheles gambiae* s.l. in Togo and genetic mechanisms involved, during 3-year survey: is there any need for resistance management? *Malar J* 18:117
5. Aplin KP, Suzuki H, Chinen AA, Chesser RT, ten Have J, Donnellan SC, Austin J, Frost A, Gonzalez JP, Herbreteau V et al (2011) Multiple geographic origins of commensalism and complex dispersal history of black rats. *PLoS ONE* 6:e26357
6. Balvin O, Booth W (2018) Distribution and frequency of pyrethroid resistance-associated mutations in host lineages of the bed bug (Hemiptera: Cimicidae) across Europe. *J Med Entomol* 55:923–928
7. Baltzegar J, Vella M, Gunning C, Vasquez G, Astete H, Stell F, Scott TW, Lenhart A, Lloyd AL, Morrison A, Gould F (2021) Rapid evolution of knockdown resistance haplotypes in response to pyrethroid selection in *Aedes aegypti*. *Evol. Appl.* <https://doi.org/10.1111/eva.13269>
8. Beans C (2019) Cities serve as testbeds for evolutionary change. *Proc. Natl. Acad. Sci. U.S.A.* 116:2787–2790
9. Biduda S, Lin C, Saleh F, Konradsen F, Hansson H, Schiøler KL, Alifrangis M (2019) Temporal pattern of mutations in the knockdown resistance (*knr*) gene of *Aedes aegypti* mosquitoes sampled from southern Taiwan. *Am J Trop Med Hyg* 101:973–975
10. Boase C (2001) Bedbugs—back from the brink. *Pestic. Outlook*. August:159–162
11. Booth W, Schal C, Vargo EL (2018) Population Genetics. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd. Oxford, UK, pp 173–182
12. Booth W, Saenz VL, Santangelo RG, Wang C, Schal C, Vargo EL (2012) Molecular markers reveal infestation dynamics of the bed bug (Hemiptera: Cimicidae) within apartment buildings. *J Med Entomol* 49:535–546
13. Booth W, Balvin O, Vargo EL, Vilimova J, Schal C (2015) Host association drive genetic divergence in the bed bug, *Cimex lectularius*. *Mol Ecol* 24:980–992
14. Bouvier J, Bues R, Boivin T, Boudinhon L, Beslay D, Sauphanor B (2001) Deltamethrin resistance in codling moth (Lepidoptera: Tortricidae): inheritance and number of genes involved. *Heredity* 87:456–462
15. Brenner RJ (1995) Economics and medical importance of German cockroaches. In: Rust MK, Owens JM, Reiersen DA (eds) *Understanding and Controlling the German Cockroach*. Oxford University Press, Oxford, pp 77–92
16. Busvine JR (1958) Insecticide-resistance in bed-bugs. *Bull World Health Organ* 19:1041–1052
17. Cho S, Kim H, Chong S, Klein TA, Kwon DH, Lee SH, Kim JH (2020) Monitoring of pyrethroid resistance allele frequency in the common bed bug (*Cimex lectularius*) in the Republic of Korea. *Korean J Parasitol* 58:99–102
18. Corrigan RM (2011) Rats and mice. In: Hedges SA (ed) *The Mallis Handbook of Pest Control*, 10th edition. The Mallis Handbook and Technical Training Co., Cleveland, OH, USA. pp. 11–48
19. Davies TGE, O'Reilly AO, Field LM, Wallace BA, Williamson MS (2008) Knockdown resistance to DDT and pyrethroids: from target-site mutations to molecular modelling. *Pest Manag Sci* 64:1126–1130
20. Dang K, Toi CS, Lilly DG, Bu W, Doggett SL (2015) Detection of knockdown resistance mutations in the common bed bug, *Cimex lectularius* (Hemiptera: Cimicidae), in Australia. *Pest Manag. Sci.* 71:914–922
21. Dang K, Doggett SL, Veera Singham G, Lee C-Y (2017) Insecticide resistance and resistance mechanisms in bed bugs, *Cimex spp.* (Hemiptera: Cimicidae). *Parasit Vectors* 10:1–31
22. Doggett SL, Geary MJ, Russell RC (2004) The resurgence of bed bugs in Australia, with notes on their ecology and control. *Environ Health* 4:30–38
23. Doggett SL, Miller DM, Vail K, Wilson MS (2018) Fiscal Impacts. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd. Oxford, UK, pp 139–146
24. Donihue CM, Lambert MR (2015) Adaptive evolution in urban ecosystems. *AMBIO* 44:194–203
25. Dong K, Du Y, Rinkevich F, Nomura Y, Xu P, Wang L, Silver K, Zhorov BS (2014) Molecular biology of insect sodium channels and pyrethroid resistance. *Insect Biochem Mol Biol* 50:1–17
26. Durand R, Cannet A, Berdjane Z, Bruel C, Haouchine D, Delaunay P, Izri A (2012) Infestation by pyrethroids resistant bed bugs in the suburb of Paris. *France Parasite* 19:381–387
27. Foster GM, Coleman M, Thomsen E, Ranson H, Yangalbé-Kalnone E, Moundai T, Kodindo ID, Nakebang A, Mahamat A, Peka M et al (2016) Spatial and temporal trends in insecticide resistance among malaria vectors in Chad highlight the importance of continual monitoring. *PLoS ONE* 11:e0155746
28. Fountain T, Butlin RK, Reinhardt K, Otti O (2014) Outbreeding effects in an inbreeding insect, *Cimex lectularius*. *Ecol Evol* 5:409–418
29. Francis RA, Chadwick MA (2015) Urban invasions: non-native and invasive species in cities. *Geography* 100:144–151

30. Freeman JC, Ross DH, Scott JG (2019) Insecticide resistance monitoring of house fly populations from the United States. *Pestic Biochem Physiol* 158:61–68
31. Gordon JR, Goodman MH, Potter MF, Haynes KF (2014) Population variation in and selection for resistance to pyrethroid-neonicotinoid insecticides in the bed bug. *Sci Rep* 4:3836
32. Gordon JR, Potter MF, Haynes KF (2015) Insecticide resistance in the bed bug comes with a cost. *Sci Rep* 5:10807
33. Holleman JG, Robison GA, Bellovich IJ, Booth W (2019) Knockdown resistance-associated mutations dominate populations of the common bed bug (Hemiptera: Cimicidae) across the south-central United States. *J Med Entomol* 56:1678–1683
34. Homer C, Dewitz J, Jin S, Xian G, Costello C, Danielson P, Gass L, Funk M, Wickham J, Stehman S et al (2020) Conterminous United States land cover change patterns 2001–2006 from the 2016 National Land Cover Database. *ISPRS J. Photogramm. Remote Sens.* 162:184–199
35. Hwang SJE, Doggett SL, Fernandez-Penas P (2018) Dermatology and Immunology. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd., Oxford, pp 107–114
36. Ismail BA, Kafy HT, Sulieman JE, Subramaniam K, Thomas B, Mnzava A, Kassim NFA, Ahmad AH, Knox TB, Kleinschmidt I et al (2018) Temporal and spatial trends in insecticide resistance in *Anopheles arabiensis* in Sudan: outcomes from an evaluation of implications of insecticide resistance for malaria vector control. *Parasit Vectors* 11:122
37. Johnson MTJ, Munshi-South J (2017) Evolution of life in urban environments. *Science* 358:1–11
38. Lee C-Y, Miller DM, Doggett SL (2018) Chemical control. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd., Oxford, pp 285–305
39. Linss JGB, Brito LP, Garcia GA, Araki AS, Bruno RV, Lima JBP, Valle D, Martins AJ (2014) Distribution and dissemination of the Val1016Ile and Phe1534Cys *kdr* mutations in *Aedes aegypti* Brazilian natural populations. *Parasit Vectors* 7:25
40. Lilly DG, Zalucki MP, Orton CJ, Russell RC, Webb CE, Doggett SL (2015) Confirmation of insecticide resistance in *Cimex lectularius* Linnaeus (Hemiptera: Cimicidae) in Australia. *Austral Entomol* 54:96–99
41. Lilly DG, Dang K, Webb CE, Doggett SL (2017) Are Australian field-collected strains of *Cimex lectularius* and *Cimex hemipterus* (Hemiptera: Cimicidae) resistant to deltamethrin and imidacloprid as revealed by topical assay? *Austral Entomol* 57:77–84
42. Mandeng SE, Awono-Ambene HP, Bigoga JD, Ekoko WE, Binyang J, Piameu M, Mbakop LR, Fesuh BN, Mvondo N, Tabue R et al (2019) Spatial and temporal development of deltamethrin resistance in malaria vectors of the *Anopheles gambiae* complex from North Cameroon. *PLoS ONE* 14:e0212024
43. Martinez-Torres D, Chandre F, Williamson MS, Darriet F, Bergé JB, Devonshire AL, Guillet P, Pasteur N, Pauron D (1998) Molecular characterization of pyrethroid knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* s.s. *Insect Mol. Biol.* 7:179–184
44. Mathias DK, Ochomo E, Atieli F, Ombok M, Bayoh MN, Olang G, Muhia D, Kamau L, Vulule JM, Hamel MJ et al (2011) Spatial and temporal variation in the *kdr* allele L1014S in *Anopheles gambiae* s.s. and phenotypic variability in susceptibility to insecticides in Western Kenya. *Malar J* 10:10
45. Matowo NS, Munhenga G, Tanner M, Coetzee M, Feringa WF, Ngowo HS, Koekemoer LL, Okumu FO (2017) Fine-scale spatial and temporal heterogeneities in insecticide resistance profiles of the malaria vector, *Anopheles arabiensis*, in rural south-eastern Tanzania. *Wellcome Open Res* 2:96
46. Milani R (1956) Comportamento mendeliano della resistenza alla ziaione abbattente del DDT e correlazione tra abbattimento e mortalità in *Musca domestica* L. *Rend Ist Sup Sanit* 19:1107–1143
47. Palenchar DJ, Gellatly KJ, Yoon KS, Mumcuoglu KY, Shalom U, Clark JM (2015) Quantitative sequencing for the determination of *kdr*-type resistance allele (V419L, L925I, I936F) frequencies in common bed bug (Hemiptera: Cimicidae) populations collected from Israel. *J Med Entomol* 52:1018–1027
48. Perron S, Hamelin G, Kaiser D (2018) Mental Health Impacts. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd. Oxford, UK, pp 127–131
49. Polanco AM, Brewster CC, Mille DM (2011) Population growth potential of the bed bug, *Cimex lectularius* L.: a life table analysis. *Insects* 2:173–185
50. Potter MF, Rosenberg B, Henriksen M (2010) Bugs without borders: defining the global bed bug resurgence. *Pestworld*. Sept./Oct:8–20
51. Potter MF (2011) The history of bed bug management—with lessons from the past. *Am Entomol* 57:14–25
52. Potter MF (2018) Bed bugs through history. In: Doggett SL, Miller DM, Lee C-Y (eds) *Advances in the Biology and Management of Modern Bed Bugs*. John Wiley & Sons Ltd. Oxford, UK, pp 9–23
53. Puckett EE, Park J, Combs M, Blum MJ, Bryant JE, Caccone A, Costa F, Deinum EE, Esther A, Himsforth CG et al (2016) Global population divergence and admixture of the brown rat (*Rattus norvegicus*). *Proc. R. Soc. Lond. [Biol]* 283:1841
54. Raab RW, Moore JE, Vargo EL, Rose L, Raab J, Culbreth M, Burzumato G, Koyee A, McCarthy B, Raffaele J et al (2016) New introductions, spread of existing matrilines, and high rates of pyrethroid resistance result in chronic infestations of bed bugs (*Cimex lectularius* L.) in lower-income housing. *PLoS ONE* 11:e0117805
55. Reed EMX, Serr ME, Maurer AS, Burford Reiskind MO (2020) Gridlock and beltways: the genetic context of urban invasions. *Oecol* 192:615–628
56. Reinhardt K, Harder A, Holland S, Hooper J, Leake-Lyall C (2008) Who knows the bed bug? Knowledge of adult bed bug appearance increases with people's age in three counties of Great Britain. *J Med Entomol* 45:956–958
57. Rinkevich FD, Hedtke SM, Leichter CA, Harris SA, Su C, Brady SG, Taskin V, Qui X, Scott JG (2012) Multiple origins of *kdr*-type resistance in the house fly, *Musca domestica*. *PLoS ONE* 7:e52761
58. Romero A, Potter MF, Potter DA, Haynes KF (2007) Insecticide resistance in the bed bug: a factor in the pest's sudden resurgence? *J Med Entomol* 44:175–178
59. Rose RI (2001) Pesticides and public health: integrated methods of mosquito management. *Emerg Infect Diseases* 7:17–23

60. Saenz VL, Booth W, Schal C, Vargo EL (2012) Genetic analysis of bed bug populations reveals small propagule size within individual infestations but high genetic diversity across infestations from the eastern United States. *J Med Entomol* 49:865–875
61. Seong KM, Lee D-Y, Yoon KS, Kwon DH, Kim HC, Klein YA, Clark JM, Lee SH (2010) Establishment of quantitative sequencing and filter contact vial bioassay for monitoring pyrethroid resistance in the common bed bug. *Cimex lectularius* *J Med Entomol* 47:592–599
62. Soderlund DM (2008) Pyrethroids, knockdown resistance and sodium channels. *Pest Manag Sci* 64:610–616
63. Soderlund SM, Knipple DC (2003) The molecular biology of knockdown resistance to pyrethroid insecticides. *Insect Biochem Mol Biol* 33:563–577
64. Tang Q, Bourguignon T, Willenmse L, De Coninck E, Evans T (2019) Global spread of the German cockroach, *Blattella germanica*. *Biol Invasions* 21:693–707
65. Tancredi A, Papandrea D, Marconcini M, Carballar-Lejarazu R, Casas-Martinez M, Lo E, Chen X, Malacrida AR, Bonizzoni M (2020) Tracing temporal and geographic distribution of resistance to pyrethroids in the arboviral vector *Aedes albopictus*. *PLoS Negl Trop Dis* 14:e0008350
66. Tawatsin A, Thavara U, Chompoonsri J, Phusup Y, Jonjang N, Khumsawads C, Bhakdeenuan P, Sawanpanyalert P, Asavadachanukorn P, Mulla MS et al (2011) Insecticide resistance in bedbugs in Thailand and laboratory evaluation of insecticides for the control of *Cimex hemipterus* and *Cimex lectularius* (Hemiptera: Cimicidae). *J Med Entomol* 48:1023–1030
67. Tomita T, Komagata O, Kasai S, Itokawa K, Watanabe M, Yaguchi N, Adachi M, Yoshida M, Kimura G, Kobayashi M (2012) Nationwide survey on pyrethroid-susceptibility of the bed bug, *Cimex lectularius*. *Jpn Soc Med Entomol Zool* 63:85
68. Vander Pan A, Kuhn C, Schmolz E, von Samson-Himmelstjerna G, Krücken J (2020) Detection of target-site and metabolic resistance to pyrethroids in the bed bug *Cimex lectularius* in Berlin. Germany *Int J Parasitol Drugs Drug Resist* 14:274–283
69. Verhaeghen K, Bortel WV, Roelants P, Okello PE, Talisuna A, Coosemans M (2010) Spatio-temporal patterns in *kdr* frequency in permethrin and DDT resistant *Anopheles gambiae* s.s. from Uganda. *Am J Trop Med Hyg* 82:566–573
70. Wada-Katsumata A, Silverman J, Schal C (2013) Changes in taste neurons support the emergence of an adaptive behavior in cockroaches. *Science* 340:972–975
71. Wang C, Wen X (2011) Bed bug infestations and control practices in China: implications for fighting the global bed bug resurgence. *Insects* 2:83–95
72. Winchell KM, Reynolds RG, Prado-Irwin SR, Puente-Rolón AR, Revell LJ (2016) Phenotypic shifts in urban areas in the tropical lizard *Anolis cristatellus*. *Evolution* 70:1009–1022
73. Yoon KS, Kwon DH, Strycharz JP, Hollingsworth CS, Lee SH, Clark JM (2008) Biochemical and molecular analysis of deltamethrin resistance in the common bed bug (Hemiptera: Cimicidae). *J Med Entomol* 45:1092–1101
74. Zalucki MP, Furlong MJ (2017) Behavior as a mechanism of insecticide resistance: evaluation of the evidence. *Curr Opin Insect Sci* 21:19–25
75. Zhu F, Wigginton J, Romero A, Moore A, Ferguson K, Palli R, Potter MF, Haynes KF, Palli SR (2010) Widespread distribution of knockdown resistance mutations in the bed bug, *Cimex lectularius* (Hemiptera: Cimicidae), populations in the United States. *Arch Insect Biochem Physiol* 73:245–257

Figures




Figure 1

(a) Genotype distribution of *kdr*-associated mutations from *Cimex lectularius* sampled across the U.S. Pies represent genotype frequencies within each state and the number below each indicates the number of populations sampled. (b) Change in genotype frequencies over time of three homozygous genotypes – wild type, 925, 925&419. Colors correspond to genotypes in Fig. 1a. 2007-2008 data from Zhu et al. (2010), 2014–2015 from Holleman et al. (2018).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supp.mat.Tables.docx](#)