Predefined and data driven CT densitometric features predict critical illness and hospital length of stay in COVID-19 patients

Tamar Shalmon

University Health Network

Pascal Salazar

Vital Images

Miho Horie

University Health Network

Kate Hanneman

University Health Network

Mini Pakkal

University Health Network

Vahid Anwari

University Health Network

Jennifer Fratesi (✉ Jennifer.fratesi@uhn.ca)

University Health Network

Research Article

Keywords: density, data, patient, driven, histograms, hospital

Posted Date: September 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-891706/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The aim of this study was to compare predefined and data-driven parameters of whole lung CT density histograms to predict critical illness outcome and hospital length of stay in a cohort of 80 COVID-19 patients. CT chest images on segmented lungs were retrospectively analyzed. Functional Principal Component Analysis (FPCA) was used to find the main modes of variations on CT density histograms (F1,F2,F3,F4) in the whole patient cohort. The data driven and a priori CT density features, the CT severity score, the COVID-GRAM score and the patient clinical data were assessed for predicting the patient outcome using logistic regression models stratified for contrast enhanced CT and non-enhanced CT, and survival analysis. ROC analysis identified as best predictors of critically ill status: 87.5th percentile CT density (Q875) - AUC: 0.88 95%CI (0.79 0.94), F1-CT - AUC: 0.87 (0.77 0.93) Standard Deviation (SD-CT)- AUC: 0.86 (0.73, 0.93). Multivariate models combining CT-density predictors and Neutrophil-Lymphocyte Ratio showed the highest accuracy with cross-validated AUCs in the 0.91–0.92 range for contrast CT and 0.82–0.88 range for non-contrast CT. SD-CT, Q875 and F1 score were significant predictors of hospital length of stay while controlling for hospital death using competing risks models. Predefined and data-driven parameters of lung CT density histograms can predict critical illness and length of stay to guide management and resources. FPCA method can be used to interpret the CT density histogram variation in a patient cohort and to extract predictive features with minimal a priori knowledge.

Introduction

Qualitative and semi-quantitative scoring methods using high resolution computed tomography (HRCT) have been increasingly applied for diagnostic, severity assessment or prognosis of interstitial lung diseases such as idiopathic lung fibrosis (IPF), chronic obstructive pulmonary diseases (COPD), and more recently for severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS)^{1–3} and COVID-19 pneumonia^{4,5}. However, the intra and inter-reader variability remain a substantial limitation. Consequently, alternative objective quantitative methods have been actively explored. Volumetric quantitative CT have been used to predict lung fibrosis outcome^{6–8} for patient stratification or prognosis in COPD, systemic sclerosis, early chronic lung allograft dysfunction in lung transplant patients⁹, ARDS, and recently in COVID-19 patients^{10–13}. Two main approaches coexist in Quantitative volumetric CT densitometry: 1. The (“first order”) radiomic method uses the whole lung CT density histogram to extract statistical features such as mean lung attenuation (MLA) standard deviation, skewness and kurtosis, quantile predictors (median, 75th percentile density, etc.) or more advanced features such as entropy¹⁴. 2. The multi-threshold method uses the whole lung divided in regions of increasing CT density ranges with predefined cutoff values. CT density ranges represent either functional versus non-functional lung regions, or regions associated with emphysema, ground glass opacification, consolidation, etc. Derived features are volumes or volume ratios of different CT density ranges^{10,15}.

Both first order radiomic and multi-thresholds methods for CT densitometry have inherent limitations: in the radiomic method, the predefined features mean lung attenuation (MLA), standard deviation,

skewness, kurtosis, or entropy were originally defined for simple formal probability distributions and they are crude descriptors of the often-complex multi-peak CT histograms of the lungs. For example, the kurtosis of a CT density histogram represents both its 'peakedness' and the thickness of the histogram left and right tails, making kurtosis hardly interpretable. In the multi-threshold method, multiple non-arbitrary cutoff values are hard to establish. As an example, it took many years to reach a consensus for a single cutoff for emphysema low attenuation area percentage (LAA%) in different CT acquisition conditions^{16,17}.

Instead of using predefined formal lung CT density features such as MLA, skewness, or any quantile measurement, it would be ideal to interrogate the whole sample of CT histograms of patient lungs with the disease of interest (e.g., Covid-19) and see how the CT histograms vary in the patient cohort and eventually, which combinations of modes of variation are associated with the severity of the disease or the patient outcome. In the present study, we are using a statistical method called Functional Principal Component Analysis (FPCA) to explore the modes of variation of the lung CT density histograms in Covid-19 patients having either non-enhanced CT or contrast-enhanced CT and to extract new data-driven features for the prediction of patients critical-illness status, hospital length-of-stay and mortality. Performances were compared with a priori methods using CT attenuation quantile values from Q50 (median density) to Q875 (87.5th percentile density), mean lung attenuation (MLA), standard deviation, skewness, and kurtosis.

Results

Patient Characteristics, Clinical, and Laboratory Findings

Patient characteristics and clinical and laboratory findings are reported in *Supplementary Table S1 online*. The study included 80 patients (median age 63.5 years, 37 of whom were female). Most patients had at least one comorbidity (71%), from which 30% had 1 comorbidity 49% had 2–3 comorbidities and 21% had 4–5 comorbidities. The mean time between CT and critical illness was 3.7 days (SD: 2.7). The mean death time was 24 days (SD: 23 days). Critical illness status was defined as meeting one or more of these patient conditions: admission to ICU, requirement for mechanical ventilation, extracorporeal membrane oxygenation (ECMO) or death within 1 month of first presentation to hospital.

The critically ill group included 35 of the 80 patients (44%), with 32 (40%) in ICU, 25 (31%) requiring mechanical ventilation or ECMO and 15 (19%) who died in hospital. More men were critically ill compared to females (23/35, [66%], $p = 0.073$), with no significant difference in patient age or comorbidities. The critically ill group had a higher mean lactate dehydrogenase value (382 vs 291 U/L, $P = 0.003$), and a higher mean Neutrophil-Lymphocyte Ratio (16.8 vs 5.71; $P = 0.0001$).

CT density curves analysis

The Functional Principal Component Analysis (FPCA) resulted in four Functional Principal Components FPCs explaining respectively 76.7% (F1) 13.5% (F2), 3.8% (F3) and 2.6% (F4) of the variability of lung CT

density curves in the Covid-19 patients (See Fig. 1). F1 represents the main mode of variation of the lung CT density curves, from homogeneous low CT density (10th percentile curve) consistent with normal lungs density to the heterogeneous distribution of CT density with high densities (above - 100 HU) (90th percentile curve associated with extensive lungs consolidation and ground glass opacification) see Fig. 1a. Two-way Anova shows F1 significantly larger for both the ‘critically-ill’ outcome $F(1,76) = 51.2 - P < 0.001$ and the contrast-CT condition $F(1,76) = 6.70 - P = 0.012$. We expect the high F1 values to be strongly associated with unfavorable outcome (“critically ill” condition) in models stratified by contrast CT conditions.

The second mode of variation F2 represents different degrees of shift in CT density from normal homogeneous low lung CT density (90th percentile curve) toward a larger density range around - 800HU to -400HU (10th percentile curve) see Fig. 1b. Interpretation: high F2 values seem associated with normal lungs while small F2 values seem associated with extensive lungs ground glass opacification but limited consolidation. We expect that the low values of F2 with extensive GGO but no significant consolidation to be moderately associated with unfavorable outcome. The third and fourth modes of variation are associated with small shifts toward higher CT densities: F3 + 46HU (25th percentile curve to 75th percentile curve) and F4 + 61HU (25th percentile curve to 75th percentile curve). See Fig. 1c and 1d.

Four score values called F1, F2, F3 and F4 quantifying for each patient the lung density curve (once for each mode of variation) were added to the list of existing features for predictive modeling.

CT Density Features Analysis

The CT analysis findings (including a priori CT features and data driven feature F1) are summarized in *Supplementary Table S2* (descriptive statistics), *Table 1* (univariate analysis) and *Table 2* (multivariate analysis).

Table 1: Univariate (ROC) Analysis

Variables	Unfavorable “Critically Ill” Outcome			
	Cutoff point	AUC (95% CI)	Sensitivity	Specificity
Best variables - All CT Cases (N=80)				
Q875 (HU)	>-380	0.88 (0.79 0.94)	85.7%	82.2%
F1	>0.099	0.87 (0.77 0.93)	82.9%	82.2%
SD-CT (HU)	>213.8 HU	0.86 (0.73 0.93)	82.86%	84.44%
Best variables - Contrast CT Cases (N=54)				
Q875 (HU)	>-335	0.89 (0.77 0.96)	82.6%	87.1%
C cases (N=54)				
SD-CT (HU)	>247	0.89 (0.74 0.96)	78.3%	96.8%
C cases (N=54)				
F1	>0.330	0.88 (0.77 0.95)	73.9%	96.8%
C cases (N=54)				
Best variables - Non-Contrast CT Cases (N=26)				
Q875 (HU)	>-589	0.87 (0.68 0.97)	91.7%	78.6%
NC cases (N=26)				
F1	>-0.255	0.84 (0.64 0.95)	91.7%	78.6%
SD-CT (HU)	> 144	0.80 (0.57 0.94)	83.3%	71.4%
NC cases (N=26)				
Other Noticeable Variables - all cases (N=80)				
CT Mean (HU)	> -646 HU	0.84 [0.73 0.91]	74.29%	84.44%
Skewness	≤1.451	0.83 (0.73 0.90)	88.6%	64.4%
Kurtosis	≤4.680	0.85 (0.75 0.92)	85.7%	77.8%
Age (years)	≤48	0.52 (0.39 0.64) *	34.29%	80.00%
Lung volume - log (ml)	≤3.43 (2670 ml)	0.59 (0.47 0.71) *	74.29%	46.67%
Best Clinical Variable – C & NC cases				
Neutrophil Lymphocyte Ratio	> 4.9	0.74 (0.60 0.85)	81.82%	61.29%
C cases (N=53) **				
Neutrophil Lymphocyte Ratio	> 4.9	0.77 (0.56 0.91)	81.82%	78.57%

Best variables (all cases, and cases with and without contrast) and other noticeable variables (Age, Lung volume and neutrophil lymphocyte ratio NLR). Cutoff points were calculated using Youden index. 95% confidence intervals. *: Non-significant P-value with test on AUC for continuous variables. **: 1 outlier removed. NC=non contrast; C=contrast. ROC= Receiving-Operating Characteristic.

Table 2
Multivariate models for critically ill outcome

Variables	Unfavorable "Critically Ill" Outcome			
	Cutoff point	AUC (95% CI)	Sensitivity	Specificity
Best variables - All CT Cases (N = 80)				
Q875 (HU)	>380	0.88 (0.79 0.94)	85.7%	82.2%
F1	> 0.099	0.87 (0.77 0.93)	82.9%	82.2%
SD-CT (HU)	> 213.8 HU	0.86 (0.73 0.93)	82.86%	84.44%
Best variables - Contrast CT Cases (N = 54)				
Q875 (HU) C cases (N = 54)	>335	0.89 (0.77 0.96)	82.6%	87.1%
SD-CT (HU) C cases (N = 54)	> 247	0.89 (0.74 0.96)	78.3%	96.8%
F1 C cases (N = 54)	> 0.330	0.88 (0.77 0.95)	73.9%	96.8%
Best variables - Non-Contrast CT Cases (N = 26)				
Q875 (HU) NC cases (N = 26)	>589	0.87 (0.68 0.97)	91.7%	78.6%
F1	>0.255	0.84 (0.64 0.95)	91.7%	78.6%
SD-CT (HU) NC cases (N = 26)	> 144	0.80 (0.57 0.94)	83.3%	71.4%
Other Noticeable Variables - all cases (N = 80)				
CT Mean (HU)	> -646 HU	0.84 [0.73 0.91]	74.29%	84.44%
Skewness	≤ 1.451	0.83 (0.73 0.90)	88.6%	64.4%
Kurtosis	≤ 4.680	0.85 (0.75 0.92)	85.7%	77.8%
Age (years)	≤ 48	0.52 (0.39 0.64) *	34.29%	80.00%
Lung volume - log (ml)	≤ 3.43 (2670 ml)	0.59 (0.47 0.71) *	74.29%	46.67%

The models with the highest AUC values included: reader-1 subjective Covid score in the enhanced CT group, mean CT density + standard deviation CT density in the enhanced CT group and Standard deviation CT density + NLR in the enhanced CT group. (95% CI = 95% confidence intervals).

Variables	Unfavorable “Critically Ill” Outcome			
	Cutoff point	AUC (95% CI)	Sensitivity	Specificity
Best Clinical Variable – C & NC cases				
Neutrophil Lymphocyte Ratio C cases (N = 53) **	> 4.9	0.74 (0.60 0.85)	81.82%	61.29%
Neutrophil Lymphocyte Ratio NC cases (N = 25) **	> 4.9	0.77 (0.56 0.91)	81.82%	78.57%
The models with the highest AUC values included: reader-1 subjective Covid score in the enhanced CT group, mean CT density + standard deviation CT density in the enhanced CT group and Standard deviation CT density + NLR in the enhanced CT group. (95% CI = 95% confidence intervals).				

Patients with critical illness had significantly different CT density features (all $P < 0.0001$) with higher CT mean density (MLA), higher CT SD, higher Q875 (87.5th percentile), higher F1 (mean: 0.326 vs. -0.254), lower skewness and lower kurtosis (See *Supplementary Table S2 online*). *Table 1* summarizes all the univariate feature performances (AUC) with optimal cutoff points using the Youden index criteria. Results for the three best CT predictors are stratified into contrast CT and non-contrast CT groups. Notice that for each of these features, the AUCs are slightly higher for the contrast enhanced group than the non-contrast group (non-significant differences). ROC analysis identified as best predictors of critically ill status: Q875 - AUC: 0.88 (0.79 0.94), F1 - AUC: 0.87 (0.77 0.93), SD-CT - AUC: 0.86 (0.73, 0.93). Remarkably, the frequently used feature trio Mean CT (MLA), skewness and kurtosis showed lower performances than the three previous features: Mean CT - AUC: 0.84 (0.73 0.91), skewness - AUC: 0.83 (0.73 0.90), Kurtosis - AUC: 0.84 (0.73, 0.91) and were not retained in the final multivariate models.

Among the clinical variables shown in *Table 1*, the Neutrophil-Lymphocyte Ratio (NLR) appears the best predictor with AUC: 0.74 (0.60 0.85) for the contrast cases and AUC: 0.77 (0.56 0.91) for the non-contrast cases and was further successfully included in the final multivariate models. Main univariate ROC curves and AUCs are shown in *Supplementary Fig. S1*.

Odds Ratios

Odds Ratios (OR) for different quantitative, clinical, and subjective variables are presented in *Fig. 2*. Whole lungs F1, SD-CT, Mean-CT and Q875 measurements were associated with critical illness: OR from 21.75 (5.63,83.96) to 8.31 (3.9,23.1) - all $p < 0.0001$. CT severity scores had high OR values as well. OR: 31.4 (9.2,107.4) $p < 0.0001$. Neutrophil-Lymphocyte Ratio (NLR) and Lactate dehydrogenase OR values also showed an association with critical illness.

Multivariate model performances

Using multivariate logistic regression models stratified for IV contrast vs. no contrast to predict critical illness, model 1 with the subjective Covid score alone had the best predictive value: AUC: 0.92 (0.83, 0.98) for IV contrast group and AUC: 0.87(0.68,0.97) for the non-contrast group. See Table 2. Considering quantitative models, the model 2 combining SD-CT and Neutrophil-Lymphocyte ratio (NLR) predictors had an AUC of 0.92 (0.81,0.97) with IV contrast compared to AUC: 0.82 (0.61,0.94) the non-contrast group. Model 3 combining F1 and NLR showed an accuracy AUC: 0.91 (0.80,0.97) with IV contrast and AUC: 0.88 (0.68,0.97) for the non-contrast group. The separation of critically ill vs. non-critically ill true cases with the model 3 and its predicted probabilities is shown in *Supplementary Fig. S2*. Model 4 combining Q875 and NLR predictive accuracy (AUC) showed the highest overall performance among the quantitative models: 0.92 (0.81,0.98) for the IV contrast group and 0.87 (0.68,0.97) for the non-contrast group.

Combined Length-of-stay (LOS) and in-hospital mortality assessment

For SD-CT, Q875 and F1 predictors, the patient cohort was stratified in two groups using optimal cutoff points for combined contrast and non-contrast studies found in the univariate analysis: SD-CT $\geq$ 213.8HU, Q875>380HU and F1>0.099.

Using Q875 feature (see Fig. 3 – Dashed lines) the cumulative incidence LOS at 30 days was: 89% (80%, 98%) for low Q875 group compared to 40% (24%, 56%) for high Q875 group. the Q875 based cumulative incidence for Death at 30 days was: 2.4% (0%, 7%) for low Q875 group 1 compared to 21% (8%, 34%) for high Q875 group 2. Both inter-group differences in cumulative incidences for LOS ($P < 0.0001$) and Death ($P = 0.027$) were significant.

Using F1 feature (Fig. 3 - solid lines) the cumulative incidence LOS at 30 days was: 81% (69% 93%) for low F1 group compared to 47% (31% 63%) for high F1 group. The F1-based cumulative incidence for Death at 30 days was: 4.6% (0% 11%) for low F1 group 1 compared to 19% (6% 31%) for high F1 group 2. Inter-group difference in cumulative incidence for LOS ($p < 0.0001$) was significant while those for Death did not reach significance ($P = 0.09$).

Similarly, the patients of the low SD-CT group had a probability of being discharged before or at 30 days of 89.7% (80.4% 99.1%) compared to 36.8% (20.9% 52.7%) for the high SD-CT group ($p < 0.0001$) and a mortality risk at 30 days of 2.3% (0.0% 6.6%) ($p = 0.022$) compared to 22.4% (8.7% 36.0%) for the high SD-CT group ($p = 0.022$).

Semi-quantitative CT severity score and COVID-GRAM Scores

The CT severity score was higher in critically ill patients vs. non critically ill patients with a combined mean score amongst both readers of 21 vs 15 ($p = 0.002$). Univariate AUC was 0.91 (0.80, 0.96) with Reader 1 and AUC: 0.83 (0.73 to 0.91) with Reader 2.

The COVID-GRAM score performed poorly with an AUC value of 0.64 (0.52, 0.74). In fact, 79/80 (99%) of our patients were predicted to have medium or high risk for critical illness based on the clinical variables from their medical records compared to the actual value of 35/80 (44%).

Interrater reliability

Interrater reliability was tested by having a second investigator perform separate Covid CT severity scoring, manual corrections on the lung segmentation (as needed) and measurements on a sample of 20 randomly selected patients.

Interrater reliability for the CT severity score measured with intra-class correlation (ICC) was: 0.90 (0.85 0.94), indicating a good agreement between both readers. Similarly, the quantitative features SD-CT ICC: 0.98 (0.95 0.99), Q875 ICC: 0.99 (0.97 0.99) and F1 ICC: 0.99 (0.96 0.99) showed an excellent interrater agreement.

Discussion

Parsimonious models combining only one quantitative lung CT density parameter (either SD-CT, Q875, or F1) and one clinical parameter (Neutrophil-Lymphocyte Ratio) allowed accurate prediction of critical illness (ICU, mechanical ventilation/ECMO and/or death) in Covid-19 patients with accuracy (AUC) ranging from 0.82 to 0.92, and prediction of hospital length-of-stay while controlling for the mortality risk. Remarkably, the performances were not adversely affected by the presence of IV contrast in the CT images and were even slightly better in the contrast enhanced group, although the lack of randomization for the IV contrast precludes evaluation of whether this factor leads to better predictions.

The three best CT parameters were Q875, F1 and SD-CT. SD-CT is a well-established a priori radiomic global feature related to the spread of the CT histogram largely available in commercial lung imaging software. In this study, the frequently seen parameters MLA, Skewness and Kurtosis⁶ performed slightly worse than either SD-CT, Q875, or F1 (see *Table 1*) and were excluded from the final models. Q875 is another radiomic parameter related to the HU value reached when 87.5% of the lung voxels have been counted (starting with the lowest densities). Q875 is the counterpart for the high CT densities of the 15th percentile density index (PD15) used to quantify the severity of emphysema in lung CT densitometry. F1 parameter is a result of the histogram functional principal component analysis (FPCA) in the patient cohort. Without a-priori knowledge or information about the patient outcome, FPCA extracts the main modes of variation in the sample of CT histograms for the patient cohort. F1 (score) values represent the different degrees of CT histograms shift from homogeneous low lung densities (better outcome) toward heterogeneous much higher densities (worse outcome) see Fig. 1. In the current study, other modes of variations (F2, F3, etc.) were not predictive of the patient outcome. F2 seems to reflect the transition from normal lung densities to extended ground glass opacifications (about -800HU to -600HU) and it was not a significant predictor of critically ill status (AUC: 0.53, P:0.60). In this study, the overall results using FPCA are consistent with previous ones for pulmonary disease subtyping^{18,19} or patient neurologic outcome prediction²⁰ confirming the value of the FPCA method: first as a non-specific data driven exploration tool,

it offers interpretable modes of variations of the CT histograms in the patient whole cohort. Second, it is a generic method giving accurate predictors related to histogram variations without a priori knowledge or delicate radiomic high dimensional parameter selection. All three CT predictors are highly correlated: Spearman Rho Q875 vs. F1: 0.97, Q875 vs. SD-CT, Rho: 0.90, F1 vs. SD-CT: Rho: 0.92 and practically exchangeable. However, the data driven FPCA approach offers a unique data analysis tool of the CT histograms in the whole patient cohort. Current machine learning research is actively extending the FPCA method with supervised FPCA²¹, multivariate FPCA²² (neuroimaging data), robust FPCA²³, etc. offering a rich toolbox for future medical imaging studies. See also Pratt et al.²⁴ for a recent application in pulmonology.

The good performances of CT density features for patient outcome prediction in COVID patients are concordant with results from a few previous studies^{10–12} including Lanza et al.¹¹ who showed that COVID patients requiring oxygenation and ventilation had higher amounts of compromised lung volumes (-500-100 HU), statistically significant at 6–23% and greater than 23% respectively. Another large study by Colombi et al. showed that a percentage of well aerated lung on CT calculated by software of 71% (OR, 3.8, 95% CI: 1.9, 7.5, $P < .001$) or less was associated with ICU admission or death¹².

Lung volume is known to affect the lung CT density in a complex way: First, the optimal lung inflation is difficult to obtain in severe acute lung disease and spirometry-controlled lung CT is often not feasible. So, partially inflated lungs may increase the apparent lung CT density. Bressen et al.¹⁰ have mentioned the potential confounding effect of the lung volume variation among patients when using CT density as a biomarker in Covid-19 patients. Second, the lung tissue density is increased with disease severity associated with extended GGO and consolidations. Third, lung CT density appears to be lower in subjects with larger lungs because of greater air spaces²⁵. In this study, adding the lung volume feature did not improve the performances of the predictive models. However, a moderate but significant correlation has been observed for the CT parameters: Q875 - Rho: -0.60 (-0.76 -0.38), F1 - Rho: -0.54 (-0.72 -0.30), Mean CT: Rho: -0.68 (-0.81 -0.48) but not SD-CT (Rho: -0.30 (-0.49 -0.08) $P = 0.0072$, for the non-critically ill patient group, in agreement with the research literature. The *supplementary Fig. S3* shows the relationship between Q875 and lung volume for both patient outcome groups. The linear relationship between Q875 and log. Volume in the non-critically ill group and for a large range of lung volumes may be best explained with Robert et al. hypothesis on lung CT density change with normal lung growth²⁵.

Moreover, the performance of the clinical predictor: Neutrophil-Lymphocyte Ratio (NLR) in either the univariate analysis (*Table 1*) or in the multivariate best logistic regression models (*Table 2 – Models 2-3-4*) supports the conclusion of the recent meta-analysis from Li et al.²⁶ pointing out the value of this biomarker to predict disease severity and patient mortality in Covid-19 patients.

Our quantitative CT density features were compared with both the COVID-GRAM score and CT severity score to predict critical illness. The CT severity score (Reader-1) alone performed well with AUC: 0.91 (0.80 0.96) and intra-class correlation (ICC): 0.90 (0.85 0.94) as previously shown in prior studies^{4,27}

However, Reader-2 CT severity score was suboptimal and illustrated the inter-reader variability of subjective features. See for example Fig. 2 (*Odd Ratios*). The COVID-GRAM score performed poorly in our study to predict critical illness with AUC of 0.64 (0.52, 0.74) 95% CI. A possible explanation of this poor performance compared to the original Chinese study to develop the model by Liang et al.⁵ is the presence of older patients (60.8 vs 48.9 years) and a higher prevalence of one of more pre-existing comorbidities (71.3% vs 25.1%) in our study. Remarkably, Al Hassan et al.²⁸ recently reported similar findings with an AUC of 0.64 for COVID-GRAM score for risk stratification with Covid-19 patients.

Hospital length of stay (LOS) and hospital mortality are mutually related and thus require a competing risks method for proper assessment of the cumulative incidence of each event of interest (discharge or death). Using this method, our study showed that the patient groups with Q875>380 HU, F1 > 0.099 or SD-CT > 213.8 HU were all associated with significantly higher cumulative incidences for longer length of stays while controlling for the hospital mortality. This information is valuable in capacity planning to provide accurate predictions of the number of beds required at each level of care.

This study has several limitations: it is retrospective and has a modest sample size resulting in larger confidence intervals or suboptimal statistical power when considering subgroup analysis (such as IV contrast vs. Non-Contrast CT) and prevents us to draw conclusions about the in-hospital mortality due to the low number of death events (14/80). The predictive accuracy results were computed with cross-validation correcting performance for overfitting. However, future work involving multiple sites would be necessary for testing the performances in a fully separated testing dataset.

Another limitation is that CT chest protocols varied based on the clinical indication with almost twice as many pulmonary angiogram studies as non-contrast studies, preventing us to better understand the role of CT contrast in the predictive performances.

Finally, the methods discussed in this study are focused on a global lung CT histogram analysis. Multi-threshold lung density analysis methods such as those described in already mentioned studies^{10,12,15} or more advanced CT density/texture methods based on local lung pattern classification²⁹ were not tested and should deserve future attention.

In conclusion, the extensive and diffuse changes in lung CT density affecting the whole lungs in COVID-19 pneumonia patients offered the opportunity to compare predefined and data-driven imaging features related to the lungs CT density histograms. All SD-CT, Q875 and F1 features could accurately predict both critical patient illness and hospital length-of-stay. Combined models with one of these features and the biomarker for inflammation Neutrophil-Lymphocyte Ratio gives the highest predictive performance. This application of CT densitometry provided similar results for both Non-enhanced CT group and the contrast enhanced group. The FPCA method allowed the unsupervised analysis of the lung density histograms in the whole patient cohort to extract interpretable CT density features with high predictive values. This approach may be considered for other predictive models with diffuse lung diseases.

Methods

Study Population

This study was approved by the UHN Coordinated Approval Process for Clinical Research (CAPCR) ethics committee for human research at our home hospital. All methods were conducted in accordance with guidelines outlined by this committee. Patient consent was waived due to the retrospective collection of patient data, and this was approved by the CAPCR UHN ethics committee. Inclusion criteria were adult patients ≥ 18 years of age with real-time reverse transcriptase polymerase chain reaction (RT-PCR) confirmed COVID-19 (positive after 1–3 tests) who had undergone a CT chest within 24 hours of admission to hospital between March 1 and December 15, 2020 and who were not in the ICU or mechanically ventilated at the time of the CT study. The indication for CT included ruling out suspected COVID-19 with a non-enhanced CT chest and to assess for a pulmonary embolism with an enhanced CT pulmonary angiogram study in patients with confirmed COVID-19. Exclusion criteria were known malignancy with pulmonary nodules on CT (added density), incomplete clinical data and those with a known superimposed bacterial pneumonia. 502 patients were retrieved, with 87 of these patients with confirmed COVID-19 by RT-PCR. Two patients with known lung cancer and lung nodules, three patients with both COVID-19 and a superimposed bacterial pneumonia and two patients with incomplete medical history and blood work were excluded from the study (see *Supplementary Fig. S4*).

Patient characteristics on admission

Clinical and laboratory data documented at admission included age, sex, symptoms, blood work (neutrophils, lymphocytes, lactate dehydrogenase, bilirubin) and comorbidities (chronic obstructive pulmonary disease, hypertension, diabetes, coronary disease, heart disease, malignancy, kidney disease, cerebral vascular disease, hepatitis B and immunodeficiency).

Outcomes

The primary outcome was critical illness, defined as one or more of admission to ICU, requirement for mechanical ventilation or extracorporeal membrane oxygenation and death within 1 month of first presentation to hospital. Secondary outcomes were hospital length of stays and mortality.

Computed tomography imaging protocol

We analyzed both non-contrast low dose and normal dose CT chest studies and contrast enhanced CT pulmonary angiogram CT studies from three hospitals in our institution. All patients were examined with either 64-CT Aquilion or 320-CT Aquilion-One scanner (Canon Medical Systems, Otawara, Japan). Chest CT acquisitions parameters were 120 kV and 20–100 mA (low dose), 120 kV and 40–150 mA (normal dose) and 100–120 kV and 0–250 mA (contrast enhanced pulmonary angiograms) according to our hospital protocols. All images were reviewed in lung windows (width: 1200 HU, level: -700 HU) and mediastinum windows (width: 350 HU, level: 40 HU) with 1–3 mm slice thickness. CT pulmonary

angiogram studies administered 70 mL of iodinated contrast (Iopromide 370 mg I/mL) at a rate of 5 mL/sec using a bolus tracking technique triggered to 250 HU in the main pulmonary artery.

COVID Gram Score

Clinical and laboratory data at time of hospital presentation was collected from the patients' electronic medical records to calculate the COVID-GRAM score⁵. These 10 parameters include age, dyspnea, conscious, hemoptysis, history of malignancy, number of comorbidities, x-ray abnormality, Neutrophil-Lymphocyte Ratio, lactate dehydrogenase and direct bilirubin. An online calculator was used to calculate a risk score and percentage and place the patient in a low, medium, or high-risk group to predict critical illness³⁰.

CT Severity Score

Subjective assessment of the percentage of ground glass opacities and consolidations on CT chest was performed by two radiologists, one with 4 years of clinical experience as a staff thoracic radiologist, and one a cardiothoracic imaging fellow with 5 years of radiology residency experience. The lungs were evaluated as per the CT Severity Score guidelines⁴ by assigning a score of 0–2 (0 = no opacity, 1 = < 50% opacity and 2 = $\geq$ 50% opacity) in each of the 10 segments in both lungs out of a total score of 40.

CT Density Analysis

Lung density measurements were performed using Vitrea Advanced Visualization version 7.14 (Vital Images, Minnetonka, USA) software to perform automatic lung segmentation and calculate a mean CT density (Mean-CT) and standard deviation (SD-CT) density for both lungs combined (See Fig. 4 & Fig. 5). Large pulmonary vessels, airways, mediastinum structures and pleural effusions are excluded from segmentation, and lung parenchyma, interstitial structures and segmental vessels and bronchi were included. Manual correction of the lung segmentation was applied when needed. The primary density analysis was performed by a cardiothoracic radiology fellow, and a second analysis to determine inter-rater reliability on a random sample of 20 cases from the data set was performed by a clinical research analyst with 4 years of experience in cardiothoracic radiology.

CT density curves analysis

Each CT density histogram for both lungs and each lung were converted in smooth curves defined between - 1000HU and 500HU using Ramsey's smoothing method for frequency distributions³¹ and previously described²⁰. Quantiles values on the CT histograms from median (50th percentile) to 85.7th percentile were computed and added to the feature list. A Functional Principal Component Analysis (FPCA) was applied to the lung CT density curves following Petersen & Müller's method for frequency distributions³². FPCA is a data driven approach akin to the Principal Component analysis (PCA) to explore and quantify the main modes of variation of a sample of curves. The resulting functional principal component scores (FPCs) for the lung region were added to the list of candidate predictors of the patient outcome including lung volume, demographic information and a priori CT density-based

features: mean lung CT attenuation, standard deviation, skewness, kurtosis and eight quantile-based features.

Statistical analysis

Statistical analysis was performed using R statistical programming and MedCalc software. A p-value of < 0.05 was considered statistically significant. Continuous variables were described using mean and standard deviation or median and interquartile range and categorical variables using numbers and percentage. Mann-Whitney tests and Fisher exact tests were used to compare continuous and binary variables. Variance ratio F-test were used to compare inter-group variance differences. Inter-rater agreement was assessed using intra-class correlation (ICC). Logistic regression models were applied for prediction of critical illness. A univariate analysis of the predictors for critical illness was performed using a receiver operating characteristic (ROC) area under the curve (AUC) metric for continuous predictors and odds ratio (OR) for binary predictors. First quartile to third quartile differences were used for defining OR in non-categorical variables. Optimal cutoff points on ROC curves were determined using the Youden index method.

In the multivariate analysis, predictive accuracy (AUC) of the final models were corrected for overfitting using a Bootstrap cross-validation method. The length-of-stay estimate for Covid patients have recently been addressed with numerous methods³³. In this study, the hospital discharge time was used as primary end point for hospital length-of-stay (LOS) and the in-hospital death was considered as the competing risk, following Brock et al. approach and publicly available R code³⁴. A cumulative incidence plot of hospital discharge was computed in R-programming language using the Aalen-Johansen estimator. The Greenwood-type method was used to estimate the standard errors and confidence intervals of our cumulative incidence plots.

Declarations

Data availability

The R-code for the CT density histogram smoothing and for the functional principal component analysis used in the current study are available from the corresponding author.

Acknowledgments

We would like to thank our colleagues at the Toronto General hospital for their support, patient care and encouragement during this pandemic.

Contributions

J.F, T.S, M.H. worked on the imaging analysis of the lungs and CT histograms, collected the imaging, clinical and pathology data, wrote and edited the manuscript.

P.S. performed the statistical analysis of the project, interpreted the results of the statistical analysis and wrote and edited the manuscript.

M.P. and K.H. helped with the study design and edited the manuscript.

V.H. helped with the study design and software functioning.

Competing interests

J.F., T.S., M.H., K.H., M.P., V.H. and P.S. declare no competing interests. P.S. is employee of Vital Images.

References

1. Wong, K. T. *et al.* Thin-section CT of severe acute respiratory syndrome: evaluation of 73 patients exposed to or with the disease., **228**, 395–400 (2003).
2. Kang, H. *et al.* Computed tomography findings of influenza A (H1N1) pneumonia in adults: pattern analysis and prognostic comparisons. *J Comput Assist Tomogr*, **36**, 285–290 (2012).
3. Das, K. M. *et al.* CT correlation with outcomes in 15 patients with acute Middle East respiratory syndrome coronavirus. *AJR Am J Roentgenol*, **204**, 736–742 (2015).
4. Yang, R. *et al.* Chest CT Severity Score: An Imaging Tool for Assessing Severe COVID-19. *Radiol Cardiothorac Imaging*, **2**, e200047 (2020).
5. Liang, W. *et al.* Development and Validation of a Clinical Risk Score to Predict the Occurrence of Critical Illness in Hospitalized Patients With COVID-19. *JAMA Intern Med*, **180**, 1081–1089 (2020).
6. Best, A. C. *et al.* Idiopathic Pulmonary Fibrosis: Physiologic Tests, Quantitative CT Indexes, and CT Visual Scores as Predictors of Mortality., **246**, 935–940 (2008).
7. Iwasawa, T. *et al.* Assessment of prognosis of patients with idiopathic pulmonary fibrosis by computer-aided analysis of CT images. *J Thorac Imaging*, **24**, 216–222 (2009).
8. Rea, G. *et al.* Comparative analysis of density histograms and visual scores in incremental and volumetric high-resolution computed tomography of the chest in idiopathic pulmonary fibrosis patients. *Radiol Med*, **126**, 599–607 (2021).
9. Horie, M. *et al.* Lung Density Analysis Using Quantitative Chest CT for Early Prediction of Chronic Lung Allograft Dysfunction., **103**, 2645–2653 (2019).
10. Bressem, K. K. *et al.* Is lung density associated with severity of COVID-19? *Pol J Radiol*, **85**, e600–e606 (2020).
11. Lanza, E. *et al.* Quantitative chest CT analysis in COVID-19 to predict the need for oxygenation support and intubation. *Eur Radiol*, **30**, 6770–6778 (2020).
12. Colombi, D. *et al.* Well-aerated Lung on Admitting Chest CT to Predict Adverse Outcome in COVID-19 Pneumonia., **296**, E86–E96 (2020).
13. Park, B. *et al.* Prognostic Implication of Volumetric Quantitative CT Analysis in Patients with COVID-19: A Multicenter Study in Daegu, Korea. *Korean J Radiol*, **21**, 1256 (2020).
14. Lubner, M. G., Smith, A. D., Sandrasegaran, K., Sahani, D. V. & Pickhardt, P. J. CT Texture Analysis: Definitions, Applications, Biologic Correlates, and Challenges. *RadioGraphics*, **37**, 1483–1503 (2017).

15. Rorat, M., Jurek, T., Simon, K. & Guziński, M. Value of quantitative analysis in lung computed tomography in patients severely ill with COVID-19. *PLoS One*, **16**, e0251946 (2021).
16. Wang, Y. *et al.* Temporal Changes of CT Findings in 90 Patients with COVID-19 Pneumonia: A Longitudinal Study., **296**, E55–E64 (2020).
17. Cao, X., Jin, C., Tan, T. & Guo, Y. Optimal threshold in low-dose CT quantification of emphysema. *European Journal of Radiology*, **129**, 109094 (2020).
18. Oikonomou, A. *et al.* Histogram-based models on non-thin section chest CT predict invasiveness of primary lung adenocarcinoma subsolid nodules. *Sci Rep*, **9**, 6009 (2019).
19. de Margerie-Mellon, C. *et al.* Assessing invasiveness of subsolid lung adenocarcinomas with combined attenuation and geometric feature models. *Sci Rep*, **10**, 14585 (2020).
20. Salazar, P. *et al.* Exploration of Multiparameter Hematoma 3D Image Analysis for Predicting Outcome After Intracerebral Hemorrhage. *Neurocrit Care*, **32**, 539–549 (2020).
21. Li, G., Shen, H. & Huang, J. Z. Supervised Sparse and Functional Principal Component Analysis. *Journal of Computational and Graphical Statistics*, **25**, 859–878 (2016).
22. Happ, C. & Greven, S. Multivariate Functional Principal Component Analysis for Data Observed on Different (Dimensional) Domains. *Journal of the American Statistical Association*, **113**, 649–659 (2018).
23. Boente, G. & Salibián-Barrera, M. Robust functional principal components for sparse longitudinal data. *METRON*, **79**, 159–188 (2021).
24. Pratt, J., Su, W., Hayes, D., Clancy, J. P. & Szczesniak, R. D. An Animated Functional Data Analysis Interface to Cluster Rapid Lung Function Decline and Enhance Center-Level Care in Cystic Fibrosis. *Journal of Healthcare Engineering*, **2021**, 1–13 (2021).
25. Robert, H. B., Robert, A. W., Kirk, G., Drummond, M. B. & Mitzner, W. Lung density changes with growth and inflation., **148**, 995–1002 (2015).
26. Li, X. *et al.* Predictive values of neutrophil-to-lymphocyte ratio on disease severity and mortality in COVID-19 patients: a systematic review and meta-analysis. *Crit Care*, **24**, 647 (2020).
27. Lieveld, A. W. E. *et al.* Chest CT in COVID-19 at the ED: Validation of the COVID-19 Reporting and Data System (CO-RADS) and CT Severity Score: A Prospective, Multicenter, Observational Study., **159**, 1126–1135 (2021).
28. Al Hassan, H., Cocks, E., Jesani, L., Lewis, S. & Szakmany, T. Clinical Risk Prediction Scores in Coronavirus Disease 2019: Beware of Low Validity and Clinical Utility. *Crit Care Explor*, **2**, e0253 (2020).
29. Ohno, Y. *et al.* Machine learning for lung CT texture analysis: Improvement of inter-observer agreement for radiological finding classification in patients with pulmonary diseases. *European Journal of Radiology*, **134**, 109410 (2021).
30. Liang, W., Walker, G., COVID-GRAM Critical Illness Risk Score & At <https://www.mdcalc.com/covid-gram-critical-illness-risk-score> (2020).

31. Ramsay, J. O. & Silverman, B. W. *Functional data analysis* (Springer, 2005).

32. Petersen, A. & Müller, H. G. Functional data analysis for density functions by transformation to a Hilbert space. *The Annals of Statistics*, **44**, 183–218 (2016).

33. Rees, E. M. *et al.* COVID-19 length of hospital stay: a systematic review and data synthesis. *BMC Med*, **18**, 270 (2020).

34. Brock, G. N., Barnes, C., Ramirez, J. A. & Myers, J. How to handle mortality when investigating length of hospital stay and time to clinical stability. *BMC Med Res Methodol*, **11**, 144 (2011).

Figures

Figure 1

First four modes of variation of the lung CT density curves (smoothed CT histograms). A. F1 explains 76.7% of the fraction of variance explained (FVE). The curves vary from homogeneous low CT density (10 percentile curve) to heterogeneous higher CT density (90 percentile curve). B. F2 explains 13.5% of the FVE. The lung CT density vary from homogeneous low density (90 percentile curve) to include mid-low density in the -700HU to -350HU range (10 percentile curve). C. F3 explains 3.8% of the FVE. This mode of variation represents a redistribution of CT lung densities from (-700HU to -600HU) to about (-600HU to -300HU). D. F4 explains 2.6% of the FVE. F4 represents a small shift of about 140HU from low homogeneous CT densities (10 percentile curve) toward higher CT densities (90 percentile curve).

Figure 2

Odd ratios with 95% confidence intervals for critically ill outcome. The continuous variables are split into two groups below / above their median value. Significant p-values ($P<0.05$) are bold. Variables with P-

values >0.5 are not shown. Notice that 6/10 of the highest ranked predictors are lung CT density quantitative features.

Figure 3

Cumulative incidence plot for hospital length-of-stay (LOS) and in-hospital deaths in Covid-19 patients stratified in Low Q875 (resp. F1) patients and high Q875 (resp. F1) patients using an optimal cutoff point for combined contrast and non-contrast studies found in the univariate analysis: Q875>380 HU (resp. F1>0.099). Both F1 and Q875 features give similar predictions of cumulative incidences for LOS and Death during the follow-up period.

Figure 4

Non-contrast chest CT images in a 28-year-old man with mild COVID-19 pneumonia. The patient was discharged 3 days later without ventilation or ICU admission. a. CT images show small, rounded ground-glass opacities in the upper lobes. b. 3D view with volume. c. Histogram of CT density values for both lungs shows most lung voxels in a normal lung density distribution and peak at about -800 HU. (perc. = percentile). Mean lung CT density: -786.1HU (21st perc.), SD-CT: 131.9HU (29th perc.), Skewness: 2.58 (97th perc.), Kurtosis: 13.40 (92nd perc.), Q875: -664HU (19th perc.), F1: -0.43 (19th perc.), F2: -0.05 (39th perc.).

Figure 5

Non-contrast low dose chest CT images in a 62-year-old woman with severe COVID-19 pneumonia. The patient was admitted to ICU, required ventilation, and died in hospital. a. CT images show bilateral extensive ground glass opacities and septal thickening. b. 3D view with volume. c. Histogram of CT density values demonstrates 2 peaks, the first peak at -800 HU representing normal well aerated lung and the second peak at 0 HU representing the lung consolidations and ground glass opacities. Mean lung density: -375.4 HU (98th percent.), SD-CT: 348.9 HU (94th perc.), skewness: -0.02 (7th perc.), kurtosis: 1.53 (0.6th perc.), Q875: 36 HU (88th perc.), F1: 0.77 (98th perc.), F2: -0.33 (9th perc.).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement.docx](#)