Seven Things to Know About The Homicides of Doctors. A Review of 30 Years From Italy.

Liliana Loretto

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Alessandra MA Nivoli

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Irma Daga

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Paolo Milia

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Cristiano Depalmas

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Giancarlo Nivoli

Sassari University Hospital: Azienda Ospedaliero Universitaria di Sassari

Saverio Bellizzi (✉ saverio.bellizzi@gmail.com)

Medical Epidemiologist, independent consultant <https://orcid.org/0000-0002-4119-0803>

Research article

Keywords: Homicides of doctors, stalking, workplace violence

Posted Date: October 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-89499/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Public Health on July 5th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-11404-5>.

Abstract

Background. Healthcare workers have a 16 times greater risk of suffering workplace violence than workers in other sectors and around 50% experience workplace violence in the course of their career. The objective of this study is to explore the characteristics and circumstances of work-related killings of doctors.

Methods. Work-related homicides of doctors over the period 1988-2019 were identified retrospectively through the Italian national statistical agencies. Variables such as perpetrator, motive and location of the crime were obtained through forensic psychiatric work. After classification, the absolute and percent values of the main characteristics of the homicides were calculated.

Results. Over the period considered, 21 doctors were killed in Italy in connection with their professional activity. In 52% (n=11) of cases, the killer was one of the doctor's patients, in 29% (n=6) of cases it was a patient's relative, in 19% (n=4) an occasional patient (first consultation). The location of the homicide was a community clinic in 48% (n=10) of cases, the street in 19% (n=4) of cases, the doctor's home in 14% (n=3), the hospital in 14% (n=3) and the patient's home in 5% (n=1). In 57% (n=12) of cases the perpetrator was not affected by any mental disorders. The motive for the homicide was revenge without previous threats in 52.4% (n=11) of cases; in 28.6% (n=6) the revenge was preceded by stalking.

Conclusions. Doctors should be aware that the risk of being killed is not limited to hospital settings and that their patients' family members might also pose a threat to them.

Background

Workplace violence is defined as any event that results in harm caused by work-related assaults, threats and abuse, whose impact on the victim may lead to a deterioration in health, safety and well-being (1). Workplace violence (WPV) is often associated with the type of occupation, with higher incidence among professions involving interactions with many individuals; therefore WPV is a matter of considerable concern for the health sector.

A US report notes that on average, 20 workers are murdered and 18,000 are assaulted each week while at work (2); similar figures are provided by European reports (3). Forty-eight per cent of non-fatal workplace violence incidents take place in the healthcare sector (4). About 50% of healthcare workers experience workplace violence in the course of their career (5, 6). Healthcare workers have a 16 times greater risk of suffering workplace violence than workers in other sectors (7). Nurses are more at risk (8, 9) and female workers, both nurses and doctors, are at even higher risk (10). In a sample of 1826 health professionals, about 11% had suffered physical assault, 5% on more than one occasion, while 64% had received threats or verbal abuse (11). Saeki et al (12) report a prevalence of 15%.

Data from the US National Crime Victimization Survey for the period 2005–2009 show a rate of workplace violence of 5.1/1000 for all occupations, 10.1/1000 for physicians and 8.1/1000 for nurses.

For mental health workers the violent victimisation rate was 20.5/1000, second only to the rate of law enforcement officers (47.7/1000) (13).

In Belgium, a study on patient-physician aggressions (14) conducted by means of an online questionnaire which was completed by 4930 participants, found that, in the preceding 12 months, 37% had been the victim of aggression: 33% verbal aggression, 30% psychological, 14% physical and 10% sexual. Psychiatric and emergency departments were the settings where violence most commonly occurred.

In Israel, Carmi-Iluz (15) compared the risk of violence between a sample of hospital-based physicians and a sample of community-based physicians, finding a substantially comparable risk.

With regard to the authors of violent acts in the workplace, Rippon (16) identifies several categories: patients, family members, visitors and co-workers. In Turkey, a study found that 64.5% of the attacks were carried out by the patients' family members (17)

A particular form of violence, homicide, is infrequent but extremely disturbing.

In 2006, a leading schizophrenia specialist, W. S. Fenton, was killed at his office by one of his patients (18).

In the healthcare professions, homicide is a malicious and intentional event and is a very rare violent circumstance.

A study in the USA found that between 1980 and 1990, 106 healthcare workers were killed at work (19). Of these, 26 were physicians. The US BLS (Bureau of Labour Statistics) (OSHA 4) reported that 69 HCWs were killed between 1996 and 2000. In Italy, in the period between 1988 and 2010, 17 physicians (20) were killed at work.

Nielssen and Large (21) have analysed homicides by psychiatric patients in psychiatric hospitals and proposed a classification into three patient categories: acute psychiatric patients soon after hospital admission, patients not receiving medication with a history of serious violence, and patients with dementia or intellectual disability, held in low-security inpatient settings in contact with vulnerable patients (victims of the homicide). In this study, the victims were both healthcare workers and fellow patients.

A study by Knable (22) describes a sample of mental health workers who were killed by patients in the US. The study highlights the low frequency of these events (about one case per year) and describes the characteristics of victims and perpetrators. The victims were most likely young female caseworkers, with little work experience, killed during unaccompanied visits to residential treatment facilities. The perpetrators were mostly male, diagnosed with schizophrenia, with poor adherence to medication.

We have carried out a retrospective analysis, to explore the main features of all the reported cases of work-related homicides of physicians in Italy over the past 32 years.

Methods

We relied on the main national statistical databases (ISTAT, EURISPES, EU-RES) (23, 24, 25) and on PubMed to trace all cases of work-related homicides of doctors in Italy from 1988 to 2019. The group was completed using the documentation from the forensic psychiatric work of the Psychiatric Clinic of the University of Sassari.

While the literature mostly focuses on workplace violence, particularly in hospital settings, and is often primarily centred on psychiatric patients, our study has broadened the scope beyond hospital settings and psychiatric patients to include all cases of work-related doctor homicides.

Specific variables were extracted from each dataset and their value as a percentage of the total was calculated.

The victims' and the perpetrators' ages were grouped in 10-year intervals while the geographical location was divided into "Northern Italy" "Central Italy" and "Southern Italy and the Islands". The method of killing has been classified as "cold weapons", "firearms" and "other", while the locations of the homicide have been classified as "community-based clinics", "hospitals", "street", "perpetrator's home" and "victim's home". In addition to their gender (male/female), the offenders have also been classified according to whether they were psychiatric patients or not. A further distinction concerned the type of patient ("regular patient", "occasional patient", or "family member") and motive ("revenge", "crime of passion", "other").

Results

From 1988 to 2019, 21 physicians were killed in work-related circumstances in Italy.

The victims were more often male ($n = 15.7\%$), with a higher concentration (Fig. 1) in age group 50–60 ($n = 7$; 33.3%).

Geographically, the homicides were more prevalent in Southern Italy (South and Islands) with 57.1% of cases ($n = 12$), while only 9.5% were recorded in Central Italy ($n = 2$) and the remaining 33.3% in Northern Italy ($n = 7$).

As to method, 47.6% of the homicides were committed with firearms ($n = 10$) and 42.9% with sharp instruments/knives ($n = 9$). In one case (4.8%) a blunt weapon was used; in another case, the method was poisoning (4.8%).

With regard to location (Fig. 2) about half of the homicides took place outside hospitals, mainly at GPs' surgeries or out-of-hours primary care services. The most common locations, accounting for 28.6% ($n = 6$) of cases, were GPs' and out-of-hours surgeries. Next come mental health outpatient clinics, making up

19.0% (n = 4) of locations. A further 19.0% of murders occurred in the street (n = 4) while 14.3% of cases occurred at the victim's home (n = 3). Three cases took place in hospitals (including adjoining locations such as the car park). One homicide took place at the offender's home (4.8%).

With regard to the victims' medical specialties, the most common was psychiatry with an incidence of 28.6% (n = 6). This is followed by 4 murders of GPs (19%), 3 of primary care doctors in the out-of-hours service (14.3%) and 2 murders of urologists (9.5%). The remaining cases involved a forensic doctor, an anatomopathologist, a general surgeon, a gynaecologist, an oncologist and a neurosurgeon.

In 52.4% (n = 11) of cases, the perpetrators were patients under the victim's care, while 19.0% (n = 4) were occasional patients. The remaining 28.6% (n = 6) were family members of the victim's patients. These latter six cases were all driven by revenge, specifically 4 of the perpetrators who were family members (19.0% of the total) sought revenge for the patient's death. In 2 out of these latter 4 cases, the perpetrators were two fathers seeking revenge for the death of their little girls.

In 52.4% of the cases recorded (n = 11), the motive for the murder was revenge without previous threats. In 2 cases, the motive was a worsening of the murderer's health. A further 2 cases were crimes of passion, while in 3 other cases the revenge was for an erroneous diagnosis made by the victim. In only one case did the claimed motive originate after the victim had stopped treating the perpetrator. In another case, the motive for the murder is not entirely clear even though the offender, a psychiatric patient on probation, blamed both his victim and all the other doctors at the mental health clinic for the regime he was subjected to (i.e. for being required to report to the MHC every three days to receive treatment). In the last case, the murderer was a medical psychologist who was both a patient and a colleague of the victim, and who took revenge on his colleague for having been subjected to two compulsory mental health hospitalisation orders at the healthcare facility where he had previously been employed.

In 6 cases (28.6%), the motive was revenge preceded by the offence of stalking/threats. Specifically, 2 cases were passion-related: the victims were two women doctors who had long been subjected to threats and stalking by their patients. In one case the murder was preceded by a caution issued by the police.

However, 23.8% of cases (n = 5) are not explained by either passion or revenge. Indeed, 3 murders were committed in the out-of-hours general medical facilities by occasional patients (2 drug addicts and 1 inebriated person) while 2 other cases took place at Mental Health Centre outpatient clinics.

Analysis of the data collected also shows that in 14.3% (n = 3) of cases the homicide was followed by the offender's suicide and in 9.5% (n = 2) the perpetrator killed more than one person.

Lastly, 42.8% (n = 9) of the perpetrators had a psychiatric disorder while the remaining 57.1% (n = 12) had no diagnosed disorders at the time of the homicides (Fig. 3).

Discussion

Based on analysis of the data, some remarks can be made on the characteristics of the phenomenon and possible preventive measures.

Firstly, the literature and statistical data confirm that the medical profession is a dangerous one. In addition to the constant risk of professional liability due to an increasing number of malpractice claims, doctors are exposed to the risk of physical assaults at work and, in extreme and fortunately rare cases, of being killed in connection with their profession.

The literature highlights the aggressiveness and violence of patients in hospitals and particularly the risk associated with psychiatric patients. (21, 22, 26, 27, 28)

We have analysed the cases of physician homicides linked to the doctor-patient professional relationship, both within and outside clinical settings. The sample we examined supports more complex reflections.

The first reflection concerns the **location of the homicide**: in most cases it was not the hospital. Out of 21 cases, only 3 homicides were committed in the hospital or in nearby places (e.g. car park), while the other murders took place outside the hospital. In particular, a significant number of murders occurred in outpatient settings (community clinics), while still others occurred at the patient's home, at the doctor's home or in casual places (on the street). In a particularly striking instance, the patient took a diabolical revenge on the doctor: he sent a poison-laced bottle of wine as a Christmas gift to the victim's home for the Christmas holidays, causing the doctor's death. This unusual and horrifying case confirms that the homicidal intent may play out beyond healthcare settings and reach the victims elsewhere, including in their own home. The significant number of homicides in places other than hospitals calls into question the limited scope of preventive measures, which so far have focused on hospital settings. Some authors (29, 30) have reported a sharp decline in violent behaviour against healthcare workers after the introduction of specific security measures in hospitals. Similar systems would certainly also be effective in community clinics, where a certain number of homicides have taken place; these outpatient settings as a rule have insufficient security arrangements and would require the implementation of security protocols. However, security measures seem hardly of any use in cases where the homicide takes place in settings other than hospital and community healthcare facilities.

Another observation concerns the type of offender. In our sample, the perpetrators included not only the doctor's regular patients, but also patients' family members and occasional (drop-in) patients. This data also exposes the limitations of violent behaviour risk assessment and management systems in reducing the risk of violent behaviour by patients (27, 31). While these tools are certainly useful when dealing with patients registered with healthcare facilities, they may fail to prevent homicide in cases where the perpetrator is a patient's family member or an occasional/drop-in patient, as these types of individuals cannot be subjected to risk assessment. In the sample analysed, almost half of the perpetrators of homicide were patients' family members or occasional patients.

The sample reveals a "wide scope of danger" both in terms of location, which extends beyond the hospital setting and may also include the doctor's or the patient's home, and in terms of offenders, who

may also be patients' family members.

Some remarks can also be made about the motives for the murder in the sample. In a large number of cases the motive was revenge against the doctor for a claimed error in diagnosis, surgery or treatment causing harm to the patient. In a number of cases the murder did not occur as an escalation of an outburst of violence but was the outcome of planned and premeditated revenge against the doctor. Often the revenge was preceded by threats.

In some cases the murder was preceded by stalking. While the instances of stalking seem to be relatively few, their number may be underestimated also due to the fact that this behaviour has been recognised as an offence only in recent years. Therefore, it would be appropriate to highlight the role of stalking as a risk factor for homicide, and to raise public awareness of this behaviour to improve its management and prevent its escalation into violent acts. While of course stalking behaviour is not necessarily a precursor of murder, a useful recommendation is to report all stalking behaviours and take specific precautionary measures.

Finally, psychiatric patients warrant specific remarks. Nielssen's study (21) focused specifically on homicides committed by patients in psychiatric hospitals. In the sample we analysed, 43% of the perpetrators had a diagnosed psychiatric disorder, while 57% did not. We do not have data on how many offenders with psychiatric disorders were found to be not prosecutable due to mental impairment.

The debate about the link between mental illness and violent behaviour is still open and very controversial. In accordance with the literature and in light of the data examined here, we recommend that psychiatric patients be assessed for their risk of violent behaviour, particularly if they have a history of violent behaviour, current substance use and non-compliance with their medication. (27, 32, 33, 34, 35, 36).

However, it should also be noted that individuals (patients or family members) who are not mentally ill can also be offenders.

Conclusions

This paper is significantly limited by the small size of the sample group, which makes it difficult to draw general conclusions.

However, with this caveat, we can highlight seven things to know about the homicides of physicians.

1. The medical profession is a dangerous one. Murder is a rare event, but has devastating impacts on the morale of healthcare workers.
2. Physicians may be attacked not only in hospital settings but in other locations too, even at home. Isolated outpatient clinics are at high risk and would benefit from improved surveillance and security systems.

3. The perpetrator is not necessarily a patient, as doctors may be targeted by a family member of the patient seeking revenge.
4. Many cases stem from allegations of medical malpractice against the doctor. Conflict mediation systems should be set up for the parties' use. In Italy, the Gelli-Bianco Law, on the safety of healthcare and the professional liability of healthcare providers, has introduced compulsory mediation before legal action can be taken.
5. Stalking is a risk factor that should not be underestimated; doctors should be aware that it might precede violent behaviours. Physicians who are victims of stalking should always contact the police.
6. Psychiatric patients, especially those with a history of violent behaviour, substance use and poor adherence to their medication regimen should undergo a specific risk assessment for violent behaviour. However, the sample analysed suggests that the threat is not limited to psychiatric patients.
7. Physicians should receive specific training in effective communication skills and conflict management with patients and their family members, including non-violent communication techniques, listening skills and conflict identification and recognition.

While each case in our sample has unique characteristics, each can be considered a piece of a broader puzzle, which requires more studies in order to move beyond prejudice and silence.

Declarations

Ethics approval and consent to participate Not applicable

Consent for publication Not applicable

Availability of data and material All data analysed in this study are publicly available

Competing interests The authors declare that they have no competing interests

Funding No funding were used for this report

Authors' contribution LR, AMAN, ID, PM and CD analysed data. LR, AMAN, GN and SB interpreted data. LR, AMAN and GN wrote the first draft of the manuscript. LR, ID, PM, CD and SB revised and finalized the final manuscript. All authors read and approved the final manuscript.

Acknowledgments Not applicable

References

1. Di Martino V. Workplace violence in the health sector. Country case studies. Geneva: International Labour Office. International Council of Nurses, World Health Organization and Public Services International, 2002

2. Jenkins L. Current Intelligence Bulletin 57 – violence in the workplace. Risk factors and prevention strategies. The National Institute for Occupational Safety and Health (NIOSH). 1996.
3. *Sixth European Working Conditions Survey* (Luxembourg: Publications Office of the European Union). 2018. <https://www.eurofound.europa.eu/fr/publications/report/2016/working-conditions/sixth-european-working-conditions-survey-overview-report>
4. OSHA (Occupational Safety and Health Administration) Guidelines for preventing workplace violence for health care & social service workers. www.osha.gov. US Department of Labor. 2004.
5. Schulte JM, Nolt BJ, Williams RL, Spinks CL, Hellsten JJ. Violence and threats of violence experienced by public health field-workers. JAMA. 1998;280(5):439-442. doi:10.1001/jama.280.5.439
6. Al-Sahlawi KS, Zahid MA, Shahid AA, Hatim M, Al-Bader M. Violence against doctors: 1. A study of violence against doctors in accident and emergency departments. Eur J Emerg Med. 1999;6(4):301-304. doi:10.1097/00063110-199912000-00005
7. Elliott PP. Violence in health care. What nurse managers need to know. Nurs Manage. 1997;28(12):38-42.
8. Kingma M. Workplace violence in the health sector: a problem of epidemic proportion. Int Nurs Rev. 2001;48(3):129-130. doi:10.1046/j.1466-7657.2001.00094.x
9. Lee DT. Violence in the health care workplace. Hong Kong Med J. 2006;12(1):4-5.
10. Boyle M, Koritsas S, Coles J, Stanley J. A pilot study of workplace violence towards paramedics. Emerg Med J. 2007;24(11):760-763. doi:10.1136/emj.2007.046789
11. Gascón S, Martínez-Jarreta B, González-Andrade JF, Santed MA, Casalod Y, Rueda MA. Aggression towards health care workers in Spain: a multi-facility study to evaluate the distribution of growing violence among professionals, health facilities and departments. Int J Occup Environ Health. 2009;15(1):29-35. doi:10.1179/107735209799449707
12. Saeki K, Okamoto N, Tomioka K, et al. Work-related aggression and violence committed by patients and its psychological influence on doctors. J Occup Health. 2011;53(5):356-364. doi:10.1539/joh.11-0108-0a
13. Harrell E. Workplace Violence, 1993-2009. National Crime Victimization Survey and the Census of Fatal Occupational Injuries. Office of Justice Programs: Bureau of Ju. 2011.
14. De Jager L, Deneyer M, Buyl R, Roelandt S, Pacqueu R, Devroey D. Cross-sectional study on patient-physician aggression in Belgium: physician characteristics and aggression types. BMJ Open. 2019;9(12):e025942. Published 2019 Dec 18. doi:10.1136/bmjopen-2018-025942
15. Carmi-Iluz T, Peleg R, Freud T, Shvartzman P. Verbal and physical violence towards hospital- and community-based physicians in the Negev: an observational study. BMC Health Serv Res. 2005;5:54. Published 2005 Aug 15. doi:10.1186/1472-6963-5-54
16. Rippon TJ. Aggression and violence in health care professions. J Adv Nurs. 2000;31(2):452-460. doi:10.1046/j.1365-2648.2000.01284.x

17. Erkol H, Gökdoğan MR, Erkol Z, Boz B. Aggression and violence towards health care providers—a problem in Turkey?. *J Forensic Leg Med*. 2007;14(7):423-428. doi:10.1016/j.jflm.2007.03.004
18. Insel TR. Tribute to Wayne Fenton, MD (1953-2006). *Schizophr Res*. 2006;88(1-3):1-2. doi:10.1016/j.schres.2006.09.019
19. Goodman RA, Jenkins EL, Mercy JA. Workplace-related homicide among health care workers in the United States, 1980 through 1990. *JAMA*. 1994;272(21):1686-1688.
20. Loretto L, Falchi L, Nivoli FL, Milia P, Nivoli G, Nivoli AM. L'omicidio del medico [Murder of the doctor]. *Riv Psichiatr*. 2015;50(4):175-180. doi:10.1708/2002.21646
21. Nielssen O, Large MM. Homicide in psychiatric hospitals in Australia and New Zealand. *Psychiatr Serv*. 2012;63(5):500-503. doi:10.1176/appi.ps.201100290
22. Knable MB. Homicides of Mental Health Workers by Patients: Review of Cases and Safety Recommendations. *Psychiatr Ann*. 2017;47(6):325-334
23. <https://www.istat.it/it/archivio/omicidi>
24. <https://eurispes.eu/ricerca-rapporto/2019-2/>
25. EU- RES. <https://www.eures.it/>
26. O'Rourke M, Wrigley C, Hammond S. Violence within mental health services: how to enhance risk management. *Risk Manag Healthc Policy*. 2018;11:159-167. Published 2018 Sep 12. doi:10.2147/RMHP.S131834
27. Loretto L, Nivoli AMA,, Milia P, Nivoli G. Violence Risk Assessment in Mental Health. In *Violence and Mental Disorders* (Ed B.Carpiniello, A.Vita, C.Mencacci). Springer International Publishing, January 2020
28. Nivoli GC, Loretto L, Nivoli FL, Nivoli AM. Sulla psicopatologia e sulla valutazione del rischio dell'omicidio compiuto dal paziente con disturbo schizofrenico [Psychopathology and risk of homicide committed by a patient with schizophrenia]. *Riv Psichiatr*. 2011;46(2):140-147. doi:10.1708/626.7314
29. Ordog GJ, Wasserberger J, Ordog C, Ackroyd G, Atluri S. Weapon carriage among major trauma victims in the emergency department. *Acad Emerg Med*. 1995;2(2):109-114. doi:10.1111/j.1553-2712.1995.tb03170.x
30. Joint Commission Resources. [Preventing workplace violence: Tips for safety in emergency departments and psychiatric hospitals](#). *Environment of Care News*. 2008; 11:4.
31. Large M, Nielssen O. The limitations and future of violence risk assessment. *World Psychiatry*. 2017;16(1):25-26. doi:10.1002/wps.20394
32. Elbogen EB, Johnson SC. The intricate link between violence and mental disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2009;66(2):152-161. doi:10.1001/archgenpsychiatry.2008.537
33. Van Dorn R, Volavka J, Johnson N. Mental disorder and violence: is there a relationship beyond substance use?. *Soc Psychiatry Psychiatr Epidemiol*. 2012;47(3):487-503. doi:10.1007/s00127-011-

34. Nivoli G, Loretto L, Milia P, Nivoli A. La valutazione clinica del paziente con disturbo mentale e comportamento violento sulle persone. *Giornale Italiano di Psicopatologia*. 2008. 14;(4):1592-1107.
35. Fazel S, Långström N, Hjern A, Grann M, Lichtenstein P. Schizophrenia, substance abuse, and violent crime. *JAMA*. 2009;301(19):2016-2023. doi:10.1001/jama.2009.675
36. Friedman RA. Violence and mental illness—how strong is the link?. *N Engl J Med*. 2006;355(20):2064-2066. doi:10.1056/NEJMp068229

Figures

Figure 1

Age distribution of doctors murdered and their perpetrators in Italy, from 1988 to 2019.

Figure 2

Proportional distribution of murders of doctors by location in Italy, from 1988 to 2019

Figure 3

Clinical features of perpetrators of murdering of doctors in Italy, from 1988 to 2019