Effect of Short- and Long-term Melatonin Treatments on the Reproductive Activity of the Tropical Damselfish *Chrysiptera Cyanea*

Satoshi Imamura

University of the Ryukyus: Ryukyu Daigaku

Sung-Pyo Hur

Korea Institute of Ocean Science & Technology

Yuki Takeuchi

Okinawa Institute of Science and Technology: Gakko Hojin Okinawa Kagaku Gijutsu Daigakuin Daigaku Gakuen

Md. Badruzzaman

Bangabandhu Sheikh Mujibur Rahman Agricultural University

Angka Mahardini

Korea Institute of Ocean Science & Technology

Dinda Rizky

University of the Ryukyus: Ryukyu Daigaku

Akihiro Takemura (✉ takemura@sci.u-ryukyu.ac.jp)

University of the Ryukyus <https://orcid.org/0000-0002-5024-4930>

Research Article

Keywords: Aralkylamine N-acetyltransferase , GnRH , Gonadotropin , Kisspeptin , Photoperiod , Reproduction

Posted Date: September 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-895393/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Fish Physiology and Biochemistry on January 31st, 2022. See the published version at <https://doi.org/10.1007/s10695-022-01051-x>.

Effect of short- and long-term melatonin treatments on the reproductive activity of the tropical
damselfish *Chrysiptera cyanea*

Satoshi Imamura · Sung-Pyo Hur · Yuki Takeuchi¹ · Md. Badruzzaman · Angka Mahardini ·
Dinda Rizky · Akihiro Takemura

S. Imamura · Y. Takeuchi · M. Badruzzaman · A. Mahardini · D. Rizky · A. Takemura
Department of Chemistry, Biology and Marine Science, University of the Ryukyus, 1 Senbaru,
Nishihara, Okinawa 903-0213, Japan
e-mail address: takemura@sci.u-ryukyu.ac.jp (✉)

S.P. Hur · A. Mahardini
Jeju International Marine Science Research & Education Center, Korea Institute of Ocean
Science & Technology, Jeju Special Self-Governing Province 63349, South of Korea

Y. Takeuchi
Okinawa Institute of Science and Technology Graduate School, 1919-1 Tancha, Onna, Okinawa
904-0495, Japan

M. Badruzzaman
Department of Biochemistry and Molecular Biology, Bangabandhu Sheikh Mujibur Rahman
Agricultural University, Salna, Gazipur-1706, Bangladesh

Abstract

Photoperiod plays a role in controlling the initiation and termination of reproduction in fish. Melatonin is an internal transducer of environmental photoperiod and is involved in regulating reproduction. The present study aimed to examine how melatonin impacts the transcript levels of kisspeptin (*kiss1* and *kiss2*), gonadotropin-releasing hormones (*gnrh1*), and the β -subunit of gonadotropins (*fsh β* and *lh β*) in the brain of the sapphire devil, a tropical damselfish with long photoperiod preference. Feeding mature females with melatonin-containing pellets inhibited increases in the transcript levels of *kiss1*, *gnrh1*, and *lh β* within 3 h. Continuous melatonin treatment for 1 week resulted in oocyte regression and downregulation of *kiss2*, *gnrh1*, *fsh β* , and *lh β* . When the transcript levels of *kiss1* and *gnrh1* were measured at 4-h intervals in the brain of sapphire devil, a day–low/night–high fluctuation was observed. The hypothalamic–pituitary–gonadal (HPG) axis may be influenced by melatonin, exerting a negative effect at night because the transcript levels of aralkylamine *N*-acetyltransferase (*aanat2*) increased during the scotophase. The expression of *aanat2* was higher under short-day than long-day conditions, suggesting that there is a seasonal change in melatonin levels at night. It was concluded that change in photoperiod becomes a proximal factor for controlling the hormone synthesis in the HPG axis through melatonin.

Keywords

Aralkylamine *N*-acetyltransferase · GnRH · Gonadotropin · Kisspeptin · Photoperiod · Reproduction

Introduction

Melatonin is an indoleamine synthesized mainly in the pineal organ and the retina and plays a crucial role in various physiological processes including development (Danilova et al. 2004), growth (Falcón et al. 2003), and reproduction (Amano et al. 2000, 2004b; Vera et al. 2007). Melatonin levels fluctuate following a day–low/night–high cycle, which is controlled by the rate-limiting enzyme aralkylamine *N*-acetyltransferase (AANAT), which has a photodegradable characteristic and is affected by clock genes (Coon et al. 1999; Falcón et al. 2007). In addition to a daily cycle, certain fish also exhibit seasonal changes to melatonin secretion according to day length (García-Allegue et al. 2001; Bromage et al. 2001). Seasonal change in melatonin secretion is likely an endogenous cue for the initiation and termination of seasonal reproduction in fish since melatonin levels increase during the winter months, corresponding to the longer length of the night (Porter 2000; Porter et al. 2001; Bromage et al. 2001). Melatonin has previously been reported as a suppressor of reproductive activity in long-day breeders. For example, the gonadal development of the spotted snakehead *Channa punctatus* was inhibited after immersion in water containing melatonin, or after injection of melatonin (Renuka and Joshi 2010). On the other hand, melatonin acts as a stimulator of reproductive activity in short-day breeders. In underyearling precocious male masu salmon *Oncorhynchus masou*, testicular development was stimulated when fish were fed with melatonin-containing pellets, mimicking short-day conditions (Amano et al. 2000). However, the role of melatonin in reproductive activity may not be generalizable because of contradicting reports of the effect of melatonin. In the zebrafish *Danio rerio*, a long-day breeder, melatonin had a stimulatory effect on ovarian development when fish were reared in melatonin-containing water for 10 days (Carnevali et al. 2011). Interestingly, low-dose melatonin administration showed an opposite effect (stimulation) on testicular maturation in the masu salmon (Amano et al. 2000). Melatonin may impact

neurons synthesizing gonadotropin-releasing hormone (GnRH) in the brain as well as cells synthesizing gonadotropin (GtH) in the pituitary gland (Amano et al. 2000, 2004a; Ghosh and Nath 2005; Carnevali et al. 2011). Since the timing and amplitude of the changes in melatonin levels differ among fish species (Bromage et al. 2001), it is hypothesized that there is diverse adaptation in melatonin action on reproduction. In addition, it is not known how fish inhabiting low-latitude shallow waters can utilize melatonin for regulation of reproduction because little change in day length as well as water temperature occurs at these latitudes.

The sapphire devil *Chrysiptera cyanea* is a tropical damselfish that is widely found in shallow reefs of the Indo-West Pacific region. Previous reports showed that this species adapted to subtropical environments undergoes active spawning from May through August (Bapary et al. 2009) and that it can utilize changes in day length and food availability for the initiation and termination of seasonal reproduction (Bapary and Takemura 2010; Bapary et al. 2012). Immersing fish in melatonin-containing seawater inhibited the ovarian development of the sapphire devil (Badruzzaman et al. 2013), although how melatonin acts on the hypothalamic–pituitary–gonadal (HPG) axis remains unclear. The present study aimed to evaluate the short-term (days) and long-term (weeks) effects of melatonin treatment on mRNA levels of kisspeptin (*kiss1* and *kiss2*), GnRH (*gnrh1*), and the β -subunit of GtH (*fsh β* and *lh β*) in the brain of the sapphire devil. Diurnal profiles of *kiss1*, *gnrh1*, and *aanat2* in the brain were also studied to confirm the relationship between their abundance and the effect of melatonin. The present study focused on *aanat2* because it expresses specifically in the pineal organ of fish (Coon et al. 1999; Isorna et al. 2011).

Materials and methods

Animals

98

99 Mature sapphire devils (1.35–2.83 g body weight) were collected on coral reefs around Sesoko
100 Island, Okinawa, Japan, using a hand net during the daytime at low tide. The fish were
101 acclimated in stock tanks with running seawater and ambient aeration under natural photoperiod
102 and ambient water temperature at Sesoko Station, Tropical Biosphere Research Center,
103 University of the Ryukyus, Okinawa, Japan. Fish were fed daily at 0900 h with commercial
104 pellets (EP1; Nisshin-Marubeni, Tokyo, Japan).

105

106 Melatonin treatment

107

108 Melatonin was purchased from Sigma-Aldrich (St. Luis, MO, USA) and dissolved with
109 absolute ethanol. Melatonin-containing pellets were prepared by spraying pellets with a
110 melatonin-containing ethanol solution, as previously described (Badruzzaman et al. 2013). The
111 final concentration of melatonin was adjusted to 500 µg/g dry pellets. Vehicle without melatonin
112 was sprayed onto pellets as a control (melatonin-free pellets).

113 The experiments were conducted to evaluate the effects of short- and long-term treatment
114 with melatonin on the mRNA levels of reproduction-related genes (*kiss1*, *kiss2*, *gnrh1*, *fshβ*,
115 and *lhβ*) in the brain of the sapphire devil. The experiments were carried out in May and June
116 when this species undergoes active reproduction (Bapary et al. 2009). Fish were randomly taken
117 from the stock tanks and transferred to four glass aquaria (60 L capacity) equipped with running
118 seawater and ambient aeration under natural photoperiod and water temperature. Plastic pipes
119 were provided as shelter and spawning nest and were placed on the bottom of each aquarium.
120 The two aquaria contained 20 females and 5 males per tank, and the treatment tanks contained
121 16 females and 4 males per tank. After acclimation for several days, fish were fed with pellets
122 twice per day, at 0900 h and 1500 h. The daily ration of food was adjusted to 2.0% of body

weight throughout the experimental period.

For the short-term melatonin treatment, females ($n = 8$) were sampled from the aquarium of the control group at 0900 h (initial control), and then from the aquariums of both groups at 1200 h and 1500 h. For the long-term melatonin treatment, females ($n = 8$) in each aquarium were sampled at 1200 h after 1 and 2 weeks. Following anesthetization with 2-phenoxyethanol (Kanto Chemical, Tokyo, Japan) and euthanization by decapitation, body mass and ovarian mass of all individuals were recorded. Then, pieces of the ovary were fixed in Bouin's solution for subsequent histological observation. Gonadosomatic index (GSI) was calculated using the following formula: $GSI = (\text{ovarian mass} / \text{body mass}) \times 100$. The whole brain was excised, immediately frozen in liquid nitrogen, and stored at -80°C until extraction of total RNA from the tissue samples using TriPure Isolation Reagent (Roche Diagnostics, Indianapolis, IN, USA) according to the manufacturer's instructions.

Daily variation of sapphire devil *kiss1* and *gnrh1*

Females ($n = 42$) were housed in filtered aquaria (60 L capacity) with ambient aeration under natural photoperiodic conditions (long-day; 0538 – 1926 h). After acclimation, fish ($n = 6$ per sampling time) were taken from the aquarium at 4-h intervals. After anesthesia, the whole brain was excised, immediately frozen in liquid nitrogen, and stored at -80°C until total RNA extraction. Sample collections during the scotophase were performed under dim red light.

Daily variation of sapphire devil *aanat2* under programmed photoperiod

This experiment was conducted in February using females ($n = 72$) with body mass ranging from 1.25 to 1.85 g, which were housed in filtered aquaria with ambient aeration under short-

day (LD = 10:14; light-on from 0800 – 1800 h) and long-day (LD = 14:10; light-on from 0600 – 1800 h), which were set using a 20 W fluorescent light (1200 lx at water surface). After acclimation under these conditions for at least 1 week, fish (n = 5 per sampling time) were taken from aquaria at 4-h intervals. The following procedures were followed by the abovementioned methods.

Histological procedures

Following dehydration with a series of ethanol and permutation with xylene, ovaries were embedded in histoparaffin (m.p. 56–58°C; Merck, Darmstadt, Germany), serially sectioned at 5 µm, and then stained with hematoxylin and eosin for microscopic observation. According to oocyte staging of the white-spotted spinefoot *Siganus canaliculatus* (Hoque et al. 1998), oocytes were classified into the following six stages: the peri-nucleolus stage (PNS), oil droplet stage (ODS), yolk vesicle stage (YVS), primary yolk stage (PYS), secondary yolk stage (SYS), and tertiary yolk stage (TYS). Atretic oocyte (AO) was also observed in the present study.

Real-time quantitative PCR (qPCR)

The transcript levels of *kiss1*, *kiss2*, *gnrh1*, *fshβ*, *lhβ*, and *aanat2* in the brain of the sapphire devil were assayed using the CFX96 Real Time System (Bio-Rad, Hercules, CA, USA) and GoTaq qPCR Master Mix (Promega, Madison, WI, USA). Total RNA was extracted from the whole brain of the sapphire devil. The first strand of cDNA was reverse-transcribed from 1 µg of total RNA using the PrimeScript RT Reagent Kit with gDNA Eraser (Takara Bio, Otsu, Japan). Specific qPCR primers were designed based on the full-length of the cDNA fragments (Table 1), which were previously cloned and characterized (Imamura et al. 2020). Each PCR

was carried out in a final volume of 10 µl, containing 5 µl of 2× GoTaq qPCR Master Mix, 0.3 µl of forward and reverse primers, 2.4 µl of nuclease-free water, and 2 µl of cDNA template. The qPCR cycling conditions were 95°C for 2 min, then 40 cycles at 95°C for 15 sec and 60°C for 1 min. Plasmid DNA as standard was produced using the gene-specific primers. cDNA at 10-fold dilution was used for qPCR to construct the standard curve. The transcript levels of the target genes were normalized to those of elongation factor 1 alpha (*ef1α*), which served as an internal control.

Statistical analyses

All data are expressed as the mean ± standard error of the mean (SEM). Changes in GSI, as well as mRNA levels of reproduction-related genes (*kiss1*, *kiss2*, *gnrh1*, *fshβ*, and *lhβ*), were compared by two-way analysis of variance (ANOVA) followed by Tukey's test. Diurnal variation of *kiss1*, *gnrh1*, and *aanat* were compared by one-way ANOVA followed by Tukey's test using $P < 0.05$ as the level of statistical significance.

Results

Effects of melatonin treatment on ovarian development and reproduction-related genes

The mature female sapphire devils were fed with melatonin-containing pellets (treatment group) or melatonin-free pellets (control group) at 0900 h and sacrificed at 0, 3, and 6 h after treatment. Females of the initial control had high GSI values. This value did not change within 6 hours after treatment (Fig. 1A). Histological observation revealed that ovaries of both groups contained vitellogenic oocytes at TYS (Supplementary data 1). The transcript levels of

reproduction-related genes (*kiss1*, *kiss2*, *gnrh1*, *fshβ*, and *lhβ*) were determined in the brain of the sapphire devil (Fig. 1B – F). The transcript levels of *kiss1* and *gnrh1* increased significantly ($P < 0.05$) in the brain of the control fish at 3 h. The transcript levels of *kiss1*, *gnrh1*, and *lhβ* decreased significantly in the melatonin-treated fish at 3 h ($P < 0.05$). However, there was little difference in the transcript levels of these genes between the groups at 6 h (Fig. 1B and D). Melatonin treatment did not affect the transcription levels of *kiss2* and *fshβ* in the brain of sapphire devil females (Fig. 1C and E).

Feeding with melatonin-containing or melatonin-free pellets lasted for 2 weeks (Fig. 2). High GSI was maintained in the control group throughout the experimental period. When fish were treated with melatonin, GSI significantly decreased within 2 weeks (Fig. 2A). Melatonin treatment had an impact on the transcript levels of *kiss2*, *gnrh1*, *fshβ*, and *lhβ*, but not *kiss1*. Significantly lower mRNA levels of *kiss2* at 1 week, and *gnrh1*, *fshβ*, and *lhβ* at 1 and 2 weeks were observed (Fig. 2B – F). Histological observation revealed that all ovaries at 1 and 2 weeks after treatment contained vitellogenic oocytes at TYS. AO and immature oocytes at PNS were observed in ovaries of the melatonin-treated fish (Supplementary data 2).

Diurnal variation of *kiss1* and *gnrh1* mRNA during spawning season

Fish were sampled at 4-h intervals to evaluate the daily profiles of *kiss1* and *gnrh1* in the brain (Fig. 3). Both genes showed clear daily variation with an increase during photophase and a decrease during scotophase. A peak of mRNA abundance occurred at 1200 h for *kiss1* (Fig. 3A) and 1600 h for *gnrh1* (Fig. 3B).

Diurnal variation of *aanat2* mRNA under long- and short-day conditions

Fish were reared under short- and long-day conditions to investigate the daily profile of *aanat2* in the brain. The transcript levels of *aanat2* fluctuated with an increase toward the start of scotophase in both conditions. A peak level occurred at Zeitgeber time (ZT)13 for short-day conditions (Fig. 4A) and ZT17 for long-day conditions (Fig. 4B).

Discussion

Administration of melatonin has previously been reported to affect—either positively or negatively—the gonadal activity of fishes (Urasaki 1972; Sundararaj and Keshavanath 1976; Ghosh and Nath 2005; Renuka and Joshi 2010). However, the results obtained may be due to intraspecies differences in the effect of melatonin and differences in experimental design. For example, intraperitoneal injections of melatonin inhibited the ovarian development of the female walking catfish *Clarias batrachus* (Ghosh and Nath 2005) and medaka *Oryzias latipes* (a long-day spawner) (Urasaki 1972). On the other hand, ovarian development of the zebrafish was stimulated when fish were reared in melatonin-containing water under long-day conditions (Carnevali et al. 2011). Our results support an inhibitory effect of melatonin treatment on ovarian development including vitellogenesis of the sapphire devil because feeding with melatonin-containing pellets for 1 to 2 weeks resulted in the induction of atresia and the arrest of recruitment in vitellogenic oocytes in the ovaries, as previously reported (Badruzzaman et al. 2013). Therefore, it is likely that melatonin acts on the synthesis and/or secretion of hormones in the HPG axis because the direct action of melatonin is exerted through the binding of G-protein-coupled melatonin receptors (Dubocovich et al. 2010), which are expressed in the diencephalon (Ikegami et al. 2009), pituitary gland (Gaildrat and Falcón 2000), and ovary of fish (Jin et al. 2013; Chai et al. 2013; Hong et al. 2014). The present study also revealed that treatment with melatonin for at least 1 week is needed to retract ovarian features, because no

change was observed in vitellogenic oocytes after melatonin treatment for 3 or 6 h.

The present study revealed that there is a negative impact of melatonin on the transcript levels of certain reproduction-related genes in the HPG axis. Unlike GSI and histological features of the ovary, changes in the transcript levels of *kiss1*, *gnrh1*, and *lhβ*, but not *kiss2* and *fshβ*, occurred within 3 h of melatonin treatment. These results demonstrate that melatonin suppresses a diurnal increase in the mRNA levels of melatonin-treated fish, suggesting that melatonin acts as a diurnal driver of *kiss1* and *gnrh1* transcription. Previous studies have reported diurnal variation of *kiss1* and *gnrh1* transcription levels in the brain of certain teleosts; sbGnRH decreased during the dark period in the pituitary of the European seabass *Dicentrarchus labrax* reared under both long-day and short-day conditions (Bayarri et al. 2004). The transcript levels of *kiss2* and *gnrh1* had a marked day–night difference with an increase at 1200 h in the brain of orange-spotted grouper *Epinephelus coioides* in developing and mature stages (Chai et al. 2013). In the latter case, the melatonin receptor (MT1) had a reciprocal relationship in regulating the day–night variation of *gnrh1* expression directly or indirectly through Kiss2 (Chai et al. 2013). A similar regulation is likely to occur in the sapphire devil because a day–high/night–low fluctuation of *kiss1* and *gnrh1* transcription was observed in the brain of female sapphire devils (Fig. 3). An inverse profile of *aanat2* (Fig. 4) also supports the involvement of melatonin in the downregulation of *kiss1* and *gnrh1*, since AANAT is the rate-limiting enzyme in melatonin synthesis in the pineal organ in fish (Coon et al. 1999; Isorna et al. 2011). The expression of *lhβ* was also downregulated in the brain of the sapphire devil, suggesting that *lhβ* transcription is a downstream target of GnRH1. In addition, there is likely a direct action of melatonin through its receptor in the pituitary gland because the expression of melatonin receptors was evidenced in certain teleosts including the pike *Esox lucius* (Gaildrat et al. 2002), the European sea bass (Sauzet et al. 2008), and the grass puffer *Takifugu niphobles* (Ikegami et al. 2009). Little difference in the transcript levels of *kiss1*, *gnrh1*, and *lhβ* was

observed in the brain of the sapphire devil at 6 h after melatonin treatment, suggesting clearance of exogenous melatonin and decrement of its effectiveness. In this regard, melatonin was reported to be eliminated within 2 h from the blood circulation of the threespot wrasse *Halichoeres trimaculatus* when melatonin was injected by intraperitoneal injection (Takemura et al. 2010).

We also evaluated the effect of weekly treatment with melatonin on ovarian development and gene expression in the sapphire devil. This experiment was founded on the previous finding that vitellogenic oocytes could be induced in the ovary when fish out of the breeding season were reared under long-day conditions within a defined range of water temperature (Bapary and Takemura 2010). The present study revealed that, in agreement with the regression and immature features of the ovary, melatonin treatment decreases *kiss2*, *gnrh1*, *fsh β* , and *lh β* transcription levels in the brain of the sapphire devil. Thus, long-term treatment with melatonin has a negative impact on the transcript levels of reproductive genes in the HPG axis. Similar findings have been shown in certain fish species undergoing seasonal reproduction (Ghosh and Nath 2005; Sébert et al. 2008; Fukunaga et al. 2019). For example, daily injections of melatonin for 2 weeks decreased the plasma levels of GtH in the walking catfish *C. batrachus* (Ghosh and Nath 2005), and an implanted osmotic pump releasing physiological levels of melatonin downregulated *lh β* in the pituitary of the honeycomb grouper *Epinephelus merra* (Fukunaga et al. 2019). Since the honeycomb grouper and the sapphire devil are both tropical species with long-day preference, it is likely that a decrease in photoperiod (an increase in melatonin effect at night) is related to the termination of breeding activity. It was previously reported that immersing female sapphire devils in estradiol-17 β (E2)-containing water resulted in increased expression levels of *kiss2* and *gnrh1* in the brain, suggesting that E2 has a positive feedback on kiss-expressing neurons in the diencephalon, and in turn, upregulates *gnrh1* in the preoptic region (Imamura et al. 2020). The inhibition of this melatonin feedback control may accelerate

regression of gonadal development in the sapphire devil.

The present study showed that higher transcription of *aanat2* was maintained in the brain of the sapphire devil when fish were reared under short-day conditions. This result indicates that the period of melatonin synthesis is longer under short-day conditions than long-day conditions since melatonin synthesis is dependent upon the expression of *aanat2* in the pineal gland (Herrera-Pérez et al. 2011; McStay et al. 2014). Since melatonin levels in the pineal organ vary according to photoperiodic conditions, seasonal changes in melatonin may be related to the recognition of the season (García-Allegue et al. 2001).

In conclusion, the ovarian development of the sapphire devil is influenced by melatonin, which has a negative impact on hormone synthesis at the HPG axis. Gradual increases in the nocturnal duration of melatonin synthesis in the seasonal progression toward winter may trigger the termination of the spawning season in long-day spawners, including the sapphire devil. Further studies would be needed to clarify how melatonin is involved in the initiation and termination of ovarian development through the circadian system in the brain.

Acknowledgements

We gratefully thank to staff of Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus, Okinawa, Japan, for use of facilities.

Author Declarations

Funding

This study was supported in part by a Grant-in-Aid for Scientific Research (B) (KAKENHI, Grant number 16H05796) from the Japan Society for the Promotion of Science (JSPS) to AT

323 and Heiwa Nakajima Foundation to AT.

324

325 **Conflicts of interest**

326 The authors declare that they have no conflict of interest.

327

328 **Ethical approval**

329 All experiments were conducted in compliance with the guidelines of the Animal Care and Use

330 Committee of the University of the Ryukyus and the regulations for the care and use of

331 laboratory animals in Japan.

332

333 **Consent to participate**

334 Not applicable.

335

336 **Consent for publication**

337 Not applicable.

338

339 **Availability of data and material**

340 All data related to this study are included in the manuscript.

341

342 **Code availability**

343 Not applicable

344

345 **Author's contribution**

346 All authors contributed to the design of this research. SI, SPH, and MB collected fish from coral

347 reefs and maintained them under experimental conditions. SI and YT sampled tissues and

carried out molecular analyses. SI, AM, DR analyzed data statistically and discussed about the results under supervision of AT. All authors contributed to the manuscript preparation. All authors revised and commented on all versions of the manuscript, and then approved the final manuscript.

References

- Amano M, Iigo M, Ikuta K, et al (2004a) Disturbance of plasma melatonin profile by high dose melatonin administration inhibits testicular maturation of precocious male masu salmon. *Zoolog Sci* 21:79–85. [https://doi.org/10.2108/0289-0003\(2004\)21\[79:DOPMPB\]2.0.CO;2](https://doi.org/10.2108/0289-0003(2004)21[79:DOPMPB]2.0.CO;2)
- Amano M, Iigo M, Ikuta K, et al (2000) Roles of melatonin in gonadal maturation of underyearling precocious male masu salmon. *Gen Comp Endocrinol* 120:190–197. <https://doi.org/10.1006/gcen.2000.7547>
- Amano M, Okubo K, Yamanome T, et al (2004b) Changes in brain GnRH mRNA and pituitary GnRH peptide during testicular maturation in barfin flounder. *Comp Biochem Physiol Part B Biochem Mol Biol* 138:435–443. <https://doi.org/10.1016/j.cbpc.2004.05.005>
- Badruzzaman M, Bapary MAJ, Takemura A (2013) Possible roles of photoperiod and melatonin in reproductive activity via changes in dopaminergic activity in the brain of a tropical damselfish, *Chrysiptera cyanea*. *Gen Comp Endocrinol* 194:240–247. <https://doi.org/10.1016/j.ygcen.2013.09.012>

369 Bapary MAJ, Fainuulelei P, Takemura A (2009) Environmental control of gonadal
 370 development in the tropical damselfish *Chrysiptera cyanea*. Mar Biol Res 5:462–469.
 371 <https://doi.org/10.1080/17451000802644722>
 372 Bapary MAJ, Nurul Amin M, Takemura A (2012) Food availability as a possible determinant
 373 for initiation and termination of reproductive activity in the tropical damselfish
 374 *Chrysiptera cyanea*. Mar Biol Res 8:154–162.
 375 <https://doi.org/10.1080/17451000.2011.605146>
 376 Bapary MAJ, Takemura A (2010) Effect of temperature and photoperiod on the reproductive
 377 condition and performance of a tropical damselfish *Chrysiptera cyanea* during different
 378 phases of the reproductive season. Fish Sci 76:769–776. [https://doi.org/10.1007/s12562-](https://doi.org/10.1007/s12562-010-0272-0)
 379 [010-0272-0](https://doi.org/10.1007/s12562-010-0272-0)
 380 Bayarri MJ, Rodríguez L, Zanuy S, et al (2004) Effect of photoperiod manipulation on the daily
 381 rhythms of melatonin and reproductive hormones in caged European sea bass
 382 (*Dicentrarchus labrax*). Gen Comp Endocrinol 136:72–81
 383 Bromage N, Porter M, Randall C (2001) The environmental regulation of maturation in farmed
 384 finfish with special reference to the role of photoperiod and melatonin. Aquaculture
 385 197:63–98. <https://doi.org/10.1016/B978-0-444-50913-0.50008-4>
 386 Carnevali O, Gioacchini G, Maradonna F, et al (2011) Melatonin induces follicle maturation in
 387 *Danio rerio*. PLoS One 6:e19978

388 Chai K, Liu X, Zhang Y, Lin H (2013) Day-night and reproductive cycle profiles of melatonin
 389 receptor, kiss , and gnrh expression in orange-spotted grouper (*Epinephelus coioides*). Mol
 390 Reprod Dev 80:535–548. <https://doi.org/10.1002/mrd.22191>
 391 Coon SL, Begay V, Deurloo D, et al (1999) Two arylalkylamine N-acetyltransferase genes
 392 mediate melatonin synthesis in fish. J Biol Chem 274:9076–9082
 393 Danilova N, Krupnik VE, Sugden D, Zhdanova I V. (2004) Melatonin stimulates cell
 394 proliferation in zebrafish embryo and accelerates its development. FASEB J 18:751–753.
 395 <https://doi.org/10.1096/fj.03-0544fje>
 396 Dubocovich ML, Delagrange P, Krause DN, et al (2010) International Union of Basic and
 397 Clinical Pharmacology. LXXV. Nomenclature, classification, and pharmacology of G
 398 protein-coupled melatonin receptors. Pharmacol Rev 62:343–380.
 399 <https://doi.org/10.1124/pr.110.002832>
 400 Falcón J, Besseau L, Fazzari D, et al (2003) Melatonin modulates secretion of growth hormone
 401 and prolactin by trout pituitary glands and cells in culture. Endocrinology 144:4648–4658.
 402 <https://doi.org/10.1210/en.2003-0707>
 403 Falcón J, Besseau L, Sauzet S, Boeuf G (2007) Melatonin effects on the hypothalamo–pituitary
 404 axis in fish. Trends Endocrinol Metab 18:81–88.
 405 <https://doi.org/10.1016/j.tem.2007.01.002>
 406 Fukunaga K, Yamashina F, Ohta N, et al (2019) Involvement of melatonin in transducing

407 moon-related signals into the reproductive network of the female honeycomb grouper
 408 *Epinephelus merra*. Gen Comp Endocrinol 282:113211.
 409 <https://doi.org/10.1016/j.ygcen.2019.113211>
 410 Gaildrat P, Becq F, Falcón J (2002) First cloning and functional characterization of a melatonin
 411 receptor in fish brain: A novel one? J Pineal Res 32:74–84. [https://doi.org/10.1034/j.1600-](https://doi.org/10.1034/j.1600-079x.2002.1817.x)
 412 079x.2002.1817.x
 413 Gaildrat P, Falcón J (2000) Melatonin receptors in the pituitary of a teleost fish: mRNA
 414 Expression, 2-[¹²⁵I]iodomelatonin binding and cyclic AMP response. Neuroendocrinology
 415 72:57–66. <https://doi.org/10.1159/000054571>
 416 García-Allegue R, Madrid JA, Sánchez-Vázquez FJ (2001) Melatonin rhythms in European sea
 417 bass plasma and eye: influence of seasonal photoperiod and water temperature. J Pineal
 418 Res 31:68–75. <https://doi.org/10.1034/j.1600-079X.2001.310110.x>
 419 Ghosh J, Nath P (2005) Seasonal effects of melatonin on ovary and plasma gonadotropin and
 420 vitellogenin levels in intact and pinealectomized catfish, *Clarias batrachus* (Linn). Indian
 421 J Exp Biol 43:224–232
 422 Herrera-Pérez P, Servili A, Rendón MC, et al (2011) The pineal complex of the European sea
 423 bass (*Dicentrarchus labrax*): I. Histological, immunohistochemical and qPCR study. J
 424 Chem Neuroanat 41:170–180. <https://doi.org/10.1016/j.jchemneu.2011.01.006>
 425 Hong LY, Hong WS, Zhu WB, et al (2014) Cloning and expression of melatonin receptors in

426 the mudskipper *Boleophthalmus pectinirostris*: Their role in synchronizing its semilunar
 427 spawning rhythm. Gen Comp Endocrinol 195:138–150.
 428 <https://doi.org/10.1016/j.ygcen.2013.11.004>
 429 Hoque M, Takemura A, Takano K (1998) Annual changes in oocyte development and serum
 430 vitellogenin level in the rabbitfish *Siganus canaliculatus* (Park) in Okinawa, Southern
 431 Japan. Fish Sci 64:44–51. <https://doi.org/10.2331/fishsci.64.44>
 432 Ikegami T, Motohashi E, Doi H, et al (2009) Synchronized diurnal and circadian expressions
 433 of four subtypes of melatonin receptor genes in the diencephalon of a puffer fish with
 434 lunar-related spawning cycles. Neurosci Lett 462:58–63.
 435 <https://doi.org/10.1016/j.neulet.2009.06.076>
 436 Imamura S, Hur S-P, Takeuchi Y, et al (2020) The mRNA expression patterns of kisspeptins,
 437 GnRHs, and gonadotropins in the brain and pituitary gland of a tropical damselfish,
 438 *Chrysiptera cyanea*, during the reproductive cycle. Fish Physiol Biochem 46:277–291.
 439 <https://doi.org/10.1007/s10695-019-00715-5>
 440 Isorna E, Aliaga-Guerrero M, M'Rabet A El, et al (2011) Identification of two arylalkylamine
 441 N-acetyltransferase 1 genes with different developmental expression profiles in the flatfish
 442 *Solea senegalensis*. J Pineal Res 51:434–444. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-079X.2011.00907.x)
 443 [079X.2011.00907.x](https://doi.org/10.1111/j.1600-079X.2011.00907.x)
 444 Jin YH, Park JW, Kim J, Kwon JY (2013) The expression pattern of melatonin receptor 1a gene

445 during early life stages in the Nile tilapia (*Oreochromis niloticus*). Dev Reprod 17:45–53.

446 <https://doi.org/10.12717/DR.2013.17.1.045>

447 McStay E, Migaud H, Vera LM, et al (2014) Comparative study of pineal clock gene and

448 AANAT2 expression in relation to melatonin synthesis in Atlantic salmon (*Salmo salar*)

449 and European seabass (*Dicentrarchus labrax*). Comp Biochem Physiol - A Mol Integr

450 Physiol 169:77–89. <https://doi.org/10.1016/j.cbpa.2013.12.011>

451 Porter M (2000) Environmental influences on melatonin secretion in Atlantic cod (*Gadus*

452 *morhua* L.) and their relevance to commercial culture. Fish Physiol Biochem 23:191–200

453 Porter MJR, Duncan N, Handeland SO (2001) Temperature, light intensity and plasma

454 melatonin levels in juvenile Atlantic salmon. J Fish Biol 58:431–438.

455 <https://doi.org/10.1006/jfbi.2000.1455>

456 Renuka K, Joshi BN (2010) Melatonin-induced changes in ovarian function in the freshwater

457 fish *Channa punctatus* (Bloch) held in long days and continuous light. Gen Comp

458 Endocrinol 165:42–46. <https://doi.org/10.1016/j.ygcen.2009.05.020>

459 Sauzet S, Besseau L, Herrera Perez P, et al (2008) Cloning and retinal expression of melatonin

460 receptors in the European sea bass, *Dicentrarchus labrax*. Gen Comp Endocrinol 157:186–

461 195. <https://doi.org/10.1016/j.ygcen.2008.04.008>

462 Sébert M-E, Legros C, Weltzien F-A, et al (2008) Melatonin activates brain dopaminergic

463 systems in the eel with an inhibitory impact on reproductive function. J Neuroendocrinol

464 20:917–929. <https://doi.org/10.1111/j.1365-2826.2008.01744.x>

465 Sundararaj BI, Keshavanath P (1976) Effects of melatonin and prolactin treatment on the

466 hypophyseal-ovarian system in the catfish, *Heteropneustes fossilis* (Bloch). Gen Comp

467 Endocrinol 29:84–96. [https://doi.org/10.1016/0016-6480\(76\)90010-1](https://doi.org/10.1016/0016-6480(76)90010-1)

468 Takemura A, Uchimura M, Shibata Y (2010) Dopaminergic activity in the brain of a tropical

469 wrasse in response to changes in light and hydrostatic pressure. Gen Comp Endocrinol

470 166:513–519. <https://doi.org/10.1016/j.ygcen.2010.01.001>

471 Urasaki H (1972) Effects of restricted photoperiod and melatonin administration on gonadal

472 weight in the Japanese killifish. J Endocrinol 55:619–620.

473 <https://doi.org/10.1677/joe.0.0550619>

474 Vera LM, De Oliveira C, López-Olmeda JF, et al (2007) Seasonal and daily plasma melatonin

475 rhythms and reproduction in Senegal sole kept under natural photoperiod and natural or

476 controlled water temperature. J Pineal Res 43:50–55. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-079X.2007.00442.x)

477 [079X.2007.00442.x](https://doi.org/10.1111/j.1600-079X.2007.00442.x)

478

Table 1. Real-time PCR Primer sets used in the present study

Primer	Sequence
<i>gnrh1</i> -Forward	5'-GGGCAACATAGCTGAAGGATT-3'
<i>gnrh1</i> -Reverse	5'-GTTCGGTGTCCATTTTCCTTG-3'
<i>kiss1</i> -Forward	5'-TCAGCAGAAGTCCACACCAG-3'
<i>kiss1</i> -Reverse	5'-ATCTTGGCGATCCACCATC-3'
<i>kiss2</i> -Forward	5'-CGTGGTTGTTGTGTGTTCG-3'
<i>kiss2</i> -Reverse	5'-GAACAGAGAAAACGGGGAGA-3'
<i>fshβ</i> -Forward	5'-AGCGTGGAGTCGGTGTTCA-3'
<i>fshβ</i> -Reverse	5'-CCGCTGAAGTGTTTGACCTC-3'
<i>lhβ</i> -Forward	5'-TTGAGTTTGTGTCTGGGAGCT-3'
<i>lhβ</i> -Reverse	5'-CTCAGAGCCACTGGGTAGGT-3'
<i>aanat2</i> -Forward	5'-CGTCTTGCCTGTTGAAGTGA-3'
<i>aanat2</i> -Reverse	5'-AGGGTTTGAGGAACGGTGAG-3'
<i>eflα</i> -Forward	5'-ACGTGTCCGTCAAGGAAATC-3'
<i>eflα</i> -Reverse	5'-GGGTGGTTCAGGATGATGAC-3'

Figure legends

- Fig. 1. Short-term effect of melatonin treatment on gonadosomatic index (GSI) (A) and relative expression levels of reproduction-related genes (*kiss1*, *kiss2*, *gnrh1*, *fsh β* , and *lh β*) (B – F) in the female sapphire devil. Fish were fed with melatonin-free pellets or melatonin-containing pellets at 0900 h. Ovary and brain were taken from fish at 0, 3, and 6 h after feeding. White and black columns indicate the data from the fish feeding with melatonin-free pellets and melatonin-containing pellets, respectively. Each value represents mean $\pm$ SEM. Different letters indicate significant difference at $P < 0.05$.
- Fig. 2. Long-term effect of melatonin treatment on gonadosomatic index (GSI) and relative expression levels of reproduction-related genes (*kiss1*, *kiss2*, *gnrh1*, *fsh β* , and *lh β*) in the female sapphire devil. Fish were fed with melatonin-free pellets or melatonin-containing pellets at 0900 h. Ovary and brain were taken from fish at 0, 1, and 2 weeks after feeding. White and black columns indicate the data from the fish feeding with melatonin-free pellets and melatonin-containing pellets, respectively. Each value represents mean $\pm$ SEM. Different letters indicate significant difference at $P < 0.05$.
- Fig. 3. Daily changes in mRNA abundance of *kiss1* (A) and *gnrh1* (B) in the brain of female sapphire devil under natural conditions. Each value represents mean $\pm$ SEM. Horizontal bar with white and black bars in the figure indicates light and dark phase, respectively. Different letters indicate significant difference at $P < 0.05$.
- Fig. 4. Daily changes in mRNA abundance of *aanat2* in the brain of female sapphire devil under conditions of long-day (A; LD = 14:10) and short-day (B; LD = 10:14). Each value represents mean $\pm$ SEM. Horizontal bar with white and black bars in the figure indicates light and dark phase, respectively. Different letters indicate significant difference at $P < 0.05$.

Supplementary data 1. Effect of single treatment with melatonin on ovarian histology of the

sapphire devil. (A) Cross section of an ovary (CSO) of fish at the onset of the experiment, (B) CSO of the fish 3 h after feeding with melatonin-free pellets, (C) CSO of the fish 3 h after feeding with melatonin-containing pellets, (D) CSO of the fish 6 h after feeding with melatonin-containing pellets. TYS; tertiary yolk stage. Scale bar = 200 μm .

Supplementary data 2. Effect of daily treatment with melatonin on ovarian histology of the

sapphire devil. (A) Cross section of an ovary (CSO) of fish at the onset of the experiment, (B) CSO of the fish fed with melatonin-free pellets for 2 weeks (C) CSO of the fish fed with melatonin-containing pellets for 1 week, (D) CSO of the fish fed with melatonin-containing pellets for 2 weeks. AO, atretic oocyte; PNS, peri-nucleous stage; TYS, tertiary yolk stage. Scale bar = 200 μm .

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary data 1