Exposure Levels and Contributing Factors of Various Arsenic Species and Their Health Effects on Korean Adults

Seul-Gi Lee

Chung-Ang University

Ingu Kang

Chung-Ang University

Mi-Na Seo

Chung-Ang University

Jung-Eum Lee

Chung-Ang University

Sang-Yong Eom

Chungbuk National University

Myung-Sil Hwang

National Institute of Food and Drug Safety Evaluation

Kyung Su Park

Korea Institute of Science and Technology

Byung-Sun Choi

Chung-Ang University

Ho-Jang Kwon

Dankook University

Young-Seoub Hong

Dong-A University

Heon Kim

Chungbuk National University

Jung-Duck Park (✉ jdpark@cau.ac.kr)

Chung-Ang University College of Medicine <https://orcid.org/0000-0003-0667-4674>

Research Article

Keywords: Arsenic species, Urine, Seafood, Rice, Human health

Posted Date: September 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-895767/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Archives of Environmental Contamination and Toxicology on February 7th, 2022. See the published version at <https://doi.org/10.1007/s00244-022-00913-y>.

Abstract

Arsenic is a human carcinogen. Data on urinary arsenic species analyses of Koreans is limited. This study evaluated the arsenic exposure level, contributing factors, and health effects in Korean adults. Dietary intake information and urine samples were obtained from 2,044 participants. Arsenic exposure was assessed based on urinary concentrations of arsenic species, such as inorganic arsenic, As(III) and As(V), monomethylarsonic acid (MMA), dimethylarsinic acid (DMA), and arsenobetaine (AsB), using high-performance liquid chromatography with inductively coupled plasma mass spectrometry, followed by determination of biomarkers, malondialdehyde and c-peptide. The geometric mean concentrations were 30.9 µg/L for the sum of inorganic arsenic and their metabolites, and 84.7 µg/L for the total sum of arsenic measured. Urinary concentrations of arsenic species were influenced by age, inhabitant area (inland or coastal), and seafood intake, which was positively correlated with inorganic arsenic, DMA, and AsB. Rice intake was positively correlated with inorganic arsenic and its metabolites but not with AsB. Additionally, malondialdehyde and c-peptide levels were significantly associated with urinary concentrations of various arsenic species. Seafood and rice are major sources of organic/inorganic arsenic exposure in Korean adults; however, it is necessary to evaluate whether their overconsumption could have a potentially detrimental effect on human health.

Introduction

Arsenic (As), which is ubiquitously distributed in the environment, is one of the major environmental pollutants. Arsenic has been originated naturally from soil, rock, and volcanic eruptions and from anthropogenic sources, such as mining, industries including copper smelter and wood preservative facilities, and agricultural sources. However, the majority of human arsenic exposure is primarily from geogenic arsenic contamination of food and water (ATSDR, 2007). There are several forms of arsenic, and the toxicities of arsenic are quite different based on their chemical form and valence; inorganic arsenic is generally more toxic than organic arsenic, and trivalent arsenic is more toxic than pentavalent arsenic. Inorganic arsenic, namely, arsenate and arsenite, is metabolized to monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA) through reduction and methylation processes in the body, which are more toxic than organoarsenics, such as arsenobetaine (AsB), arsenocholine (AsC), and arsenosugars (AsS). MMA(III), which is a methylated intermediate metabolite of inorganic arsenic, is the most toxic arsenic species (Petrick et al., 2001; Stýblo et al., 2002).

Inorganic arsenic is known to cause non-carcinogenic diseases, including skin pigmentation and keratosis, diabetes, cardiovascular diseases, and peripheral neuropathy, as well as various cancers of the skin, lungs, liver, and bladder (ATSDR, 2007; Schuhmacher-Wolz et al., 2009). Human exposure to arsenic can be estimated from arsenic levels in blood, hair, nails, and urine. Arsenic is metabolized relatively quickly in the body and is excreted mainly in the urine. Blood arsenic level rapidly decreases within several hours after exposure, which makes it a relatively poor exposure marker. Therefore, urinary arsenic concentration is the most valuable biomarker that reflects arsenic exposure in the past several days (Buchet et al., 1981; Link et al., 2007). Arsenic in the hair and nails is a biomarker of chronic and relatively older arsenic exposure (Hindmarsh, 2002; Middleton et al., 2016). Additionally, several forms of arsenic, such as organoarsenic as

well as inorganic arsenic and their metabolites, are excreted in the urine. Therefore, it is necessary to determine the concentrations of urinary arsenic species for the risk assessment of human arsenic exposure.

Several studies reported that seafood consumption increased DMA excretion in urine and suggested that organoarsenic in seafood could be metabolized to DMA after consumption (Ma and Lee, 1998; Choi et al., 2010; Molin et al., 2014). It is questionable whether overconsumption of seafood could be hazardous to human health. The general population, who are not occupationally exposed to arsenic, is mainly exposed to arsenic through ingestion of arsenic contaminated water, grains harvested from arsenic contaminated fields, and seafood. Associations between arsenic intake from food, especially seafood and rice, and urinary arsenic concentration in the general population have been reported (Navas-Acien et al., 2011; Wei et al., 2014; Bae et al., 2017; Signes-Pastor et al., 2017). Seafood is one of the favorite foods, and rice is a staple food in Korea. However, limited data are available on urinary arsenic species analyses in the Korean population (Choi et al., 2010; Park et al., 2016; Bae et al., 2017). Additionally, it is not fully understood whether environmental arsenic exposure could detrimentally influence the health of the general Korean population.

Thus, we performed speciation analysis of urinary arsenic, analyzed the relations between the consumption of food groups and the urinary arsenic species, and measured possible health effectors, namely, malondialdehyde (MDA) and c-peptide, for a risk assessment of arsenic exposure in the Korean adults.

Materials And Methods

Study population

This cross-sectional study included a total of 2,044 study subjects, 888 males and 1,156 females, who were 19 years old or older. They had not been occupationally exposed to arsenic. Study subjects were sampled using the multistage and probability sampling method and were stratified by sex and age from 102 sampling sites during 2010–2011 in Korea as described previously (Eom et al., 2014; Lim et al., 2015). In briefly, the 102 sampling sites were distributed throughout the Korea, and included 15 metropolitans and provinces, excluding Jeju province. We selected 34 cities and counties from 15 metropolitans and provinces, followed by sampling of 102 towns and townships from these 34 cities and counties. The number of study subject from each site was allocated in proportion to the square root of the population size of the district. Because we decided that having stable sample was very important in this small-sample study. Written informed consent was obtained from all the study subjects. Analyses of various arsenic species in the urine were performed in this study. The study protocol of 2010–2012 was approved by the Chung-Ang University Ethical Committee for Medical Research and Other Studies Involving Human Subjects, and the study protocol for additional analyses of arsenic species was approved by the Institutional Review Board of Chung-Ang University.

Personal interview and urine sampling

We conducted personal interviews with study subjects individually to obtain information about demographic characteristics, such as sex, age, smoking, alcohol consumption, education level, monthly income, type of

drinking water, pesticides used, residential area size, and inhabitant area (coastal or inland). Coastal or inland area was categorized whether each study site included a seashore or not. Additionally, we asked about seafood intake within 72 hours before the start of this study. All urine excreted by the study subjects was collected as an aggregated sample starting from post-dinner until the next morning's interview, also included the first-void sample of the next morning. This corresponds to urine collected for approximately 15–18 h and averaged about 1.1 L. Urine samples were refrigerated during the sample collection time, frozen after dispensing, and subsequently stored at -80°C in the laboratory without thawing before analysis after sample collection.

Estimation of daily food consumption

Previously, we estimated daily food consumption during the last 24 h before the interview (Seo et al., 2016). A 24-h recall method has a limitation which could not reflect long-term, usual intakes and may be underestimated (Tucker, 2007), although this method could provide more detailed information on dietary intake in a population study. A diet study was carefully conducted with a prepared questionnaire by well-trained personnel to minimize recall bias. In this study, we included 138 specific food items (16 food groups) that are frequently and largely consumed in Korea.

Analyses of arsenic species

Speciation analyses of various arsenic species in urine were performed using high-performance liquid chromatography (HPLC, PerkinElmer Series 200, Shelton, CT) coupled to inductively coupled plasma mass spectrometry (ICP-MS, PerkinElmer NEXION 300S, Concord, Ontario, Canada). We determined five different arsenic species, namely, arsenite [As(III)], arsenate [As(V)], MMA, DMA, and AsB, using CAPCELL PAK, 4.6 $\times$ 250 μm , 5 μm C18 columns (Shiseido, Japan) with a mobile phase of 10 mM sodium 1-butanedisulfonate, 4 mM malonic acid, and 4 mM tetramethylammonium hydroxide pentahydrate (pH 2.5). The concentrations of urinary arsenic are expressed as inorganic arsenic [InAs, As(III)+As(V)], MMA, DMA, AsB, the sum of inorganic arsenic and their metabolites (TmetAs, InAs+MMA+DMA), and the total sum of arsenic measured (TsumAs, InAs +MMA+DMA+AsB). The analytical method for various arsenic species was validated for linearity using each corresponding standard solution, such as As(III) (Inorganic Ventures, Christiansburg, VA), As(V) (Inorganic Ventures), MMA (TCLC, Japan), DMA (TCLC), and AsB (TCLC). The calibration curves had a linearity of $r^2 > 0.99$. The standard reference material from the National Institute of Standards Technology (NIST), NIST 2669 I&II (USA), was used to validate the accuracy and precision of the experimental method. The recovery of NIST 2669 levels I and II were 99.7% and 100.7% for As(III), 102.1% and 98.6% for As(V), 99.5% and 103.2% for MMA, 100.8% and 99.7% for DMA, and 101.0% and 99.4% for AsB, respectively, and the coefficient of variations were 4.5% and 4.0% for As(III), 6.4% and 5.3% for As(V), 11.3% and 7.2% for MMA, 6.4% and 1.6% for DMA, and 7.8% and 3.5% for AsB, respectively. The limits of detection were 0.026, 0.049, 0.012, 0.030, and 0.059 $\mu\text{g/L}$ for As(III), As(V), MMA, DMA, and for AsB, respectively. The concentrations were below the detection limits in 242, 206, 70, 0, and 11 samples for As(III), As(V), MMA, DMA, and AsB, respectively. More than 80% of the samples had detectable values in the various arsenic species. The levels of arsenic below the detection limits were assigned to values of detection limits divided by a square root of two (Hornung and Reed, 1990).

Measurements of malondialdehyde and c-peptide

Urinary MDA and c-peptide levels were measured as possible indicators of oxidative stress and endogenous endocrine response, respectively. The urinary concentration of MDA was determined using HPLC with a fluorescence detector (RF-10AxL, Shimadzu, Kyoto, Japan) as described previously by Agarwal and Chase (2002). The urinary concentration of c-peptide was measured using electrochemiluminescence immunoassay (Cobas 8000 e602, Roche, Germany).

Statistical analyses

Statistical analyses were performed with SAS version 9.2 (SAS Institute Inc., Cary, NC, USA). The urinary arsenic level was presented as the arithmetic mean, geometric mean, median, and the value at the 95th percentile. The urinary arsenic concentrations were distributed log-normally rather than as normal distribution, which were log-transformed for statistical analyses. The comparisons of means were analyzed using a two-tailed Student's *t*-test or analysis of variance following multiple comparison test using Duncan's method. Contributing factors to the urinary arsenic levels were determined using multiple regression analyses. The relationship between urinary arsenic concentrations and consumption of each food group was analyzed using Spearman's rank correlation coefficients after adjusting for potential confounding variables, such as sex, age, inhabitant area, and body weight. Statistical significance was set at $p < 0.05$ or $p < 0.01$.

Results

The arithmetic and geometric means and median of various urinary arsenic species in this study subjects are presented in Table 1, and individual levels of various arsenic species are plotted in Fig. 1. The urinary concentrations (geometric means) of different arsenic species were 3.0 $\mu\text{g/L}$ of InAs, As(III) + As(V), 1.4 $\mu\text{g/L}$ of MMA, 24.7 $\mu\text{g/L}$ of DMA, 41.4 $\mu\text{g/L}$ of AsB, 30.9 $\mu\text{g/L}$ of TmetAs, and 84.7 $\mu\text{g/L}$ of TsumAs in Korean adults. The values of the 95th percentiles were 19.1, 5.6, 83.1, and 338.0 $\mu\text{g/L}$ for InAs, MMA, DMA, and AsB, respectively. No statistically significant difference in geometric means was observed between males and females, except AsB. The proportion of AsB was 56.7% of TsumAs, and the sum of InAs and their metabolites (TmetAs) accounted for 43.3% of TsumAs. DMA was the major species accounting for 80.8% of TmetAs, followed by InAs and MMA (Fig. 2).

Table 1
Mean concentrations of various arsenic (As) species in urine (µg/L)

		InAs	MMA	DMA	AsB	TmetAs	TsumAs
Male (n = 888)	AM ± SD	5.4 ± 7.6	2.4 ± 2.1**	32.5 ± 27.6	100.8 ± 181.9	40.3 ± 33.6	141.0 ± 194.3
	GM (GSD)	3.0 (3.2)	1.5 (3.6)	24.2 (2.2)	45.6 (3.9)**	30.4 (2.2)	87.7 (2.6)
	Median	3.0	1.9	25.2	48.1	32.1	87.9
	P95	18.2	5.8	81.2	363.4	99.6	434.3
Female (n = 1,156)	AM ± SD	6.5 ± 17.0	2.1 ± 2.1	35.3 ± 46.4	96.1 ± 226.7	43.9 ± 61.0	140.1 ± 240.6
	GM (GSD)	3.0 (3.5)	1.4 (3.4)	25.1 (2.3)	38.4 (4.5)	31.3 (2.2)	82.4 (2.7)
	Median	2.9	1.6	25.3	41.3	31.8	79.9
	P95	20.4	5.5	86.0	314.6	104.1	385.4
Total (n = 2,044)	AM ± SD	6.0 ± 13.8	2.2 ± 2.1	34.1 ± 39.4	98.2 ± 208.4	42.4 ± 51.0	140.5 ± 221.6
	GM (GSD)	3.0 (3.3)	1.4 (3.5)	24.7 (2.3)	41.4 (4.2)	30.9 (2.2)	84.7 (2.6)
	Median	3.0	1.7	25.3	43.8	31.9	82.9
	P95	19.1	5.6	83.1	338.0	100.9	407.1
InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, AM: arithmetic mean, SD: standard deviation, GM: geometric mean, GSD: geometric standard deviation, P95: value at the 95th percentile. Student's t-test was performed to compare means (arithmetic and geometric) of various arsenic species between males and females. ** p < 0.01 from the Student's t-test							

The DMA and AsB levels increased with age and peaked in the forties; however, no significant difference was observed for urinary InAs and MMA levels by age group. Urinary arsenic levels did not differ generally in regard to smoking and alcohol consumption as well as education level, income, drinking water, pesticides used, and residential area size. However, the concentrations (geometric mean) of most arsenic species, except MMA, were higher in coastal area inhabitants than in inland inhabitants and were higher in those who had consumed seafood during the past 3 days than in those who had not (Table 2). In multiple regression analysis, factors contributing to the concentrations of urinary arsenic species, except MMA, were generally determined based on age, living in coastal areas, and seafood consumption (Table 3).

Table 2

Mean urinary concentrations of various arsenic (As) species based on demographic characteristics (µg/L)

	Class	N	InAs	MMA	DMA	AsB	TmetAs	TsumAs
Age (years)	19–29	355	2.8 (3.3)	1.5 (3.1)	20.6 (2.4) ^a	29.1 (4.8) ^a	26.6 (2.3) ^a	66.4 (2.7) ^a
	30–39	360	3.1 (3.1)	1.3 (3.8)	23.7 (2.3) ^b	37.8 (4.0) ^b	30.1 (2.2) ^b	78.7 (2.6) ^b
	40–49	450	3.2 (3.1)	1.6 (3.3)	26.8 (2.2) ^c	47.1 (4.0) ^c	33.2 (2.1) ^b	92.9 (2.6) ^c
	50–59	475	2.8 (3.3)	1.5 (3.3)	26.2 (2.1) ^{bc}	46.7 (4.0) ^c	32.3 (2.1) ^b	92.2 (2.6) ^c
	≥ 60	404	3.0 (3.9)	1.3 (4.0)	25.6 (2.3) ^{bc}	45.8 (4.1) ^{bc}	31.7 (2.3) ^b	91.4 (2.6) ^c
	F-value		1.24	1.63	6.75 ^{**}	8.05 ^{**}	4.88 ^{**}	8.78 ^{**}
Smoking	Non-smoker	1302	3.0 (3.4)	1.4 (3.4)	25.4 (2.3)	39.8 (4.4)	31.7 (2.2)	84.4 (2.6)
	Smoker	740	2.8 (3.3)	1.4 (3.7)	23.6 (2.3)	44.3 (3.9)	29.5 (2.2)	85.1 (2.7)
	t-value		1.16	-0.03	1.93	1.62	1.96 [*]	-0.18
Alcohol drinking	Non-drinker	496	2.9 (3.6)	1.5 (3.2)	25.3 (2.3)	37.9 (4.2)	31.7 (2.2)	81.7 (2.7)
	Drinker	1538	3.0 (3.2)	1.4 (3.6)	24.6 (2.2)	42.6 (4.2)	30.7 (2.2)	85.8 (2.6)
	t-value		-0.58	1.42	0.73	-1.57	0.74	-0.97
Education	<High school	615	2.8 (3.6)	1.5 (3.5)	26.3 (2.2) ^a	42.4 (4.4)	32.5 (2.2) ^a	89.2 (2.6)
	=High school	687	3.1 (3.2)	1.4 (3.2)	25.0 (2.2) ^{ab}	44.0 (3.8)	31.3 (2.1) ^{ab}	86.6 (2.6)
	>High school	742	2.9 (3.2)	1.4 (3.7)	23.2 (2.3) ^b	38.2 (4.4)	29.3 (2.2) ^b	79.5 (2.7)
	F-value		1.12	0.66	4.20 [*]	1.88	3.04	2.67

InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, Data are presented as geometric mean and geometric standard deviation, [†]: Coastal or inland area was categorized to specify whether each study site included a seashore. [‡]: seafood intake during the 3 days before this study. ^{a,b,c} Duncan grouping, * p < 0.05, ** p < 0.01

	Class	N	InAs	MMA	DMA	AsB	TmetAs	TsumAs
Income (US\$/month)	< 1,599	827	2.8 (3.5)	1.4 (3.7)	24.9 (2.3)	40.8 (4.5)	30.9 (2.2)	85.1 (2.7)
	1,600-3,199	794	3.0 (3.2)	1.5 (3.3)	24.9 (2.2)	42.0 (4.2)	31.2 (2.1)	85.4 (2.6)
	≥ 3,200	305	3.2 (3.5)	1.4 (3.3)	25.6 (2.4)	43.2 (3.9)	32.3 (2.3)	87.6 (2.6)
	F-value		1.55	0.19	0.18	0.20	0.38	0.11
Drinking water	Self-tap water	209	3.0 (3.0)	1.4 (4.2)	24.6 (2.5)	41.6 (4.1)	30.8 (2.3)	82.9 (2.7)
	Tap water	672	3.0 (3.3)	1.4 (3.6)	25.0 (2.2)	42.6 (4.3)	31.2 (2.2)	86.3 (2.6)
	Bottled water	135	2.6 (3.7)	1.2 (4.8)	24.7 (2.2)	37.6 (5.4)	30.4 (2.2)	82.7 (2.7)
	Filtered water	795	3.0 (3.4)	1.5 (3.1)	24.3 (2.3)	41.2 (4.1)	30.5 (2.2)	84.3 (2.7)
	Others	214	2.6 (3.4)	1.5 (3.3)	25.3 (2.2)	40.6 (4.0)	31.3 (2.1)	84.2 (2.4)
	F-value		1.17	0.79	0.17	0.23	0.10	0.11
Pesticide	No	1720	3.0 (3.4)	1.4 (3.4)	24.8 (2.2)	41.2 (4.2)	31.0 (2.2)	84.9 (2.6)
	Yes	323	3.0 (3.2)	1.5 (3.9)	24.4 (2.3)	42.6 (4.0)	30.4 (2.3)	84.1 (2.7)
	t-value		0.02	-0.59	0.35	-0.40	0.46	0.15
Residence area size	Metropolitan	815	2.9 (3.8)	1.4 (3.5)	25.8 (2.2)	40.3 (4.5)	32.3 (2.2)	87.0 (2.6)
	Urban	817	3.1 (3.1)	1.5 (3.2)	23.9 (2.3)	42.2(3.7)	30.1 (2.2)	82.8 (2.5)
	Rural	412	2.8 (2.9)	1.3 (4.0)	24.2 (2.2)	41.8 (4.6)	29.8(2.2)	83.9 (2.8)
	F-value		0.85	1.68	2.02	0.23	2.23	0.57
Inhabitant area [†]	Inland	1541	2.8 (3.3)	1.5 (3.3)	23.9 (2.2)	35.2 (4.1)	29.9 (2.1)	76.1 (2.5)

InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, Data are presented as geometric mean and geometric standard deviation, [†]: Coastal or inland area was categorized to specify whether each study site included a seashore. [‡]: seafood intake during the 3 days before this study. ^{a,b,c} Duncan grouping, * p < 0.05, ** p < 0.01

	Class	N	InAs	MMA	DMA	AsB	TmetAs	TsumAs
	Coastal	503	3.6(3.3)	1.3 (3.9)	27.5 (2.4)	67.5 (4.0)	34.3 (2.3)	117.3 (2.9)
	t-vale		-4.41**	1.69	-3.25**	-9.00**	-3.27**	-8.25**
Seafood intake [‡]	No	734	2.6 (3.4)	1.3 (3.8)	21.0 (2.3)	29.2 (4.5)	26.6 (2.2)	65.6 (2.6)
	Yes	1295	3.2 (3.3)	1.5 (3.3)	27.2 (2.2)	50.7 (3.9)	33.7 (2.2)	98.1 (2.6)
	t-value		3.51**	1.40	6.98**	8.26**	6.60**	9.17**
InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, Data are presented as geometric mean and geometric standard deviation, [†] : Coastal or inland area was categorized to specify whether each study site included a seashore. [‡] : seafood intake during the 3 days before this study. ^{a,b,c} Duncan grouping, * p < 0.05, ** p < 0.01								

Table 3

Contributing factors with significant coefficients of various urinary arsenic (As) species were determined using multiple regression analysis

Variable	InAs	MMA	DMA	AsB	TmetAs	TsumAs
Age	-	-	0.003**	0.007**	0.002**	0.005**
Residence area size	-	-	-	-	-0.025*	-
Education level	-	-0.048*	-	-	-	-
Inhabitant area	0.127**	-	0.050**	0.259**	0.053**	0.170**
Seafood intake	0.073**	-	0.109**	0.217**	0.099**	0.160**
InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, * $p < 0.05$, ** $p < 0.01$						

The daily mean food intake of study subject was estimated previously at $1,373.6 \pm 652.4$ g and 22.0 ± 10.5 g/kg body weight by Seo et al. (2016). Generally, no significant correlation coefficients were observed between total food consumption and urinary arsenic species, such as InAs, DMA, AsB, TmetAs, and TsumAs, concentrations (Table 4). However, the consumption of specific food groups significantly correlated with urinary arsenic levels. Namely, the consumption of fish&shellfish and seaweeds positively correlated with the concentrations of InAs ($r = 0.108$, $p < 0.01$ for fish&shellfish; $r = 0.234$, $p < 0.01$ for seaweeds) and DMA ($r = 0.167$, $p < 0.01$ for fish&shellfish; $r = 0.178$, $p < 0.01$ for seaweeds), but not with the MMA concentration (Table 4). The concentration of AsB showed a significantly high association with the consumption of fish&shellfish ($r = 0.234$, $p < 0.01$) but was not related to seaweeds consumption (Table 4). Furthermore, dose-dependent increases of InAs, DMA, and AsB levels were observed according to the amount of fish&shellfish consumed. However, dose-dependent increases based on the level of seaweeds consumption were observed for InAs and DMA only, and AsB levels did not significantly differ statistically based on the amount of seaweeds consumed (Table 5). Daily intake of grain, which included 17 kinds of food items including rice, wheat, noodles, and so on, as described in a previous study (Seo et al., 2016), was positively correlated with the concentrations of MMA and DMA but not with those of InAs and AsB. Specifically, the amount of rice intake was statistically significantly correlated with the concentrations of InAs, MMA, DMA, and TmetAs, but not of AsB (Tables 4 and 5). Flavorings also correlated with various urinary arsenic species, except for MMA (Table 4). Moreover, the urinary concentrations of MDA and c-peptide significantly increased according to the levels of all arsenic species as well as TmetAs and TsumAs in a dose-dependent manner (Table 6).

Table 4

Daily food intakes and Spearman's correlation coefficients, adjusted for sex, age, inhabitant area, and body weight, in different concentrations of urinary arsenic (As) species and daily consumption from each food group

Group	Food intake (g/day,%) [‡]	InAs	MMA	DMA	AsB	TmetAs	TsumAs
Grains	390.6 ± 351.7 (28.4%)	0.035	0.078**	0.063**	-0.016	0.058**	0.005
Potatoes	35.9 ± 86.6 (2.6%)	-0.008	0.009	-0.041	-0.075**	-0.035	-0.067**
Sugars	10.5 ± 9.9 (0.8%)	0.030*	-0.014	0.020	-0.004	0.021	0.008
Pulse	35.0 ± 64.2 (2.5%)	0.024	0.031	0.005	-0.032	0.010	-0.014
Seeds	0.8 ± 3.2 (0.1%)	0.020	-0.012	0.014	0.001	0.013	0.001
Vegetables	288.7 ± 201.2 (21.0%)	0.011	-0.035	0.026	0.035	0.019	0.031
Mushrooms	3.5 ± 15.4 (0.3%)	0.021	0.016	0.006	-0.001	0.008	0.000
Fruits	138.8 ± 252.9 (10.1%)	-0.032	-0.029	-0.017	-0.014	-0.023	-0.021
Meats	64.1 ± 98.0 (4.7%)	-0.006	-0.048*	-0.037	-0.011	-0.034	-0.022
Eggs	25.3 ± 40.7 (1.8%)	0.011	-0.026	-0.013	-0.020	-0.012	-0.021
Fish/Shellfish	59.0 ± 82.0 (4.3%)	0.108**	0.032	0.167**	0.234**	0.160**	0.236**
Seaweeds	1.8 ± 9.0 (0.1%)	0.234**	0.039	0.178**	0.037	0.197**	0.108**
Milks	76.9 ± 148.5 (5.6%)	0.016	-0.007	0.003	-0.004	0.006	-0.001
Oils	11.6 ± 9.4 (0.8%)	0.065**	-0.025	0.027	-0.024	0.034	-0.003
Beverage	197.1 ± 346.3 (14.4%)	-0.023	-0.114**	-0.045*	0.023	-0.050*	0.002
Flavorings	34.1 ± 33.9 (2.5%)	0.045*	-0.022	0.050*	0.067**	0.046*	0.063**
Total	1373.6 ± 652.4 (100%)	-0.003	-0.044*	0.018	0.021	0.009	0.016
	(22.0 ± 10.5) ^a						
InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB, [‡] : Seo et al. (2016), ^a : Values per kg body weight basis (g/kg body weight/day); data are presented as mean and standard deviation, * p < 0.05, ** p < 0.01							

Table 5

Mean concentrations of various urinary arsenic (As) species (µ/L) according to the consumption amounts of fish/shellfish, seaweeds, and rice

	Group	N	InAs	MMA	DMA	AsB	TmetAs	TsumAs
Fish/shellfish consumption (g)	below 50	1263	2.7 (3.4) ^a	1.4 (3.6)	22.5 (2.2) ^a	32.7 (4.1) ^a	28.3 (2.2) ^a	71.1 (2.5) ^a
	50–99	358	3.3 (2.9) ^b	1.5 (3.0)	26.4 (2.2) ^b	50.4(4.1) ^b	33.0 (2.1) ^b	98.3 (2.5) ^b
	100–199	310	3.4 (3.4) ^b	1.4 (3.5)	28.5 (2.3) ^b	60.5(3.9) ^b	35.2 (2.2) ^b	111.0 (2.7) ^b
	≥ 200	131	3.8 (3.2) ^b	1.7 (2.9)	35.6 (2.3) ^c	91.9 (3.7) ^c	43.2 (2.2) ^c	153.8 (2.8) ^c
	F-value		6.55 ^{**}	1.54	18.68 ^{**}	35.87 ^{**}	17.26 ^{**}	43.25 ^{**}
Seaweeds consumption (g)	below 1.0	1325	2.5 (3.2) ^a	1.4 (3.4)	22.4 (2.2) ^a	39.4 (4.2)	27.9 (2.2) ^a	78.2 (2.6) ^a
	1.0–1.9	174	3.2 (3.1) ^b	1.5 (3.2)	25.3 (2.2) ^{ab}	45.2 (4.7)	31.7 (2.1) ^a	92.7 (2.8) ^b
	2.0–2.9	163	3.6 (3.3) ^b	1.5 (3.5)	28.8 (2.1) ^{bc}	43.8 (3.9)	36.2 (2.00) ^b	93.3 (2.4) ^b
	≥ 3.0	400	4.8 (3.3) ^c	1.5 (3.9)	31.5 (2.2) ^c	45.4 (4.1)	40.2 (2.2) ^b	100.8 (2.6) ^b
	F-value		34.01 ^{**}	0.28	20.70 ^{**}	1.36	25.86 ^{**}	8.29 ^{**}
Rice consumption (g)	below 100	437	2.5 (3.5) ^a	1.3 (3.2) ^a	22.3 (2.3) ^a	37.0 (4.0)	28.0 (2.2) ^a	75.7 (2.6) ^a
	100–199	809	3.0 (3.3) ^{ab}	1.4 (3.4) ^{ab}	23.8 (2.2) ^{ab}	40.4 (4.1)	29.8 (2.2) ^a	82.3 (2.6) ^{ab}
	200–299	641	3.2 (3.3) ^b	1.6 (3.5) ^{ab}	26.9 (2.2) ^{bc}	44.3(4.4)	33.6 (2.1) ^b	92.0 (2.6) ^b
	≥ 300	143	3.2 (3.2) ^b	1.6 (4.4) ^b	28.5 (2.3) ^c	48.0 (4.0)	35.2 (2.2) ^b	94.3 (2.9) ^b
	F-value		3.46 [*]	2.65 [*]	6.91 ^{**}	1.99	6.74 ^{**}	4.38 ^{**}

InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs + MMA + DMA, TsumAs (total sum of arsenic measured): InAs + MMA + DMA + AsB; data are presented as geometric mean and geometric standard deviation, ^{a,b,c} Duncan grouping, ^{*} p < 0.05, ^{**} p < 0.01

Table 6

Mean urinary concentrations of MDA ($\mu\text{mol/g}$ creatinine) and c-peptide ($\mu\text{g/day}$) according to the urinary arsenic (As) levels in study subjects

	As Level	MDA		c-peptide	
		Mean $\pm$ SD	F-value	Mean $\pm$ SD	F-value
InAs	G1	1.389 $\pm$ 1.308 ^a	8.54 ^{**}	34.048 $\pm$ 29.756 ^a	5.43 ^{**}
	G2	1.676 $\pm$ 1.491 ^b		40.629 $\pm$ 37.066 ^b	
	G3	1.753 $\pm$ 1.476 ^b		40.990 $\pm$ 32.899 ^b	
	G4	1.826 $\pm$ 1.628 ^b		41.305 $\pm$ 34.604 ^b	
MMA	G1	1.330 $\pm$ 1.282 ^a	33.04 ^{**}	34.137 $\pm$ 30.989 ^a	10.69 ^{**}
	G2	1.448 $\pm$ 1.141 ^a		36.354 $\pm$ 33.159 ^a	
	G3	1.697 $\pm$ 1.495 ^b		41.790 $\pm$ 34.088 ^b	
	G4	2.168 $\pm$ 1.812 ^c		44.712 $\pm$ 35.821 ^b	
DMA	G1	1.267 $\pm$ 1.182 ^a	28.13 ^{**}	33.406 $\pm$ 29.838 ^a	10.35 ^{**}
	G2	1.509 $\pm$ 1.132 ^b		37.426 $\pm$ 32.140 ^a	
	G3	1.815 $\pm$ 1.513 ^c		44.082 $\pm$ 38.254 ^b	
	G4	2.049 $\pm$ 1.883 ^d		42.046 $\pm$ 33.450 ^b	
AsB	G1	1.370 $\pm$ 1.318 ^a	13.59 ^{**}	33.509 $\pm$ 30.228 ^a	13.68 ^{**}
	G2	1.568 $\pm$ 1.443 ^b		36.374 $\pm$ 31.655 ^a	
	G3	1.790 $\pm$ 1.551 ^c		41.070 $\pm$ 35.443 ^b	
	G4	1.914 $\pm$ 1.573 ^c		46.004 $\pm$ 36.231 ^c	
TmetAs	G1	1.271 $\pm$ 1.197 ^a	27.87 ^{**}	33.416 $\pm$ 29.749 ^a	10.98 ^{**}
	G2	1.522 $\pm$ 1.212 ^b		37.019 $\pm$ 31.129 ^a	
	G3	1.783 $\pm$ 1.394 ^c		44.131 $\pm$ 39.042 ^b	

SD: standard deviation, InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs: InAs + MMA + DMA, TsumAs: InAs + MMA + DMA + AsB. The levels of various As species were divided into quartiles, such as G1, G2, G3 and G4; Urine As level criteria are as follows: P25 = 1.539, P50 = 2.949, P75 = 6.205 $\mu\text{g/L}$ for InAs; P25 = 1.033, P50 = 1.728, P75 = 2.773 $\mu\text{g/L}$ for MMA; P25 = 15.199, P50 = 25.286, P75 = 42.207 $\mu\text{g/L}$ for DMA; P25 = 19.867, P50 = 43.820, P75 = 99.757 $\mu\text{g/L}$ for AsB; P25 = 18.766, P50 = 31.875, P75 = 51.141 $\mu\text{g/L}$ for TmetAs; P25 = 45.460, P50 = 82.854, P75 = 157.769 $\mu\text{g/L}$ for TsumAs, Data are presented as mean and standard deviation, ^{a,b,c,d} Duncan grouping, ^{**} $p < 0.01$

	G4	2.065 ± 1.916 ^d		42.396 ± 33.514 ^b	
TsumAs	G1	1.249 ± 1.119 ^a	24.22 ^{**}	32.132 ± 29.243 ^a	15.39 ^{**}
	G2	1.638 ± 1.563 ^b		37.195 ± 33.626 ^b	
	G3	1.732 ± 1.276 ^b		42.283 ± 34.185 ^c	
	G4	2.022 ± 1.800 ^c		45.352 ± 36.316 ^c	
SD: standard deviation, InAs: inorganic As [As(III) + As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs: InAs + MMA + DMA, TsumAs: InAs + MMA + DMA + AsB. The levels of various As species were divided into quartiles, such as G1, G2, G3 and G4; Urine As level criteria are as follows: P25 = 1.539, P50 = 2.949, P75 = 6.205 µ/L for InAs; P25 = 1.033, P50 = 1.728, P75 = 2.773 µ/L for MMA; P25 = 15.199, P50 = 25.286, P75 = 42.207 µ/L for DMA; P25 = 19.867, P50 = 43.820, P75 = 99.757 µ/L for AsB; P25 = 18.766, P50 = 31.875, P75 = 51.141 µ/L for TmetAs; P25 = 45.460, P50 = 82.854, P75 = 157.769 µ/L for TsumAs, Data are presented as mean and standard deviation, ^{a,b,c,d} Duncan grouping, ^{**} p < 0.01					

Discussion

In our study, the geometric mean concentrations of various arsenic species in urine were 3.0, 1.4, 24.7, and 41.4 µ/L of inorganic arsenic, MMA, DMA, and AsB, respectively; therefore, TmetAs and TsumAs were 30.9 and 84.7 µ/L, respectively. The urinary arsenic level in the general adult population of Korea was considerably higher than that of the United States (7.3 µ/L < no seafood > and 24.5 µ/L < with seafood > of the median total arsenic, Navas-Acien et al., 2011; 4.76 µ/L of the median TmetAs and 5.79 µ/L of the median AsB, Gilbert-Diamond et al., 2013), France (3.75 µ/L of of the geometric mean TmetAs and 13.42 µ/L of the geometric mean total arsenic, Saoudi et al., 2012), Germany (4.9 µ/L of the arithmetic mean TmetAs and 10.8 µ/L of the arithmetic mean TsumAs, Heitland and Köster, 2008), and United Kingdom (3.6 µ/L of the arithmetic mean TmetAs and 33.9 µ/L of the arithmetic mean TsumAs, Morton and Leese, 2011), but was similar to or lower than that of Taiwan (86.08 µ/L of the arithmetic mean TmetAs and 267.05 µ/L of the arithmetic mean total arsenic in a previously contaminated area, Hsueh et al., 1998; 57.08 µ/L of the arithmetic mean TmetAs in a previously contaminated area, Huang et al., 2009; 20.94 µ/g creatinine of the arithmetic mean TmetAs, Huang et al., 2012), Japan (141.3 µ/L of the median total arsenic, Hata et al., 2007; 132.2 µ/L of the median total arsenic, Suzuki et al., 2009), and China (28.3 µ/L of the arithmetic mean TmetAs of the control, Wen et al., 2011; 56 µ/L of the arithmetic mean TmetAs, Cui et al., 2013). In addition, approximately 44.0% of this study subjects (900/2,044) had urinary TmetAs over 35 µ/L of the biological exposure index (BEI, ACGIH, 2014); approximately 26.3% of study subjects (537/2,044) had urinary TmetAs exceeding 50 µ/L of the biological limit value (BLV, DFG, 2017).

Diet, especially seafood, was the main source of arsenic exposure in the general population who lived in arsenic non-polluted areas. Previously, several studies reported that seafood intake was associated with urinary arsenic concentrations (Navas-Acien et al., 2011; Bae et al., 2017; Signes-Pastor et al., 2017). In our study, urinary concentrations of various arsenic species, such as InAs, DMA, and AsB, were significantly higher in people who ate seafood, including fish/shellfish and seaweeds, during the 3 days before the

personal interview than in those who did not. Additionally, the urinary concentrations of arsenic were higher in inhabitants of the coastal area than in those living inland, highlighting that seafood may be a major source of arsenic exposure in the general population (Luvonga et al., 2020). In our study, the distributions of urinary arsenic profiles were quite different from those in Western countries, including the United States. The relative proportions of AsB and TmetAs in relation to TsumAs were approximately 57% and 43%, respectively, in our study population. In contrast, the relative proportions of AsB in relation to TsumAs or the levels of AsB in urine were much lower in the general populations of Western countries, such as the United States, Germany, and France, than in Koreans (Heitland and Köster, 2008; Caldwell et al., 2009; Saoudi et al., 2012). Thus, the Korean general population is exposed to a higher arsenic level than the populations of Western countries, and AsB was a dominant contributor to the total arsenic exposure.

AsB is essentially a non-toxic and rapidly excreted compound, with a relatively high concentration in seafood (Wolle and Conklin, 2018). In our previous study, total urinary arsenic concentrations measured using ICP-MS were associated with the amount of seafood consumption (Bae et al., 2017). Furthermore, the speciation analyses of urinary arsenic indicated a major source of organic arsenic, namely, AsB, which was significantly associated with the consumption of fish/shellfish but not of seaweeds. Moreover, the urinary AsB concentration increased with the amount of fish/shellfish consumption in a dose-dependent pattern; seaweeds intake positively correlated with urinary InAs and DMA concentrations, and dose-dependent increases of InAs and DMA were observed according to the seaweeds consumption. However, seaweeds intake was not statistically associated with urinary concentrations of AsB. These findings indicate that seaweeds might afford inorganic arsenic rather than organic species (from fish/shellfish) in the Korean population. Nonetheless, the urinary concentrations of InAs and DMA positively correlated with fish/shellfish intake and increased according to the amount of fish/shellfish consumption, which might indicate that fish/shellfish is a part of InAs exposure source in Koreans.

However, the contributing mechanism remains to be understood as to whether a little amount of InAs in fish/shellfish, which is much consumed favorite food in Korea (Sirot et al., 2009; Seo et al., 2016), any kind of labile organic arsenic in fish/shellfish (Choi et al., 2010; Luvonga et al., 2020) as well as both and other foods might be source of arsenic exposure.

Inhabitants in arsenic contaminated area could be exposed to arsenic mainly through contaminated drinking water or harvested crops in the contaminated soil, which has previously caused the so-called “black foot disease” in Taiwan, Bangladesh, and elsewhere (Tseng, 1977; Smith et al., 2000; Nordstrom, 2002; Sun et al., 2007). Others could be exposed through inhalation or ingestion in the industry or accidentally (Morton and Mason, 2006; Heitland and Köster, 2008; Wen et al., 2011). However, urinary arsenic profiles show a different pattern according to the exposure source of arsenic. Our data present a high proportion of AsB (56.7%) and a relatively low proportion of TmetAs (43.3%) in TsumAs, but a relatively high proportion of DMA (80.8%) in TmetAs with low proportions and concentrations of InAs and MMA. Despite a relatively low concentration of AsB and high concentration of TmetAs, a relatively low concentration of DMA with a high concentration of InAs and MMA in TmetAs were observed in inhabitants of arsenic-contaminated areas or in workers occupationally exposed to arsenic in industries (Morton and Mason, 2006; Sun et al., 2007; Wen et al., 2011). Differences in urinary arsenic profiles by arsenic exposure levels could be explained based on a previous

study by Huang et al. (2009), which presented changes in urinary arsenic profiles, such as the decreased proportion of InAs (−4.9%) and MMA (−6.8%) and increased DMA (11.7%), after cessation of arsenic ingestion for 15 years in people residing in the arsenic-contaminated area of Taiwan. Therefore, the speciation analyses of arsenic are essential for evaluating the health risk from arsenic exposure, especially in countries where populations mainly consume seafood, such as Korea.

In our previous study, increased urinary excretion of DMA was observed after seafood consumption in volunteers (Choi et al., 2010). Thus, a labile organoarsenic such as arsenosugar and arsenolipid could be metabolized to DMA, which is more toxic than the original form of organoarsenic (Molin et al., 2014; Luvonga et al., 2020). Furthermore, this study showed significantly higher urinary DMA concentrations in subjects who consumed seafood during the 3 days before the personal interview than in those who did not. Seafood is known as a healthy food, especially for growing children, pregnant women, and the elderly, as it is a rich source of essential amino acids, unsaturated fatty acids (omega 3 & 6), vitamins, and minerals (Mozaffarian and Rimm, 2006; Venugopal and Gopakumar, 2017). However, it remains to be elucidated whether overconsumption of seafood may increase exposure to hazardous arsenic species and result in toxicological implications.

Human exposure to the most hazardous metals, such as lead, mercury, and cadmium, are generally influenced by individual lifestyles, such as smoking and alcohol consumption, socioeconomic status, as well as sex and age in the general population (McKelvey et al., 2007; Eom et al., 2018). However, there were no observed prominent differences in urinary arsenic levels based on sex, smoking and alcohol consumption, economic and educational levels, residential area size, and pesticide used in our study. The factors affecting arsenic exposure in humans were quite different from those for other metals. Diet, especially seafood, and residential area (inland or coastal) mainly contributed to determining the human exposure levels to various arsenic species in our study population. Additionally, the amount of rice intake associated with the urinary levels of inorganic arsenic and its metabolites. Our finding is consistent with previous reports where rice consumption contributed to inorganic arsenic exposure (Wei et al., 2014; Signes-Pastor et al., 2017). Rice is a staple food and one of the major sources of inorganic arsenic exposure in Koreans (Seo et al., 2016). However, the levels of human arsenic exposure among individuals or countries could be affected by several factors, including lifestyles, dietary habits, and geological contamination (Vahter et al., 2000; Mandal and Suzuki, 2002; Minatel et al., 2018). Moreover, it is well known that drinking water is a principal contributing factor to arsenic exposure (Smith et al., 2000; Sun et al., 2007; Huang et al., 2009). In this study, there was no difference in the urinary concentrations of various arsenic species in the study subjects according to the type of drinking water used. The concentration of arsenic in drinking water is well regulated, with a standard limit of $< 10 \text{ } \mu\text{g/L}$ in Korea. Water supply is available for 99.3% of the population, the mean arsenic concentration in the water supply is $< 1 \text{ } \mu\text{g/L}$, and only 3 times were reported as exceeding a standard limit in the supplied water during the last 10 years (MOE, 2021). Nevertheless, as the concentration of arsenic in drinking water was not analyzed in this study, our findings do not suggest that groundwater is safe from arsenic contamination. In a previous nationwide survey of arsenic concentrations in groundwater, about 98% of 722 groundwater had $< 10 \text{ } \mu\text{g/L}$ (Park et al., 2016).

Taken together, the big difference in the urinary concentrations of TmetAs, which is a toxicologically relevant arsenic species, between Korean (30.9 µg/L) and European/American (3–5 µg/L) populations could be possibly explained mainly by their dietary habits. Particularly, the amount of rice consumption is much higher in Koreans, at 62 g/person/year, than in Europeans, at 6 g/person/year (OECD/FAO, 2021). Moreover, Koreans eat higher amounts of seafood and fish, at 56 g/person/year, than Europeans, at 23 g/person/year (OECD/FAO, 2021). The significant portion of the seafood consumed by Koreans is made up of crustacean and mollusk which contain relatively high levels of inorganic arsenic and DMA compared to fish (Sioen et al., 2009; Taylor et al., 2017). Seaweeds, such as dried tangle, dried laver, and kelp, contain high levels of inorganic arsenic (Seo et al., 2016). Koreans consume considerably more seaweeds, at 33 kg/person/year, than Europeans/Americans, who consume very little seaweeds, if at all (FAO, 2021). Nevertheless, it is necessary a more comprehensive study to assess the risk conferred by contributing factors to the arsenic exposure of Koreans and to develop effective exposure-reduction measures.

Previous epidemiologic studies suggested that chronic arsenic exposure may induce metabolic syndrome, diabetes, atherosclerosis and cancer, which could be associated with the increased oxidative stress (De Vizcaya-Ruiz et al., 2009; James et al., 2015; Kuo et al., 2017; Spratlen et al., 2018). C-peptide is a small peptide by-product of insulin synthesis from proinsulin that may be associated with metabolic disease (Suzuki et al., 1997; Kim and Lee, 2017; Yaribeygi et al., 2019). In our study, the increase of MDA and c-peptide concentrations was shown to occur in a dose-dependent pattern according to the urinary concentrations of various arsenic species. These findings suggest that environmental arsenic exposure might be a potential cause of metabolic diseases, such as diabetes mellitus and atherosclerosis, through oxidative stress and endogenous endocrine effects; however, further studies are needed to improve our understanding and to protect the general public from environmental pollutants.

In summary, urinary arsenic concentrations in the adult population in the Republic of Korea was similar to or lower than those in other Asian countries but higher than those in Western countries, including the United States. Overall, our findings suggest that seafood and rice are the main sources of arsenic exposure in Korean adults. Furthermore, overconsumption of seafood might be the main source of exposure to organic arsenic also additional exposure source to inorganic arsenic, which primarily exposed by rice. All of this might have a potentially detrimental effect on human health.

Declarations

Acknowledgements

This study was supported by a Grant (14162MFDS654) from the Ministry of Food and Drug Safety in 2014.

Conflicts of interest

The authors declare no conflicts of interest.

Authors' contributions

Seul-Gi Lee: Writing – Original Draft, Investigation. **Ingu Kang:** Investigation, Methodology. **Mi-Na Seo:** Investigation, Resources. **Jung-Eum Lee:** Formal Analysis, Data Curation. **Sang-Yong Eom:** Resources, Formal Analysis, Software. **Myung-Sil Hwang:** Formal Analysis, Data Curation. **Kyung Su Park:** Methodology, Validation. **Byung-Sun Choi:** Methodology, Resources, Formal Analysis, Data Curation. **Ho-Jang Kwon:** Conceptualization, Resources, Data Curation. **Young-Seoub Hong:** Resources, Methodology. **Heon Kim:** Conceptualization, Resources, Writing - Review & Editing. **Jung-Duck Park:** Conceptualization, Writing - Original Draft, Writing - Review & Editing, Supervision.

References

1. ACGIH (2014) TLV® and BEIs based on the documentation of the threshold limit values for chemical substances and physical agents & biological exposure indices. ACGIH, Cincinnati
2. Agarwal R, Chase SD (2002) Rapid, fluorimetric-liquid chromatographic determination of malondialdehyde in biological samples. *J Chromatogr B Analyt Technol Biomed Life Sci* 775:121–126
3. ATSDR (2007) Toxicological profile for arsenic. Agency for Toxic Substances and Disease Registry. Division of Toxicology, Atlanta
4. Bae HS, Kang IG, Lee SG, Eom SY, Kim YD, Oh SY, Kwon HJ, Park KS, Kim H, Choi BS, Yu IJ, Park JD (2017) Arsenic exposure and seafood intake in Korean adults. *Hum Exp Toxicol* 36:451–460
5. Buchet JP, Lauwerys R, Roels H (1981) Comparison of the urinary excretion of arsenic metabolites after a single oral dose of sodium arsenite, monomethylarsonate, or dimethylarsinate in man. *Int Arch Occup Environ Health* 48:71–79
6. Caldwell KL, Jones RL, Verdon CP, Jarrett JM, Caudill SP, Osterloh JD (2009) Levels of urinary total and speciated arsenic in the US population: National Health and Nutrition Examination Survey 2003–2004. *J Expo Sci Environ Epidemiol* 19:59–68
7. Choi BS, Choi SJ, Kim DW, Huang M, Kim NY, Park KS, Kim CY, Lee HM, Yum YN, Han ES, Kang TS, Yu IJ, Park JD (2010) Effects of repeated seafood consumption on urinary excretion of arsenic species by volunteers. *Arch Environ Contam Toxicol* 58:222–229
8. Cui J, Shi J, Jiang G, Jing C (2013) Arsenic levels and speciation from ingestion exposures to biomarkers in Shanxi, China: implications for human health. *Environ Sci Technol* 47:5419–5424
9. De Vizcaya-Ruiz A, Barbier O, Ruiz-Ramos R, Cebrian ME (2009) Biomarkers of oxidative stress and damage in human populations exposed to arsenic. *Mutat Res* 674:85–92
10. Deutsche Forschungsgemeinschaft (DFG) (2017) List of MAK and BAT Values 2017. Permanent Senate Commission for the Investigation of Health Hazards of Chemical Compounds in the Work Area. Report 53. WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
11. Eom SY, Choi SH, Ahn SJ, Kim DK, Kim DW, Lim JA, Choi BS, Shin HJ, Yun SW, Yoon HJ, Kim YM, Hong YS, Yun YW, Sohn SJ, Kim H, Park KS, Pyo HS, Kim H, Oh SY, Kim J, Lee SA, Ha M, Kwon HJ, Park JD (2014) Reference levels of blood mercury and association with metabolic syndrome in Korean adults. *Int Arch Occup Environ Health* 87:501–513

12. Eom SY, Lee YS, Lee SG, Seo MN, Choi BS, Kim YD, Lim JA, Hwang MS, Kwon HJ, Kim YM, Hong YS, Sohn SJ, Park KS, Pyo HS, Kim H, Kim H, Park JD (2018) Lead, mercury, and cadmium exposure in the Korean general population. *J Korean Med Sci* 33:e9
13. FAO (2021) New Food Balances (database). FAOSTAT. <http://www.fao.org/faostat/en/#data/FBS>. Accessed 13 May 2021
14. Gilbert-Diamond D, Li Z, Perry AE, Spencer SK, Gandolfi AJ, Karagas MR (2013) A population-based case-control study of urinary arsenic species and squamous cell carcinoma in New Hampshire, USA. *Environ Health Perspect* 121:1154–1160
15. Hata A, Endo Y, Nakajima Y, Ikebe M, Ogawa M, Fujitani N, Endo G (2007) HPLC-ICP-MS speciation analysis of arsenic in urine of Japanese subjects without occupational exposure. *J Occup Health* 49:217–223
16. Heitland P, Köster HD (2008) Fast determination of arsenic species and total arsenic in urine by HPLC-ICP-MS: Concentration ranges for unexposed German inhabitants and clinical case studies. *J Anal Toxicol* 32:308–314
17. Hindmarsh JT (2002) Caveats in hair analysis in chronic arsenic poisoning. *Clin Biochem* 35:1–11
18. Hornung RW, Reed LD (1990) Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg* 5:46–51
19. Hsueh YM, Huang YL, Huang CC, Wu WL, Chen HM, Yang MH, Lue LC, Chen CJ (1998) Urinary levels of inorganic and organic arsenic metabolites among residents in an arseniasis-hyperendemic area in Taiwan. *J Toxicol Environ Health Part A* 54:431–444
20. Huang CY, Su CT, Chung CJ, Pu YS, Chu JS, Yang HY, Wu CC, Hsueh YM (2012) Urinary total arsenic and 8-hydroxydeoxyguanosine are associated with renal cell carcinoma in an area without obvious arsenic exposure. *Toxicol Appl Pharmacol* 262:349–354
21. Huang YK, Huang YL, Hsueh YM, Wang JTJ, Yang MH, Chen CJ (2009) Changes in urinary arsenic methylation profiles in a 15-year interval after cessation of arsenic ingestion in Southwest Taiwan. *Environ Health Perspect* 117:1860–1866
22. James KA, Byers T, Hokanson JE, Meliker JR, Zerbe GO, Marshall JA (2015) Association between lifetime exposure to inorganic arsenic in drinking water and coronary heart disease in Colorado residents. *Environ Health Perspect* 123:128–134
23. Kim MS, Lee DY (2017) Urinary glucose screening for early detection of asymptomatic type 2 diabetes in Jeonbuk province Korean schoolchildren. *J Korean Med Sci* 32:985–991
24. Kuo CC, Moon KA, Wang SL, Silbergeld E, Navas-Acien A (2017) The Association of arsenic metabolism with cancer, cardiovascular disease, and diabetes: a systematic review of the epidemiological evidence. *Environ Health Perspect* 125(8):087001
25. Lim JA, Kwon HJ, Ha M, Kim H, Oh SY, Kim JS, Lee SA, Park JD, Hong YS, Sohn SJ, Pyo H, Park KS, Lee KG, Kim YD, Jun S, Hwang MS (2015) Korean research project on the integrated exposure assessment of hazardous substances for food safety. *Environ Health Toxicol* 30:e2015004
26. Link B, Gabrio T, Piechotowski I, Zöllner I, Schwenk M (2007) Baden-Wuerttemberg environmental health survey (BW-EHS) from 1996 to 2003: toxic metals in blood and urine of children. *Int J Hyg Environ*

27. Luvonga C, Rimmer CA, Yu LL, Lee SB (2020) Organoarsenicals in seafood: occurrence, dietary exposure, toxicity, and risk assessment considerations – a review. *J Agric Food Chem* 68:943–960
28. Ma M, Le XC (1998) Effect of arsenosugar ingestion on urinary arsenic speciation. *Clin Chem* 44:539–550
29. Mandal BK, Suzuki KT (2002) Arsenic round the world: a review. *Talanta* 58:201–235
30. McKelvey W, Gwynn RC, Jaffery N, Kass D, Thorpe LE, Garg RK (2007) A biomonitoring study of lead, cadmium and mercury in the blood of New York city adults. *Environ Health Perspect* 115:1435–1441
31. Middleton DRS, Watts MJ, Hamilton EM, Fletcher T, Leonardi GS, Close RM, Exley KS, Crabbe H, Polya DA (2016) Prolonged exposure to arsenic in UK private water supplies: toenail, hair and drinking water concentrations. *Environ Sci Process Impacts* 18:562–574
32. Minatel BC, Sage AP, Anderson C, Hubaux R, Marshall EA, Lam WL, Martinez VD (2018) Environmental arsenic exposure: From genetic susceptibility to pathogenesis. *Environ Int* 112:183–197
33. Ministry of Environment (MOE) (2021) National Waterworks Information System. <https://www.waternow.go.kr/web/lawData/search>. Accessed 26 May 2021
34. Molin M, Ulven SM, Dahl L, Goessler W, Fliegel D, Holck M, Sloth JJ, Oshaug A, Alexander J, Meltzer HM, Ydersbond TA (2014) Urinary excretion of arsenicals following daily intake of various seafoods during a two weeks intervention. *Food Chem Toxicol* 66:76–88
35. Morton J, Leese E (2011) Arsenic speciation in clinical samples: urine analysis using fast micro-liquid chromatography ICP-MS. *Anal Bioanal Chem* 399:1781–1788
36. Morton J, Mason H (2006) Speciation of arsenic compounds in urine from occupationally unexposed and exposed persons in the U.K. using a routine LC-ICP-MS method. *J Anal Toxicol* 30:293–301
37. Mozaffarian D, Rimm EB (2006) Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA* 296:1885–1899
38. Navas-Acien A, Francesconi KA, Silbergeld EK, Guallar E (2011) Seafood intake and urine concentrations of total arsenic, dimethylarsinate and arsenobetaine in the US population. *Environ Res* 111:110–118
39. Nordstrom DK (2002) Public health. Worldwide occurrences of arsenic in ground water. *Science* 296:2143–2145
40. OECD/FAO (2021) OECD-FAO Agricultural Outlook (Edition 2020). OECD Agriculture Statistics (database). <https://doi.org/10.1787/4919645f-en>. Accessed 13 May 2021
41. Park JD, Choi SJ, Choi BS, Lee CR, Kim H, Kim YD, Park KS, Lee YJ, Kang S, Lim KM, Chung JH (2016) Arsenic levels in the ground waters of Korea and the urinary excretion among contaminated area. *J Expo Sci Environ Epidemiol* 26:458–463
42. Petrick JS, Jagadish B, Mash EA, Aposhian HV (2001) Monomethylarsonous acid (MMA(III)) and arsenite: LD(50) in hamsters and in vitro inhibition of pyruvate dehydrogenase. *Chem Res Toxicol* 14:651–656
43. Saoudi A, Zeghnoun A, Bidondo ML, Garnier R, Cirimele V, Persoons R, Fréry N (2012) Urinary arsenic levels in the French adult population: the French National Nutrition and Health Study, 2006–2007. *Sci*

44. Schuhmacher–Wolz U, Dieter HH, Klein D, Schneider K (2009) Oral exposure to inorganic arsenic: evaluation of its carcinogenic and non-carcinogenic effects. *Crit Rev Toxicol* 39:271–298
45. Seo MN, Lee SG, Eom SY, Kim J, Oh SY, Kwon HJ, Kim H, Choi BS, Yu IJ, Park JD (2016) Estimation of total and inorganic arsenic intake from the diet in Korean adults. *Arch Environ Contam Toxicol* 70:647–656
46. Signes-Pastor AJ, Vioque J, Navarrete-Muñoz EM, Carey M, García de la Hera M, Sunyer J, Casas M, Riaño-Galán I, Tardón A, Llop S, Amorós R, Amiano P, Bilbao JR, Karagas MR, Meharg AA (2017) Concentrations of urinary arsenic species in relation to rice and seafood consumption among children living in Spain. *Environ Res* 159:69–75
47. Sioen I, De Henauw S, Van Camp J, Volatier JL, Leblanc JC (2009) Comparison of the nutritional–toxicological conflict related to seafood consumption in different regions worldwide. *Regul Toxicol Pharmacol* 55:219–228
48. Sirot V, Guérin T, Volatier JL, Leblanc JC (2009) Dietary exposure and biomarkers of arsenic in consumers of fish and shellfish from France. *Sci Total Environ* 407:1875–1885
49. Smith AH, Lingas EO, Rahman M (2000) Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. *Bull World Health Organ* 78:1093–1103
50. Spratlen MJ, Grau-Perez M, Best LG, Yracheta J, Lazo M, Vaidya D, Balakrishnan P, Gamble MV, Francesconi KA, Goessler W, Cole SA, Umans JG, Howard BV, Navas-Acien A (2018) The association of arsenic exposure and arsenic metabolism with the metabolic syndrome and its individual components: prospective evidence from the Strong Heart Family Study. *Am J Epidemiol* 187:1598–1612
51. Stýblo M, Drobná Z, Jaspers I, Lin S, Thomas DJ (2002) The role of biomethylation in toxicity and carcinogenicity of arsenic: a research update. *Environ Health Perspect* 110(Suppl 5):767–771
52. Sun G, Xu Y, Li X, Jin Y, Li B, Sun X (2007) Urinary arsenic metabolites in children and adults exposed to arsenic in drinking water in Inner Mongolia, China. *Environ Health Perspect* 115:648–652
53. Suzuki Y, Iizuka T, Kobayashi T, Nishikawa T, Atsumi Y, Kadowaki T, Oka Y, Kadowaki H, Taniyama M, Hosokawa K, Asahina T, Matsuoka K (1997) Diabetes mellitus associated with the 3243 mitochondrial tRNA^{Leu(UUR)} mutation: insulin secretion and sensitivity. *Metabolism* 46:1019–1023
54. Suzuki Y, Shimoda Y, Endo Y, Hata A, Yamanaka K, Endo G (2009) Rapid and effective speciation analysis of arsenic compounds in human urine using anion-exchange columns in HPLC-ICP-MS. *J Occup Health* 51:380–385
55. Taylor V, Goodale B, Raab A, Schwerdtle T, Reimer K, Conklin S, Karagas MR, Francesconi KA (2017) Human exposure to organic arsenic species from seafood. *Sci Total Environ* 580:266–282
56. Tseng WP (1977) Effects and dose-response relationships of skin cancer and blackfoot disease with arsenic. *Environ Health Perspect* 19:109–119
57. Tucker KL (2007) Assessment of usual dietary intake in population studies of gene-diet interaction. *Nutr Metab Cardiovasc Dis* 17:74–81

58. Vahter M, Concha G, Nermell B (2000) Factors influencing arsenic methylation in humans. *J Trace Elem Exp Med* 13:173–184
59. Venugopal V, Gopakumar K (2017) Shellfish: nutritive value, health benefits, and consumer safety. *Compr Rev Food Sci Food Saf* 16:1219–1242
60. Wei Y, Zhu J, Nguyen A (2014) Rice consumption and urinary concentrations of arsenic in US adults. *Int J Environ Health Res* 24:459–470
61. Wen W, Wen J, Lu L, Liu H, Yang J, Cheng H, Che W, Li L, Zhang G (2011) Metabolites of arsenic and increased DNA damage of p53 gene in arsenic plant workers. *Toxicol Appl Pharmacol* 254:41–47
62. Wolle MM, Conklin SD (2018) Speciation analysis of arsenic in seafood and seaweed: Part I-evaluation and optimization of methods. *Anal Bioanal Chem* 410:5675–5687
63. Yaribeygi H, Maleki M, Sathyapalan T, Sahebkar A (2019) The effect of c-peptide on diabetic nephropathy: A review of molecular mechanisms. *Life Sci* 237:116950

Figures

Figure 1

Distribution of urinary concentrations of various arsenic species according to age in Korean adults

Figure 2

The relative proportions of TmetAs and AsB in relation to TsumAs (A), and the relative proportions of InAs and their metabolites (MMA and DMA) in relation to TmetAs (B). InAs: inorganic As [As(III)+As(V)], MMA: monomethylarsonic acid, DMA: dimethylarsinic acid, AsB: arsenobetaine, TmetAs (sum of inorganic arsenic and their metabolites): InAs+MMA+DMA, TsumAs (total sum of arsenic measured): InAs+MMA+DMA+AsB