Identification of Nucleus Sterility Candidate Genes using a Resequencing Technique in Sweet Pepper Sterile Line AB91

Yaning Meng

Hebei Academy of Agriculture and Forestry Sciences <https://orcid.org/0000-0002-2326-2156>

Yanqin Fan (✉ nkyfanyq@126.com)

<https://orcid.org/0000-0001-5724-0097>

libin Yan

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

hongxiao zhang

Hebei Academy of Agriculture and Forestry Sciences

Research article

Keywords: sweet pepper, nuclear male sterile line, genome-wide resequencing technique, candidate gene

Posted Date: December 10th, 2019

DOI: <https://doi.org/10.21203/rs.2.18377/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Agricultural Sciences on January 1st, 2021.
See the published version at <https://doi.org/10.4236/as.2021.122010>.

Abstract

Background: Breeding hybrids with nuclear male sterile lines is an important method for the cross breeding of sweet peppers. To date, few reports have been published on the nuclear male sterility gene of sweet pepper. Yet, there are approximately 20 pepper nuclear male sterility lines in the world.

Results: Using the self-developed testing material, sweet pepper nuclear male sterile dual-purpose line AB91, the genome-wide resequencing technique was applied to firstly find that the mutation site causing the abortion of sweet pepper nuclear male sterility AB91 is on chromosome #5. The mutation gene Capana05g000747 was filtered out and validated by the flight mass spectrometry genotyping method and determined to be the gene causing the abortion of sweet pepper nuclear male sterility AB91. The gene Capana05g000747 contains eight exons and seven introns, and its mutation site is a non-synonymous mutation site located at the 6th exon; the base C mutated into A, and the amino acid changed from alanine to serine. Sequence alignment analysis showed that the gene Capana05g000747 has a similar function to gene At2g02148.

Conclusion: The gene At2g02148 contains a pentatricopeptide repeat protein which has important physiological functions in the gene expression process of organelles and is closely related to the performance of male sterility genes. Therefore, Capana05g000747 was selected as an important candidate gene for sweet pepper nuclear male sterile testing material AB91.

Background

Sweet pepper (*Capsicum annuum* var. *grossum*) is a one-year or perennial crop in the Solanaceae and a variant of pepper. The heterosis of sweet pepper has high utilization value and using male infertility to breed hybrids is an important approach to solving the problem of artificial emasculation. Therefore, the research topic of male infertility has drawn a great deal of attention of domestic and international research scholars. Due to the lack of recovery genes for male sterility in the sweet pepper nucleus and the difficulty of finding excellent recovery lines, the breeding and application of cytoplasmic male sterility in sweet pepper hybrids is limited. However, the genetics of sweet pepper nucleus male sterility is relatively simpler and the number of recovery sources is relatively large, which are a great advantage inbreeding sweet pepper hybrids. Martin and Crawford^[1] first reported capsicum sterility materials controlled by a recessive single-gene. Since then, new genes have been discovered in the pepper male sterile material. To date, nearly 20 nuclear sterility genes have been discovered in peppers^[2-4]. However, there are very few reports on the pepper nuclear male sterility genes. Fan et al.^[5] discovered a natural sterile sweet pepper plant. After years of research, improvement, and breeding, the sweet pepper nuclear male sterile dual-purpose line AB91 was bred, and its sterility feature was controlled by a pair of recessive nuclear genes named msc2^[6]. The sterile line AB91 is completely aborted and stable in sterility with no adverse cytoplasmic effect and has excellent agronomic traits and wide recovery sources. Usually, inbred lines can easily become their own recovery lines, the degree of freedom for association is large, and they are more likely to obtain advantageous combinations. So far, 14 sweet pepper hybrids have been bred using

AB91 and have been widely planted^[7]. We have found that the abortion of sterile plants occurs after the development of quarter microspores at the cytological level; the reasons for microspores abortion are disintegration of the tetraploid enamel wall, tapetum cell dysplasia, and delayed disintegration^[8]; however, there are no reports of co-segregation markers or candidate genes for this gene site.

In recent years, with rapid development of modern molecular biology technology in the field of plants, male infertility research has risen to the level of genetic engineering. At present, genes including *coxII*^[9], *atp6*^[10], *orf456*^[11], *PPR*^[12] and *atp6-2*^[13] are the genes that are connected to the sterility traits for the pepper sterile lines. Ma et al.^[14] and Li et al.^[15] discovered the gene *PAP3* in peppers, which was found to cause atrophy of pollen grains leading to male sterility. Two functional genes of the *MADS-box* gene family are *APETALA3* (*AP3*)/*DEFICIENS* (*DEF*) and *GLOBOSA* (*GLO*)/*PISTILLATA* (*PI*), which are transcription indexes controlling petals and stamens^[16-17]. Fan et al.^[18] found a new gene *CaCTS* from the pepper male sterile line by cloning, which is highly expressed in flowers and seeds, moderately expressed in the placenta and pericarp, and weakly expressed in stems and leaves. Deng et al.^[19] expanded out the malate dehydrogenase gene by reverse transcription PCR. For the *MDH* gene, the performance level is low in abortion and the performance trend is not stable, which may interfere with the energy metabolism balance of the sterile line. Guo et al.^[20] discovered the protein *CaAMS*, the down-regulation of which causes partial filament shortening, withering, stamen non-cracking, and pollen abortion. *CaAMS* plays an important role in the development of pepper tapetum and pollen through a complex regulatory network. Deng et al.^[21] expanded the complete coding sequence for the triose phosphate isomerase (*TPI*) gene by reverse transcriptase PCR. In the abortion process, the activity and performance level of *TPI* in the anthers of male sterile lines were significantly decreased, and the levels of *TPI* in F1 hybrids and maintainer lines remained normal, which indicated that the stable *TPI* transcripts maintained the energy metabolism at a normal level. Qing et al.^[22] performed genomic resequencing and comparison analysis for the male sterile line and fertile line of capsicum and selected Capana02g002096 as a candidate gene for the *msc1* site according to genetic variation and annotation. Capana02g002096 encodes a homologous gene of *AtDYT1*, which is a bHLH transcription index involved in the early development of the tapetum. In addition, a deletion of 7bp was found in the exon of Capana02g002096, which leads the codon to terminate prematurely, resulting in infertility due to a function loss. These prior genetic studies related to infertility have provided important insights for the further study of the mechanism of male infertility, and they also provided a theoretical foundation for related research.

Based on the studies mentioned above, our study used sweet pepper nuclear male sterile line AB91 as the testing material. According to the pepper genome information published^[23], the difference sites and differential candidate genes of fertile and sterile plants for sweet pepper nuclear infertility AB91 were obtained through whole genome resequencing technology, and the genome resequencing results were analyzed by bioinformatics methods. The candidate gene sites were filtered out and validated by a mass spectrometry genotyping method, and the candidate gene of male sterility for AB91 was finally identified, which provides a theoretical and technical foundation for further cloning as well as transformation and utilization of the gene.

Results

Fertility results and statistical analysis

As Figure 1 shows for the male sterile dual-purpose line of the sweet pepper AB91, the anthers of the fertile plant are full and bright yellow, the stigma is lower than the anther, and the whole anther is covered in pollen; however, the sterile plant anther is small and lavender, the stigma is higher than the anther, the anther is shriveled, and there is no pollen. A total of 473 strains of self-crossing F2 generation of the sweet pepper AB91 dual-purpose fertility strain (Msms) were planted in a field, and fertility identification was performed at the flowering stage. As the investigation results for the F2 groups implies, there were 359 fertile plants and 114 sterile plants; the ratio between fertile and sterile plants was in line with the expected theoretical value 3:1. The fertile plants (Msms, MsMs) of the male sterile F2 generation of sweet peppers were singly seeded and planted separately; the group size for each single plant ranges from 30 to 40. The fertile plant (Msms) and pure dominant fertile plant (MsMs) of sweet pepper nuclear male sterile hybrid were filtered out according to the fertility separation of groups of strains. There were 110 pure dominant fertile plants and 249 heterozygous fertile plants in F2.

Bulked-segregant analysis and data analysis

MsMs and msms gene banks were sequenced by the bulked-segregant analysis sequencing (BSA-Seq) technique; 417.04 million and 467.1 million filtered reads were obtained from the MsMs and msms pools, respectively. The MsMs-pool comparison ratio was 99.31%, the average coverage depth for the reference genome (excluding the N region) was 19.26X, and the 1X coverage degree (at least one base got covered) was 92.78%. The msms-pool comparison ratio was 99.03%, the average coverage depth for the reference genome (excluding the N region) was 21.55X, and the 1X coverage degree (at least one base got covered) was 92.95%. According to the results in Table 1, the comparison results are normal and can be used for subsequent mutation detection and correlation analysis.

Table 1. Sequencing depth and coverage ratio statistics

Sample	Mapped reads	Total reads	Mapping rate (%)	Average depth (X)	Coverage at least 1X (%)
MsMs	414,182,342	417,041,448	99.31	19.26	92.78
msms	462,562,325	467,111,904	99.03	21.55	92.95

In order to reduce the impact of sequencing errors and comparison errors, the SNP index and InDel index polymorphism sites were filtered out, and the number of polymorphic marker sites was 11,348,482 after filtration. By making a difference between the SNP index and InDel index for two descendants by choosing a 95% confidence level. The window larger than the threshold was selected as the candidate

interval, and the numbers of polymorphism marker sites selected out for SNP and InDel were 27,541 and 1,865 respectively. For the candidate site extraction ANNOVAR annotation results, stop loss, and stop gain were selected with the priority, non-synonymous mutation, or alternative splicing site selected as the candidate gene. The number of SNP candidate genes filtered out from the fertile and sterile gene pools was 35(InDel has no annotated candidate genes above),and 33 candidate genes of them located at chromosome #5 (Figure 2).

Functional annotations of genes

The 33 genes in the correlated region are compared to the databases of NR, SwissProt^[24], GO^[25] and KEGG^[26] used BLAST software to identify 10 candidate genes and 13 candidate sites connecting to sweet pepper nuclear infertility, respectively, and they are speculated to be related to putative pentatricopeptide repeat-containing protein., galactinol synthase, glucosyltransferase, zeatin O-xylosyltransferase, threonine-protein kinase, histone acetyltransferase, U-box domain-containing protein, cell division cycle, and apoptosis regulator protein. All these candidate gene sites are located at chromosome #5. Hence, the candidate gene controlling the sweet pepper nuclear male sterile AB91 is determined at chromosome #5(Table 2).

Table 2. Candidate genes comments

TransID	Pos	Subject Description
Capana05g000747	28594037 28599926	PREDICTED: uncharacterized protein At2g02148 isoform X1[Solanum lycopersicum]
Capana05g000767	30024258 30026988	PREDICTED: pentatricopeptide repeat-containing protein At1g03100, mitochondrial-like isoform X1[Solanum tuberosum]
Capana05g000608	18775367 18776004	PREDICTED: galactinol synthase 2-like [Nicotiana tomentosiformis]
Capana05g000611	18835766	PREDICTED: putative glucosyltransferase [Solanum lycopersicum]
Capana05g000612	18852022	PREDICTED: zeatin O-xylosyltransferase-like [Solanum lycopersicum]
Capana05g000617	19194137	
Capana05g000657	22059680	PREDICTED: probable LRR receptor-like serine/threonine-protein kinase At1g34110-like [Solanum tuberosum]
Capana05g000711	25862908	PREDICTED: cell division cycle and apoptosis regulator protein 1-like [Solanum tuberosum]
Capana05g000715	26374526	PREDICTED: histone acetyltransferase HAC1-like isoform X1 [Solanum tuberosum]
Capana05g000896	34599677	PREDICTED: U-box domain-containing protein 21-like [Solanum tuberosum]

Mass spectrometry analysis

A mass spectrometry genotyping technique was used to detect the 13 mutation sites obtained by BSA-sequencing a population of 222 self-crossed F2 generation of sweet pepper nuclear male sterile dual-purpose AB91 fertile strain (Msms). Both sites 19194137 and 34599677 did not exhibit complete cosegregation with msc2. Therefore, the two corresponding genes Capana05g000617 and Capana05g000896 were excluded. The remaining 11 locations showed co-segregation, but it was slightly different from the msc2 phenotype detection in the field; the mutation site 28594037 had the highest similarity with msc2, and the accuracy rate (the number consistent with the msc2 phenotype detection in the field/F2 generation total population size) was 99.5%, and the corresponding gene for this site is Capana05g000747. Hence, the Capana05g000747 gene was determined to be the most powerful candidate gene for msc2 (Figure 3).

Sequence analysis of male sterility gene msc2 for sweet pepper line AB91

The gene *msc2* of sweet pepper male sterile AB91 is composed of eight exons and seven introns. The cDNA sequence is 1350bp in length and encodes 450 amino acids. The mutation site of the sterility gene is located at the 6th exon of Capana05g000747, the base C is mutated into A, and the amino acid is changed from alanine to serine, which causes a change in the fertility expression of the sterile plant of sweet pepper nuclear male sterile dual-purpose line AB91, leading to male sterility(Figure 4).

Homology of nuclear male sterility gene *msc2* in sweet pepper AB91 with other species

For a further functional study of *msc2*, we performed a sequence alignment and homology relationship analysis for the gene *msc2* and other species, which showed that *msc2* gene was at the same branch as tomato, potato, and tobacco, indicating that the their homology relationship is very close and *Msc2* is more likely to have the same effect as a close homology. However, the function of gene *msc2* in tomato, potato, and tobacco sequences is predicted, and its function was found to be similar to the At2g02148 gene, which encodes the pentatricopeptide repeats (PPR) protein. This finding preliminarily implies that the *msc2* gene has a connection with the PPR protein(Figure 5).

Discussion

Employing male sterility to breed advantageous hybrids is the most economical and efficient approach. To date, two types of recessive nuclear male sterility resources for sweet peppers have been discovered in China; one is the pepper male sterile material discovered by Shizhou Yang with the nuclear sterility gene named *msc1*, and the other is the sweet pepper male sterile line AB91 used in this study with its nuclear sterility gene named *msc2*. To date, sweet pepper nuclear male sterile line AB91 has been used to breed five national recognized vegetable varieties (JiYan5, JiYan6, JiYan15, JiYan16, JiYan108) and nine provincial recognized vegetable varieties (JiYan8, JiYan12, JiYan13, JiYan19, JiYan4, JiYan105, JiYan102, JiYan20, JiYan28), which are widely planted throughout China. Therefore, it is of great meaning to study the sterility gene *msc2* of the sweet pepper male sterile line AB91. However, the research on the infertility mechanism of the *msc2* gene has not been reported yet. Whole genome sequencing technology (WGS) is currently the most effective method for group function genetic mining and has many advantages such as its ability to contain comprehensive, efficient, and accurate information. WGS is a bioinformatics technique that performs differential sequence analysis of individuals or groups at the genomic level to explore species and filter out functional genes^[27]. Currently, WGS has been widely used in rice^[28], cucumber^[29], potato^[30], watermelon^[31], sorghum^[32], and other crops. Therefore, this study carried out in-depth research to analyze the reason for abortion caused by gene *msc2* using genetic sequencing and genotyping technology.

Material with prominent and genetically stable objective traits

The sweet pepper male sterile dual-purpose line AB91 is a sister cross-hybrid whose agronomic traits within groups have remained stable after years of breeding and whose sterility is controlled by a pair of recessive nuclear genes. The pure dominant fertile plant has a genetic background that is consistent with that of the pure recessive sterile plant, which avoids the wrong interference analysis caused by the genetic background difference between fertility materials.

Capana05g000747 is an important candidate gene of Msc2

In this study, the the mutation site that caused the abortion of sweet pepper male sterile AB91, which is on chromosome #5, was first discovered through WGS technology. Flight mass spectrometry genotyping was used to verify the differential gene and filter out the gene Capana05g000747, which is most likely to be the important gene leading to the abortion of sweet pepper male sterile AB91. For the gene Capana05g000747 including eight exons and seven introns, the mutation site 28549037 is a non-synonymous mutation site with its position at the 6th exon of Capana05g000747; the features with base C mutated into A and the amino acid changed from alanine to serine are what causes the male sterility of sweet pepper malesterile line AB91.

Sequence analysis of the Capana05g000747 gene

For a further function of the gene Capana05g000747, we performed a sequence alignment analysis that showed that the gene homology is close to tomato, potato, and tobacco, but their functional annotations were all predicted and are similar to that of the gene At2g02148. At2g02148 encodes a PPR gene protein, which is encoded by a nuclear gene and consists of 35 degenerate amino acids in a series of repeating units; most of the protein is transported into organelles to fulfill their functions^[33]. The PPR gene protein has important physiological functions in the gene performance process of organelles and is involved in almost all stages of gene performance, including transcription^[34], RNA splicing^[35], RNA editing^[36], translation^[37], and RNA stability maintenance^[38]. Previous studies have shown that the PPR gene protein is dispersed all over the entire pepper genome^[39-40], while most of the Rf candidate genes are on chromosome #6^[41-42] and the sterility performance is realized by controlling the male sterility related genes. However, our research shows that the candidate gene protein PPR of the nuclear male sterility line AB91 on chromosome #5 has a base mutation, which causes a protein function change, resulting in abortion microspores, but the molecular mechanism of the PPR gene protein in the sweet pepper nuclear male sterility AB91 is still unclear and its verification will require further research.

Materials And Methods

Plant materials

The sweet pepper recessive male sterile dual-purpose line AB91 was provided by the Sweet Pepper Research Group from the Institute of Cash Crops of Hebei Academy of Agriculture and Forestry Sciences.

Fifty percent of the sweet pepper recessive male sterile dual-purpose line AB91 groups are fertile plants and the remainder are sterile plants. The sterility trait, which is controlled by a pair of recessive nuclear genes, is genetically stable.

Basic group construction: F2 generation isolated groups containing recessive sterile plants (msms) and fertile plants (Msms, MsMs) were obtained from the self-crossing of fertile plants (Msms) in the sweet pepper male sterile line AB91. In order to determine the genotype of the fertile plants, the fertile plants in the F2 plants were self-crossed, and the homozygous dominant fertile plants (MsMs) and hybrid plants (Msms) were separated and identified according to offspring fertility to complete the basic group construction.

Gene bank construction and Illumina sequencing

The homozygous fertile plants (MsMs) and recessive sterile plants (msms) in the F2 generation isolated group of sweet pepper AB91 were used as testing materials, and the DNA of the young leaves of plants was extracted by the CTAB method^[43] to construct the MsMs and Msms gene banks. The DNA sample was fragmented into a size of 350bp by sonication, then the DNA fragments were finally polished, A-tailed, and ligated with the full-length adapter for Illumina sequencing with further PCR expansion. At last, the PCR products were purified (AMPure XP system). Then, the size distribution of gene banks was analyzed by an Agilent2100 Bioanalyzer (Aligent, Santa Clara, CA, USA). Quantified analysis was performed using real-time PCR. The genebanks constructed above were sequenced by an Illumina HiSeq4000 platform (Illumina, San Diego, CA, USA) and 150bp paired-end reads were generated with an insert size of approximately 350bp.

Data analysis

The raw reads from the msms and MsMs pools were filtered and aligned to the pepper sequence genomics (<http://peppersequence.genomics.cn/page/species/download.jsp>) using the Burrows Wheeler BWA alignment tool^[44]. GATK software was used to detect single-nucleotide polymorphisms (SNPs) and insertion-deletion (InDels)^[45]. The reads depth information for the above homozygous SNPs/InDels in the offspring pools was obtained to calculate the SNP/InDel index^[46]. We filtered out those points that had an SNP/InDel index in both pools of less than 0.3. The sliding window method was used to present the SNP/InDel index of the whole genome. The average of all SNP/InDel indexes in each window was as the SNP/InDel index for this window. Usually, we use a window size of 1 Mb and step size of 10 Kb as default settings. The difference of the SNP/InDel index of the two pools was calculated as the delta SNP/InDel index. The differential candidate genes for male sterile and fertile plants of sweet pepper male sterile line AB91 are identified by calculating the indexes of the SNP and SNP/InDel index. The differential candidate genes of male sterile and female fertile AB91 nuclear male sterile lines were determined by calculating the values of SNP/InDel index and $\Delta(\text{SNP/InDel index})$.

Mass spectrometry method identification

The self-crossing F2 generation of the sweet pepper male sterile line AB91 fertile line (Msms) was tested by the flying mass spectrometry Sequenom platform typing technique^[47], and the testing results were read in real-time and completely analyzed by the software Mass ARRAY® (Agena Bioscience Inc., San Diego, CA, USA). The detection for all samples was repeated to verify accuracy. The test results were compared with the test results with the results of phenotypic identification in the field to calculate the accuracy rate (the number of test results consistent with the phenotype identification in the field divided by the total number of F2 generation groups), and filter out the candidate genes for sweet pepper nuclear male sterility msc2 according to the accuracy rate.

Declarations

Conflicts of interest The authors declare that they have no competing interests.

Funding This work was supported by National Key Research and Development Projects (No. 2016YFD0101704) and Hebei Science and Technology Support Program Project (16226308D-9).

Author Contributions Yanqin Fan and Libin Yan designed the project. Yaning Meng performed the experiments, analyzed the data and wrote the manuscript. All authors contributed to the revision of this manuscript and approved the final manuscript.

Acknowledgments We thank LetPub (www.letpub.com) for its linguistic assistance during the preparation of this manuscript.

References

1. Martin J.A., and Crawford J.H. (1951) Several types of sterility in *Capsicum frutescens*. Hort. Sci. 57:335-338.
2. Wang D.Y., Yang F.M., Li Y., Wang H.M. (2008) Advances in research on male sterility genes in pepper. China Vegetables 9:40-43.
3. Bartoszewski G., Waszczak C., Gawronski P., Stepień I., Bolibok B.H., Palloix A., Lefebvre V., Korzeniewska A., Niemirowicz S.K. (2012) Mapping of the ms8 male sterility gene in sweet pepper (*Capsicum annuum* L.) on the chromosome P4 using PCR based markers useful for breeding programmes. Euphytica, 186(2):453-461.
4. Lee J., Han J.H., An C.G., Lee W.P., Yoon J.B. (2010a) A CAPS marker linked to a genic male-sterile gene in the colored sweet pepper 'Paprika' (*Capsicum annuum* L.). Breed Sci. 60(1): 93-98.
5. Fan Yanqin, Guo Jingyin (1993) A Preliminary Report on Breeding of Male-sterile Lines AB91 in Sweet Pepper. Acta agriculture universitatis pekinensis 19:118-122.
6. Wang D.Y., Yang F.M., Li Y., Wang H.M. (2008) Advances in research on male sterility genes in pepper. China Vegetables 9:40-43.

7. Fan Yanqin,Liu Yun,Yan Libin(2003)Breeding and Application of Male-sterile Lines in Sweet Pepper. *Acta Agriculturae Boreali Sinica* 18:81-83.
8. Luo Shuangxia, Fan Yanqin, Shen Shuxing, Chen Xueping, Liu Yun, Yan Libin (2006) Observation on Microspore Development of Genetic Male-sterile Pepper. *Acta Agriculturae Boreali-Sinica* 21(2):99-103.
9. Kim D,Kang J G,Kim S,et al(2001)Identification of *coxII* and *atp6* regions as associated to CMS in *Capsicum annuum* by using RFLP and long accurate PCR.Korean Society for Horticultural Science 42(2):121-127
10. Kim D H,Kim B D(2006)The organization of mitochondrial *atp6* gene region in male fertile and CMS lines of pepper(*Capsicum annuum* L.).*Curr Genet* 49(1):59-67
11. Kim D H,Kang J G,Kim B D(2007)Isolation and characterization of the cytoplasmic male sterility-associated *orf456* gene of chili pepper(*Capsicum annuum* L.).*Plant Mol Biol* 63(4):519-532
12. Derek W,Barchenger,Joseph I(2018)Genome wide identification of chile pepper pentatricopeptide repeat domains provides insight into fertility restoration. *Journal of the American Society for Horticultural Science* 143(6):418-429.
13. Ji JiaoJiao,Huang Wei,Li DaWei,Yin YanXu,Chai WeiGuo,Gong ZhenHui (2014)A CMS-Related Gene *atp6-2* Causes Increased ATP Hydrolysis Activity of the Mitochondrial F1F0-ATP Synthase and Induces Male Sterility in Pepper (*Capsicum annuum* L.).*Plant Molecular Biology Reporter* 32(4):888-899.
14. Ma Ning,Liu Chen,Yang Wencai,Shen Huolin(2016)*PAP3* regulates stamen but not petal development in *Capsicum annuum* L.*Horticultural Plant Journal* 2(2):91-96.
15. Li Xin,Liu Chen,Da Fengjiao,Ma Ning,Shen Huolin(2013)Expression pattern of class B gene *PAP3* in flower development of pepper.*International Journal Of Molecular Sciences* 14(12):24643-24655.
16. BeckerA.,Theissen G(2003)The major clades of *MADS-box* genes and their role in the development and evolution of flowering plants.*Phylogenet Evol* 29:464-489.
17. Whipple C.J.,Ciceri P,Padilla C.M.,Ambrose B.A.,Bandong S.L.,Schmidt R.J.(2004) Conservation of B-class floral homeotic gene function between maize and Arabidopsis.*Development* 131:6083-6091.
18. Fan Shi,Deng Minghua,et al(2016)Isolation and characterization of citrate synthase gene *CaCTS* in pepper cytoplasmic male sterility.*Research Journal Of Biotechnology* 11(10):9-15.
19. Deng Minghua,Lv Junheng,Zhao Kai(2017)Molecular cloning sequence characterization of a novel pepper gene *MDH* and its effect on cytoplasmic male sterility.*Research Journal Of Biotechnology* 12(4):62-68.
20. Guo Jinju,Liu Chen,Wang Peng,Cheng Qing,Sun Liang,Yang Wencai,Shen Huolin(2018)The Aborted Microspores (AMS)-Like Gene Is Required for Anther and Microspore Development in Pepper(*Capsicum annuum* L.).*International Journal Of Molecular Sciences* 19(5):12-20.
21. Deng Minghua, Wen Jinfen, Huo Jinlong,et al(2012)cDNA cloning sequence characterization and expression analyses of a novel gene-*TPI* from pepper cytoplasmic male sterility(CMS),*Journal of Medicinal Plants Research* 6(10):1997-2003.

22. Qing Cheng,Peng Wang,Jinqiu Liu,et al(2018)Identification of candidate genes underlying genic male-sterile msc1 locus via genome resequencing in *Capsicum annuum* L.Theoretical and Applied Genetics 131:1861-1872.
23. Cheng Qin,Changshui Yu,Yaou Shen,Xiaodong Fang,Lang Chen,et al(2014)Whole-genome sequencing of cultivated and wild peppers provides insights into *Capsicum* domestication and specialization.Proceedings of the National Academy of Sciences 111(14):5135-5140.
24. Yip YL,Famiglietti M,Gos A,Duek PD,David FPA,Gateau A,Bairoch A(2008)Annotating single amino acid polymorphisms in the UniProt Swiss-Prot knowledgebase.Hum Mutat 29(3):361-6.
25. Sherlock G.(2009)Gene ontology: tool for the unification of biology.Canadian Ins J Food Scie Tech.22(4):415.
26. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M(2016)KEGG as a reference resource for gene and protein annotation. Nucleic Acids Res.44:457-62.
27. Pan Z Y,He X Y,Liu Q Y,Guo X F,Cao X H,Hu W P,Di R,Wang X Y,Chu M X(2016)Application of Whole Genome Sequencing(WGS) in Population Evolution and Gene Discovery of Domestic Animals.Journal of Agricultural Biotechnology 24(12): 1945-1954.
28. Dayong lia,Zhiyuan Huangb,Shuhui Song,et al(2016)Integrate analisys of phenome genome and transcriptome of hybrid rice uncovered multiple heterosis-related loci for yield increase.PNAS 113(41):E6026-6035.
29. Lu H,Lin T,Klein J,et al(2014)QTL-seq indentifies an early flowering QTL located near flowering locus T in cucumber.Theoretical and applied genetics 127(7):1491-1499.
30. Zhang G D,Liu B I,Li X B,SI HuaiJ,Li X Q(2016)Effects of potato cultivars and tuber sections on the sequencing depth pattern in genome.Journal of Gansu Agricultural University. 51(3): 43-48.
31. Guo S G,Zhang J G,Sun H H,Salse J,Lucas W J,Zhang H Y,Zheng Y,Mao L Y,Ren Y,Wang Z W(2013)The draft genome of water melon(*Citrullus lanatus*)and resequencing of 20 diverse accessions.Nat Genet 45: 51-58.
32. Zheng L Y,Guo X S,He B,Sun L J,Peng Y,Dong S S,Liu T F,Jiang SRamachandran S,Liu C M,Jing H C(2011)Genome wide patterns of genetic variation in sweet and grain sorghum (*Sorghum bicolor*).Genome Biol 12:R114.
33. Ma Ning,Liu Chen,Yang Wencai,Shen Huolin(2016)*PAP3* regulates stamen but not petal development in *Capsicum annuum* L.Horticultural Plant Journal 2(2):91-96.
34. Hashimoto M,Endo T,Peltier G,et al(2003)A nucleus encoded factor CRR2 is essential for the expreesion of chloroplast ndhB in Arabidopsis.The Plant Journal 36(4):541-549.
35. Ichinose M,Tasak I E,Sugita C,et al(2012)A PPR-DYW protein is required for splicing of a group intron of cox1 pre-mRNA in *Physcomitrella patens*.The Plant Journal 70(2):271-278.
36. Rudinger M,Volkmar U,Lenz H,et al(2012)Nuclear DYW-type PPR gene families diversify with increasing RNA editing frequencies in liverwort and moss mitochondria.Journal of Molecular Evolution.74(1-2):37-51.

37. Tavares C F,Camacho V Y,Zamudio O,et al(2007)The pentatricopeptide repeats present in PET 309 are necessary for translation but not for stability of the mitochondrial COX1 mRNA in yeast.Journal of Biological Chemistry.283(3):1472-1479.
38. Yamazaki H,Tasaka M,Shinkanai T(2004)PPR motifs of the nucleus encoded factor PGR3 fuction in the selective and distinct steps of chloroplast gene expression in Arebidopsis.The Plant Journal.38(1):152-163.
39. Lurin, C., C. Andres, S. Aubourg, M. Bellaoui, et al(2004)Genome-wide analysis of Arabidopsis pentatricopeptide repeat proteins reveals their essential role in organelle biogenesis. Plant Cell 16:2089–2103.
40. Wang,Z.,Y.Zou,X.Li,Q.Zhang,L.Chen,H.Wu,D.Su,Y.Chen,J.Guo,D.Luo,Y.Long,Y.Zhong,and Y.Liu(2006)Cytoplasmic male sterility of rice with boro II cytoplasm is caused by a cytotoxic peptide and is restored by two related PPR motif genes via distinct modes of mRNA silencing.Plant Cell 18:676–687.
41. Jo Y.D., Kim Y.M., Park M.N.,Yoo J.H.,Park M.,Kim B.D.,Kang B.C.(2010) Development and evaluation of broadly applicable markers for Restorer of fertility in pepper.Mol.Breeding 25:187-201.
42. Ding A M,Li F X,Chen Y Q,et al(2014)Homology based cloning and expression analysis of Rf genes encoding PPR-containing proteins in tobacco.Genet Mol Res 13: 2310-2322.
43. Healey A, Furtado A, Cooper T, Henry RJ(2014)Protocol: a simple method for extracting next-generation sequencing quality genomic DNA from recalcitrant plant species.Plant Methods 10:21.
44. Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357–359.
45. McKenna A, Hanna M, Banks E, et al(2010)The Genome Analysis Toolkit: a MapReduce framework for analyzing next generation DNA sequencing data.Genome research 20(9): 1297-1303.
46. Takagi H,Abe A,Yoshida K,et al(2013)QTL-seq: rapid mapping of quantitative trait loci in rice by whole genome resequencing of DNA from two bulked populations.The Plant Journal 74(1):174-183.
47. Schaeffeler E,Zanger U M,Eichelbaum M,et al(2008)Highly multiplexed genotyping of thiopurine smethyl transferase variants using malditof mass spectrometry:Reliable genotyping in different ethnic groups.ClinChem 54(10):1637-1647

Figures

Figure 1

Morphological identification of sterile and fertile plants of the sweet pepper AB91 male sterile dual-purpose line: (A) The sterile plant anther is small and lavender, and the stigma is higher than the anther; (B) fertile plant is full of anthers and bright yellow, and the stigma is lower than the anther; (C) sterile plant anther is shriveled and has no pollen; (D) pollen covers entire anther.

Figure 2

Distribution of msms and MsMs-bulk ΔAll -index at chromosome for sweet pepper male sterile Horizontal axis: chromosome length (Mb); vertical axis: Δ (All-index); and the blue horizontal line is the threshold at the 95% confidence level.

Figure 3

Mass spectrometry genotyping test results for sweet pepper nuclear male sterile AB91:(A)mass spectrometry detection result of a high accuracy candidate site; (B) accuracy histograms of sweet pepper nuclear male sterile AB91F2 generation for 11 candidate sites.

Figure 4

Sequence information of the MSMS gene and the msms gene in sweet pepper nuclear male sterile line AB91

Figure 5

Homologous relationship tree for gene msc2 between sweet pepper nuclear male sterility AB91 and other species; unknown represents annotated with gene msc2