Construction of high copy number and highly stable *Escherichia coli*-*Bacillus subtilis* shuttle plasmids based on pWB980

XingYa Zhao

University of the Chinese Academy of Sciences

JianYong Xu

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

Ming Tan

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

Jie Zhen

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

WenJu Shu

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

ShiBin Yang

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

YanHe Ma

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

Hongchen Zheng

Tianjin Institute of Industrial Biotechnology Chinese Academy of Sciences

Hui Song (✉ song_h@tib.cas.cn)

Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences <https://orcid.org/0000-0003-2642-1850>

Research

Keywords: Expression vectors, pUC980, *Bacillus subtilis*, Alkaline pectate lyase, Alkaline protease, Pullulanase

Posted Date: December 10th, 2019

DOI: <https://doi.org/10.21203/rs.2.18382/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background: pWB980 derived from pUB110 is a promising expression vector in *Bacillus* for its high copy number and high stability. However, the low transformation rate of recombinant plasmids to the wild cells limited the application of it. On the basis of pWB980, constructing an *E. coli* - *B. subtilis* shuttle plasmid could facilitate the transformation rate to *Bacillus* cells. Because the insertion site for *E. coli* replication origin sequence (ori) is not unique in pWB980, in order to investigate the best insertion site, eight shuttle plasmids (pUC980-1~pUC980-8) containing all possible insertion sites and directions were constructed.

Results: The results showed that all the selected insertion sites could be used to construct shuttle plasmid but some sites required a specific direction. And different insertion sites led to different properties of the shuttle plasmids. The best shuttle plasmids pUC980-1 and pUC980-2, which showed copies more than 450 per cell and segregational stabilities up to 98%, were selected for heterologous expressions of an alkaline pectate lyase gene *pelN*, an alkaline protease *spro1* and a pullulanase gene *pulA11*, respectively. The highest extracellular activities of *PelN*, *Spro1* and *PulA11* were upto 5200 U/mL, 21537 U/mL and 504 U/mL correspondingly after 54 h, 60 h and 48 h fermentation in a 10 L fermentor. Notably, *PelN* and *Spro1* showed remarkably higher yields in *Bacillus* than previous reports.

Conclusion: In this work we can conclude that the optimum ori insertion site was the upstream region of BA3-1 in pWB980 which resulted in shuttle plasmids with higher copy numbers and higher stabilities. The novel shuttle plasmids pUC980-1 and pUC980-2 will be promising expression vectors in *B. subtilis*. And the ori insertion mechanism revealed in this work could provide theoretical guidance for further studies of pWB980 and constructions of other shuttle plasmids.

Background

Bacillus subtilis possessing non-pathogenic and strong protein secretion ability is a promising safe host strain for bio-industrial productions^[1–3]. In the last 30 years, *B. subtilis* expression systems have been well developed. Many foreign proteins have been produced in *B. subtilis* cells successfully by using different kinds of expression plasmids^[4–6]. However, compared to *Escherichia coli* expression system, relatively difficult in plasmid transformation, instable of plasmids, and lack of diversity for available plasmids became the main obstacles of the development of *B. subtilis* expression system^[7, 8].

At present, the majority of available *B. subtilis* expression plasmids are derived from pC194^[9–11], pLS1^[12, 13] and pUB110^[3, 14]. In order to improve transformation efficiency of plasmids, a lot of *E. coli*-*B. subtilis* shuttle plasmids such as pHCMC05^[15] and pHT43^[16] have been constructed. However, the available plasmids in recent researches remain difficult to combine high copy number and high stability in the fermentation for producing heterologous proteins. As a delegate of high copy number plasmids, pWB980^[17] derives from plasmid pUB110 has been doing well in heterologous gene expression in *B. subtilis*^[14, 18, 19]. However, the low transformation rate of the ligation products between pWB980 and target genes became a barrier to the widely use of it. On the purpose of solving this problem, many

shuttle plasmids were constructed based on pWB980^[19, 20]. But so far, there is still a lack of systematical studies about the manipulations of pWB980.

In this work, the *ori* region where plasmid-replication is originated in *E. coli* was inserted into any possible sites of pWB980 to produce a series of *E. coli*-*B. subtilis* shuttle plasmids. Because the replication gene insertions may usually cause the changes of copy number and stability of the plasmids^[21, 22], we focused on detecting the copy number and stability of the recombinant plasmids and found some promising *E. coli*-*B. subtilis* shuttle plasmids with high copy numbers and good stabilities. Moreover, an alkaline pectate lyase gene *pelN* from *Paenibacillus* sp. 0602^[23], an alkaline protease *spo1* from alkaliphilic *Bacillus* sp. 221^[24] and a pullulanase gene *pulA11* from *Anoxybacillus* sp. LM18-11^[25] were improving expressed with the specific plasmids in *B. subtilis* WB600. Thus, we believe that these plasmids will be a wider range of uses in further research of *B. subtilis* expression system.

Results

Construction of *E. coli*-*B. subtilis* shuttle plasmids

In general, smaller size of plasmid usually cause easier to be manipulated. On the purpose of shortening pWB980 before construction, the more commonly used gene *kan* was kept while the *bleoR* was deleted from it (Fig 1). As the co-existence of selective genes *bleoR* and *kan* were proved to be unnecessary in other studies and our previous work^[18, 19]. The truncated plasmid pWB980-DB was successfully obtained in this work (Fig 2). It was found to have the same stability with pWB980 along with a higher copy number 272 than that (134) of pWB980 as shown in Fig 3.

Then eight *E. coli*-*B. subtilis* shuttle plasmids derived from pWB980-DB were constructed successfully according to the design in Fig 1. pWB980-DB was in the size of 3388 bp and the eight shuttle plasmids were all in the size of 4276 bp. In order to verify the plasmids clearly, double enzyme-digestions with *EcoRI*/*HindIII* were carried out and the results were shown in Fig 2. All of them replicated normally both in *E. coli* and *B. subtilis* except pUC980-8 which could not replicate in *B. subtilis* host cells. This might be due to the interruption of reverse *ori* to the replication of the *rep* gene.

Copy number of recombinant shuttle plasmids in *B. subtilis* 168

The copy numbers of all new plasmids were detected and stated in Fig 3. Shuttle plasmids pUC980-1, pUC980-2 and pUC980-6 obtained high copy numbers of 584, 492 and 487 correspondingly. pUC980-1 had the highest copy number, which was over four times than that of pWB980 and twice than that of pWB980-DB. The copy numbers of plasmids pUC980-3 (305), pUC980-4 (295) and pUC980-5 (286) were similar to that of pWB980-DB. Only plasmid pUC980-7 with 107 copies per cell had a lower copy number than that of pWB980. pUC980-8 was not counted due to its un-replicability in *B. subtilis*. Therefore, the shuttle plasmids were classified into three groups (C1~C3) according to their copy number levels from high to low as shown in Fig 3B. Plasmids in C1 were determined to be high copy plasmids had copy

numbers higher than 400, including pUC980-1, pUC980-2 and pUC980-6. C2 is the middle copy plasmids group included plasmids pUC980-3, pUC980-4, and pUC980-5 with copy numbers from 200 to 400. Plasmid pUC980-7 was ranked into C3 with relatively low copy numbers from 100 to 200.

Stability of recombinant shuttle plasmids in *B. subtilis* 168

Plasmid stability is consist of structural stability and segregation stability. All plasmids are kept stable in structures after 30-days culturing as they were enzymatic digested into expected fragments consistent with Fig 2. The good structural stabilities of these plasmids were further verified by sequencing (data not shown). pWB980 and pWB980-DB were stable during passages with segregation stabilities of 100% (Fig 3A). The pUC980-series shuttle plasmids could be clarified into three levels based on their segregation stabilities from high to low (Fig 3B). Plasmids pUC980-1(99%), pUC980-2 (100%) and pUC980-5 (99%) were ranged into group S1 (high stable plasmid group). They acted well with more than 98% transformants retained after 30-days incubation in LB medium showing the similar stability with pWB980 and pWB980-DB. pUC980-4 were included in group S2 (low stable plasmid group) with a segregation stability of 39%. Plasmids in group S3 (unstable plasmids group) contained segregation stabilities lower than 10%, including pUC980-3 (2%), pUC980-6 (4%) and pUC980-7 (0).

Synthetically, all plasmids mentioned above were arranged and stated in Fig 3B combining their ranks in copy number (C1 to C3) and segregation stability (S1 to S3). pUC980-1 and pUC980-2 were clarified both in C1 and S1. Plasmid pUC980-6 in C1 obtained high copy number but low segregation stability in S3. Plasmid pUC980-5 in C2 were listed in S1, having relatively lower copy number than plasmids in C1. pUC980-4 performed moderately both in C2 and S2, while plasmid pUC980-3 in C2 was so unstable that ranked in S3. pUC980-7 was not only the most unstable plasmid but also had the lowest copy number, listed in C3 and S3. Based on the classification, plasmids pUC980-1 and pUC980-2 performing well both in copying and passaging were supposed to be the optimal expression vectors. Plasmids pUC980-5 and pUC980-6 only showed high stability and high copy number respectively. To further test the application potential of these new plasmids in production of industrial enzymes, three different industrial enzymes were selected for heterologous expression using these plasmids in *B. subtilis* WB600.

Alkaline pectate lyase expression in *B. subtilis* WB600

The alkaline pectate lyase gene *pelN* was expressed in these plasmids mentioned above separately. All plasmids functioned normally with extracellular secretory expressions after 48 h culturing in shake flasks. The results indicated obvious differences in expression levels as presented in Fig 4A. The extracellular activities of PelN were more than 1500 U/mL when using pUC980-1, pUC980-2, pUC980-5 and pWB980-DB as expression vectors (Fig 4A). The highest activity was 2738 U/mL with pUC980-2-*pelN*, which was nearly 3 times than that of pWB980. It also was the highest expression level in *Bacillus*^[26, 27]. It is worth noting that the plasmids pUC980-3, pUC980-4, pUC980-6 produced activities lower than 200 U/mL. The lowest PelN activity was produced by pUC980-7 (4 U/mL) that was nearly the same to the background of the host cells.

In addition, comparative analysis of these expressions were carried through SDS-PAGE as shown in Fig 4B. The SDS-PAGE analysis were sustained and correlated well with the PleN activities. The proportions of PelN protein in the supernatants at 48 h culturing with plasmids pWB980, pWB980-DB, pUC980-1 to pUC980-7 were estimated as 48%, 63%, 78%, 80%, 4%, 12%, 65%, 5% and 2% correspondingly (Fig 4B).

Alkaline protease expression in *B. subtilis* WB600

As stated in the expression of *peIN*, plasmids pUC980-1 and pUC980-2 have shown their accomplishments during foreign expression. They were further used to express alkaline protease gene *spro1* in *B. subtilis* WB600, employing pWB980 and pWB980-DB as controls. After culturing in shake-flask for 48 h, the extracellular activities of protease expressed with pUC980-1-*spro1* and pUC980-2-*spro1* were 4590 U/mL and 4818 U/mL (Fig 5A), which were 2.3 times and 2.4 times than that with pWB980-*spro1* (1991 U/mL), respectively. The proportions of the Spro1 in the supernatants were 72%, 77%, 92% and 95% in which expressed with plasmids pWB980, pWB980-DB, pUC980-1 and pUC980-2, respectively (Fig 5B).

Pullulanase expression in *B. subtilis* WB600

Meanwhile, plasmids pUC980-1 and pUC980-2 were used to express a pullulanase gene *pulA11* in WB600 in shake flask culturing for 48 h. Enzyme activities of plasmids pUC980-1-*pulA11* and pUC980-2-*pulA11* in the supernatants were 173 U/mL and 187 U/mL respectively, which were 1.9 times and twice than pWB980 (92 U/mL) correspondingly as stated in Fig 6A. The percentages of PulA11 proteins in the supernatants after 48 h fermentation of the recombinant strains containing plasmids pWB980-*pulA11*, pWB980-DB-*pulA11*, pUC980-1-*pulA11*, and pUC980-2-*pulA11* were 6%, 8%, 27% and 30%, respectively (Fig 6B).

Scale up fermentation of the optimum recombinant strains in a 10 L fermenter

The above results indicated that the recombinant strains containing pUC980-2-*peIN*, pUC980-2-*spro1*, and pUC980-2- *pulA11* showed the highest expression level in shake flask culturing basing on the remarkable properties of pUC980-2. In order to verify the stable production capacity of these strains in scale up fermentation, we performed batch-feed fermentations in a 10 L fermenter. The highest APL activity reached to 5200 U/mL after 54 h fermentation of the recombinant strain *B. subtilis* WB600 (pUC980-2-*peIN*). The curves for cell-growth and wet weight were shown in Fig 7A. The cells grown into the steady stage at 30 h with OD_{600nm} in 34.1 and 66.3 g/L of wet cell weight. The fermentation of this strain in a 10 L fermenter were repeat 20 times and the activities fluctuated little as shown in Fig 7B. Furthermore, the highest production of Spro1 was 21537 U/mL at 60 h and the highest production of pullulanase was 504 U/mL at 48 h (Fig S1). So far, the extracellular yields of PelN and Spro1 by the recombinant strains *B. subtilis* WB600 (pUC980-2-*peIN*) and *B. subtilis* WB600 (pUC980-2-*spro1*) showed the highest level in *Bacillus subtilis*^[28, 29]

Discussion

The properties of pUB110-derived plasmid pWB980 were regulated by the replicase encoding gene rep and membrane-binding region BA3-1 according to previous studies [30–33]. The rep gene, encoding the replicase protein, was fully remained to keep the plasmid function during construction of pWB980 [17, 34–36] and the BA3-1 region was related with plasmid divisions in daughter cells [30, 37–39]. It has long been suspected that plasmid replication origin ori might be important for plasmids replications and divisions [40, 41]. On these basis, the rep gene and the membrane-binding region have been manipulated with ori insertion separately.

Before ori insertions, the plasmid size was reduced for easier manipulation. The occasionally used resistance gene bleoR was deleted from pWB980, resulting plasmid pWB980-DB with a few higher copy numbers than pWB980 (Fig. 3A). The improvement of plasmid copying might be due to the hamper elimination of the bleoR gene to the initial signal in BA3-1 region for lagging strand synthesis. Using pWB980-DB as the original plasmids, ori was inserted into any possible positions upstream and downstream the rep gene and the BA3-1 region.

As for shuttle plasmids derived from pWB980-DB, ori insertions at certain sites influenced the properties of each shuttle plasmid differently. Firstly, all ori insertions flanking the BA3-1 improved the plasmids copies in host cells concluded from the enhancements of copy numbers of plasmids pUC980-1, pUC980-2, pUC980-3 and pUC980-4. Moreover, insertions upstream BA3-1 effected the plasmid-segregations less than down-stream ones. While insertions downstream BA3-1 depressed the plasmid-stabilities sharply. These depressions strongly suggested the importance of this area to the distribution of plasmids in the cell division. In addition, ori insertions upstream the rep gene in plasmids pUC980-5 and pUC980-6 were conducive to plasmid copying. While there was a direction bias in plasmid segregations as the forward insertion was largely more stable than the reverse one. Furthermore, at the position downstream the rep gene, the obstructions of ori insertions to plasmid properties were presented by the resulted plasmids pUC980-7 and pUC980-8 with poor performances in copy number and stability.

All plasmids (pUC980-1 ~ pUC980-7) kept structural stabilities in *B. subtilis*. They were systematically classified based on their copy numbers and segregation stabilities and further used to produce heterogenous proteins in *B. subtilis* WB600. Plasmids with good segregation stabilities dramatically outperformed the unstable ones, stated by the disparity between S1 plasmids and others. Moreover, the production of heterogenous proteins increased along with the copy numbers when the plasmids had similar stabilities. This conclusion was approved by the gradually improved expressions along the copy numbers of plasmids in group S1. And it was also supported by the plasmids in group S3. As for plasmids with similar copies, improvements in segregation stabilities enhanced the heterogenous productions of heterogenous proteins notably, corroborated by plasmids in every segregation stability levels.

Plasmids pUC980-1 and pUC980-2 performed well in expressing the tested heterogenous proteins, especially plasmid pUC980-2, with which the highest productions of alkaline pectate lyase PelN and alkaline protease Spro1 were obtained in *B. subtilis*. The high producing levels of foreign proteins have

proved the high efficiency and universality of these two plasmids and stated that they were capable of expressing many other proteins. Furthermore, the practicability of pUC980-2 in industrial applications were strongly illustrated by the stable expressions of heterogenous proteins with pUC980-2 in larger scale fermentations.

As for the relatively lower expressions of PulA11 than the other two proteins, the result indicated that the foreign gene itself also make important effect in the final expression level. Meanwhile, the expression levels of protein PulA11 with plasmid pUC980-2 in *B. subtilis* was still promoted to a certain extent compared with our previous studies in *E. coli* expression system^[42–44]. These improvements further confirmed the availability of pUC980-2 in production of heterogenous proteins.

Conclusion

In this work, some novel rules about the constructions of *E. coli*-*B. subtilis* shuttle plasmids based on pWB980 were concluded: 1) The deletion of bleoR gene improved the plasmid copying of pWB980; 2) The ori insertion was generally beneficial to plasmids copying in *B. subtilis*, while its impacts on plasmids segregation stabilities differed with insertion sites; 3) The site upstream the membrane binding region BA3-1 was the optimum position for plasmids manipulations with improvements both in copy number and stability. Based on these construction rules, two plasmids pUC980-1 and pUC980-2 with high copy numbers and high stabilities were selected out and performed well in tested protein productions. The highest production levels in *B. subtilis* of pectate lyase PelN and alkaline protease Spro1 were achieved by using plasmid pUC980-2 as expressing vector through 10 L fermentations. Plasmid pUC980-2 was proved to be capable and valuable in further industrial productions by these promising results.

Methods

Bacterial strains, plasmids and materials

The bacterial strains and plasmids used in this study are listed in Table 1. *E. coli* DH5α was used in plasmids construction and amplification. *B. subtilis* 168 were used in the determinations of plasmids properties. *B. subtilis* WB600 was employed in gene expressions. The sequences of the alkaline pectate lyase from *Paenibacillus* sp. 0602 (*pelN*, GenBank: KC351190.1), the alkaline protease gene *spro1* from alkaliphilic *Bacillus* sp. 221 (*spro1*, Sequence ID: D13157.1) and the pullulanase gene *pulA11* from *Anoxybacillus* sp. LM18-11 (GeneBank ID: HQ844266.1) were deposited in NCBI. The restriction endonucleases and DNA polymerase were commercially supplied by Thermo Fisher Scientific Co., Ltd. All other enzymes, chemicals and reagents were purchased from TaKaRa Biotechnology (Dalian, China) Co., Ltd.

Cultivation conditions

All cells were routinely grown at 37 °C and 200 rpm in Luria-Bertani (LB) medium or fermentation medium. LB medium (1 L) consisted of tryptone 10 g, yeast extract 5 g and NaCl 5 g. 1 L of fermentation

medium consisted of tryptone 16 g, soluble starch 37 g, CaCl_2 1.5 g, NaCl 2 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.2 g, KH_2PO_4 1 g, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.05 g and $(\text{NH}_4)_2\text{SO}_4$ 1.5 g. Different antibiotics (50 $\mu\text{g}/\text{mL}$ kanamycin and 25 $\mu\text{g}/\text{mL}$ kanamycin) were added in the medium for the relevant recombinant strains.

B. subtilis WB600 strains containing recombinant plasmids were cultured in Shake flask (SHUNIU, GG-17, Sichuan SHUBO Co., LTD, China) or in a 10-L fermenter (5BG, bxbio, China) to produce foreign proteins. The shake-flask culturing was performed as followed. A 5 mL aliquot of LB medium supplemented with 25 $\mu\text{g}/\text{mL}$ kanamycin was inoculated with a frozen glycerol stock of recombinant strain (20 μL), and then incubated for up to 14 h at 37 °C and 200 rpm in a rotary shaker (ZQWY-200G, Shanghai Zhichu Instruments Co., Ltd., China). An aliquot of this preculture (0.5 mL) was transferred into a 250 mL shake-flask that was loaded with 50 mL of LB medium supplemented with 25 $\mu\text{g}/\text{mL}$ kanamycin, which was then shaken in the rotary shaker (200rpm) at 37 °C. After culturing for 48 h, samples of each culture were collected and analyzed for enzyme activities. The culture was harvested by centrifugation at 12,000 $\times g$ for 10 min at 4 °C to obtain the culture supernatant. Enzyme activities in the supernatant were determined with corresponding assays. Then the culture supernatant was analyzed with SDS-PAGE. The percentages of the produced extracellular proteins were calculated by a software Gel-Pro analyzer 4.0 (Media Cybernetics, CA, USA).

Fermentation of the recombinant *B. subtilis* WB600 strain was performed in a 10-L fermenter. The seed culture was prepared as described in the above section and then used to inoculate an initial batch of fermentation medium supplemented with 25 $\mu\text{g}/\text{mL}$ kanamycin. The biomass and enzyme activities were analyzed at designated time intervals. The cell density was determined by measuring the $\text{OD}_{600\text{nm}}$ with a UV-1800/PC spectrophotometer (Epoch 2, BioTek, USA). To determine the wet cell weight (WCW), a 2-mL sample of the culture broth were centrifuged at 12,000 $\times g$ for 10 min at 4 °C. The resulting pellet was collected and weighed. The enzyme activities in the culture supernatant were further determined with corresponding methods. The whole fermentation process was stopped and finished when continuous reductions of biomass and enzyme activities were observed.

Shuttle plasmids construction

Construction of shuttle plasmids is schematically presented in Fig.1. All plasmids were constructed through the method of MEGAWHOP^[45]. All of the specific primers used for PCR amplification were synthesized by the Beijing Genomics Institute (BGI) and listed in Table 2 and the isolation and manipulation of recombinant DNA were performed according to the previously published protocol^[46]. 1.5 mL of culture was poured into a 2 mL EP tube and centrifuged at 12,000 $\times g$ for 10 min at 4 °C. The supernatant was discarded and the bacterial pellet was collected. The bacterial samples were treated with solution I (100 mL of solution I consisted of 1 M pH 8.0 Tris-HCl, 2.5 mL; 0.5 M EDTA 2 mL; ddH₂O 91 mL; 20% sterile glucose 4.5 mL), solution II (100 mL of solution II consisted of 10% SDS 50 mL; 2 M NaOH 50 mL), solution III (500 mL of solution III consisted of KAc 147 g; HAc 57.5 mL, ddH₂O was added to 500 mL), mixture of phenol: chloroform (volume ration 1:1) and 70% ethanol successively according to

the protocol. The DNA sample was dissolved with 80 μ L ddH₂O and stored in -20 °C for further experiments.

See Table 2

According to the sequencing map, 169 bp of the membrane-binding region BA3 (515 bp) originated from pUB110 was deleted during the construction of pWB980. In order to be mentioned conveniently, the remaining 346 bp fragment of BA3 was named BA3-1 in this work as shown in Figure 1. pWB980 is consist of five parts: BA3-1, P43 promoter cohered with a signal-peptide sequence, the replicase-coding gene *rep* in *B. subtilis*, the kanamycin-resistance gene *kanR* and the bleomycin gene *bleoR* gene (Fig 1). In this work, the *bleoR* was deleted with primers DB-1/DB-2. The plasmid replication origin *ori* (888 bp) from pUC19 was inserted at the four sites up- and down-stream BA3-1 region and *rep* gene in pWB980-DB (3388 bp) separately. With primer pairs P1-S/P1-A and P2-S/P2-A, *ori* integrated into the site (659 bp) upstream the BA3-1 with forward and reverse directions. The recombinant plasmids were named as pUC980-1 and pUC980-2 respectively. At the site (169 bp) downstream the BA3-1, primers P3-S/P3-A and P4-S/P4-A were used in *ori*-insertions, resulting plasmids pUC980-3 and pUC980-4 in forward and reverse directions. Upstream the *rep* gene, the *ori* was inserted at the 3118 site using primers P5-S/P5-A and P6-S/P6-A. The recombinant plasmids were named as pUC980-5 and pUC980-6 with forward and reverse *ori*-insertions respectively. *Ori* was put into the 1722 site downstream the *rep* by using primers P7-S/P7-A and P8-S/P8-A. The plasmid with forward *ori* insertion was pUC980-7 and the other one was pUC980-8.

With primer pair 62-S/62-A, the alkaline pectate lyase gene *pelN* was inserted into the sites after the P43 promoter of plasmids pWB980, pWB980-DB and all pUC980-serial plasmids except pUC980-8. The recombinant plasmids were pWB980-*pelN*, pWB980-DB-*pelN*, pUC980-1-*pelN*, pUC980-2-*pelN*, pUC980-3-*pelN*, pUC980-4-*pelN*, pUC980-5-*pelN*, pUC980-6-*pelN* and pUC980-7-*pelN* correspondently.

Determination of plasmid copy number by quantitative real time PCR (qPCR)

The quantitative real-time PCR (qPCR) was performed using iQ™ SYBR® Green Supermix (Bio-Rad) as described by Turgeon^[47]. The *dnaN* and *kanR* genes were chosen as a single copy reference gene on the *B. subtilis* chromosome and the test gene in plasmids correspondingly^[48-50]. Specific primer pairs dnaNS/dnaNA and kanS/kanA (Table 2) were designed to generate products of approximately 150 bp. The standard curves were prepared based on the pMD-18-T vector. The PCR mixtures (total volume 20 μ L) contained 10 μ L 2xTakara SYBR Green Real-Time PCR Master Mix, 1.5 μ L forward and reverse primers (10 μ M) and 1 μ L diluted (10^{-1} to 10^{-4}) sample DNA. The reaction condition was as followed: 95 °C for 10 min, 45 cycles of 95 °C for 10 s, 62 °C for 20 s, and 72 °C for 30 s, followed by a gradient temperature from 55 °C to 95 °C. The calculated mean of the *kanR* gene in each compared plasmid was divided by the *dnaN* gene (single-copy reference), resulting the copy number. All experiments were performed with three independent biological replicates.

Plasmid stability assay

Plasmid segregational stability in *B. subtilis* 168 was detected according to the method of Bron and Luxen^[38] with minor modifications. Single colonies on kanamycin selective (25 µg/mL) LB agar plate were used to inoculate 5 mL kanamycin selective (25µg/mL) LB medium. After 24 h culturing at 37 °C, the cultures were diluted 1:1000 in fresh LB media without antibiotics and incubated at 37 °C. This iterative subculturing process was repeated every 24 h for 30 consecutive days. Samples taken at appropriate intervals were checked for the fraction of plasmid-containing cells by replica plating on LB agar plates containing 25µg/mL kanamycin.

N: Total amount of colonies on the selectable plate

Plasmids were extracted from host cells after 30-days continuously culturing and detected the structural stability by *EcoRI/HindIII* enzyme-digestion verifications.

Alkaline pectate lyase assay

The pectate lyase activity was measured using the method described previously by Wang et al.^[51] 20µL of the diluted enzyme solutions were added to 400µL of 0.2% pectin in 50 mmol/L glycine-NaOH buffer (with 50µmol/L CaCl₂) at pH 9.0. The reaction mixture was incubated at 65 °C for 10 min, and the reaction was terminated by adding 580µL of 30 mmol/L H₃PO₄. The product yields were detected using a spectrophotometer at 235 nm (Epoch 2, BioTek, USA). One standard enzyme unit was defined as the yield of 1µmol unsaturated pectin per minute under the above-mentioned conditions.

Alkaline protease assay

The alkaline protease activity was measured using the modified method described by Tjalsma et al.^[27] One standard enzyme unit was defined as the amount of enzyme that hydrolyzes 1 µg casein minute under the situation of pH 9.0, 50°C. 250µL of the diluted enzyme solution were mixed with 250 µL solution containing 2% casein in pH 9.0 buffer. The mixture was incubated at 50 °C for 10 min, and the reaction was terminated by adding 500µL 0.4 M TCA. The productions were detected using a spectrophotometer (Epoch 2, BioTek, USA) at 280 nm. Casein with concentrations from 0 to 500 µg/mL were used as standard.

Pullulanase assay

Pullulanase activity was measured using the modified dinitrosalicylic acid (DNS) method^[52]. One standard enzyme unit was defined as the amount of enzyme that releases 1 µmol reducing sugar per minute under the assay conditions. Glucose with concentrations from 0 to 400 mg/mL were used as standard. 50 µL of the diluted enzyme solutions were added into the 450 µL reaction mixture containing 5% pullulan solutions and the pH6.0 buffer at the volume ratio of 1:8. After incubations at pH 6.0, 60°C for 30 min, 500 µL DNS were used to stop the reactions. The productions were detected using a spectrophotometer (Epoch 2, BioTek, USA) at 540 nm. The amount of reducing sugar released during incubations were tested as the enzyme-index.

SDS-PAGE

B. subtilis WB600 transformant cells harboring recombinant plasmids were cultured in LB medium containing 25 µg/ml kanamycin at 37 °C for 48 h. The cultures (0.5 mL respectively) were centrifuged at 5000 rpm, 4 °C for 10 min and the supernatant solutions were reserved. The cell pellets were re-suspended in 0.5 mL ddH₂O, followed by sonication to release intracellular content. Extracellular proteins were precipitated with 13% (w/v) TCA for 30 min on ice, then the precipitates which were collected by 10 min centrifugation at 4 °C and 13,000×g, were carefully washed with ice-cold acetone and dried under vacuum. The collective proteins were dissolved in 20µL urea (6M) and mixed with appropriate volume of 4× SDS loading buffer. Samples were then heated in boiling water bath for 10 min and centrifuged at 13,000×g for 10 s. 12 µL of the samples were loaded onto 10% SDS-PAGE and run at 20 mA for around 1h. Gel was stained using Coomassie brilliant blue dye.

Declarations

Ethics approval and consent to participate

This article does not contain any studies with human participants or animals performed by any of the authors.

Availability of data and material

We declared that materials described in the manuscript, including all relevant raw data, will be freely available to any scientist wishing to use them for non-commercial purposes, without breaching participant confidentiality.

Competing interests

The authors declare that they have no conflict of interest.

Funding

This work was supported by the National Natural Science Fund of China (Grant 31701534), the Tianjin outstanding talent training program, the Tianjin Science & Technology Planning Project (Grant 14ZCZDSY00157 and 15PTCYSY00020) and Yantai Marine economy innovation development demonstration project (Grant YHCX-SW-L-201703).

Authors' contributions

XYZ and JYX designed the experiments together. XYZ led the performance of the experiments, analysis of the data and writing of the paper. MTJJZ, WJS and SBY participated in experiments and analysis. HCZ, HS and YHM participated in editing the paper. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

Abbreviations

SDS-PAGE: sodium dodecyl sulfate-polyacrylamide gel electrophoresis; BGI: the Beijing Genomics Institute; qPCR: the quantitative real-time PCR.

References

1. Dong H, Zhang D: Current development in genetic engineering strategies of *Bacillus* species. *Microbial Cell Factories* 2014, 13:63.
2. Zhang K, Su L, Duan X, Liu L, Wu J: High-level extracellular protein production in *Bacillus subtilis* using an optimized dual-promoter expression system. *Microbial Cell Factories* 2017, 16:32.
3. Ji S, Li W, Rasheed Baloch A, Wang M, Li H, Cao B, Zhang H: Efficient biosynthesis of a Cecropin A-melittin mutant in *Bacillus subtilis* WB700. *Scientific Reports* 2017, 7:40587.
4. Marcus S, Ajay S, Ward OP: Developments in the use of *Bacillus* species for industrial production. *Canadian Journal of Microbiology* 2004, 50:1.
5. Hallberg ZF, Su Y, Kitto RZ, Ming CH: Engineering and in vivo applications of riboswitches. *Annual Review of Biochemistry* 2017, 86:515-539.
6. Willenbacher J, Mohr T, Henkel M, Gebhard S, Mascher T, Syltatk C, Hausmann R: Substitution of the native *srfA* promoter by constitutive *P_{veg}* in two *B. subtilis* strains and evaluation of the effect on Surfactin production. *Journal of Biotechnology* 2016, 224:14-17.
7. Maeda T, Sanchez-Torres V, Wood TK: Hydrogen production by recombinant *Escherichia coli* strains. *Microbial Biotechnology* 2012, 5:214-225.
8. Lu Z, Yang S, Yuan X, Shi Y, Ouyang L, Jiang S, Yi L, Zhang G: CRISPR-assisted multi-dimensional regulation for fine-tuning gene expression in *Bacillus subtilis*. *Nucleic Acids Research* 2019, 47:e40-e40.
9. Iannelli F, Santagati M, Santoro F, Oggioni MR, Stefani S, Pozzi G: Nucleotide sequence of conjugative prophage Φ 1207.3 (formerly Tn1207.3) carrying the *mef(A)/msr(D)* genes for efflux resistance to macrolides in *Streptococcus pyogenes*. *Frontiers in Microbiology* 2014, 5:687.
10. Xu H, Xiao T, Chen CH, Li W, Liu SX: Sequence determinants of improved CRISPR sgRNA design. *Genome Research* 2015, 25:1147-1157.
11. Gros MF, Riele H, Te, Ehrlich SD: Rolling circle replication of single-stranded DNA plasmid pC194. *the EMBO Journal* 1987, 6:3863-3869.
12. Pluta R, Espinosa M: Antisense and yet sensitive: copy number control of rolling circle-replicating plasmids by small RNAs. *WIREs RNA* 2018, 9:e1500.
13. Liu M, M Kwong S, Jensen S, Brzoska A, Firth N: Biology of the staphylococcal conjugative multiresistance plasmid pSK41. *Plasmid* 2013, 70:42-51.

14. Yihan L, Fuping L, Guanqun C, Snyder CL, Jing S, Yu L, Jianling W, Jing X: High-level expression, purification and characterization of a recombinant medium-temperature alpha-amylase from *Bacillus subtilis*. *Biotechnology Letters* 2010, 32:119-124.
15. Nguyen HD, Nguyen QA, Ferreira RC, Ferreira LC, Tran LT, Schumann W: Construction of plasmid-based expression vectors for *Bacillus subtilis* exhibiting full structural stability. *Plasmid* 2005, 54:241-248.
16. Phan TT, Nguyen HD, Schumann W: Novel plasmid-based expression vectors for intra- and extracellular production of recombinant proteins in *Bacillus subtilis*. *Protein Expression and Purification* 2006, 46:189-195.
17. Wu SC, Wong SL: Development of improved pUB110-based vectors for expression and secretion studies in *Bacillus subtilis*. *Journal of Biotechnology* 1999, 72:185-195.
18. Reinhard B, Sui-Lam W, Dieter J: Production of PHA depolymerase A (PhaZ5) from *Paucimonas lemoignei* in *Bacillus subtilis*. *FEMS Microbiology Letters* 2002, 209:237-241.
19. Lin B, Li Z, Zhang H, Wu J, Luo M: Cloning and expression of the γ -polyglutamic acid synthetase gene pgsBCA in *Bacillus subtilis* WB600. *Biomed Research International* 2016, 2016:1-7.
20. Sun TY, Shao JX, Gao YY, Chen LM, Gao ZQ, Lin WP, Fu-Ping LU: Establishment of high-efficiency transformation and expression of the gene encoding alkaline protease PB92 in *Bacillus subtilis* DB104. *Jiangsu Journal of Agricultural Sciences* 2009, 25:534-537.
21. Kramer MG, Espinosa M, ., Misra TK, Khan SA: Characterization of a single-strand origin, ssoU, required for broad host range replication of rolling-circle plasmids. *Molecular Microbiology* 2010, 33:466-475.
22. Gryczan TJ, Dubnau D, . Construction and properties of chimeric plasmids in *Bacillus subtilis*. *Proceedings of the National Academy of Sciences of the United States of America* 1978, 75:1428-1432.
23. Zhou Z, Liu Y, Chang Z, Wang H, Leier A, Marquez-Lago TT, Ma Y, Li J, Song J: Structure-based engineering of a pectate lyase with improved specific activity for ramie degumming. *Applied Microbiology & Biotechnology* 2016, 101:1-11.
24. van der Laan JC, Gerritse G, Mulleners LJ, van der Hoek RA, Quax WJ: Cloning, characterization, and multiple chromosomal integration of a *Bacillus* alkaline protease gene. *Applied and environmental microbiology* 1991, 57:901-909.
25. Xu J, Ren F, Huang CH, Zheng Y, Zhen J, Sun H, Ko TP, He M, Chen CC, Chan HC, et al: Functional and structural studies of pullulanase from *Anoxybacillus* sp. LM18-11. *Proteins* 2014, 82:1685-1693.
26. Zhou M, Wu J, Wang T, Gao L, Yin H, Lü X: The purification and characterization of a novel alkali-stable pectate lyase produced by *Bacillus subtilis* PB1. *World Journal of Microbiology and Biotechnology* 2017, 33:190.
27. Tjalsma H, Koetje EJ, Kiewiet R, Kuipers OP, Kolkman M, Laan J, Daskin R, Ferrari E, Bron S: Engineering of quorum-sensing systems for improved production of alkaline protease by *Bacillus subtilis*. *Journal of Applied Microbiology* 2004, 96:569-578.

28. Feto NA: *Bacillus spp. and their biotechnological roles in green industry*.2016.
29. Liu L, Liu Y, Shin HD, Chen RR, Wang NS, Li J, Du G, Chen J: Developing *Bacillus* spp. as a cell factory for production of microbial enzymes and industrially important biochemicals in the context of systems and synthetic biology. *Applied Microbiology and Biotechnology* 2013, 97:6113-6127.
30. Bron S, Luxen E, Swart P: Instability of recombinant pUB110 plasmids in *Bacillus subtilis*: Plasmid-encoded stability function and effects of DNA inserts. *Plasmid* 1988, 19:231-241.
31. Maciag IE, Viret JF, Alonso JC: Replication and incompatibility properties of plasmid pUB110 in *Bacillus subtilis*. *Molecular & General Genetics* 1988, 212:232-240.
32. Tanaka T, Sueoka N: Site-specific in vitro binding of plasmid pUB110 to *Bacillus subtilis* membrane fraction. *Journal of Bacteriology* 1983, 154:1184-1194.
33. Viret JF, Alonso JC: A DNA sequence outside the pUB110 minimal replicon is required for normal replication in *Bacillus subtilis*. *Nucleic Acids Research* 1988, 16:4389-4406.
34. Thomas CD, Jennings LJ: RepD/D*: a protein-DNA adduct arising during plasmid replication. *Biochemical Society Transactions* 1995, 23:442S.
35. Alonso JC, Viret JF, Tailor RH: Plasmid maintenance in *Bacillus subtilis* recombination-deficient mutants. *Molecular Genetics and Genomics* 1987, 208:349-352.
36. Noirot-Gros MF, Ehrlich SD: Change of a catalytic reaction carried out by a DNA replication protein. *Science* 1996, 274:777-780.
37. Alonso JC, Stiege CA, Tailor RH, Viret JF: Functional analysis of the dna (Ts) mutants of *Bacillus subtilis* : Plasmid pUB110 replication as a model system. *Molecular & General Genetics* 1988, 214:482-489.
38. Bron S, Luxen E: Segregational instability of pUB110-derived recombinant plasmids in *Bacillus subtilis*. *Plasmid* 1985, 14:235-244.
39. Wassenaar TM, Cabal A: The mobile dso-gene-sso element in rolling-circle plasmids of staphylococci reflects the evolutionary history of its resistance gene. *letters in applied microbiology* 2017, 65:192-198.
40. Costa A, Hood IV, Berger JM: Mechanisms for initiating cellular DNA replication. *Annual Review of Biochemistry* 2013, 82:25-54.
41. Bleichert F, Botchan MR: Mechanisms for initiating cellular DNA replication. *Science* 2017, 355.
42. Li SF, Xu JY, Bao YJ, Zheng HC, Song H: Structure and sequence analysis-based engineering of pullulanase from *Anoxybacillus* sp. LM18-11 for improved thermostability. *Journal of Biotechnology* 2015, 210:8-14.
43. Zeng Y, Zheng H, Shen Y, Xu J, Tan M, Liu F, Song H: Identification and analysis of binding residues in the CBM68 of pullulanase Pula from *Anoxybacillus* sp. LM18-11. *Journal of Bioscience and Bioengineering* 2019, 127:8-15.
44. Zeng Y, Xu J, Fu X, Tan M, Liu F, Zheng H, Song H: Effects of different carbohydrate-binding modules on the enzymatic properties of pullulanase. *International Journal of Biological Macromolecules*

- 2019, 137:973-981.
45. Miyazaki K: MEGAWHOP cloning: a method of creating random mutagenesis libraries via megaprimer PCR of whole plasmids. *Methods in Enzymology* 2011, 498:399-406.
 46. Green MR, Sambrook J: Molecular Cloning: a laboratory manual (fourth edition): three-volume set. *Cold Spring Harbor Laboratory Press* 2012, 1:part1.
 47. Turgeon N, Laflamme C, Ho J, Duchaine C: Evaluation of the plasmid copy number in *B. cereus* spores, during germination, bacterial growth and sporulation using real-time PCR. *Plasmid* 2008, 60:118-124.
 48. Friehs K: Plasmid copy number and plasmid stability. *Advances in Biochemical Engineering Biotechnology* 2004, 86:47.
 49. Maidak BL, Larsen N, Mccaughey MJ, Overbeek R, Olsen GJ, Fogel K, Blandy J, Woese CR: The ribosomal database project. *Nucleic Acids Research* 1994, 22:3485-3487.
 50. Providenti MA, O'Brien JM, Ewing RJ, Paterson ES, Smith ML: The copy-number of plasmids and other genetic elements can be determined by SYBR-Green-based quantitative real-time PCR. *Journal of Microbiological Methods* 2006, 65:476-487.
 51. Wang H, Li X, Ma Y, Song J: Process optimization of high-level extracellular production of alkaline pectate lyase in recombinant *Escherichia coli* BL21 (DE3). *Biochemical Engineering Journal* 2015, 93:38-46.
 52. He XS, Shyu YT, Nathoo S, Wong SL, Doi RH: Construction and use of a *Bacillus subtilis* mutant deficient in multiple protease genes for the expression of eukaryotic genes. *Annals of the New York Academy of Sciences* 2010, 646:69-77.

Tables

Table 1 Strains and plasmids used in this study

strains and plasmids	characteristics	reference
strains		
<i>E. coli</i> DH5α	F-, φ80dlacZΔM15, Δ(<i>lacZYA-argF</i>)U169, <i>deoR</i> , <i>recA1</i> , <i>endA1</i> , <i>hsdR17</i> (rk-,mk+), <i>phoA</i> , <i>supE44</i> , λ-, <i>thi-1</i> , <i>gyrA96</i> , <i>relA1</i>	Takara
<i>B. subtilis</i> 168 ^a	<i>trpC2</i>	ATCC
<i>B. subtilis</i> WB600	<i>apr</i> , <i>nprA</i> , <i>epr</i> , <i>bpf</i> , <i>mpr</i> , <i>nprB</i> , <i>trpC2</i>	Our lab. preservation
plasmids		
pMD-18-T	Standard vector for plasmid copy number determination, Amp ^r	Takara
pUB110	Kan ^r , <i>mob</i> region, membrane biding region BA3	Our lab. preservation
pWB980	pUB110 derivative, destroyed <i>mob</i> region, Kan ^r , Bleo ^r , P43 promoter, part BA3 region(BA3-1)	Our lab. Preservation
pWB980-DB	pWB980 derivate, <i>bleoR</i> was deleted, Kan ^r , P43 promoter, BA3-1 region	this work
pUC980-1	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between BA3-1 region and <i>kanR</i> gene in homodromous direction	this work
pUC980-2	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between BA3-1 region and <i>kanR</i> gene in reversed direction	this work
pUC980-3	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> region was inserted between BA3-1 region and P43 promoter in forward direction	this work
pUC980-4	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> region was inserted between BA3-1 region and P43 promoter in reversed direction	this work
pUC980-5	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between P43 promoter and <i>rep</i> in homodromous direction	this work
pUC980-6	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between P43 promoter and <i>rep</i> in reverse direction	this work
pUC980-7	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between <i>rep</i> and <i>kanR</i> in forward direction	this work
pUC980-8	<i>B.subtilis-E.coli</i> shuttle expression vector, Kan ^r , P43 promoter, <i>ori</i> was inserted between <i>rep</i> and <i>kanR</i> in reverse direction	this work
pWB980- <i>peIN</i>	Kan ^r , Bleo ^r , P43 promoter, signal peptide <i>sacB</i> , control plasmid in Alkaline pectate lyase <i>peIN</i> expression in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work
pWB980-DB- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C2 and segregation stability of S1	this work
pUC980-1- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , with both high copy number and segregation stability, <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C1 and segregation stability of S1	this work

pUC980-2- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , with both high copy number and segregation stability, <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work
pUC980-3- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C2 and segregation stability in S3	this work
pUC980-4- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C2 and segregation stability in S2	this work
pUC980-5- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , control plasmid with low copy number and high segregation stability, <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C2 and segregation stability in S1	this work
pUC980-6- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , control plasmid with high copy number and low segregation stability, <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C1 and segregation stability in S3	this work
pUC980-7- <i>peIN</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , control plasmid with high copy number and low segregation stability, <i>peIN</i> expression plasmid in <i>B. subtilis</i> with copy number in C3 and segregation stability in S3	this work
pWB980- <i>spro1</i>	Kan ^r , Bleo ^r , P43 promoter, control plasmid in alkaline protease <i>spro1</i> expression in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work
pWB980-DB- <i>spro1</i>	Kan ^r , P43 promoter, control plasmid in alkaline protease <i>spro1</i> expression in <i>B. subtilis</i> with copy number in C2 and segregation stability of S1	this work
pUC980-1- <i>spro1</i>	Kan ^r , P43 promoter, with both high copy number and segregation stability, <i>spro1</i> expression plasmid in <i>B. subtilis</i> with copy number in C1 and segregation stability of S1	this work
pUC980-2- <i>spro1</i>	Kan ^r , P43 promoter, with both high copy number and segregation stability, <i>spro1</i> expression plasmid in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work
pWB980- <i>pulA11</i>	Kan ^r , Bleo ^r , P43 promoter, signal peptide <i>sacB</i> , control plasmid in pullulanase gene <i>pulA11</i> expression in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work
pWB980-DB- <i>pulA11</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , <i>pulA11</i> expression plasmid in <i>B. subtilis</i> with copy number in C2 and segregation stability of S1	this work
pUC980-1- <i>pulA11</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , with both high copy number and segregation stability, <i>pulA11</i> expression plasmid in <i>B. subtilis</i> with copy number in C1 and segregation stability of S1	this work
pUC980-2- <i>pulA11</i>	Kan ^r , P43 promoter, signal peptide <i>sacB</i> , with both high copy number and segregation stability, <i>pulA11</i> expression plasmid in <i>B. subtilis</i> with copy number in C3 and segregation stability of S1	this work

^a*Bacillus subtilis* 168 (ATCC23857) was provided by the American Type Culture Collection (ATCC).

Table 2 Primers used in this study

primers	sequence(5'-3')
DB-1	AATCTATTATTAATCTGTTCAGCAATC
DB-2	TCTCACGCATAAAATCCCC
P1-S	GGGGATTTTATGCGTGAGACTGTCAGACCAAGTTTACTCAT
P1-A	GATTGCTGAACAGATTAATAATAGATTAGCGGTATCAGCTCACTC
P2-S	GATTGCTGAACAGATTAATAATAGATTCTGTCAGACCAAGTTTACTCAT
P2-A	GGGGATTTTATGCGTGAGAAGCGGTATCAGCTCACTC
P3-S	ATTGAACATACGGTTGATTTAATAACTGACTGTCAGACCAAGTTTACTCAT
P3-A	GCTTTAGCAAGAGGGTGATGTTTGAGCGGTATCAGCTCACTC
P4-S	GCTTTAGCAAGAGGGTGATGTTTGCTGTCAGACCAAGTTTACTCAT
P4-A	GAACATACGGTTGATTTAATAACTGAAGCGGTATCAGCTCACTC
P5-S	CAAGAAAAACACGATTTAGAACCTGTCAGACCAAGTTTACTCAT
P5-A	GAGTTAGTTCAAATTCGTTCTTTTTAAGCGGTATCAGCTCACTC
P6-S	GAGTTAGTTCAAATTCGTTCTTTTTACTGTCAGACCAAGTTTACTCAT
P6-A	CAAGAAAAACACGATTTAGAACCGCGGTATCAGCTCACTC
P7-S	TGCTTAGGAAGACGAGTTATTAATACTGTCAGACCAAGTTTACTCAT
P7-A	GAATATTTGGAGAGCACCGTTCTTATTCAGCAGCGGTATCAGCTCACTC
P8-S	GAATATTTGGAGAGCACCGTTCTTATTCAGCCTGTCAGACCAAGTTTACTCAT
P8-A	TGCTTAGGAAGACGAGTTATTAATAAGCGGTATCAGCTCACTC
62-S	CATAAAAAAGGAGACATGAACGATGGCGGGCAATGCAGATTAC
62-A	CCCCGGGTACCGAGCTCGATTAATAGCTCGTCTTCAGCCAGTTGTCC
spro-S	GCGCAACTCAAGCTTTTGCCATGAAGCTTAAGAAACCGTTG
spro-A	CCCGGGTACCGAGCTCGATTAGCGTGTTGCCGCTTC
pulA-S	GAGGCGCAACTCAAGCTTTTGCCCCC AAAACAACAGTCGTTTGAAG
pulA-A	GATCCCCGGGTACCGAGCTCGATCAACATTGAATTAATACCCACGCAC
kanS	CGGATATTGAGATGATGTGTGTCATGTC
kanA	GACCATGTGTAAGCGGCCAATC
dnaNS	GCACTTGCCGCAGATTGA
dnaNA	AATGCAAGACGGTGGCTATC