Impacts of the COVID-19 lockdown on air quality and its association with human mortality trends in megapolis Mexico City

Gurusamy Kutralam-Muniasamy

Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional

Fermín Pérez-Guevara

Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional

Priyadarsi D. Roy

Instituto de Geología, Universidad Nacional Autónoma de México (UNAM)

I. Elizalde-Martínez

Instituto Politécnico Nacional (IPN), Centro Mexicano para la Producción más Limpia (CMP+L)

V.C. Shruti (✉ shrutifmd@gmail.com)

Instituto Politécnico Nacional (IPN), Centro Mexicano para la Producción más Limpia (CMP+L)

Research Article

Keywords: Air Pollution, NO₂, PM_{2.5}, O₃, Traffic, Temperature

Posted Date: October 12th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-90079/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Air Quality, Atmosphere & Health on October 28th, 2020. See the published version at <https://doi.org/10.1007/s11869-020-00960-1>.

[Click here to view linked References](#)

Impacts of the COVID-19 lockdown on air quality and its association with human mortality trends in megapolis Mexico City

Gurusamy Kutralam-Muniasamy¹, Fermín Pérez-Guevara^{1,2}, Priyadarsi D. Roy³, I. Elizalde-Martínez⁴, V.C. Shruti^{4*}

¹Department of Biotechnology and Bioengineering, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Ciudad de México, México

²Nanoscience & Nanotechnology Program, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Ciudad de México, México

³Instituto de Geología, Universidad Nacional Autónoma de México (UNAM), Ciudad Universitaria, Del. Coyoacán, C.P. 04510, Ciudad de México, Mexico

⁴Instituto Politécnico Nacional (IPN), Centro Mexicano para la Producción más Limpia (CMP+L), Av. Acueducto s/n, Col. Barrio la Laguna Ticomán, Del Gustavo A. Madero, C.P. 07340, México City, México

*Corresponding Author Mailing Address: Instituto Politécnico Nacional (IPN), Centro Mexicano para la Producción más Limpia (CMP+L), Av. Acueducto s/n, Col. Barrio la Laguna Ticomán, Del Gustavo A. Madero, C.P. 07340, México, D.F., México.

E-mail address: shrutifrnd@gmail.com Tel.: 57296000 Extn- 52612

Keywords: Air Pollution; NO₂; PM_{2.5}; O₃; Traffic; Temperature.

Abstract

Mexico City is the second most populated city in Latin America, and it went through two partial lockdowns between April 1 and May 31, 2020 for reducing the COVID-19 propagation. The present study assessed air quality and its association with human mortality rates during the lockdown by estimating changes observed in air pollutants (CO, NO₂, O₃, SO₂, PM₁₀ and PM_{2.5}) between the lockdown (April 1 - May 31) and pre-lockdown (January 1 – March 31) periods, as well as by comparing the air quality data of lockdown period with the same interval of previous five-years (2015-2019). Concentrations of NO₂ (-29%), SO₂ (-55%) and PM₁₀ (-11%) declined and the contents of CO (+1.1%), PM_{2.5} (+19%) and O₃ (+63%) increased during the lockdown compared to the pre-lockdown period. This study also estimated that NO₂, SO₂, CO, PM₁₀ and PM_{2.5} reduced by 19-36%, and O₃ enhanced by 14% compared to the average of 2015-2019. Reduction in traffic as well as less emission from vehicle exhausts led to remarkable decline in NO₂, SO₂ and PM₁₀. The significant positive associations of PM_{2.5}, CO and O₃ with the numbers of COVID-19 infections and deaths, however, underscored the necessity to enforce air pollution regulations to protect human health in one of the important cities of the northern hemisphere.

Introduction

Geographical setting (elevated basin surrounded by mountain ridges), climatological conditions and emission of higher amounts of pollutant have led to poor air quality of Mexico City, the second most populated city in Latin America (SMA-GDF 2016; Calderon-Garcidueñas et al. 2015). Different air pollutants are associated with anthropogenic activities like transportation, industry and construction (SMA-GDF 2016; Table 1) and the elevated concentrations of particulate matter (PM₁₀ and PM_{2.5}), carbon monoxide (CO) and nitrogen oxides (nitric oxide, NO_x and nitrogen dioxide, NO₂) during the spring-summer of every year give rise to implementation of the no circulation program (Benítez-García 2014; Molina et al. 2019). Similarly, the concentrations of ozone (O₃) remain more than the World Health Organization (WHO) recommended 100 microgram per cubic meter in about 79% of every year. Negative health effects due to air pollution also include specific effects associated with fine particles (Gold et al. 1999; Osornio-Vargas et al. 2003; Calderon-Garcidueñas et al. 2015). The government of Mexico City has been implementing the program of No-drive days since 1989 to restrict the circulation of 50% of privately-owned vehicles and reduce almost 20% of air pollutants for mitigating the deteriorating effects of air pollution on health (Nava Escudero 2008; Davis 2017). Even with the restrictions, the environmental contingency, because of higher levels of PM₁₀, PM_{2.5}, and O₃, has been regular in the months of April to June with additional air pollution coming through forest fire and the agricultural waste combustion in the surrounding states, as well as new vehicles introduced to the system every year (Molina et al. 2019).

The rapid spread of COVID-19 to more than 100 countries within three months of its discovery in China prompted the WHO to declare the outbreak as pandemic in March 2020 (WHO 2020). Recent reports from several countries like China (Shakoor et al. 2020; Gautam

2020), India (Agarwal et al. 2020; Panda et al. 2020; Sharma et al. 2020), Morocco (Otmani et al. 2020), Brazil (Nakada and Urban 2020), Saudi Arabia (Anil and Alagha 2020), and US (Pata 2020; Doğan et al. 2020) have assessed improvement in the air quality from an appreciable decline in many anthropogenic pollutants in response to restrictions imposed on traffic and industrial activity during the novel coronavirus (SARS-CoV-2) pandemic. In this context, we believe that the estimation of changes in air pollutants from Mexico City is necessary for a global comparison. The new data might even provide important clues for implementing new measures or modification of existing strategies for improving the air quality of one of the polluted cities in Latin America.

By August 2020, the COVID-19 death toll from Mexico was the third highest in the world, next to US and Brazil, with the highest confirmed cases (n=95185) and deaths (n=8301) in Mexico City among all the federal states (<https://coronavirus.gob.mx/datos/>). In Mexico, the first case of infection was registered on February 28, 2020 and subsequently, all 32 federal states of Mexico presented COVID-19 cases by March 17, 2020 (Mexico Health Ministry 2020). On March 31, the government of Mexico City announced Sanitary Emergency starting on April 1 as the number of cases exceeded one thousand by March 30, 2020 (Mexico Health Ministry 2020). The first partial lockdown was imposed until April 30 and all non-essential activities in the public and private sector, schools and universities were suspended as part of the contention measures. The shopping malls, bars, clubs, zoos, saunas, gyms, and cinemas remained closed and all the individuals over 60 years of age and those with pre-existing medical conditions were instructed to follow the stay-at-home measures. Circulation restrictions (No-drive days) are generally on place all year long, however, on April 23, 2020 an extraordinary scheme was implemented, and the vehicular circulation decreased by 62% due to the mandatory controls (GDF 2020). In addition, about 20% of the

metro train, metro bus and light rail stations were closed from April 23, 2020. The government extended the restriction by imposing another period of lockdown until May 31 as the number of confirmed cases increased to 5,847 (Mexico Health Ministry 2020). There is no available information about the effects of anthropogenic activities and other sources on overall air quality of Mexico City amid the COVID-19 pandemic. Thus, the main objective of this paper is: (1) to assess the impact of COVID-19 measures on air quality by evaluating the levels of air pollutants, i.e. CO, NO₂, O₃, SO₂, PM₁₀ and PM_{2.5}, between January 1 – May 31, 2020 and comparing the air quality of the lockdown interval with the same period of previous five years, (2) to evaluate their associations with mobility trends and (3) to empirically analyse whether or not the concentration level of air pollutants can contribute to the outbreak of COVID-19. Therefore, this very first report, as per our knowledge, evaluated the effect of reduced human activities through COVID-19 lockdown on the air quality of Mexico City megapolis.

Materials and methods

Mexico City is the second most populated city of Latin America with >21 million people, divided into 16 municipalities, and circulation of an estimated 9.6 million vehicles (CONAPO 2019). An atmospheric monitoring system, known as SIMAT, continuously measures O₃, SO₂, NO₂, CO, PM₁₀ and PM_{2.5} through the manual and automated atmospheric monitoring networks (REDMA and RAMA), atmospheric deposit network (REDDA) and meteorology and solar radiation network (REDMET). The hourly data from RAMA follows the United States Environmental Protection Agency. The daily concentrations (24h) of SO₂, NO₂, CO, PM₁₀, PM_{2.5} and O₃ and meteorological data (i.e. temperature, relative humidity, wind speed and precipitation) were acquired from the official website of atmospheric monitoring system (<http://www.aire.cdmx.gob.mx>) for the months of January, February,

March, April and May of 2015, 2016, 2017, 2018, 2019 and 2020. We calculated the average of each pollutant ($\mu\text{g}/\text{m}^3$, ppm) for each month.

We considered the levels of air pollutants between January 1 - March 31, 2020 to represent the air quality of pre-lockdown period. Concentrations of air pollutants of the lockdown period between April 1 - May 31, 2020 were compared with the pre-lockdown concentrations to estimate the short-term changes in air quality during the lockdown period. The lockdown, however, was divided into different periods: PL1 between April 1 - April 30, 2020 and PL2 between May 1-31, 2020. The Phase I represents changes (%) occurred in the level of air pollutants of the first period of lockdown, PL1 (i.e. April 1 - April 30, 2020), with respect to pre-lockdown (i.e. January 1 - March 31, 2020). Phase II represents the changes (%) in air pollutants of the second period of the lockdown, PL2 (May 1 – May 31, 2020) with respect to the first period (PL1) of lockdown. The average concentrations in April and May of 2015-2019 were considered as historical trend and they were compared with air pollutants of the entire lockdown period of 2020 to estimate the long-term changes.

To better understand the effects of traffic/mobility on air quality, we acquired online data of average traffic count (https://www.tomtom.com/en_gb/traffic-index/mexico-city-traffic/) for Mexico City by collecting data from TomTom. It represents change in vehicle movements for each day during the COVID-19 pandemic. Basically, the traffic index data is built from users who use TomTom tech in navigation devices, in-dash systems and smartphones. The collected datasets were used to understand the potential relationship between traffic measures and the concentration of air pollutants. We retrieved data of confirmed COVID-19 cases and deaths in Mexico City during the lockdown period from the official website of the Government of Mexico (<https://coronavirus.gob.mx/datos/>) to find the association between COVID-19 mortality and air pollutants between April 1, 2020 to May

31, 2020. Statistica (Version 8) was used to process the data, to perform correlation tests ($p < 0.05, 0.01$ and 0.001) and to determine the relationship between variables.

Results and discussion

Meteorological parameters of 2015-2020

As the meteorological conditions impact the air quality of a region, we carefully considered the variables of temperature, precipitation, relative humidity and wind speed of the studied interval (i.e. January-May) during 2015-2020 (e.g. Chen et al. 2019). Most of the meteorological conditions of the lockdown period of 2020 and the same interval of previous years were almost similar (Table 2). The average temperature ($20-21^{\circ}\text{C}$) of April-May of 2020 remained almost similar to the average temperature of the previous five years for the same interval ($18-22^{\circ}\text{C}$). Average precipitation of the lockdown period (27 mm) was slightly lower compared to the same months in previous years (34 mm). The average temperature of the lockdown interval (April-May), however, increased by 4°C compared to previous months (January-March) in all the six years. The average temperature of the pre-lockdown period (i.e. January-March between 2015-2020) remained $15-17^{\circ}\text{C}$, and the average temperature of lockdown period (i.e. April-May between 2015-2020) increased to $19-21^{\circ}\text{C}$. The average relative humidity (42%) and wind speed (7.1 mph) of the lockdown period exhibited non-significant variations in comparison with the pre-lockdown period (42% of relative humidity and 7 mph of wind speed). The amount of precipitation during lockdown period, however, was higher compared to the pre-lockdown period (Table 2).

Air pollutants during COVID-19 pandemic

Figure 1 shows the mean levels of PM_{10} , $\text{PM}_{2.5}$, NO_2 , CO, SO_2 , and O_3 during pre-lockdown and lockdown periods in Mexico City. The concentrations did not exceed environmental quality standards for air, specified by the Mexican Government (for $\text{PM}_{2.5}$ =

166 $45 \mu\text{g}/\text{m}^3$, $\text{PM}_{10} = 75 \mu\text{g}/\text{m}^3$, $\text{SO}_2 = 290 \text{ ppm}$, $\text{NO}_2 = 400 \text{ ppm}$ based on 24-hour average, CO
167 $= 12.5 \text{ mg}/\text{m}^3$ and $\text{O}_3 = 137 \text{ ppm}$ based on 8-h average) during the lockdown period.
168 Furthermore, all criteria pollutants before lockdown were also within the Mexican air quality
169 standard limits. Our study revealed important changes in concentrations for all air pollutants
170 due to drastic measures adopted to limit the spread of coronavirus during the pandemic.
171 Contents of NO_2 , SO_2 , and PM_{10} declined during the lockdown period compared to pre-
172 lockdown (Fig. 1). On the other hand, the concentrations of O_3 increased and both CO and
173 $\text{PM}_{2.5}$ fluctuated (Fig. 1).

174 Changes in air quality (percentage) for the period of assessment i.e., lockdown period
175 (April 1 – May 31, 2020) vs pre-lockdown period (January 1 – March 31, 2020) estimated
176 the effectiveness of lockdown on air quality (Fig. 2). SO_2 concentration showed a maximum
177 reduction of 55%, and we detected 29% and 11% reductions in NO_2 and PM_{10} , respectively.
178 By contrast, the concentrations of O_3 , $\text{PM}_{2.5}$ and CO increased during the lockdown period.
179 Most notably, we observed higher augmentation in O_3 (63%), and the concentrations of CO
180 and $\text{PM}_{2.5}$ showed increments of 1.1% and 19%, respectively.

181 Furthermore, we analysed changes in the air quality between two periods of lockdown
182 (Phase I and Phase II) to better understand the influence of COVID-19 pandemic restrictions
183 on air pollutants (Fig. 2). Phase I showed much better air quality with significant reduction
184 in concentrations of pollutants during the first period of lockdown (PL1) with respect to pre-
185 lockdown. In addition, the reduction (Phase II) in second period of lockdown (PL2) with
186 respect to the first period (PL1) indicated that second period of lockdown was also successful
187 with significant improvement in the air quality. For instance, SO_2 and NO_2 registered 55%
188 and 20% reductions in Phase I, and both showed further reduction of $<10\%$ in Phase II. In
189 case of PM_{10} , the Phase I registered 1% reduction and the reduction was still higher in Phase

II (18%). However, a fluctuating pattern of PM_{2.5} showed increase of 21% in Phase I and decline of 5% in Phase II. Additionally, we also noted 32% decline of CO in Phase I and a hike of 98% during the Phase II.

Data in Table 3 shows the change (%) of air pollutants concentration during the lockdown months of 2020 compared to the historical trend of same interval comprising of previous five years (2015-2019). The SO₂ concentration recorded a maximum reduction of 36%, while NO₂ concentration exhibited 33% reduction. CO and PM₁₀ concentrations presented almost similar reductions of 29% and 25%, respectively. Moreover, PM_{2.5} displayed less reductions (19%) than other air pollutants. However, the levels of O₃ increased by 14% (Table 3). While significant reductions were reported during lockdown compared to pre-lockdown period of 2020, even much higher reductions in the air pollutants concentration were observed when comparing the lockdown period with the historical trend of 2015-2019. For instance, NO₂ and PM₁₀ recorded reductions of 29% and 11% during lockdown period and pre-lockdown period, whereas they registered higher reductions of 33% and 25% in comparison with five-year trend (2015-2019). Even though CO (1.1%) and PM_{2.5} (19%) were found increased during lockdown and pre-lockdown period, a significant decrease of 29% and 19% was observed in comparison to previous five years. Overall, these results clearly indicate that most of the air pollutants significantly decreased during the pandemic compared with the previous five years.

In Mexico City, the vehicular emission is the most prominent source of air pollutants, (Table 1). According to the emission inventories of Government of Mexico, the transportation sector contributes more than 5,000 tons of PM_{2.5} into the atmosphere and 48% of PM₁₀ (SMA-GDF 2016). The vehicular emission sources account for more than 50% of air pollutants such as PM₁₀, PM_{2.5}, CO and nitrogen oxides (Table 1) and it is expected that

change in traffic patterns had an important effect on the observed concentrations. Therefore, it is necessary to understand the effect of traffic count on air quality during the COVID-19 pandemic. There were 4.6 million cars registered in Mexico City in 2017 and nearly 9.6 million in 2019 (CONAPO 2019). During the lockdown period, the vehicular circulation decreased by 62% due to the mandatory no circulation scheme (GDF 2020). In addition, about 20% of the metro train, metro bus and light rail stations were closed from April 23, 2020 (GDF 2020). It is important to mention here that the great majority of population rely heavily on public transportation to commute in Mexico City. In this regard, we assessed the relationship between the traffic count from TomTom database and concentration of air pollutants (PM_{10} , $PM_{2.5}$, NO_2 , CO , SO_2 , and O_3) during the lockdown period (Fig. 3). We observed an immediate change in the traffic volume on April 1, 2020 when the lockdown measures were implemented (Fig. 3 and 4). The daily averaged traffic volume decreased four orders of magnitude in lockdown period compared to the pre-lockdown period, indicating that only fewer people drove and took public transportation to work, while more people stayed at home. This suggests that the measures were effective and widely followed.

Furthermore, we performed correlation analysis to explore associations between the air pollutants and vehicular traffic (Fig. 4). The relationship between traffic count and PM_{10} ($r^2=0.33$), NO_2 ($r^2=0.46$) and SO_2 ($r^2=0.30$) confirms that reduced vehicular movements led to a significant reduction in the concentrations of PM_{10} (11%), NO_2 (29%) and SO_2 (55%) during lockdown period (Fig. 2). Moreover, the positive correlations among PM_{10} vs NO_2 ($r^2=0.36$) and SO_2 ($r^2=0.48$) suggest less vehicular emission as the principal reason of better air quality (Table 4). It is important to note that the reduction in SO_2 (55%) over the entire lockdown period (March 30-May 31, 2020) is comparable to the reduction (65%) obtained by limiting the use of sulphur bearing gasoline and diesel oil in 2000-2011 and reduced

emissions from combustion of fossil fuels in industry and power generation plants (Benítez-García 2014). While the traffic patterns could explain some of the trends observed in air quality, they do not explain all the features such as high concentrations of PM_{2.5}, CO and O₃ during lockdown period. In case of CO and PM_{2.5}, the lack of significant relation with traffic count indicate other sources such as electric energy production and household combustion (SMA-GDF 2016). Certainly, the usage of electric energy and household combustion augmented as people stayed at home and cooked during this global crisis. Regarding O₃, we observed negative correlation with traffic count ($r^2 = -0.21$) and strong interactions with PM_{2.5} ($r^2 = 0.43; 0.66; 0.60$) for all the three periods (Table 4). Fine aerosols (PM_{2.5}) have longer residence times in the atmosphere in comparison with the coarse fraction (PM₁₀), and their composition determines how they interact with light (Salcedo et al. 2006; Aiken et al. 2009). Increase in PM_{2.5} during the Phase I possibly had an impact on optical properties, but it is difficult to know if it was positive or negative without knowing the composition and physical property of aerosol. Similarly, the role of nitrogen oxides and other nitrogenated species in O₃ production is also complex in the urban atmosphere. In addition, the average temperature of Mexico City during the lockdown interval (April-May) increased by 4°C compared to previous months (January-March) of the last six years (Table 2). The strong solar radiations and favourable conditions for photochemical production also lead to peak in O₃ levels during the months of April and May (e.g. Chen et al. 2019). Previous intensive campaigns conducted in the Mexico City metropolitan area (i.e., IMADA-AVER, MCMA-2003 and MILAGRO; Song et al. 2010; Palomera et al. 2016) had demonstrated the complexity of atmospheric chemistry in formation of the secondary species like O₃ and secondary aerosols. Thus, the increment in both the pollutants during pandemic could be telling something important about changes at the chemistry level. The impacts on air quality due to drastic changes in urban

function during the pandemic would provide valuable information for the air quality management in Mexico City and other cities with similar photochemical pollution problems.

Association between COVID-19 and air pollutants

Air pollution is a concern in Mexico as it was responsible for one in seventeen (5.9%) deaths in the country with nearly 48,100 deaths in 2017 (Larsen 2018; Alves 2020). In a previous study in China, Cui et al. (2003) observed high mortality rates in areas with deteriorated air quality during the SARS epidemic, a virus genetically identical to COVID-19. Given that COVID-19 is a respiratory disease that could quickly spread to the community and would remain viable and infectious in aerosols for hours (van Doremalen et al. 2020), this study investigated the association between concentrations of air pollutants and reported daily number of COVID-19 confirmed cases and deaths in Mexico City. At end of the lockdown period, Mexico City reported 29,826 COVID-19 positive cases and 3,676 deaths. About 70% (n=2567) of the mortality were represented by men and the rest 30% (n=1109) were women (Government of Mexico: <https://coronavirus.gob.mx/datos/>). **Figure 3** illustrates daily variations in air pollutants and daily reported COVID-19 cases and deaths. **Table 4** summarizes the association between air pollutants and the COVID-19 cases and deaths for the entire lockdown as well as in PL1 and PL2. The correlation matrix revealed a significant association between COVID-19 cases and deaths and concentrations of CO and O₃ for all the three periods. PM_{2.5}, however, showed significant correlations only with cases and mortalities of PL1 and PL2. We observed that the increasing levels of aforementioned air pollutants equally affected the COVID-19 cases and mortality rate, and they were independent of sex. Our results show similarities with other studies conducted in other parts of the world. Most of the countries show substantial relationship between PM_{2.5} and COVID-19 mortality rate. Identical to this study, Xiao et al. (2020) found strong correlation between

both in United States and suggested that the fatality rate can increase up to 15% by increase in the concentration of PM_{2.5} (1 µg/m³). Similarly, Yao et al. (2020) identified positive association between higher PM_{2.5} and PM₁₀ and COVID-19 deaths in China. In northern Italy, Setti et al. (2020) recognized strong relationship between both the fine particulate matter and COVID-19 mortality rate. Our study also shows strong positive correlations ($r^2 = 0.86; 0.65$) of CO with COVID-19 cases and deaths in Mexico City. Pansini and Fornacca (2020) observed that the level of CO in the air was positively related with the number of infections in Italy, the USA, and China. Alike to our study, Zhu et al. (2020) reported the association between 4.76% increase in daily confirmed cases in China and O₃ (10 µg/m³) enhancement. Overall, these evidences indicate air pollutants as a potential risk factor for respiratory infections and increase in COVID-19 confirmed cases and deaths.

Limitations of the study

Among the limitations of this study, (i) the emission inventories could not be examined to evaluate the impacts from changed patterns, change in fuel consumption at industrial and domestic levels due to lack of data. A thorough analysis of air pollution with respect to emission during the lockdown would be required in the future research. (ii) We also did not take into account the contribution of air pollutants from water heaters, petrochemical solvents in open areas and emissions from a refinery from the neighbouring Hidalgo state (90 km from Mexico City), as potential sources of contamination. The activity of Popocatepetl volcano might have also played a role. (iii) We could not perform subgroup analyses based on the age specific COVID-19 confirmed cases and deaths and (iv) we did not take into account the contributions of immunology, mask-wearing, hand-washing habits, people gathering and social distancing for evaluating the relationship between daily COVID-19 confirmed cases and deaths. This could be the first study that provides evidence about the

association between air pollutants and COVID-19 mortality in Mexico City. The future studies should consider these limitations as an opportunity in order to establish mechanisms to improve the air quality.

Conclusion

This paper examined the implications of restrictions imposed during two different periods of partial lockdown to mitigate the propagation of COVID-19 on the air quality of Mexico City (Mexico). Contents of NO₂, SO₂, PM₁₀ reduced and ozone was enhanced during the lockdown period compared to the pre-lockdown period as well as historical average of the same interval of last five years. The observed variations in air pollutants did not show association with meteorological parameters and they were mainly caused by stringent restrictions on anthropogenic activities during the COVID-19 pandemic. For example, NO₂, SO₂ and PM₁₀ showed strong relationships with traffic count, indicating significant effect of vehicular movements on air quality. The air pollutants mostly diminished during the first period of lockdown (PL1) between April 1 - April 30, 2020 and the air quality further improved in second period of the lockdown (PL2). Furthermore, PM_{2.5}, CO and O₃ were significantly correlated with the confirmed COVID-19 cases and deaths. Future studies can provide insight about the inter-regional response towards COVID-19 pandemic and deliver a better understanding of air quality and the spread of COVID-19. We also assume that the air pollutants would intensify once the lockdown restrictions are lifted and this temporary improvement in air quality might help to formulate new strategies to revoke the years of damages caused by different human activities.

Acknowledgements

VCS thanks CONACYT project no. 274276 “Fase I De La Remediación de Áreas Contaminadas Con Hidrocarburos En La Refinería Gral. Lázaro Cárdenas” for Postdoctoral

fellowship. The authors would like to acknowledge all the health-care workers involved in the diagnosis and treatment of patients world-wide. We also thank Dirección de Monitoreo Atmosférico-CDMX for making air quality data open access.

References

- Agarwal A, Kaushik A, Kumar S, Mishra R (2020) Comparative study on air quality status in Indian and Chinese cities before and during the COVID-19 lockdown period. *Air Qual Atmos Health* 1-12
- Aiken AC, Salcedo D, Cubison MJ, Huffman JA, DeCarlo PF, Ulbrich IM, Docherty KS, Sueper D, Kimmel JR, Worsnop DR, Trimborn A (2009) Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (T0)–Part 1: Fine particle composition and organic source apportionment. *Atmos Chem Phys* 9:6633–6653
- Alves B (2020) Mexico: deaths from air pollution 2000-2017 *Energy & Environmental Services* <https://www.statista.com/statistics/868758/number-deaths-air-pollution-mexico/>
- Anil I, Alagha O (2020) The impact of COVID-19 lockdown on the air quality of Eastern Province, Saudi Arabia. *Air Qual Atmos Health* 1-12
- Benítez-García SE, Kanda I, Wakamatsu S, Okazaki Y, Kawano M (2014) Analysis of criteria air pollutant trends in three Mexican metropolitan areas. *Atmosphere* 5:806-829
- Calderón-Garcidueñas L, Kulesza RJ, Doty RL, D'Angiulli A, Torres-Jardón R (2015) Megacities air pollution problems: Mexico City Metropolitan Area critical issues on the central nervous system pediatric impact. *Environ Res* 137:157-169

366 Chen Z, Zhuang Y, Xie X, Chen D, Cheng N, Yang L, Li R (2019) Understanding long-term
367 variations of meteorological influences on ground ozone concentrations in Beijing
368 During 2006–2016. *Environ Pollut* 245:29-37

369 CONAPO (2019) Consejo Nacional de Población Datos Abiertos del Indice de Marginacion.
370 [http://www.conapogob.mx/es/CONAPO/Datos_Abiertos_del_Indice_de_Marginacio](http://www.conapogob.mx/es/CONAPO/Datos_Abiertos_del_Indice_de_Marginacion)
371 n

372 Cui Y, Zhang ZF, Froines J, Zhao J, Wang H, Yu SZ (2003) Air pollution and case fatality
373 of SARS in the People's Republic of China: an ecologic study. *Environ Health* 2:15

374 Davis LW (2017) Saturday driving restrictions fail to improve air quality in Mexico City. *Sci*
375 *Rep* 7:41652

376 Doğan B, Jebli MB, Shahzad K, Farooq TH, Shahzad U (2020) Investigating the Effects of
377 Meteorological Parameters on COVID-19: Case Study of New Jersey, United States.
378 *Environ Res* 110148

379 Gautam S (2020) COVID-19: air pollution remains low as people stay at home. *Air Qual*
380 *Atmos Health* 13:853–857

381 GDF (2020) Nuevas medidas para la fase 3 por COVID-9.
382 <https://www.cdmx.gob.mx/portal/articulo/nuevas-medidas-para-la-fase-3-por-covid-9>

383 Gold DR, Damokosh AI, Pope III CA, Dockery DW, McDonnell WF, Serrano P, Retama A,
384 Castillejos M (1999) Particulate and ozone pollutant effects on the respiratory
385 function of children in southwest Mexico City. *Epidemiology* 8:16

386 Larsen B (2018) Mexico Perspective: Air Pollution Post 2015 CONSENSUS
387 [https://www.copenhagenconsensus.com/sites/default/files/mexico_resources_air_poll](https://www.copenhagenconsensus.com/sites/default/files/mexico_resources_air_pollutionpdf)
388 [utionpdf](https://www.copenhagenconsensus.com/sites/default/files/mexico_resources_air_pollutionpdf)

- 389 Mexico Health Ministry (2020) Nuevo Coronavirus en el mundo COVID-19 Comunicado
390 Técnico Diario. [https://www.gobmx/salud/prensa/nuevo-coronavirus-en-el-mundo-](https://www.gobmx/salud/prensa/nuevo-coronavirus-en-el-mundo-covid-19-comunicado-tecnico-diario-236595)
391 [covid-19-comunicado-tecnico-diario-236595](https://www.gobmx/salud/prensa/nuevo-coronavirus-en-el-mundo-covid-19-comunicado-tecnico-diario-236595)
- 392 Molina LT, Velasco E, Retama A, Zavala M (2019) Experience from integrated air quality
393 management in the Mexico City metropolitan area and Singapore. *Atmosphere*
394 10:512
- 395 Nakada LYK, Urban RC (2020) COVID-19 pandemic: Impacts on the air quality during the
396 partial lockdown in São Paulo state, Brazil. *Sci Total Environ* 139087
- 397 Nava Escudero C (2008) From homo sapiens to homo automobilis: Revisiting air quality
398 management in Mexico City. *Boletín mexicano de derecho comparado* 41:245-278
- 399 Osornio-Vargas ÁR, Bonner JC, Alfaro-Moreno E, Martínez L, García-Cuellar C, Ponce-de-
400 León Rosales S, Miranda J, Rosas I (2003) Proinflammatory and cytotoxic effects of
401 Mexico City air pollution particulate matter in vitro are dependent on particle size and
402 composition. *Environ Health Perspect* 111:1289-1293
- 403 Otmani A, Benchrif A, Tahri M, Bounakhla M, El Bouch M, Krombi MH (2020) Impact of
404 Covid-19 lockdown on PM10, SO2 and NO2 concentrations in Salé City
405 (Morocco). *Sci Total Environ* 139541
- 406 Palomera MJ, Álvarez HB, Echeverría RS, Hernández EG, Álvarez PS, Villegas RR (2016)
407 Photochemical assessment monitoring stations program adapted for ozone precursors
408 monitoring network in Mexico City. *Atmósfera* 29:169-188
- 409 Panda S, Mallik C, Nath J, Das T, Ramasamy B (2020) A study on variation of atmospheric
410 pollutants over Bhubaneswar during imposition of nationwide lockdown in India for
411 the COVID-19 pandemic. *Air Qual Atmos Health* 1-12

- 1
- 2
- 3
- 4 412 Pansini R, Fornacca D (2020) COVID-19 higher induced mortality in Chinese regions with
- 5
- 6 413 lower air quality. medRxiv <https://doi.org/10.1101/2020040420053595>
- 7
- 8
- 9 414 Pata UK (2020) How is COVID-19 affecting environmental pollution in US cities? Evidence
- 10
- 11 415 from asymmetric Fourier causality test. Air Qual Atmos Health 1-7
- 12
- 13
- 14 416 Salcedo D, Onasch TB, Dzepina K, Canagaratna MR, Zhang Q, Huffman JA, DeCarlo PF,
- 15
- 16 417 Jayne JT, Mortimer P, Worsnop DR, Kolb CE (2006) Characterization of ambient
- 17
- 18 418 aerosols in Mexico City during the MCMA-2003 campaign with Aerosol Mass
- 19
- 20 419 Spectrometry: results from the CENICA Supersite Atmos Chem Phys 6:925–946
- 21
- 22
- 23 420 Setti L, Passarini F, De Gennaro G, Barbieri P, Perrone MG, Piazzalunga A, Borelli M,
- 24
- 25 421 Palmisani J, Di Gilio A, Piscitelli P, Miani A (2020) The Potential Role of Particulate
- 26
- 27 422 Matter in the Spreading of COVID-19 in Northern Italy: First Evidence-Based
- 28
- 29 423 Research Hypotheses.
- 30
- 31 424 <https://www.medrxiv.org/content/10.1101/2020.04.11.20061713v1>
- 32
- 33
- 34
- 35 425 Shakoor A, Chen X, Farooq TH, Shahzad U, Ashraf F, Rehman A, e Sahar N, Yan W (2020)
- 36
- 37 426 Fluctuations in environmental pollutants and air quality during the lockdown in the
- 38
- 39 427 USA and China: two sides of COVID-19 pandemic. Air Qual Atmos Health 1-8
- 40
- 41
- 42 428 Sharma M, Jain S, Lamba BY (2020) Epigrammatic study on the effect of lockdown amid
- 43
- 44 429 Covid-19 pandemic on air quality of most polluted cities of Rajasthan (India). Air
- 45
- 46 430 Qual Atmos Health 1-9
- 47
- 48
- 49
- 50 431 SMA-GDF (2016) Inventario de emisiones de la CDMX Contaminantes criterio, toxicos y
- 51
- 52 432 compuestos de efecto invernadero. Secretaría del Medio Ambiente Gobierno del
- 53
- 54 433 Distrito Federal, México DF
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Song J, Lei W, Bei N, Zavala M, De Foy B, Volkamer R, Cardenas B, Zheng J, Zhang R, Molina LT (2010) Ozone response to emission changes: a modeling study during the MCMA-2006/MILAGRO Campaign. *Atmos Chem Phys* 10:3827–3846

van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, (2020) Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New Engl J Med* 382:1564-1567

WHO (2020) <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>

Xiao Wu, Rachel CN, Sabath MB, Braun D, Xiao FD (2020) Exposure to air pollution and COVID-19 mortality in the United States: a nationwide cross-sectional study. medRxiv <https://doi.org/10.1017/CBO9781107415324004>

Yao Y, Pan J, Wang W, Liu Z, Kan H, Meng X, Wang W (2020) Spatial correlation of particulate matter pollution and death rate of COVID19/China/PM25. PM10/multiple linear regression medRxiv <https://doi.org/10.1101/2020.04.07.20052142>

Zhu Y, Xie J, Huang F, Cao L (2020) Association between short-term exposure to air pollution and COVID-19 infection: evidence from China. *Sci Total Environ* 727:138704

Figure legends

Figure 1 Average concentrations of air pollutants during the pre-lockdown and two different periods of partial lockdown (PL1 and PL2) in Mexico City (Mexico).

Figure 2 Bar chart representing the overall changes (%) in air pollutants of Mexico City.

Figure 3 Daily trends of air pollutants, traffic counts and COVID-19 confirmed cases and deaths since the start of restrictions in Mexico City.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 4 Traffic counts for the period of January 1 – May 31, 2020 in Mexico City from the TomTom traffic database and the correlation between traffic count and air pollutants during the lockdown period.

Table legends

Table 1 Main sources of air pollutants for Mexico City (Mexico) according to emission inventory report of SMA-GDF (2016).

Table 2 Meteorological parameters of Mexico City (Mexico) from January to May 2015-2020.

Table 3 Comparison of changes in air pollutants for April and May in 2020 (lockdown period) and the same intervals during 2015-2019 in Mexico City (Mexico).

Table 4 Correlation between daily confirmed COVID-19 cases and deaths and air pollutants in Mexico City (Mexico).

Table 1. Main sources of air pollutants for Mexico City (Mexico) according to emission inventory report of SMA-GDF (2016).

Pollutant sources	Contribution of emissions (%)				
	PM ₁₀	PM _{2.5}	SO ₂	CO	NO _x
Point source e.g. chemical, food, textile, metal and mineral based industries and electrical energy generation	8.3	12.9	9.3	0.3	3.6
Area source e.g. combustion, urban solid waste, construction, non-highway vehicles, paved roads and others	35.1	29.5	62.6	13.3	10.1
Mobile source e.g. Motorcycles, aircraft operation, SUVs, vehicles > 3.8 t, minibuses, industrial combustions, taxis, buses, tractor trucks, private cars,	52.5	55.7	28.1	86.4	86.1
Agriculture activity	4.1	1.9	N/A	N/A	0.2

Table 2: Meteorological parameters of Mexico City (Mexico) from January to May 2015-2020.

	Temperature (°C)	Relative humidity (%)	Wind speed (mph)	Precipitation (mm)
<i>2015</i>				
January	14	54	6.1	0.3
February	15	45	6.1	2.9
March	16	54	7.3	19.5
April	20	48	7.0	8.3
May	18	61	7.4	69.6
<i>2016</i>				
January	14	45	6.0	6.7
February	15	41	6.3	0.8
March	17	42	7.8	27.8
April	20	38	6.8	19
May	20	51	7.5	51.8
<i>2017</i>				
January	15	42	5.9	0
February	17	36	7.2	0
March	17	44	7.7	18.1
April	20	35	8.0	12.9
May	21	46	7.5	48.9
<i>2018</i>				
January	13	48	6.3	7.9
February	17	47	6.3	7.7
March	19	39	7.1	12.2
April	19	44	7.9	44.5
May	20	46	8.1	59.1
<i>2019</i>				
January	15	46	5.8	5.3
February	18	40	6.4	5.7
March	19	38	6.9	7.2
April	20	35	7.4	4.3
May	22	37	7.9	21.6
<i>2020</i>				
January	15	47	6.5	11.3
February	18	41	7.5	5.3
March	20	38	7.0	8.1
April	21	41	7.1	21.9
May	20	43	7.1	32.8

Table 3. Comparison of changes in air pollutants for April and May in 2020 (lockdown period) and the same intervals during 2015-2019 in Mexico City (Mexico).

Air pollutant	2015-2019	2020	Relative change (%) (2015-2019 vs 2020)
CO (ppm)	0.60	0.43	-29
NO ₂ (ppm)	0.026	0.017	-33
O ₃ (ppm)	0.038	0.044	+14
PM ₁₀ (µg/m ³)	50.89	38.17	-25
PM _{2.5} (µg/m ³)	28.82	23.34	-19
SO ₂ (ppm)	0.0041	0.0026	-36

Values in negative shows decreased contents and positive shows increased contents compared to the historical trend (2015-2019).

Table 4. Correlation between daily confirmed COVID-19 cases and deaths and air pollutants in Mexico City (Mexico).

	Total number of cases	Female cases	Male cases	Total number of mortality	Female mortality	Male mortality	PM _{2.5}	PM ₁₀	CO	O ₃	NO ₂	SO ₂
Entire lockdown period (April 1-May 31, 2020)												
Total number of cases	1.00											
Female cases	0.81**	1.00										
Male cases	0.51**	0.40*	1.00									
Total number of mortality	0.99**	0.80**	0.52**	1.00								
Female mortality	0.84**	0.76**	0.60**	0.84**	1.00							
Male mortality	0.86**	0.79**	0.55**	0.85**	0.82**	1.00						
PM _{2.5}	-	-	-	-	-	-	1.00					
PM ₁₀	-0.38*	-0.29*	-	-0.39*	-0.44**	-0.46**	0.81**	1.00				
CO	0.86**	0.84**	0.47**	0.86**	0.88**	0.84**	-	-0.39*	1.00			
O ₃	0.49**	0.56**	0.54**	0.49**	0.51**	0.48**	0.43*	-	0.53**	1.00		
NO ₂	-	-	-0.32*	-	-	-	0.48**	0.36*	-	-	1.00	
SO ₂	-	-	-	-	-	-	0.37*	0.48**	-	-	-	1.00
PL 1 (April 1-April 30, 2020)												
Total number of cases	1.00											
Female cases	0.85**	1.00										
Male cases	0.49*	-	1.00									
Total number of mortality	0.99**	0.83**	0.50*	1.00								
Female mortality	0.86**	0.76**	-	0.87**	1.00							
Male mortality	0.87**	0.75**	-	0.87**	0.67**	1.00						
PM _{2.5}	0.44*	0.41*	-	0.45*	-	0.40*	1.00					
PM ₁₀	-	-	-	-	0.29†	-	0.85**	1.00				
CO	0.65**	0.49*	-	0.65**	0.57*	0.64**	0.53*	-	1.00			
O ₃	0.44*	0.43*	-	0.42*	-	0.52†	0.66**	0.42*	-	1.00		
NO ₂	-	-	-	-	-	-	-	-	0.54†	-	1.00	
SO ₂	-	-	-	-	-	-	0.53*	0.50*	-	-	-	1.00
PL 2 (May 1-May 31, 2020)												
Total number of cases	1.00											
Female cases	0.43*	1.00										
Male cases	-	-	1.00									
Total number of mortality	0.99**	0.42*	-	1.00								
Female mortality	-	-	-	-	1.00							
Male mortality	0.46*	-	-	0.46*	-	1.00						
PM _{2.5}	0.40*	0.45*	-	0.39*	0.41*	-	1.00					
PM ₁₀	-	-	-	0.44*	-	-	0.86**	1.00				
CO	0.39*	0.53†	-	0.38*	0.38*	-	0.77**	0.72**	1.00			
O ₃	-	0.44*	0.45*	-	0.40*	-	0.60**	0.66**	0.47†	1.00		
NO ₂	-	0.64**	-0.37*	-	-	-	0.67**	0.60**	0.89**	-	1.00	
SO ₂	-	-	-	-	-	-	-	0.38*	-	-	-	1.00

 $p < 0.05^*$; $0.01^\dagger$; $0.001^\ddagger$

Figure 1

[Click here to access/download;Figure;Figure 1.jpg](#)

PRE-LOCKDOWN: January 1-March 31, 2020

PL1: April 1-April 30, 2020

PL2: May 1-May 31, 2020

Figure 2

Figure 3

[Click here to access/download;Figure;Figure 3.jpg](#)

Figure 4

Period		PM _{2.5}	PM ₁₀	CO	O ₃	NO ₂	SO ₂
Lockdown	Traffic count	0.14	0.33 [*]	-0.29 [*]	-0.21	0.46 ^{*†‡}	0.30 ^{*†‡}

$p < 0.05^*$; 0.01[†]; 0.001[‡]

Figures

Figure 1

Average concentrations of air pollutants during the pre-lockdown and two different periods of partial lockdown (PL1 and PL2) in Mexico City (Mexico).

Figure 2

Bar chart representing the overall changes (%) in air pollutants of Mexico City.

Figure 3

Daily trends of air pollutants, traffic counts and COVID-19 confirmed cases and deaths since the start of restrictions in Mexico City.

Period		PM _{2.5}	PM ₁₀	CO	O ₃	NO ₂	SO ₂
Lockdown	Traffic count	0.14	0.33*	-0.29*	-0.21	0.46*†‡	0.30*†‡

$p < 0.05^*$; $0.01^\dagger$; $0.001^\ddagger$

Figure 4

Traffic counts for the period of January 1 – May 31, 2020 in Mexico City from the TomTom traffic database and the correlation between traffic count and air pollutants during the lockdown period.