

# Prevalence of Metabolic Syndrome Among Adolescents in India: a Population-Based Study

**Sowmya Ramesh** (✉ [sramesh@popcouncil.org](mailto:sramesh@popcouncil.org))

Population Council

**Ransi Ann Abraham**

All India Institute of Medical Sciences

**Avina Sama**

Population Council

**Harshpal S Sachdev**

Sitaram Bhartia Institute of Science and Research

**Nizamuddin Khan**

Population Council

**Akash Porwal**

Population Council

**Rajib Acharya**

Population Council

**Praween K. Agrawal**

IPE Global Limited

**Sana Ashraf**

Population Council

**Lakshmi Ramakrishnan**

All India Institute of Medical Sciences

---

## Research Article

**Keywords:** Adolescents, Metabolic syndrome, CNNS, India

**Posted Date:** September 24th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-902711/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

**Background:** In India, the prevalence of overweight among adolescents is on the rise, setting the stage for an increase in metabolic syndrome (MS). This paper presents the national prevalence of MS in adolescents in India.

**Methods:** A nationally representative data of adolescents (10–19 years) from the Comprehensive National Nutrition Survey was used. MS was defined based on the NCEP–ATP III criteria for adolescents. Bivariate analysis was used to report socio-demographic differentials in prevalence and to assess interstate variability. Multivariate logistic regression model was constructed to measure the association between socio-demographic characteristics and prevalence of MS. Census data from 2011 was projected to 2017 to calculate burden.

**Results:** The prevalence of MS was 5.2% among adolescents. 11.9%, 15.4%, 26.0%, 31.9% and 3.7% had central obesity, high blood pressure, hypertriglyceridemia, low HDL-cholesterol and high fasting glucose, respectively. The prevalence was higher among males (5.7% vs. 4.7%, adjusted odds ratio (AOR): 1.3, 95% confidence interval [CI]: 1.0, 1.6), those residing in urban areas (7.9% vs 4.2%, AOR: 1.4, 95% CI: 1.1, 1.8), and from wealthier households as compared to their counterparts (8.3% vs. 2.4%, AOR: 3.4, 95% CI: 2.1, 5.5). There was wide interstate variability in the prevalence of MS (0.5% – 16.5%). In 2017, 14.2 million adolescents had MS in India.

**Conclusions:** The prevalence of MS among adolescents in India is low and clustered in urban areas and richer households. Early prevention interventions promoting a healthy lifestyle, especially in high prevalence areas, are needed to keep MS from becoming a public health issue.

## Background

Globally, the prevalence of obesity among adolescents has increased rapidly over the last decade. India is seeing a similar rising trend in the prevalence of overweight and obesity among adolescents. As per the recent National Family Health Survey – 4 (NFHS-4) conducted in 2015–16, the prevalence of overweight or obesity among adolescents aged 15 to 19 years was reported to be 4.2%, a significant increase from the reported prevalence in 2005–06 (NFHS-3) [1, 2]. As well, a systematic review conducted among adolescents aged 10 to 18 years in 2016 in India indicated increasing rates of overweight and obesity [3]. The study further highlighted that the prevalence was not increasing just among adolescents from the higher socio-economic strata but also among those from lower income groups where the prevalence of undernutrition continues to be a major public health issue in the country [3]. A growing proportion of the adolescent population with overweight and obesity sets the stage for a potential increase in metabolic syndrome (MS) as several research studies conducted across the globe indicate a connection between MS and obesity [4].

MS is a cluster of conditions that occur together that include central adiposity, dyslipidemia, hyperglycemia and hypertension. Several large-scale studies conducted across the globe such as the Framingham Offspring Study, Kuopio Ischemic Heart Disease Study, NHANES Mortality Study, Diabetes Epidemiology: Collaborative Analysis of Diagnostic Criteria in Europe, have reported a significant increase in all causes of mortality and cardiovascular complications among adults with MS [5–10]. Studies indicate that children and adolescents with MS have an increased risk of MS as adults, increasing their risk of developing type 2 diabetes mellitus and cardiovascular disease later in life [11–13]. Due to this, in the recent years, there have been increasingly more researches on children and adolescents to better understand MS' patho-physiology, risk factors and levels in different countries, in order to identify strategies to manage MS at an early age.

Different criteria have been used to define MS. Reaven and his colleagues first described MS in 1988, as a cluster of changes associated with resistance to insulin-mediated glucose uptake, including increase in plasma triglyceride, decrease in high density lipoprotein-cholesterol concentration and high blood pressure [14]. Later, the World Health

Organization (WHO) [15], National Cholesterol Education Program and Adult Treatment Panel III (NCEP-ATP III) [16, 17], American Association of Clinical Endocrinologists (AACE) [18] and International Diabetes Federation [19] published their criteria for MS for adolescents [19]. In fact, since 1988, more than 40 different definitions of MS have been in use [20]. The prevalence of MS varies by definition. A systematic review conducted among adolescents aged 10 to 19 years in low to medium-income countries reported a prevalence ranging from 4.2–15.4% among studies that used the NCEP-ATP III criteria; and 4.5–38.7% in studies where the WHO criteria was used [21]. Researchers who used the same data to estimate the prevalence of MS using different definition reported divergent prevalence estimates, ranging from 4.2% to 9.2% [22]. Recently, few smaller studies have been undertaken to estimate the prevalence of MS among adolescents in different parts of India [4, 23–26]. A study conducted in Delhi among children aged 6–18 years belonging to higher socio-economic strata, reported a prevalence of 6.5% [23]. Similarly, a study conducted by Singh *et al* in the state of Jammu and Kashmir reported that 2.6% of adolescents had MS [4]. Yet another study conducted among urban Indian adolescents reported a prevalence of 4.3% and 3.0% using NCEP ATP III and IDF criteria, respectively [25]. Another study identified MS in 4.3% of Asian Indian adolescents [26].

Estimates on the prevalence of MS and associated factors among adolescents in India are largely derived from small scale or regional studies; there are no estimates available at the national level. This study presents the national prevalence of MS in adolescents 10–19 years of age and the associated socio-demographic differentials.

## Methods

### Design, setting and sample

Data for this study was drawn from the Comprehensive National Nutrition Survey (CNNS), a nationally representative sample of children and adolescents (0–19 years), conducted across all 29 states of India and the capital Delhi in 2016–18. CNNS used a multi-stage, stratified, probability proportion to size cluster sampling to recruit participants. This study used data of adolescents aged 10–19 years. Adolescents with any current illness, chronic disease or physical deformity were excluded from participation in CNNS. Half of all adolescents from whom anthropometric measurements were taken were selected by systematic random sampling and were invited to enroll for biological sampling. Out of these, participants with data for waist circumference, blood pressure, triglycerides, HDL cholesterol and glucose were included in the analytical sample (N=8,007). Details of the survey design and sampling methodology are published elsewhere [27]. Figure 1 presents the details of the sample size for various parameters and the final analytical sample.

#### *Blood sample collection and quality control*

Trained phlebotomists collected blood samples from adolescents and accredited private reference laboratories in Mumbai, Delhi and Kolkata were utilized to conduct all tests included in the study. Rigorous quality control measures and monitoring systems were established for sample collection, transportation and testing. The laboratories had their own internal Quality Control (QC) procedures of the reference laboratory. For external quality assurance, laboratories participated in the BIORAD and US Centre for Disease Control external quality assurance scheme. A subset of samples (5%) was sent to other laboratories (All India Institute of Medical Sciences [AIIMS], Delhi, and the National Institute of Nutrition [NIN], Hyderabad) on a monthly basis for comparison testing. Technical Advisory Group (TAG) members, constituted for CNNS and quality control partners including NIN, AIIMS and CDSA made regular visits to the field and the laboratories to ensure that the Standard Operating Procedures were followed for collection, transportation and analysis of biological samples.

#### *Anthropometric data collection and quality control*

Anthropometric data was collected by trained and standardized female health investigators and a three-tier monitoring system was established for quality control of anthropometric data collection. Level 1 monitoring involved an internal quality control observer within the field team to oversee anthropometric equipment calibration and anthropometric measurements. Level 2 comprised, monitoring and supervision undertaken by an external three-member data quality assurance team in each state. The three-member team observed and re-measured at least three respondents in each PSU they visited. Level 3 involved monitoring by Postgraduate Institute for Medical Education and Research (PGIMER) Chandigarh, UNICEF and Population Council to ensure that the measurements were taken as per protocol. Further details of the quality control protocol are available elsewhere [27].

## Measures

### *Biological and anthropometric measures*

HDL cholesterol was assessed by spectrophotometry and direct measure polyethylene glycol modified cholesterol oxidase methods; triglycerides were estimated by spectrophotometry and enzymatic endpoint method and glucose was estimated using spectrophotometry, Hexokinase method. Waist circumference was measured in centimeters (to the nearest 0.1 cm), using a non-elastic fiberglass measuring tape, at the midpoint between the lowest rib and the iliac crest in the mid-axillary line at the end of normal expiration. A mean of two readings was recorded. Blood pressure was measured using an automated device. Three blood pressure readings were taken with a gap of at least two minutes and the mean of the last two readings was used for the analysis.

### *Metabolic syndrome*

The criteria considered for MS in this study was based on the NCEP ATP III criteria, defined as the coexistence of at least three of following five risk factors: high waist circumference (WC) defined as  $WC \geq 90$ th percentile for age and sex, hypertension defined as systolic blood pressure (SBP) and/or diastolic blood pressure (DBP)  $\geq 90$ th percentile for age and sex, impaired glucose tolerance defined as fasting blood glucose of 110 mg/dL or more, hypertriglyceridemia defined as triglycerides of 110 mg/dL or more and Low HDL defined as HDL cholesterol  $\leq 40$  mg/dL [17].

### *Background characteristics*

Background characteristics included in this study were sex (male, female), age (10–12 years, 13–15 years, 16–19 years), current schooling status of adolescents (Yes, No), area of residence (rural, urban), religion (Hindu, Muslim, Sikh, Christians and Others), caste (Scheduled Caste, Scheduled Tribe, Other Backward Class and Others), and household wealth (wealth index: poorest, poor, middle, rich, richest). Wealth index was created by giving scores that were derived using principal component analysis to households based on the number and kinds of consumer goods they owned and their housing characteristics [2]. Following this step, the household score was assigned to each usual household member, ranked based on their score, and then divided by the distribution into five equal categories, each with 20 percent of the population.

### *Unhealthy diet*

In addition, an attempt was made to assess the association between the prevalence of MS and adolescents' consumption of an unhealthy diet. An unhealthy diet included daily consumption of either fried Indian foods such as *pooris*, *pakor*as, *vadas*, *samosas*, *tikkis* etc.; junk foods such as burgers, pizzas, pastas, instant noodles etc. or sweets such as Indian sweets, pastries/cakes, donuts or aerated drinks.

### *Statistical analysis*

Appropriate sampling weights were used to account for non-response rates and differential probabilities for selection of participants across states. Descriptive statistics were generated for the anthropometric and biochemical data. Mean and standard deviation (SD) are presented. The summary statistics for WC, SBP, DBP, fasting glucose levels, triglyceride and HDL cholesterol level were compared between males and females using t-test and Wilcoxon rank sum test. The prevalence of individual risk factors of MS with 95% confidence intervals (CI) was estimated at the national level. The prevalence of MS with 95% CI was estimated at the national, as well as for individual states. Univariate analysis was used to present the prevalence of coexistence of different MS risk factors. The burden of MS in different states and at the national level was estimated using population projections for the year 2017, based on Government of India, Census 2001 and 2011 data. Bivariate analysis was used to report socio-demographic differentials in prevalence and to assess interstate variability. Multivariate logistic regression model was constructed to measure the association between socio-demographic characteristics and prevalence of MS. Adjusted odds ratios (AOR) and corresponding 95% CI are presented in the tables. Stata version 16.0 (College Station, TX, USA) was used for all analyses.

## Ethical considerations

The research was performed in **accordance** with the Declaration of Helsinki. Ethical approval for conducting CNNS was received from the Ethics Committee of the Postgraduate Institute for Medical Education and Research in Chandigarh, India, and the Institutional Review Board of the Population Council in New York. A comprehensive consent process was adopted. All aspects of the study were conveyed to the participants prior to taking consent, including the purpose of the study, their right to discontinue anytime during the study without any penalty, they were assured of the confidentiality of data, they were informed told that there will be no monetary benefits of participation in the study and they will not experience any serious risks due to participation in the study. For adolescents aged 11–17 years, written informed consent was obtained from the caregivers and written assent was obtained from the adolescents. Adolescents aged 18–19 years provided their own consent.

## Results

Overall, the mean WC, SBP, DBP, glucose level, triglyceride level and HDL cholesterol level among adolescents was 63.0cm, 112.1mm of Hg, 71.8 mm of Hg, 89.4mg/dL, 94.0, mg/d and 46.0mg/dL respectively (Table 1). Male adolescents had a significantly higher WC and fasting blood glucose level, compared to females. Whereas, females had significantly higher values for triglycerides and HDL-Cholesterols, compared to males. It was found that the SBP was higher among males, whereas DBP was higher among females.

The prevalence of each component of MS is presented in Table 2. Low HDL cholesterol was the most prevalent risk factor of MS among adolescents (31.9%), followed by hypertriglyceridemia (26%), hypertension (15%) and central adiposity (12%). Impaired glucose was the least prevalent risk factor (3.7%). More than three-fifth (62%) of adolescents were found to have at least one risk factor of MS (Table 3). Higher proportion of males had at least one risk factor of MS compared to females (63 % vs. 60%, p-value <0.01). Less than one percent of adolescents had four or five risk factors of MS.

Overall, the prevalence of MS was 5.2% among adolescents (Table 4). The prevalence was higher among males compared to females (5.7% vs. 4.7%, AOR: 1.3, 95% CI: 1.0, 1.6). Adolescents aged 13–15 years had higher odds of having MS as compared to those in the age group of 10–12 years (10–12 years vs. 13–15 years: 4.2% vs. 6.4%, AOR: 1.5, 95% CI: 1.1, 2.0). Adolescents who reported being in school at the time of the survey had lower odds of having MS, compared to those not in school (4.9% vs 6.1%, AOR: 0.6, 95% CI: 0.5, 0.8). A significant association was observed between the prevalence of MS and area of residence, with higher proportion of adolescents residing in urban areas having MS, compared to those in rural areas (7.9% vs. 4.2%, AOR: 1.4, 95% CI: 1.1, 1.8). Further, in terms of household

wealth, significantly higher proportion of adolescents belonging to wealthier households had a higher chance of getting diagnosed with MS as compared to those from the poorest wealth quintile (8.3% vs. 2.4%, AOR: 3.4, 95% CI: 2.1, 5.5).

Wide interstate variability was observed in the prevalence of MS, ranging from less than 1% in Punjab to as high as 17% in Manipur (Figure 2). In 15 states, the prevalence of MS was higher than the national-level prevalence. Further, in four states—Uttarakhand, Arunachal Pradesh, Manipur and Tripura—the prevalence was higher than 10%. In terms of the burden of MS, this study estimated 14.2 million adolescents (range 11.2 million–17.9 million) aged 10–19 years as having MS in 2017 in India (Table 5). Together, only six states (Uttar Pradesh, Karnataka, Gujarat, Tamil Nadu, Madhya Pradesh and West Bengal) contributed to 53% of MS in the country.

## Discussion

MS is characterised by a set of conditions associated with metabolic disorders and can be referred to as insulin resistance syndrome, dysmetabolic syndrome, syndrome X, Reaven syndrome, metabolic cardiovascular syndrome, to name a few. Several definitions of MS have been used across the globe; however, dyslipidemia (hypertriglyceridemia and low levels of high-density lipoprotein cholesterol), elevated blood pressure, impaired glucose tolerance and central adiposity are common to all. The prevalence of MS varies depending on the diagnostic criteria used. This study used the NCEP ATP III criteria and observed a prevalence of 5.2% among adolescents aged 10–19 years. Recent studies in India also reported clustering of MS risk factors among adolescents and reported similar prevalence ranging from 3.0–6.5% [4, 23–26]. Further, among adolescents observed under this study, the prevalence of MS was as high as 23.3% in those who were overweight or obese as compared to those who were not overweight or obese (4.1%) (based on body mass index [BMI] for age z score) (not presented in tabular format). These results were concurrent with previous research where a connection between overweight and MS was reported among adolescents [4]. High prevalence of MS among overweight or obese adolescents highlights the need for effective preventive strategies that focus on diet, exercise and lifestyle modification rather than medical treatment for this age group.

The present study showed a higher prevalence of MS in males (5.7%) than females (4.7%). Though these results were concurrent with some previous studies, they conflicted with the findings of others. For example, in the study undertaken by Singh *et al* in the state of Jammu and Kashmir, a significantly higher prevalence among males was reported, whereas two other studies undertaken in north India reported either no such difference or a higher prevalence among females [4, 23, 26]. The reason for these differences could be the small sample size in those studies or because they were undertaken in specific states and contextual state-level factors might have contributed to those differentials.

As per the findings of this study, adolescents who resided in urban areas had higher odds of having MS as compared to those living in rural areas (7.9 %vs 4.2%). There are conflicting observations reported in prior research, with some studies reporting findings that are similar to what was observed in the present and others studies reporting no such rural-urban differentials [4, 28]. The probable reason for observing higher prevalence in adolescents living in urban areas could be because of changes in the standards of living, unhealthy dietary habits and adoption of sedentary lifestyles. Studies have reported that low physical activity is associated with increased risk of MS [29]. Similarly, adolescents belonging to households from higher wealth quintiles had higher odds of having MS as compared to those from poorer households. Probable reason could be increased access to high-fat diet or junk food.

The most common component of MS observed in this study was low HDL-cholesterol, which is similar to what was reported in several previous studies undertaken in India and elsewhere [4, 30]. A very small percent of adolescents in this study had 4 to 5 risk factors of MS (0.5%), which is in contrast to an earlier study conducted in India where no adolescent was reported as having four or five components of MS [4], probably due to the small sample size.

This study observed a wide variability in the prevalence of MS across states and in 15 states the prevalence was higher than the national average. It was not possible to compare this with previous studies due to lack of studies reporting state level prevalence for all the states. However, looking at the trends related to overweight and obesity in previous research work in India, the probable reason for this wide geographic variability could be due to multiple factors, such as differences in lifestyles, the level of urbanisation, access to fast food outlets, etc.. In terms of number adolescents, six states (Uttar Pradesh, Karnataka, Gujarat, Tamil Nadu, Madhya Pradesh and West Bengal) contributed to more than half the burden of MS in the country.

This study had several limitations. First, there is little consensus on the definition to be used for MS. The NCEP ATP III definition for MS was used in this study and the prevalence might change if a different definition was used to measure associated conditions of MS. Second, the fasting status considered in the study was based on self-reported information and the limitation of self-reported data is widely recognized. Similarly, some of the socio-demographic variables included in this study such as the items owned by the household, which was used to derive household wealth variables, were based on self-reported responses. Finally, the findings of this study are limited to adolescents and therefore cannot be generalized to the entire population, including all the age groups. Nonetheless, these limitations do not compromise the findings of this study and to the knowledge of the research team of this study, this is the first attempt made to estimate the prevalence and distribution of MS based on a nationally representative sample of adolescents in India. Future research using various other definitions of MS will help gather a range of estimates. Further, contextual state-level factors need to be explored to better understand the geographical variations in MS, as observed by this study, to design targeted interventions promoting lifestyle changes among adolescents to prevent disease progression into adulthood.

## Conclusion

In conclusion, the findings of this study highlight MS as a growing threat to the health of people in India. It is still clustered in urban areas, among economically better off households and in certain states. There is an increasing urgency to focus on early screening for metabolic abnormalities, especially in high prevalence/burden areas and promote healthy lifestyles among adolescents, given the risk that adolescents diagnosed with MS could develop chronic diseases in adulthood if MS goes unaddressed in early stages.

## Abbreviations

**AACE:** American Association of Clinical Endocrinologists

**AIIMS:** All India Institute of Medical Sciences

**AOR:** Adjusted odds ratios

**BMI:** Body mass index

**CDSA:** Clinical Development Services Agency

**CNNS:** The Comprehensive National Nutrition Survey

**CI:** Confidence Intervals

**Gov:** Government of India

**DBP:** Diastolic blood pressure

**HDL cholesterol:** High-density lipoprotein cholesterol

**SBP:** Systolic blood pressure

**MoHFW:** Ministry of Health and Family Welfare

**MS:** Metabolic Syndrome

**NCEP-ATP III:** National Cholesterol Education Program and Adult Treatment Panel III

**NFHS:** National Family Health Survey

**NHANES:** National Health and Nutrition Examination Survey

**NIN:** National Institute of Nutrition

**PGIMER:** Postgraduate Institute for Medical Education and Research

**QC:** Quality Control

**SD:** Standard Deviation

**TAG:** Technical Advisory Group

**WHO:** The World Health Organization

**WC:** Waist circumference

## Declarations

### *Ethics approval and consent to participate*

Ethics approval for conducting CNNS was obtained from the Ethics Committee of the Postgraduate Institute for Medical Education and Research in Chandigarh, India, and the Institutional Review Board of the Population Council in New York. Participation in the study was voluntary. All aspects of the study were conveyed to the participants prior to taking consent, including the purpose of the study, their right to discontinue anytime during the study without any penalty, they were assured of the confidentiality of data, they were informed told that there will be no monetary benefits of participation in the study and they will not experience any serious risks due to participation in the study. Written informed consent for adolescents aged 11–17 years, was obtained from the caregivers and written assent was obtained from the adolescents. Adolescents aged 18-19 years provided their own consent.

***Consent for publication:***Not applicable

### ***Availability of data and materials:***

Reasonable request for data used in this article may be made to the corresponding author.

**Competing interests:** None of the authors listed on this manuscript report any competing interest.

***Funding:*** This work was supported by UNICEF under the CNNS knowledge products project and funded by Aditya and Megha Mittal

***Authors' contributions:*** SR and RAA conceptualized the manuscript. PKA designed the survey, NK computed sampling weights. SR led the statistical analyses and SA, AP and NK contributed to the analyses. RAA, AS, HPS and LR provided

expert advice for the manuscript. SR led the writing of the manuscript with inputs from RAA, AS, RA, HPS, AP, NK and PKA. All authors reviewed the manuscript.

**Acknowledgements:** *Not Applicable*

## References

1. International Institute for Population Sciences (IIPS) and ICF: **National Family Health Survey (NFHS-3), 2005-06: India.** Mumbai: IIPS; 2007.
2. International Institute for Population Sciences (IIPS) and ICF: **National Family Health Survey (NFHS-4), 2015-16: India.** Mumbai: IIPS.; 2017.
3. Ranjani H, Mehreen TS, Pradeepa R, Anjana RM, Garg R, Anand K, Mohan V: **Epidemiology of childhood overweight & obesity in India: A systematic review.** *Indian J Med Res* 2016, **143**:160–174.
4. Singh N, Parihar RK, Saini G, Mohan SK, Sharma N, Razaq M: **Prevalence of metabolic syndrome in adolescents aged 10–18 years in Jammu, J and K.** *Indian J Endocrinol Metab* 2013, **17**:133–137.
5. Halpern A, Mancini MC: **Treatment of obesity: an update on anti-obesity medications.** *Obes Rev* 2003, **4**:25–42.
6. Hu G, Qiao Q, Tuomilehto J, Balkau B, Borch-Johnsen K, Pyorala K, Group DS: **Prevalence of the metabolic syndrome and its relation to all-cause and cardiovascular mortality in nondiabetic European men and women.** *Arch Intern Med* 2004, **164**:1066–1076.
7. Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, Taskinen MR, Groop L: **Cardiovascular morbidity and mortality associated with the metabolic syndrome.** *Diabetes Care* 2001, **24**:683–689.
8. McNeill AM, Rosamond WD, Girman CJ, Golden SH, Schmidt MI, East HE, Ballantyne CM, Heiss G: **The metabolic syndrome and 11-year risk of incident cardiovascular disease in the atherosclerosis risk in communities study.** *Diabetes Care* 2005, **28**:385–390.
9. Sattar N, Gaw A, Scherbakova O, Ford I, O'Reilly DS, al. e: **Metabolic syndrome with and without C-reactive protein as a predictor of coronary heart disease and diabetes in the West of Scotland Coronary Prevention Study.** *Circulation* 2003, **108**:414–419.
10. Skilton MR, Moulin P, Sérusclat A, Nony P, Bonnet F: **A comparison of the NCEP-ATPIII, IDF and AHA/NHLBI metabolic syndrome definitions with relation to early carotid atherosclerosis in subjects with hypercholesterolemia or at risk of CVD: evidence for sex-specific differences.** *Atherosclerosis* 2007, **190**:416–422.
11. Flegal KM, Carroll MD, Kit BK, Ogden CL: **Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010.** *JAMA* 2012, **307**:491–497.
12. Morrison JA, Friedman LA, Gray-McGuire C: **Metabolic syndrome in childhood predicts adult cardiovascular disease 25 years later: the Princeton Lipid Research Clinics Follow-up Study.** *Pediatrics* 2007, **120**:340–345.
13. Morrison JA, Friedman LA, Wang P, Glueck CJ: **Metabolic syndrome in childhood predicts adult metabolic syndrome and type 2 diabetes mellitus 25 to 30 years later.** *J Pediatr* 2008, **152**:201–206.
14. Raven GM: **Banting Lecture 1988: role of insulin resistance in human diseases.** *Diabetes Care* 1988, **37**:1595–1607.
15. Avogaro P, Crepaldi G, Enzi G, Tiengo A: **Metabolic aspects of essential obesity.** *Epatologia* 1965, **11**:226–238.
16. The National Cholesterol Education Program: **Executive summary of the third report of The National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III).** *JAMA* 285:2486e2497.
17. National Heart LaBI, National Institutes of Health.: **Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report: National Cholesterol Education Program.** (September NPN- ed.: NIH; 2002.

18. Einhorn D, Reaven GM, Cobin RH, Ford E, Ganda OP, Handelsman Y, Hellman R, Jellinger PS, Kendall D, Krauss RM, et al: **American College of Endocrinology position statement on the insulin resistance syndrome.** *Endocr Pract* 2003, **9**:237–252.
19. International Diabetic Federation: **The IDF Consensus Definition of the Metabolic Syndrome In Children and Adolescents.** 2007.
20. Ford ES, Li C: **Defining the metabolic syndrome in children and adolescents: will the real definition please stand up?** *J Pediatr* 2008, **152**:160–164.
21. Moraes AC, Fulaz CS, Netto-Oliveira ER, Reichert FF: **[Prevalence of metabolic syndrome in adolescents: a systematic review].** *Cad Saude Publica* 2009, **25**:1195–1202.
22. de Ferranti SD, Gauvreau K, Ludwig DS, Neufeld EJ, Newburger JW, Rifai N: **Prevalence of the metabolic syndrome in American adolescents: findings from the Third National Health and Nutrition Examination Survey.** *Circulation* 2004, **110**:2494–2497.
23. Kapil U, Kaur S: **Prevalence of pediatrics metabolic syndrome (PMS) amongst children in the age group of 6–18 years belonging to high income group residing in national capital territory (NCT) of Delhi.** *Indian J Pediatr* 2010, **77**:1041.
24. Singh R, Bhansali A, Sialy R, Aggarwal A: **Prevalence of metabolic syndrome in adolescents from a north Indian population.** *Diabet Med* 2007, **24**:195–199.
25. Tandon N, Garg MK, Singh Y, Marwaha RK: **Prevalence of metabolic syndrome among urban Indian adolescents and its relation with insulin resistance (HOMA-IR).** *J Pediatr Endocrinol Metab* 2013, **26**:1123–1130.
26. Vikram NK, Misra A, Pandey RM, Luthra K, Wasir JS, Dhingra V: **Heterogeneous phenotypes of insulin resistance and its implications for defining metabolic syndrome in Asian Indian adolescents.** *Atherosclerosis* 2006, **186**:193–199.
27. UNICEF and Population Council: **Comprehensive national nutrition survey: anthropometric measurement manual.** March 2016.
28. Fadzlina AA, Harun F, Nurul Haniza MY, Al Sadat N, Murray L, Cantwell MM, Su TT, Majid HA, Jalaludin MY: **Metabolic syndrome among 13 year old adolescents: prevalence and risk factors.** *BMC Public Health* 2014, **14 Suppl 3**:S7.
29. Nyugen TH, Tang HK, Kelly P, van der Ploeg HP, Dibley MJ: **Association between physical activity and metabolic syndrome: a cross sectional survey in adolescents in Ho Chi Minh City, Vietnam.** *BMC Public Health* 2010, **10**:141.
30. Azizi F, Salehi P, Etemadi A, Zahedi-Asl S: **Prevalence of metabolic syndrome in an urban population: Tehran Lipid and Glucose Study.** *Diabetes Res Clin Pract* 2003, **61**:29–37.

## Tables

**Table 1: Descriptive statistics of key risk factors of metabolic syndrome among adolescents, India, CNNS 2016–18**

Variable	All		Male		Female	
	Mean	Standard Deviation	Mean	Standard Deviation	Mean	Standard Deviation
<b>Anthropometric Measurements</b>						
Waist circumference (cm) <sup>c</sup>	63.0	8.3	63.7	8.6	62.3	7.8
BMI z-score	-1.1	1.3	-1.2	1.3	-1.0	1.2
<b>Blood Pressure and Biochemical Measurements</b>						
Systolic blood pressure (mmHg) <sup>c</sup>	112.1	10.6	112.6	10.6	111.7	10.6
Diastolic blood pressure (mmHg) <sup>a</sup>	71.8	9.1	71.6	9.1	72.1	9.1
Glucose ( mg/dL ) <sup>b</sup>	89.4	10.9	89.7	11.3	89.0	10.4
Triglycerides (mg/dL) <sup>c</sup>	94.0	50.1	90.1	46.0	98.1	53.6
HDL (mg/dL) <sup>c</sup>	46.0	10.4	45.4	10.6	46.6	10.2

<sup>a</sup>p<0.05, <sup>b</sup>p<0.01, <sup>c</sup>P<0.001

**Table 2: Prevalence of individual risk factors of metabolic syndrome among adolescents, India, CNNS 2016-18**

Risk factors	All			Male			Female		
	Prevalence (%)	95% CI	Total N	Prevalence (%)	95% CI	N	Prevalence (%)	95% CI	N
Central obesity	11.9	10.1, 13.8	8,007	12.2	9.9, 15.0	4,067	11.4	9.7, 13.5	3,940
Hypertension	15.4	13.4, 17.7	8,007	15.9	13.2, 18.9	4,067	15.0	12.9, 17.4	3,940
Hypertriglyceridemia	26.0	23.3, 28.9	8,007	23.3	20.0, 26.9	4,067	28.7	25.1, 32.6	3,940
Low HDL	31.9	29.1, 34.9	8,007	34.7	30.9, 38.8	4,067	29.0	25.8, 32.4	3,940
Impaired glucose	3.7	3.0, 4.5	8,007	4.4	3.3, 5.7	4,067	3.1	2.3, 4.0	3,940

**Note:** prevalence presented here is for the sub-sample of adolescents for whom the values for all parameters were available

**Table 3: Prevalence of number of risk factors of metabolic syndrome among adolescents, India, CNNS 2016-18**

No. of risk factors of Metabolic syndrome	All		Male		Female	
	%	N	%	N	%	N
0	38.5	3,006	37.3	1,532	39.7	1,474
1	39.8	3,014	41.2	1,542	38.4	1,472
2	16.5	1,465	15.8	725	17.2	740
3	4.7	449	5.0	229	4.4	220
4	0.5	68	0.7	37	0.3	31
5	0.0	5	0.0	2	0.1	3

**Table 4: Prevalence of the metabolic syndrome among adolescents by demographic characteristics and diet, India, CNNS 2016-18**

Characteristics	Prevalence	95% CI	AOR (95% CI)	p-value	Total N
<b>Sex</b>					
Male	5.7	3.9, 8.2	1.3 (1.0, 1.6)	0.048	4,067
Female	4.7	3.7, 6.0	Reference		3,940
<b>Age (years)</b>					
10-12.	4.2	3.1, 5.6	Reference	-	2,634
13-15	6.4	3.9, 10.5	1.5 (1.1, 2.0)	0.004	2,561
16-18	5.0	3.8, 6.5	1.0 (0.7, 1.3)	0.961	2,812
<b>Currently in school</b>					
No	6.1	4.2, 8.8	Reference		1,305
Yes	4.9	3.7, 6.5	0.6 (0.5, 0.8)	<0.001	6,702
<b>Area of residence</b>					
Rural	4.2	3.4, 5.3	Reference	-	4,451
Urban	7.9	5.0, 12.3	1.4 (1.1, 1.8)	0.013	3,556
<b>Household Characteristics</b>					
<b>Religion</b>					
Hindu	5.3	4.0, 6.9	Reference		5,646
Muslim	4.9	3.4, 7.1	0.8 (0.6, 1.2)	0.288	974
Christian	4.0	1.9, 8.1	0.6 (0.3, 1.4)	0.268	856
Sikh	0.5	0.1, 2.1	0.1 (0.0, 1.1)	0.061	185
Other	9.8	3.3, 25.7	1.7 (0.7, 3.9)	0.221	346
<b>Caste/Tribe</b>					
Scheduled Caste/ Scheduled Tribe	4.8	3.5, 6.4	Reference		3,030
Other Backward Class	5.3	3.4, 8.3	1.0 (0.7, 1.3)	0.843	2,614
Other	5.6	4.1, 7.5	1.1 (0.8, 1.4)	0.769	2,363
<b>Wealth Index</b>					
Poorest	2.4	1.4, 4.1	Reference		606
Poor	4.9	3.2, 7.4	2.1 (1.3, 3.3)	0.001	1,035
Middle	3.9	2.5, 6.0	1.7 (1.1, 2.7)	0.029	1,604
Rich	6.1	4.3, 8.6	2.6 (1.6, 4.1)	<0.001	2,200
Richest	8.3	4.7, 14.1	3.4 (2.1, 5.5)	<0.001	2,562
<b>Unhealthy Diet <sup>a</sup></b>					
No	5.1	4.0, 6.5	Reference		7,242
Yes	6.6	3.2, 13.0	1.1 (0.7, 1.6)	0.657	765

NCEP ATP III criteria was used for metabolic syndrome

<sup>a</sup>Daily consumption of either fried foods (poori, pakora, vada, samosa, tikki etc.) or junk foods (burger, pizza, pasta, instant noodles) or sweets (Indian sweets, pastries/cakes, donuts) or aerated drinks


Table 5: Medium, low and high estimates of total number of adolescents with metabolic syndrome, by state in India in 2017

State	Medium	Low	High
Andhra Pradesh	409174	166632	981277
Arunachal Pradesh	44956	28474	69380
Assam	563796	236948	1247460
Bihar	495024	273494	902530
Chhattisgarh	386478	240777	611203
Goa	18242	11776	27774
Gujarat	1037415	601957	1754640
Haryana	0	0	0
Himachal Pradesh	23757	3751	143795
Jammu & Kashmir	58700	27430	137150
Jharkhand	363363	148311	881629
Karnataka	854292	567213	1284910
Kerala	349771	190028	635407
Madhya Pradesh	860693	303564	2321370
Maharashtra	707940	346393	1385570
Manipur	111464	75121	161071
Meghalaya	26770	10578	65093
Mizoram	11626	4814	26718
Nagaland	0	0	0
NCT of Delhi	123304	50328	301969
Odisha	722895	472200	1090353
Punjab	25458	5417	108332
Rajasthan	298785	71565	1180825
Sikkim	8600	5375	13655
Tamil Nadu	1151024	605138	2105377
Telangana	531673	327842	855239
Tripura	86684	48786	147066
Uttar Pradesh	1244017	342390	4279875
Uttarakhand	243027	90372	588638
West Bengal	1236328	812876	1852602
<b>India</b>	<b>14268398</b>	<b>11293520</b>	<b>17904361</b>

### Population projection for 10-19 age group using Census 2001 and 2011 data


Population projection Year 2017 (using decadal GR= $(\text{population 2011} - \text{population 2001}) / \text{population 2001} * \text{time}$ )\*population 2011+population 2011

## Figures



**Figure 1**

Sample size of various biochemical and anthropometric parameters, CNNS, 2016–18



**Figure 2**

Prevalence of metabolic syndrome among adolescents by state in India, CNNS 2016–18 NCEP ATP III criteria was used for metabolic syndrome Results for Haryana and Nagaland are not presented as the sample size was <50