

Visceral Leishmaniasis Outbreaks in Bihar: Community-level Investigations in the Context of Elimination of Kala-azar as a Public Health Problem

Khushbu Priyamvada

Care India

Joy Bindroo

Care India

Madan Prashad Sharma

Government of Bihar

Lloyd Chapman

London School of Hygiene & Tropical Medicine

Pushkar Dubey

Care India

Tanmay Mahapatra

Care India <https://orcid.org/0000-0002-5401-2045>

Allen Hightower

Independent consultant

Caryn Bern (✉ caryn.bern2@ucsf.edu)

University of California San Francisco <https://orcid.org/0000-0001-8195-7303>

Sridhar Srikantiah

Care India

Research

Keywords: visceral leishmaniasis, epidemiology, risk factors, outbreak investigation, India

Posted Date: October 15th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-90830/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 15th, 2021. See the published version at <https://doi.org/10.1186/s13071-020-04551-y>.

Abstract

Background: With visceral leishmaniasis (VL) incidence at its lowest level since the 1960s, increasing attention has turned to early detection and investigation of outbreaks.

Methods: Outbreak investigations were triggered by recognition of case clusters in the VL surveillance system established for the elimination program. Investigations included ascertainment of all VL cases by date of fever onset, household mapping and structured collection of risk factor data.

Results: VL outbreaks were investigated in 13 villages in 10 blocks of 7 districts. Data were collected for 20,670 individuals, of whom 272 were diagnosed with VL between 2012 and 2019. Risk was significantly higher among 10-19 year-olds and adults 35 or older compared to children younger than 10 years.

Outbreak confirmation triggered vector control activities and heightened surveillance. VL cases strongly clustered in tolas (hamlets within villages) in which >66% of residents self-identified as scheduled caste or scheduled tribe (SC/ST); 79.8% of VL cases occurred in SC/ST tolas whereas only 24.2% of the population resided in them. Other significant risk factors included being an unskilled non-agricultural laborer, migration for work in a brick kiln, living in a kuccha (mud brick) house, household crowding, habitually sleeping outside or on the ground, and open defecation.

Conclusions: Our data highlight the importance of sensitive surveillance with triggers for case cluster detection and rapid, careful outbreak investigations to better respond to ongoing and new transmission. The strong association with SC/ST tolas suggests that efforts should focus on enhanced surveillance in these disadvantaged communities.

Introduction

In the Indian subcontinent, visceral leishmaniasis (VL) is caused by the protozoan parasite *Leishmania donovani*, and transmitted by the sand fly, *Phlebotomus argentipes* [1]. The only proven reservoir is human, and the most severe form of the disease is called kala-azar (black fever). Female sand flies become infected when they take blood meals from patients with kala-azar or those with post-kala-azar dermal leishmaniasis (PKDL), a dermatosis that follows VL in 5 to 15% of apparently cured VL patients [2, 3].

In India, suppression of VL incidence during the malaria eradication campaigns of the 1950s-1960s was followed by three large-scale epidemic waves, peaking in 1978, 1992 and 2007 [4]. Throughout this time, Bihar has been the most severely affected state in India, with endemic districts concentrated north of the Ganges River. A regional effort by the governments of India, Bangladesh and Nepal, coordinated by the World Health Organization (WHO), aims to eliminate VL as a public health problem (defined as VL incidence below 1 per 10,000 population at the subdistrict level) [5]. The main program components comprise rapid VL diagnosis followed by effective treatment, vector control, and systematic disease surveillance [6]. CARE-India conducted a VL situation assessment in 2013 [7], followed by the refinement

and establishment of the online Kala-azar Management Information System (KAMIS) to provide ongoing surveillance under the auspices of the National Vector-Borne Disease Control Programme (NVBDCP).

Visceral leishmaniasis incidence in the Indian subcontinent is now at its lowest level since the 1960s [8]. In this setting of low incidence, increasing attention has turned to early detection of outbreaks, defined by WHO/NVBDCP as a newly reported VL or PKDL case in a village with no reported cases for at least 3 years; a cluster of 3 or more VL or PKDL cases reported in a village within 4 weeks; or an increase in case numbers in an endemic block over an unspecified outbreak threshold [9]. Outbreak investigations consist of a series of steps to confirm the outbreak and define its extent, initiate appropriate interventions, and when feasible, determine risk factors for disease [10]. We conducted a series of community-level VL outbreak investigations from 2017 to 2019. Our aims were to understand the epidemiology of VL clusters meeting the VL outbreak definition and to evaluate risk factors for VL within the outbreak communities. In coming years, prevention of VL resurgence will depend upon the ability of the public health system to anticipate and detect such outbreaks and to rapidly institute effective control measures.

Methods

Two of our investigations were conducted in blocks that had mean annual VL incidence below the elimination threshold of 1 per 10,000 population from 2009 to 2012; we considered these newly affected blocks. The other investigations occurred in eight known endemic blocks whose mean incidence was greater than 2 per 10,000 from 2009 to 2012. The first investigation was conducted in 2017 in Kosra village, Sheikhpura Block, where a large cluster of newly recognized VL cases received considerable press coverage and attention from public health authorities [11, 12]. Subsequent to this investigation, kala-azar block coordinators (KBCs) deployed by CARE throughout Bihar to assist VL elimination program operations were asked to report unexpected VL case clusters, and KAMIS data were evaluated on a regular basis to identify potential outbreaks. In all investigated villages, the field team worked with local public health officials to confirm and map VL cases, and to ensure that control program interventions were instituted. We conducted a full census of each village and collected data on all residents, to facilitate the risk factor assessment. Data included demographics, social status, house construction materials, migration history, occupation, sleeping practices and domestic animal ownership.

Kala-azar block coordinators report VL cases in KAMIS after confirmation at the primary health centre based on the nationally mandated case definition (at least 2 weeks of fever, splenomegaly and positive rK39 rapid test or biopsy)[6]. Report date reflects the date of confirmation. As part of the ongoing surveillance system, KBCs review medical records for each VL patient and collect case details using a structured data format. The collected data include determination of illness duration prior to treatment using a calendar based on locally memorable events such as festivals. Our epidemiological analyses were based on the illness onset month determined in this way.

Statistical Analysis: Epidemiological curves were constructed for each village or geographical cluster of villages, based on month and year of VL case fever onset. To provide a visual assessment of geographic

clustering, case and non-case households were mapped based on GPS readings. Potential risk factors for VL were evaluated in log-binomial regression models, with generalized estimating equations to account for intra-household correlation. Multivariable models were constructed using a stepwise selection procedure with significance level of 0.05. Analysis was performed in SAS version 9.4.

Results

Between December 2017 and April 2019, VL outbreaks were identified and investigated in 13 villages in 10 blocks of 7 districts (Table 1) (See also Additional file 1: Figure S1). Two blocks south of the Ganges River, Sheikhpura and Nawada, had mean annual VL incidence below 1 per 10,000 in 2009–2011. The other 8 blocks, all north of the Ganges River, had mean annual incidence between 2 and 7 per 10,000 population during 2009–2011. VL surveillance data are not available at the block level prior to 2009. Data were collected for 20,670 individuals in these 13 villages, of whom 272 were diagnosed with VL between 2012 and 2019 (Table 2).

Table 1

Locations in which visceral leishmaniasis (VL) outbreak investigations were conducted, Bihar 2017–2019

District	Block	VL incidence, 2009–2011 ¹	Villages investigated	Village population	
				Total N	VL cases ²
Endemic blocks			Villages in endemic blocks		
Muzaffarpur	Paroo	6.19	Pandey	3258	11
Purnia	Banmanhki	6.76	Binowagram	2971	34
Purnia	Bhawanipur	4.29	Bhamath	1379	3
Purnia	Krityanand Nagar	6.36	Pansohi	209	9
Saharsa	Mahishi	6.65	Thanwar	1757	4
Saharsa	Nauhatta	4.25	Rasalpur	2380	15
Saran	Dariyapur	2.77	Jitwarpur	3044	25
Sitamarhi	Dumra	2.01	Chakka Majhauriya, Chakka Rasalpur, Panapur	545	31
Newly affected blocks			Villages in newly affected blocks		
Nawada	Kashichak	0.05	Kashichak, Lal Bigha	2692	66
Sheikhpura	Sheikhpura	0.90	Kosra	2435	74
¹ Mean annual VL incidence in cases per 10,000 population from 2009 to 2011. Data prior to 2009 not available at the block level.					
² Cumulative number of VL cases in the study population, 2012 to 2019					

Table 2
Study population characteristics, visceral leishmaniasis (VL) outbreak investigations conducted in Bihar 2017–2019

Characteristic	Population ¹		VL ²		PKDL ³	
	N	%	N	%	N	%
Sex ⁴						
Male	10693	51.8%	165	60.7%	3	37.5%
Female	9964	48.2%	107	39.3%	5	62.5%
Age group ⁵						
<10	5297	25.7%	55	20.2%	2	25.0%
10–19	4753	23.0%	79	29.0%	2	25.0%
20–34	4828	23.4%	52	19.1%	2	25.0%
35–49	3060	14.8%	44	16.2%	1	12.5%
>=50	2699	13.1%	42	15.4%	1	12.5%
Lives in SC/ST tola ⁶						
Yes	5006	24.2%	217	79.8%	4	50.0%
No	15664	75.8%	55	20.2%	4	50.0%
¹ Total N = 20,670						
² Four subjects with VL prior to 2012 excluded from VL analyses						
³ Includes four VL patients with onset in 2012 (3) and 2014 (1) who are also listed in VL column. Four others with onset prior to 2012 are not included in VL column.						
⁴ Sex data missing for 13 individuals						
⁵ Age data missing for 33 individuals						
⁶ SC/ST, Scheduled caste/scheduled tribe; tola classification presented in detail in Additional File 2: Table S1.						

All age groups were affected by VL, but risk was significantly higher among 10–19 year-olds and adults 35 or older compared to children younger than 10 years (Table 3). In univariable analyses, adult males had significantly higher risk than adult females, but the differences by sex within age groups did not reach statistical significance, and there was no significant interaction between age and sex ($p = 0.49$ by Breslow-Day test for interaction). Age-specific patterns of VL incidence differed markedly in communities

located in historically endemic blocks compared to newly affected blocks (Fig. 1). The cumulative incidence by age group in communities in endemic blocks ranged from 0.7–1.2%, with a peak in the 10–19 year age group and low incidence in older adults. In blocks with newly recognized transmission, the cumulative incidence was substantially higher (2.1 to 4.1%) than in endemic blocks, and was highest in the oldest age group.

Table 3

Univariable analyses of risk factors for visceral leishmaniasis (VL) based on log-binomial regression models. Generalized estimating equations used to account for intra-household correlation.

Characteristic	Cumulative VL incidence n/N (%)	Univariable analyses		
		Relative risk	95% CI	P
Sex				
Male	165/10689 (1.5)	1.42	1.15, 1.76	0.0009
Female	107/9960 (1.1)			
Age group (years)				
<10	55/5296 (1.0)	Referent		
10–19	79/4751 (1.7)	1.69	1.18, 2.43	0.005
20–34	52/4827 (1.1)	1.20	0.84, 1.72	0.312
35–49	44/3059 (1.4)	1.54	1.06, 2.26	0.025
≥50	42/2696 (1.6)	1.71	1.19, 2.47	0.004
Lives in predominantly SC/ST tola ¹				
Yes	217/5004 (4.3)	12.37	8.79, 17.41	< 0.0001
No	55/15658 (0.4)			
Unskilled non-agricultural laborer				
Yes	73/1143 (6.4)	3.95	2.49, 6.26	< 0.0001
No	199/19519 (1.0)			
Ever sleeps outside				
Yes	166/7528 (2.2)	2.40	1.83, 3.14	< 0.0001
No	106/13106 (0.8)			
Sleeps on ground				
Yes	155/3918 (4.0)	4.43	3.22, 6.10	< 0.0001
No	117/16686 (0.7)			

¹SC/ST, Scheduled caste/scheduled tribe; tola classification presented in detail in Table S1.

²Data missing for 2444 respondents, including 75 VL cases.

³*Kuccha*, unfired mud brick; *pucca*, cement or fired brick; *semi-pucca*, a mixture of the two.

	Cumulative VL incidence	Univariable analyses		
Sleeps under net ²				
Ever	145/15764 (0.9)	0.49	0.31, 0.80	0.004
Never	52/2462 (2.1)			
Sleeps under net ²				
Always	71/10963 (0.7)	0.47	0.32, 0.71	0.0003
Sometimes	74/4801 (1.5)	0.73	0.43, 1.24	0.246
Never	52/2462 (2.1)	Referent		
Defecation site				
Field	245/13204 (1.9)	5.28	3.28, 8.49	< 0.0001
Latrine or toilet	27/7451 (0.4)			
Caste				
Scheduled caste/tribe	219/6310 (3.5)	9.40	6.52, 13.53	< 0.0001
Other/General	53/1432 (0.4)			
House materials ³				
Kuccha	174/9560 (1.8)	2.00	1.42, 2.79	< 0.0001
Semipucca or pucca	98/11086 (0.9)			
Crowding				
≥3 people per room	186/10675 (1.7)	1.95	1.40, 2.73	< 0.0001
<3 people per room	86/9968 (0.9)			
Owns cow				
Yes	109/11317 (1.0)	0.58	0.41, 0.80	0.0007
No	163/9336 (1.8)			
Owns goats				
Yes	132/8436 (1.6)	1.39	1.00, 1.93	0.047
¹ SC/ST, Scheduled caste/scheduled tribe; tola classification presented in detail in Table S1.				
² Data missing for 2444 respondents, including 75 VL cases.				
³ <i>Kuccha</i> , unfired mud brick; <i>pucca</i> , cement or fired brick; semi- <i>pucca</i> , a mixture of the two.				

	Cumulative VL incidence	Univariable analyses		
No	140/12217 (1.2)			
Owns pigs				
Yes	21/456 (4.6)	3.39	1.88, 6.12	< 0.0001
No	251/20197 (1.2)			
Migrated for work in past 3 years				
Yes	96/4472 (2.2)	1.47	1.11, 1.94	0.007
No	176/16190 (1.1)			
Migrated to work in brick kiln				
Yes	75/1053 (7.1)	5.06	3.30, 7.76	< 0.0001
No	197/19609 (1.0)			
¹ SC/ST, Scheduled caste/scheduled tribe; tola classification presented in detail in Table S1.				
² Data missing for 2444 respondents, including 75 VL cases.				
³ <i>Kuccha</i> , unfired mud brick; <i>pucca</i> , cement or fired brick; <i>semi-pucca</i> , a mixture of the two.				

Epidemiological curves reveal that VL cases occurred in outbreak villages over a period of several years (Figs. 2 and 3) (see also Additional File 1: Figures S2-S5). South of the Ganges, three nearby villages had VL cases occurring at a low rate from 2013 onward, with increasing transmission leading to larger clusters in Lal Bigha in 2015 and Kosra in 2016–2017 (Fig. 3). Case maps indicate spatial clustering within villages, often within a single tola (Fig. 4) (see also Additional file 1: Figures S6-S8). In all villages, confirmation of the outbreak triggered vector control activities and heightened surveillance through active case detection activities that continued for at least one year following the last detected VL case.

An array of factors, most with evident links to poverty, were associated with a significant increase in VL risk (Table 3). The strongest determinant was residence in a tola in which two-thirds or more of the residents self-identified as scheduled caste or scheduled tribe (SC/ST), associated with a relative risk of 12.4 (95% confidence interval (CI) 8.8, 17.4). VL cases strongly clustered in these tolas; 79.8% (217/272) of VL cases occurred in SC/ST tolas whereas only 24.2% (5006/20,670) of the study population resided in them (Additional File 2: Table S1). Self-identification as SC/ST was associated with a 9-fold increase in risk (95% CI 6.5, 13.5), but the individual-level variable was less strongly predictive of VL than residence location.

Other significant risk factors included being an unskilled non-agricultural laborer, migration for work in a brick kiln, living in a *kuccha* (mud brick) house, household crowding, habitually sleeping outside or on the ground, and open defecation (Table 3). Sleeping under a bed net was associated with significant

protection in the univariate analysis, with a trend toward a dose-response relationship. Unfortunately, net use data were missing for the villages investigated early in the process, which decreased statistical power for these analyses. Ownership of a cow was associated with significant decreased risk. Ownership of a pig, while rare, was associated with significant increased risk; ownership of goats was associated with increased risk that just reached statistical significance. We included questions on indoor residual insecticide spraying (IRS), but spraying occurred after illness onset for 84% of VL cases. Vector control activities occurred in response to the outbreaks and were not relevant as a predictor of disease risk. In the multivariable model, residence in an SC/ST tola remained the strongest predictor of disease, associated with a 7-fold increase in risk (95% CI 5.0, 10.6) compared to residence in other tolas (Table 4). Male sex, sleeping outside or on the ground, open defecation and migration to work in a brick kiln remained in the model with lower relative risk levels.

Table 4

Multivariable log-binomial regression model of risk factors associated with visceral leishmaniasis. The analysis dataset included 272 participants with and 20,357 participants without the disease. Model constructed using a stepwise selection procedure with significance level of 0.05, using generalized estimating equations to account for intra-household correlation.

Characteristic	Adjusted relative risk	95% Confidence Interval	p
Male sex	1.41	1.13, 1.77	0.003
Age group			
<10	Referent		
10–19	1.84	1.27, 2.66	0.002
20–49	1.38	0.99, 1.92	0.06
>=50	1.94	1.30, 2.90	0.001
Lives in predominantly SC/ST tola ¹	7.29	5.03, 10.57	< 0.0001
Ever sleeps outside	1.54	1.12, 2.11	0.008
Sleeps on ground	1.61	1.14, 2.28	0.007
Open defecation	2.06	1.29, 3.30	0.003
Migrates to work in brick kiln	1.71	1.19, 2.46	0.004
¹ SC/ST, Scheduled caste/scheduled tribe; tola classification presented in detail in Table S1.			

In Kosra, where the biggest outbreak occurred, VL case occurrence was highly clustered in a single tola (Sheikhpura tola 1; Fig. 4). In tola 1, the cumulative incidence was 12.6% (65/516) compared to 0.4% (2/459) and 0.5% (7/1460) in tolas 2 and 3. Of 76 households in tola 1, 44% (34/76) had at least one case; 17 (22%) households had more than one member with VL. By contrast, a single VL case occurred in 3.1% (2/65) and 3.7% (7/187) of households in tolas 2 and 3, respectively. Although IRS was instituted

soon after the outbreak was reported in May 2017, VL cases continued to occur, including 12 cases in 2018 and 3 cases in 2019 (Fig. 2).

The first recognized VL patient in Kosra had symptom onset in 2013 and was ill for 440 days before treatment in the private sector in 2015; this case was never reported to KAMIS (Fig. 2). The second VL patient in Kosra was the uncle of the first; he had onset in May 2016. Symptom duration was extremely prolonged in the early phase of the outbreak and shortened progressively after the outbreak was recognized and control measures instituted (median [IQR] days of illness 142 [115,177], 55.5 [39,69.5] and 20 [15, 30] for patients with onset in 2016, 2017 and 2018–2019 respectively). The mean patient age also decreased over the course of the outbreak (from 38.2 years in 2016 to 26.5 in 2017 and 20.3 in 2018/2019).

In Kosra, residence in tola 1 was associated with a relative risk of 28.2 (95% CI 14.0, 57.0); no other risk factors remained in an age-adjusted multivariable model. The strength of the tola 1 association was not due solely to poverty. Tola 2 was also SC/ST predominant and had higher rates of some poverty-associated risk factors than tola 1, including migration to work in a brick kiln (85% vs 46%; $p < 0.0001$), sleeping outside (59% vs 52%; $p < 0.05$) and sleeping on the ground (76% vs 71%; $p > 0.05$).

Discussion

Despite remarkable progress toward the elimination of VL as a public health problem in India, transmission continues and VL outbreaks are increasingly recognized [13]. Our data suggest two different epidemiological scenarios. The age patterns observed in the villages north of the Ganges are characteristic of an endemic pattern. Herd immunity built up during high incidence periods causes a subsequent fall in case numbers, but after a hiatus of 5–10 years, VL returns predominantly in younger age groups not exposed in the earlier wave [4]. By contrast, in the villages south of the Ganges, where transmission was absent or very low during the last epidemic cycle, the highest VL incidence was in adults over 50. Interestingly, the very youngest age group appeared to be spared in both epidemiological patterns in the Indian subcontinent; this may be related to lower exposure due to different sleeping habits or other behavioral patterns.

Despite these differences, the salient risk factors were similar throughout our data, and echo epidemiological studies conducted over the past 20 years [14, 15, 16, 17]. Neighborhood-level clustering results from proximity to previous VL cases, the proven human infection reservoir; such spatial clustering is a well-recognized feature of VL [15, 18, 19, 20, 21, 22]. The occurrence of clusters in SC/ST tolas has been seen in other investigations in Bihar [19], and the intense local risk reflects a matrix of factors linked to poverty [23]. The risk is not due to caste *per se*. As seen in our data, not all SC/ST tolas are associated with risk, and an early study in the Nepali lowlands bordering Bihar showed no significant association with caste [16]. However, once an infectious VL case is present, closely situated houses of vulnerable construction, intra-household crowding, environmental factors that increase local sand fly density, and behaviors that facilitate human-sand fly contact all come into play to promote transmission [17]. The

protective association with cattle has been previously reported, though the effect of their presence is variable [15, 17, 24]. Cattle are an attractive blood meal source, and may thereby increase local sand fly density. However, cattle may also divert flies away from humans, decreasing both biting and the prevalence of leishmanial infection in vectors. Poor nutritional status in the most disadvantaged groups may also increase the likelihood of an infection progressing to kala-azar and boost the ratio of symptomatic to asymptomatic infection [18, 25].

A crucial unanswered question is where people become infected. Because sand flies are active from dusk to dawn, with peak activity for *P. argentipes* recorded around midnight [26], sleeping patterns have long been assumed to alter risk [16]. Bed net use was protective in our univariate analyses, despite having data for only a portion of our study population; this finding is consistent with the protective association for both treated and untreated net use in observational data from Bangladesh and Nepal [15, 16, 27]. Nevertheless, a cluster-randomized trial of treated nets in India and Nepal failed to demonstrate protection, leading to the abandonment of nets as a major tool in the Indian VL elimination program [28]. Vector control in the Indian program relies almost exclusively on IRS, which has a direct impact only on sand flies resting inside houses [6]. Recent data suggest that *P. argentipes* may be less endophilic than previously assumed [29], with trap yields in outdoor vegetation higher than those inside human dwellings, though lower than in mixed cow-human dwellings [30]. Vector surveillance has failed to show a consistent decrease in sand fly densities inside sprayed vs unsprayed houses [31], and VL cases continue to be reported to KAMIS from sprayed villages. Sleeping outdoors was a strong risk factor in the current analysis, suggesting outdoor transmission unlikely to be controlled by IRS. The risk associated with open defecation, which remained significant in the full multivariable model, is intriguing; perhaps vegetation stands constitute a location where humans and sand flies come into contact in the early morning hours [32].

Our data highlight the importance of sensitive surveillance with triggers for case cluster detection and rapid, careful outbreak investigations to better respond to ongoing and new transmission [8, 29]. The intense efforts to control the epidemic waves of the 1970s and 1990s were not followed by comprehensive VL surveillance, and the resurgence in the early 2000s caught the subcontinent unprepared. Without close local attention, VL transmission can pass unrecognized for several years, as occurred in Kosra. The outbreak there was amplified by the long symptomatic (and likely highly infectious) periods of the early VL cases and lack of pre-existing population immunity. No other case cluster we investigated had an attack rate approaching that in Kosra.

Conclusions

In the future, as overall VL incidence falls even farther, the central challenge is to sustain sufficient levels of long-term surveillance and readiness to enable prompt detection and control of VL case clusters. The strong association with the tolas of scheduled castes and scheduled tribes suggests that efforts should focus on enhanced surveillance in these disadvantaged communities.

Abbreviations

CI: Confidence interval

IQR: Interquartile range

IRS: Indoor residual insecticide spraying

KAMIS: Kala-azar Medical Information System

KBC: Kala-azar block coordinator

NVBDCP: National Vector-Borne Disease Control Programme

PKDL: Post-kala-azar dermal leishmaniasis

SC/ST: Scheduled caste or scheduled tribe

VL: Visceral leishmaniasis

WHO: World Health Organization

Declarations

Ethics approval and consent to participate:

The study was approved by Ashirwad Ethics Committee, Ashirwad Hospital and Research Center, Ulhasnagar, Maharashtra, India. Verbal informed consent was obtained from participants and parents or guardian of participating children prior to interview.

Consent for publication:

Not applicable.

Availability of data and materials:

Due to concerns for participant confidentiality, delinked versions of the datasets used during the current study will be available from the corresponding authors upon reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

This study was financially supported by a grant from the Bill and Melinda Gates Foundation (Grant ID# OPP1196454), <http://www.gatesfoundation.org/>. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. LACC was supported by the Bill and Melinda Gates Foundation through the Setting the Post-Elimination Agenda for Kala-azar in India consortium (Grant OPP1183986).

Authors' contributions:

KP, JB, CB and SS conceived, designed and supervised the data collection process. KP conducted the field investigations. CB, KP, JB, LACC and AWH performed the data cleaning and analysis. TM, LACC, AWH, CB and SS reviewed the analytical plan. KP, LACC and CB wrote the manuscript. TM, MPS, AWH and SS reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgments:

The authors thank the KBCs who participated and the members of the communities included in the investigations, especially VL patients and their families, for their generous contribution to the effort to understand and control this disease.

References

1. Burza S, Croft SL, Boelaert M. Leishmaniasis. *Lancet*. 2018;392 10151:951-70; doi: 10.1016/S0140-6736(18)31204-2. <https://www.ncbi.nlm.nih.gov/pubmed/30126638>.
2. Mondal D, Bern C, Ghosh D, Rashid M, Molina R, Chowdhury R, et al. Quantifying the Infectiousness of Post-Kala-Azar Dermal Leishmaniasis Toward Sand Flies. *Clin Infect Dis*. 2019;69 2:251-8; doi: 10.1093/cid/ciy891. <https://www.ncbi.nlm.nih.gov/pubmed/30357373>.
3. Zijlstra EE, Alves F, Rijal S, Arana B, Alvar J. Post-kala-azar dermal leishmaniasis in the Indian subcontinent: A threat to the South-East Asia Region Kala-azar Elimination Programme. *PLoS Negl Trop Dis*. 2017;11 11:e0005877; doi: 10.1371/journal.pntd.0005877. <https://www.ncbi.nlm.nih.gov/pubmed/29145397>.
4. Courtenay O, Peters NC, Rogers ME, Bern C. Combining epidemiology with basic biology of sand flies, parasites, and hosts to inform leishmaniasis transmission dynamics and control. *PLoS Pathog*. 2017;13 10:e1006571; doi: 10.1371/journal.ppat.1006571. <https://www.ncbi.nlm.nih.gov/pubmed/29049371>.
5. WHO Regional Office for South-East Asia: Regional strategic framework for elimination of kala-azar from the South-East Asia Region (2011-2015). New Delhi, India 2012: 20.

6. National Vector Borne Disease Control Programme: Accelerated Plan for Kala-azar Elimination 2017. New Delhi, India: Ministry of Health and Family Welfare, Government of India; 2017: 80.
7. Das A, Karthick M, Dwivedi S, Banerjee I, Mahapatra T, Srikantiah S, et al. Epidemiologic Correlates of Mortality among Symptomatic Visceral Leishmaniasis Cases: Findings from Situation Assessment in High Endemic Foci in India. *PLoS Negl Trop Dis*. 2016;10 11:e0005150; doi: 10.1371/journal.pntd.0005150. <https://www.ncbi.nlm.nih.gov/pubmed/27870870>.
8. Sundar S, Singh OP, Chakravarty J. Visceral leishmaniasis elimination targets in India, strategies for preventing resurgence. *Expert Rev Anti Infect Ther*. 2018;16 11:805-12; doi: 10.1080/14787210.2018.1532790. <https://www.ncbi.nlm.nih.gov/pubmed/30289007>.
9. WHO Country Office for India: Kala-azar Outbreak Response Guidelines in India. New Delhi, India: World Health Organization; 2017: 76.
10. Murhekar M, Moolenaar R, Hutin Y, Broome C. Investigating outbreaks: practical guidance in the Indian scenario. *Natl Med J India*. 2009;22 5:252-6. <https://www.ncbi.nlm.nih.gov/pubmed/20334049>.
11. Rao M: In this Bihar village, a sudden surge of kala azar has hit Dalit migrant workers. In: *Scrollin*. India: ACK Media; 2017.
12. Kumar A, Saurabh S, Jamil S, Kumar V. Intensely clustered outbreak of visceral leishmaniasis (kala-azar) in a setting of seasonal migration in a village of Bihar, India. *BMC Infect Dis*. 2020;20 1:10; doi: 10.1186/s12879-019-4719-3. <https://www.ncbi.nlm.nih.gov/pubmed/31906924>.
13. Rijal S, Sundar S, Mondal D, Das P, Alvar J, Boelaert M. Eliminating visceral leishmaniasis in South Asia: the road ahead. *BMJ*. 2019;364:k5224; doi: 10.1136/bmj.k5224. <https://www.ncbi.nlm.nih.gov/pubmed/30670453>.
14. Picado A, Ostyn B, Singh SP, Uranw S, Hasker E, Rijal S, et al. Risk factors for visceral leishmaniasis and asymptomatic *Leishmania donovani* infection in India and Nepal. *PloS one*. 2014;9 1:e87641; doi: 10.1371/journal.pone.0087641. <http://www.ncbi.nlm.nih.gov/pubmed/24498159>.
15. Bern C, Hightower AW, Chowdhury R, Ali M, Amann J, Wagatsuma Y, et al. Risk factors for kala-azar in Bangladesh. *Emerg Infect Dis*. 2005;11 5:655-62; doi: 10.3201/eid1105.040718. <http://www.ncbi.nlm.nih.gov/pubmed/15890115>.
16. Bern C, Joshi AB, Jha SN, Das ML, Hightower A, Thakur GD, et al. Factors associated with visceral leishmaniasis in Nepal: bed-net use is strongly protective. *The American journal of tropical medicine and hygiene*. 2000;63 3:184-8.
17. Bern C, Courtenay O, Alvar J. Of cattle, sand flies and men: a systematic review of risk factor analyses for South Asian visceral leishmaniasis and implications for elimination. *PLoS Negl Trop Dis*. 2010;4 2:e599; doi: 10.1371/journal.pntd.0000599. <https://www.ncbi.nlm.nih.gov/pubmed/20161727>.
18. Bern C, Haque R, Chowdhury R, Ali M, Kurkjian KM, Vaz L, et al. The epidemiology of visceral leishmaniasis and asymptomatic leishmanial infection in a highly endemic Bangladeshi village. *The American journal of tropical medicine and hygiene*. 2007;76 5:909-14.

- http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17488915
19. Bulstra CA, Le Rutte EA, Malaviya P, Hasker EC, Coffeng LE, Picado A, et al. Visceral leishmaniasis: Spatiotemporal heterogeneity and drivers underlying the hotspots in Muzaffarpur, Bihar, India. *PLoS Negl Trop Dis*. 2018;12 12:e0006888; doi: 10.1371/journal.pntd.0006888.
<https://www.ncbi.nlm.nih.gov/pubmed/30521529>.
 20. Hasker E, Malaviya P, Cloots K, Picado A, Singh OP, Kansal S, et al. Visceral Leishmaniasis in the Muzaffarpur Demographic Surveillance Site: A Spatiotemporal Analysis. *The American journal of tropical medicine and hygiene*. 2018;99 6:1555-61; doi: 10.4269/ajtmh.18-0448.
<https://www.ncbi.nlm.nih.gov/pubmed/30298812>.
 21. Chapman LAC, Jewell CP, Spencer SEF, Pellis L, Datta S, Chowdhury R, et al. The role of case proximity in transmission of visceral leishmaniasis in a highly endemic village in Bangladesh. *PLoS Negl Trop Dis*. 2018;12 10:e0006453; doi: 10.1371/journal.pntd.0006453.
<https://www.ncbi.nlm.nih.gov/pubmed/30296295>.
 22. Chapman LAC, Spencer SEF, Pollington TM, Jewell CP, Mondal D, Alvar J, et al. Inferring transmission trees to guide targeting of interventions against visceral leishmaniasis and post-kala-azar dermal leishmaniasis. *Proc Natl Acad Sci U S A*. 2020; doi: 10.1073/pnas.2002731117.
<https://www.ncbi.nlm.nih.gov/pubmed/32973088>.
 23. Alvar J, Yactayo S, Bern C. Leishmaniasis and poverty. *Trends Parasitol*. 2006;22 12:552-7.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17023215
 24. Barnett P, Singh SP, Bern C, Hightower AW, Sundar S. Virgin soil: the spread of visceral leishmaniasis into Uttar Pradesh, India. *The American journal of tropical medicine and hygiene*. 2005;73 4:720-5.
 25. Cerf BJ, Jones TC, Badaro R, Sampaio D, Teixeira R, Johnson WD, Jr. Malnutrition as a risk factor for severe visceral leishmaniasis. *The Journal of infectious diseases*. 1987;156 6:1030-3.
 26. Dinesh DS, Ranjan A, Palit A, Kishore K, Kar SK. Seasonal and nocturnal landing/biting behaviour of *Phlebotomus argentipes* (Diptera: Psychodidae). *Annals of tropical medicine and parasitology*. 2001;95 2:197-202. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11299126
 27. Mondal D, Chowdhury R, Huda MM, Maheswary NP, Akther S, Petzold M, et al. Insecticide-treated bed nets in rural Bangladesh: their potential role in the visceral leishmaniasis elimination programme. *Tropical medicine & international health : TM & IH*. 2010;15 11:1382-9.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20946233
 28. Picado A, Singh SP, Rijal S, Sundar S, Ostyn B, Chappuis F, et al. Longlasting insecticidal nets for prevention of *Leishmania donovani* infection in India and Nepal: paired cluster randomised trial. *BMJ*. 2010;341:c6760; doi: 10.1136/bmj.c6760. <http://www.ncbi.nlm.nih.gov/pubmed/21190965>.

29. Cameron MM, Acosta-Serrano A, Bern C, Boelaert M, den Boer M, Burza S, et al. Understanding the transmission dynamics of *Leishmania donovani* to provide robust evidence for interventions to eliminate visceral leishmaniasis in Bihar, India. *Parasit Vectors*. 2016;9:25; doi: 10.1186/s13071-016-1309-8. <https://www.ncbi.nlm.nih.gov/pubmed/26812963>.
30. Poche D, Garlapati R, Ingenloff K, Remmers J, Poche R. Bionomics of phlebotomine sand flies from three villages in Bihar, India. *Journal of vector ecology : journal of the Society for Vector Ecology*. 2011;36 Suppl 1:S106-17; doi: 10.1111/j.1948-7134.2011.00119.x. <https://www.ncbi.nlm.nih.gov/pubmed/21366762>.
31. Poche DM, Garlapati RB, Mukherjee S, Torres-Poche Z, Hasker E, Rahman T, et al. Bionomics of *Phlebotomus argentipes* in villages in Bihar, India with insights into efficacy of IRS-based control measures. *PLoS Negl Trop Dis*. 2018;12 1:e0006168; doi: 10.1371/journal.pntd.0006168. <https://www.ncbi.nlm.nih.gov/pubmed/29324760>.
32. Poche DM, Poche RM, Mukherjee S, Franckowiak GA, Briley LN, Somers DJ, et al. Phlebotomine sandfly ecology on the Indian subcontinent: does village vegetation play a role in sandfly distribution in Bihar, India? *Medical and veterinary entomology*. 2017;31 2:207-13; doi: 10.1111/mve.12224. <https://www.ncbi.nlm.nih.gov/pubmed/28106262>.

Figures




Figure 1

Age-specific patterns of visceral leishmaniasis incidence in communities located in historically endemic blocks compared to newly affected blocks. The cumulative incidence in communities in endemic blocks was highest in the 10-19 year age group and low in older age groups. In blocks with newly recognized transmission, the cumulative incidence was substantially higher than in endemic blocks, and was highest in the oldest age group.




Figure 2

Visceral leishmaniasis case count by year and month of fever onset in Kosra village, Sheikhpora block, Sheikhpora district. The first recognized patient reported fever onset in 2013 but was not treated until late 2014. Interventions, including indoor residual spraying, first occurred in March 2017, but cases continued to be reported in 2018 and 2019.




Figure 3

Visceral leishmaniasis case count by year and month of fever onset in Kosra village in Sheikhpura district and Lal Bigha and Kashichak villages in Nawada district. These blocks are located south of the Ganges River and were not considered endemic for VL. Social links between villages may have facilitated low level sustained transmission prior to recognition of the large outbreak in Kosra in 2017.




Figure 4

Map of houses with (red) and without (green) visceral leishmaniasis cases from 2013 to 2019 in Kosra village, Sheikhpura block, Sheikhpura district. In Tola 1, 44% of households had at least one case.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1figuresS1toS8.pptx](#)
- [Additionalfile2TableS1.docx](#)