

Octogenarian Patients with Colon Cancer – Postoperative Morbidity and Mortality are the Major Challenges

Øystein Høydahl (✉ oystein.hoydahl@gmail.com)

Levanger Hospital, Nord-Trøndelag Hospital Trust

Tom-Harald Edna

Levanger Hospital, Nord-Trøndelag Hospital Trust

Athanasios Xanthoulis

Levanger Hospital, Nord-Trøndelag Hospital Trust

Stian Lydersen

Regional Centre for Child and Youth Mental Health and Child Welfare – Central Norway, Norwegian University of Science and Technology

Birger Henning Endreseth

Trondheim University Hospital

Research Article

Keywords: Colon cancer, Octogenarians, Treatment, Survival, Epidemiology

Posted Date: September 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-911268/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Cancer on March 21st, 2022. See the published version at <https://doi.org/10.1186/s12885-022-09384-9>.

Abstract

Background Few studies have addressed colon cancer surgery outcomes in an unselected cohort of octogenarian patients. The present study aimed to evaluate the relative survival of octogenarian patients after a major resection of colon cancer with a curative intent.

Methods All patients diagnosed with colon cancer at Levanger Hospital between 1980 and 2016 were included. We performed logistic regression to test for associations between 100-day mortality and explanatory variables. We performed a relative survival analysis to identify factors associated with short- and long-term survival.

Results Among 239 octogenarian patients treated with major resections with curative intent, the 100-day mortality was 10.1%. Among 215 patients that survived the first 100 days, the five-year relative survival rate was 99.7%. The 100-day mortality of octogenarian patients was significantly shorter than that of younger patients, but the long-term survival converged with that of younger patients. Among octogenarian patients, the incidence of colon cancer more than doubled during our 37-year observation period. The relative increase in patients undergoing surgery exceeded the increase in incidence; hence, more patients were selected for surgery over time. A high 100-day mortality was associated with older age, a high American Society of Anaesthesiologists (ASA) score, and emergency surgery. Moreover, worse long-term survival was associated with a high Charlson Comorbidity Index, a high ASA score, a worse TNM stage, emergency surgery and residual tumours. Both the 100-day and long-term survival rates improved over time.

Conclusion Among octogenarian patients with colon cancer that underwent major resections with curative intent, the 100-day mortality was high, but after surviving 100 days, the relative long-term survival rate was comparable to that of younger patients. Further improvements in survival will primarily require measures to reduce the 100-day mortality risk.

Background

Colon cancer mainly occurs among older individuals. In Nordic countries, increases have been observed in the population, life expectancy, and incidence of colon cancer over the last few decades. These trends are likely to continue; thus, the number of older patients with colon cancer will continue to increase [1, 2], and a significant proportion of these patients will be octogenarians (i.e., aged 80–89 years) [3, 4].

In Norway, a standardized evidence-based approach to assessing and treating colon cancer has been established at a national level [5]. The final treatment strategy for an individual patient should be based on an accurate staging of the disease and on patient-related factors. The national guidelines recommend that multidisciplinary teams undertake treatment decision-making. Guidelines related to adjuvant chemotherapy administration and follow-up times are recommended according to the patient's chronological age.

In the literature, the group of older patients with colon cancer is a vaguely defined term. The definition of 'older age' ranges from ≥ 65 to > 80 years [6–8]. The mainstay of treatment for colon cancer is radical surgery, and this is combined with chemotherapy in selected subgroups of patients. Previous reports have noted that both radical surgery and chemotherapy are increasingly underused with increasing patient age [7, 9]. Despite some variation, several studies have also reported that postoperative morbidity and mortality increased with increasing age [10, 11]. Although it is well known that, overall, long-term survival decreases with increasing age, estimates of long-term disease-free and relative survival rates have varied for older patients treated for colon cancer.

The present study aimed to evaluate the trends, treatments, and outcomes observed over a period of nearly four decades in patients diagnosed with colon cancer. In particular, we investigated octogenarian patients. Over time, this heterogeneous group of patients has become larger. Therefore, it is of paramount importance to raise our awareness of patient-related factors and their impact on cancer treatment outcomes in older patients. Based on this knowledge, we can establish evidence-based, individualized treatment strategies.

Methods

This study included 1530 consecutive patients admitted with colon cancer at Levanger Hospital during 1980–2016. Levanger Hospital is the primary hospital of 10 municipalities in Norway, and the catchment area remained unchanged throughout the study period. The population increased by 18%, from 83,890 inhabitants in 1980, to 99,566 inhabitants in 2016. During this period, the average age of the population also increased. In particular, the number of octogenarian inhabitants increased by 73%, from 2184 individuals in 1980, to 3800 individuals in 2016 [4].

Through the hospital administrative system, we accessed the health records for all patients that were discharged with diagnosis codes of the International Classification of Diseases, 8th revision (ICD-8) from 153.1 to 153.9, with ICD-9 codes from 153.0 to 153.9, and with ICD-10 codes from C18.0 to C19. Data on all patients were recorded, crosschecked, and confirmed with data from the Norwegian Cancer Registry, during 1980–2016. From the hospital database, we retrieved data on demographic and logistic variables, comorbidities, treatment, tumour characteristics (including histopathology), complications after treatment, and short- and long-term outcome measures.

We defined colon cancer as any tumour located above 15 cm from the anal verge. Right colon tumours were defined as tumours localized in the caecum, ascending colon, hepatic flexure, or transverse colon. Left colon tumours were defined as tumours localized in the splenic flexure, descending colon, or sigmoid colon. Tumours located within 15 cm from the anal verge were defined as rectal cancer, and we excluded these and cancers localized in the appendix.

We characterized patient comorbidity with the American Society of Anaesthesiology (ASA) score and the Charlson Comorbidity Index (CCI) [12, 13]. We defined anaemia at admission, as advocated by the World Health Organization, as blood haemoglobin levels below 13 g/dL in males and below 12 g/dL in females

[14]. We also defined “moderate to severe” anaemia as haemoglobin levels below 11 g/dL in males and 10 g/dL in females. Surgical complications were defined according to the Clavien-Dindo classification of surgical complications, grades I-V [15].

Disease stages were based on the TNM classification, sixth edition [16]. An R0 resection was defined as no detectable residual tumour postoperatively; an R1 resection was defined as a microscopic residual tumour detected in a postoperative histological examination; and an R2 resection was defined as a macroscopic residual tumour detected after surgical treatment [17]. An R0 resection was further classified into two groups: an R0 without tumour perforation and an R0 with tumour perforation. Tumour perforations included both spontaneous (12) and iatrogenic perforations (9).

Patients were categorized into five groups, according to treatment intent: (i) a major resection with curative intent (R0 and R1), (ii) a polypectomy, (iii) a major resection with non-curative intent, (iv) a bypass/stoma, and (v) best supportive care.

Emergency surgery was defined as surgery due to evidence of a large bowel obstruction or large bowel perforation. The laparoscopic colon resection technique was gradually introduced during the last part of the study period. A total of 49 patients underwent laparoscopic surgery. In ten of these patients, the procedure was converted to open surgery.

Staging varied throughout the observation period. Staging was based on complete clinical and histopathological examinations of the resected specimen in 84.9% (1299/1530) of patients; a clinical examination and histopathological examination of a tumour biopsy in 7.8% (120/1530) of patients; a pathological evaluation during an autopsy in 1.4% (21/1530) of patients, and clinical evaluations alone in 5.9% (90/1530) of patients.

Since 1993, the Norwegian national guidelines for treatment of colon cancer advocated that all patients aged 75 years or under with Stage III disease should be evaluated for adjuvant chemotherapy. Later, this recommendation was applied to selected patients with Stage II disease [5].

Follow-ups were initially conducted according to local guidelines. Starting in 1993, they were based on very similar, national guidelines [5]. The follow-up time was calculated as the patient-years at risk, starting from the date of admission. The study endpoints were: local recurrence, metastasis, or death, regardless of cause. The mean follow-up time was 6.05 years (standard deviation [SD] = 6.89, range: 0-38.7 years). The end of follow-up was December 31st, 2018.

Statistical analyses

The Exact Unconditional z-pooled test was used to compare binomial proportions; for example, the percentage of reoperations, relative to the percentage of emergency or elective primary operations. The Cochran Armitage exact trend test was used to test for trends in proportions; for example, the proportions of elective surgeries vs. emergency surgeries in different age groups. The Joncheere-Terpstra test was

used to test for the distribution of age, as a dependent variable, across 10-year age groups, as the independent variable. The five-year rates of local recurrence and metastases were estimated with the Kaplan-Meier method.

Logistic regression analyses were performed to assess associations between the 100-day mortality, as the dependent variable, and different explanatory variables. Ordinal logistic regressions were performed to analyse the associations in doubly-ordered $r \times c$ tables; for example, the ASA score stratified by age group. The resulting odds ratios (ORs) represent a common OR estimate for any 2×2 table that would occur, if the $r \times c$ table was collapsed to a 2×2 table, based on any cut-off threshold, along the columns and rows. Multinomial logistic regression analyses were performed in singly ordered $r \times c$ tables; for example, the type of treatment, stratified by age groups.

Relative survival analysis

Relative survival was defined as a measure of mortality compared to the general population. The observed survival in the group with cancer was divided by the expected survival of a comparable group in the general Norwegian population, matched by age, sex, and the calendar year of investigation. Relative survival was estimated with the Ederer II method and analysed with STATA 16 [18]. Multivariable analyses were performed with a full likelihood approach. Norwegian population survival probabilities were downloaded from the Human Mortality Database, for every year from 1980, calculated for groups stratified by sex and age [19].

Two-sided p-values < 0.05 were considered significant. Means are reported with the range (minimum to maximum) and SD, where relevant. Ninety-five percent confidence intervals (95% CI) are reported, when relevant. Analyses were carried out in Stata 16, IBM SPSS Statistics 25, and StatXact 9.

Results

Study population

Table 1 presents the characteristics of all 1530 patients admitted with colon cancer between 1980 and 2016. There were 750 males (49%) and 780 females, with mean ages of 72.3 (range: 32.9–96.1, SD: 11.1) years and 73.2 (range: 20.3–99.6, SD: 11.6) years, respectively. The mean age of the population increased from 71.5 years, in 1980–1989, to 74.5 years, in 2010–2016 ($p = 0.001$). The mean number of patients admitted per year increased by 109%, from 27.4 patients/y in 1980–1989 to 57.4 patients/y in 2010–2016. The number of octogenarian patients increased by 131%, from 6.7 to 15.5 patients admitted per year, respectively.

Table 1
 Characteristics of all patients admitted with colon cancer during the 1980–2016 study period

Characteristic	Age group (years)							P value
	Total, n = 1530	< 65, n = 353	65–74, n = 451	75–79, n = 281	80–84, n = 269	85–89, n = 124	≥ 90, n = 52	
Sex								0.031 ^a
Females	780 (51)	181 (23)	210 (27)	139 (18)	148 (19)	70 (9)	32 (4)	
Males	750 (49)	172 (23)	241 (32)	142 (19)	121 (16)	54 (7)	20 (3)	
Calendar year								< 0.001 ^b
1980–1989	274 (18)	70 (26)	89 (33)	45 (16)	49 (18)	18 (7)	3 (1)	
1990–1999	367 (24)	85 (23)	115 (31)	73 (20)	56 (15)	29 (8)	9 (3)	
2000–2009	487 (32)	129 (27)	121 (25)	85 (18)	100 (21)	33 (7)	19 (4)	
2010–2016	402 (26)	69 (17)	126 (31)	78 (19)	65 (16)	44 (11)	21 (5)	
Charlson Comorbidity Index								< 0.001 ^b
0	1076 (70)	300 (85)	321 (71)	187 (66)	165 (61)	75 (61)	28 (54)	
1–2	358 (23)	51 (14)	100 (22)	72 (26)	81 (30)	39 (32)	15 (29)	
> 2	96 (6)	2 (1)	30 (7)	22 (8)	23 (9)	10 (8)	9 (17)	
ASA score								< 0.001 ^b
1–2	832 (54)	282 (80)	284 (63)	134 (48)	94 (35)	28 (23)	10 (19)	

Values are the number of patients (%), unless otherwise indicated. ^a Cochran-Armitage exact trend test; ^b Ordinal logistic regression with age group as covariate; ^c Multinomial logistic regression with age group as covariate

^d Ordinal logistic regression with age group as covariate, for known stages; ^e Including polypectomy

Age group (years)								
3	598 (39)	66 (19)	147 (33)	131 (47)	150 (56)	76 (61)	28 (54)	
4-5	100 (7)	5 (1)	20 (4)	16 (6)	25 (9)	20 (16)	14 (27)	
Localization								< 0.001 ^a
Right colon	845 (55)	166 (47)	232 (51)	174 (62)	158 (59)	79 (64)	36 (69)	
Left colon	685 (45)	187 (53)	219 (49)	107 (38)	111 (41)	45 (36)	16 (31)	
Stage (TNM)								0.14 ^d
I	189 (12)	42 (12)	61 (14)	34 (12)	34 (13)	15 (12)	3 (6)	
II	582 (38)	127 (36)	160 (36)	117 (42)	101 (38)	47 (38)	30 (58)	
III	331 (22)	78 (22)	120 (27)	62 (22)	54 (20)	12 (11)	5 (10)	
IV	377 (25)	102 (29)	102 (23)	60 (21)	66 (25)	35 (27)	12 (23)	
Unknown	51 (3)	4 (1)	8 (2)	8 (3)	14 (5)	15 (12)	2 (4)	
Treatment intent categories								< 0.001 ^c
<i>Curative intent</i>								
Major resection	1034 (68)	239 (68)	328 (73)	204 (73)	172 (64)	67 (54)	24 (46)	
Polypectomy	38 (3)	10 (3)	11 (2)	11 (4)	6 (2)	0	0	
<i>Non-curative intent</i>								
Major resection	220 (19)	62 (18)	64 (14)	39 (14)	38 (14)	14 (11)	3 (6)	
Bypass/stoma	67 (4)	17 (5)	17 (4)	6 (2)	14 (5)	11 (9)	2 (4)	

Values are the number of patients (%), unless otherwise indicated. ^a Cochran-Armitage exact trend test; ^b Ordinal logistic regression with age group as covariate; ^c Multinomial logistic regression with age group as covariate

^d Ordinal logistic regression with age group as covariate, for known stages; ^e Including polypectomy

	Age group (years)						
<i>Best supportive care</i>	171 (11)	25 (7)	31 (7)	21 (8)	39 (15)	32 (26)	23 (44)
Surgery							0.005 ^b
Elective surgery	1081 (82)	263 (83)	339 (83)	217 (88)	176 (79)	72 (78)	14 (48)
Emergency surgery	240 (18)	55 (17)	70 (17)	32 (12)	48 (21)	20 (22)	15 (52)
Values are the number of patients (%), unless otherwise indicated. ^a Cochran-Armitage exact trend test; ^b Ordinal logistic regression with age group as covariate; ^c Multinomial logistic regression with age group as covariate							
^d Ordinal logistic regression with age group as covariate, for known stages; ^e Including polypectomy							

The mean CCI, the mean ASA score, and the proportion of patients with right-sided colon cancer increased with increasing age. We observed no differences in stages among the age groups. Over time, the percentage of patients diagnosed with stage I or II disease increased from 41%, in 1980–1989, to 58% in 2010–2016 ($p < 0.001$). The number of patients with an unknown stage declined over time and was zero in the last time period (2010–2016).

Overall, 89% (1359/1530) of all patients diagnosed with colon cancer underwent a surgical treatment, including a major resection, a polypectomy, or a palliative procedure. The rate of surgeries decreased as patient age increased. It was 93% (1008/1085) among patients younger than 80 years and 82% (316/393) among octogenarian patients. Overall, 68% (1034/1530) of patients were treated with major resections with a curative intent. Again, this rate decreased as age increased. It was 71% (771/1085) among patients younger than 80 years and 61% (239/393) among octogenarian patients. During the study period, the percentage of octogenarian patients that underwent a major resection with curative intent increased over time. It was 54% (36/67), in the first time-period (1980–1989), and 61% (66/108), in the last time-period (2010–2016).

Among all patients, 16% (240/1530) underwent emergency surgery, and this rate remained stable over time. However, emergency surgery was required more frequently as patient age increased. The rates were 16% (157/976) among patients younger than 80 years and 22% (68/316) among octogenarian patients. Over time, the proportion of emergency procedures in octogenarian patients decreased: the rates were 28.6% (16/56) during 1980-89 and 20% (17/85) during 2010–2016.

The mean hospital stay after a major resection with curative intent decreased from 17.0 days (range: 2–67, SD: 11.7) during 1980–1989 to 9.7 days (range: 4–47, SD: 6.3) during 2010–2016. The mean hospital stay for patients undergoing a major resection with non-curative intent decreased over the years.

The means were 21.6 days (range: 2-119, SD: 17.5) during 1980–1989 and 10.8 days (range: 3–24, SD: 4.7) during 2010–2016. The corresponding mean hospital stays for patients that underwent the best supportive care were 17.6 days (range: 1-102, SD: 25.4) and 6.8 days (range: 1–28, SD: 6.1), respectively.

Mortality within 100 days for all patients

Overall, the 100-day mortality rate after admission was 14.3% (219/1530). The mortality rate increased successively as patient age increased. Mortality rates were 7.4% in patients < 65 years, 23.7% in octogenarian patients, and 48.1% in patients above 90 years ($p < 0.001$). During 1980–1989, 22.6% (62/274) of all admitted patients died within 100 days. In comparison, during 2010–2016, only 11.2% (45/402) of patients died. Table 2 presents the prognostic factors we identified that were associated with mortality within 100 days after admission. The odds of death increased with increasing patient age.

Table 2
Factors associated with 100-day mortality for all patients admitted with colon cancer in 1980–2016; n = 1530

Factors	Unadjusted odds ratio (95% CI)	P value	Adjusted odds ratio (95% CI)	P value
Age (years)				
< 65	1 (reference)		1 (reference)	
65–74	1.26 (0.75 to 2.01)	0.38	1.19 (0.64 to 2.24)	0.59
75–79	1.73 (1.01 to 2.96)	0.045	1.91 (0.97 to 3.78)	0.063
80–84	3.53 (2.16 to 5.78)	< 0.001	2.59 (1.35 to 4.98)	0.004
85–89	4.75 (2.71 to 8.33)	< 0.001	2.14 (0.99 to 4.64)	0.053
≥ 90	11.65 (5.93 to 22.87)	< 0.001	6.44 (2.55 to 16.25)	< 0.001
Calendar year	0.97 (0.96 to 0.98)	< 0.001	0.94 (0.92 to 0.96)	< 0.001
Female sex	0.75 (0.56 to 0.996)	0.047	0.73 (0.50 to 1.07)	0.11
CCI	1.64 (1.39 to 1.93)	< 0.001	1.39 (1.09 to 1.76)	0.007
ASA score				
1–2	1 (reference)		1 (reference)	
3	2.76 (1.96 to 3.90)	< 0.001	1.90 (1.21 to 3.00)	
4–5	21.27 (13.11 to 34.49)	< 0.001	6.90 (3.65 to 13.05)	< 0.001
Anaemia ^a	1.47 (1.09 to 1.98)	0.012	1.21 (0.80 to 1.85)	0.37
Emergency surgery	2.98 (2.15 to 4.14)	< 0.001	4.97 (3.03 to 8.13)	< 0.001
Localization (left vs. right)	0.84 (0.63 to 1.12)	0.24	0.75 (0.50 to 1.13)	0.17
TNM-stage				
I	1 (reference)		1 (reference)	
II	2.37 (0.99 to 5.67)	0.052	1.28 (0.47 to 3.50)	0.63
III	1.55 (0.60 to 4.03)	0.37	1.07 (0.36 to 3.20)	0.91
IV	15.87 (6.85 to 36.77)	< 0.001	4.45 (1.44 to 13.78)	0.010

Results are from a logistic regression analysis, with death within 100 days as dependent variable; unadjusted: analysis performed with one covariate at a time; adjusted: analysis performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified in three levels: 0, 1 and 2+; ^aAnaemia was defined as < 13 g/dL in males and < 12 g/dL in women (based on WHO recommendations)

Factors	Unadjusted odds ratio	P value	Adjusted odds ratio	P value
Unknown	31.72 (11.89 to 84.61)	< 0.001	1.61 (0.44 to 5.85)	0.47
Treatment intent categories				
<i>Curative intent</i>				
Major resection	1 (reference)		1 (reference)	< 0.001
Polypectomy	0.56 (0.08 to 4.13)	0.57	1.36 (0.17 to 10.8)	0.77
<i>Non-curative intent</i>				
Major resection	4.99 (3.21 to 7.76)	< 0.001	4.31 (2.51 to 7.38)	< 0.001
Bypass/stoma	28.61 (16.26 to 50.36)	< 0.001	28.28 (13.00 to 61.49)	< 0.001
<i>Best supportive care</i>	21.78 (14.35 to 33.05)	< 0.001	29.99 (16.06 to 56.03)	< 0.001
Results are from a logistic regression analysis, with death within 100 days as dependent variable; unadjusted: analysis performed with one covariate at a time; adjusted: analysis performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified in three levels: 0, 1 and 2+; ^a Anaemia was defined as < 13 g/dL in males and < 12 g/dL in women (based on WHO recommendations)				

Long-term relative survival rates for all patients

Overall, the five-year relative survival rate for all patients was 58.5% (95% CI: 55.2 to 61.6). Figure 1a presents the five-year relative survival rates, stratified by age groups. Patients aged 75–79 years had the highest five-year relative survival rate, at 63.1% (95% CI: 55.2 to 70.6), compared to 55.4% (95% CI: 47.4 to 63.5) in octogenarian patients. Figure 1b presents the relative survival rates stratified by treatment intent categories. The five-year relative survival rate for the R0 resection group was 85.1% (95% CI: 81.2 to 88.7), compared to 49.1% (95% CI: 22.1 to 75.6) for the R1 resection group, and 18.3% (95% CI: 4.6 to 41.2) for the R0 resection with perforation group. Among patients < 65 years, the two-year relative survival rates were: 32.6% (95% CI: 21.3 to 44.4) after a major resection with non-curative intent, 0% after a bypass/stoma, and 12.2% (95% CI: 0.3 to 28.1) after the best supportive care. The corresponding rates in octogenarian patients were: 18.8% (95% CI: 8.8 to 32.3) after a major resection with non-curative intent, 0% after a bypass/stoma, and 18.1% (95% CI: 9.3 to 29.8) after the best supportive care.

After excluding patients that died within the first 100 days, the overall five-year relative survival rate was 67.8% (95% CI: 64.4–71.2). Patients aged 65–74 years had the lowest five-year relative survival rate, at 64.8% (95% CI: 59.1–70.1), compared to 72.2% (95% CI: 62.3 to 81.9) in octogenarian patients. Table 3 presents the prognostic factors we identified that were associated with long-term relative survival, among patients that survived 100 days after admission.

Table 3
Factors associated with long-term relative survival in 1311 patients that survived 100 days after admission

Factors	Unadjusted hazard ratio (95% CI)	<i>P</i> value	Adjusted hazard ratio (95% CI)	<i>P</i> value
Age (years)				
<65	1 (reference)		1 (reference)	
65–74	1.08 (0.83 to 1.40)	0.55	1.26 (0.96 to 1.65)	0.092
75–79	0.91 (0.65 to 1.28)	0.60	1.14 (0.81 to 1.60)	0.45
80–84	0.98 (0.67 to 1.42)	0.91	0.89 (0.61 to 1.30)	0.55
85–89	1.35 (0.81 to 2.26)	0.25	0.88 (0.53 to 1.47)	0.63
≥90	0.88 (0.20 to 3.79)	0.86	1.08 (0.46 to 2.51)	0.86
Female sex	1.17 (0.94 to 1.45)	0.16	1.31 (1.06 to 1.62)	0.014
Calendar year				
1980–1989	1 (reference)		1 (reference)	
1990–1999	0.97 (0.71 to 1.33)	0.86	0.70 (0.51 to 0.96)	0.029
2000–2009	0.69 (0.51 to 0.94)	0.020	0.55 (0.41 to 0.76)	< 0.001
2010–2016	0.61 (0.43 to 0.85)	0.004	0.45 (0.32 to 0.64)	< 0.001
CCI	1.36 (1.19 to 1.55)	< 0.001	1.24 (1.07 to 1.44)	0.004
ASA score				
1–2	1 (reference)		1 (reference)	
3	1.31 (1.05 to 1.64)	0.017	1.28 (0.99 to 1.66)	0.061
4–5	4.45 (2.89 to 6.84)	< 0.001	1.92 (1.19 to 3.10)	0.008
Emergency surgery	2.11 (1.64 to 2.72)	< 0.001	1.41 (1.08 to 1.85)	0.013
TNM-stage				
I	1 (reference)		1 (reference)	
II	1.76 (0.65 to 4.75)	0.27	1.46 (0.72 to 2.95)	0.30
III	6.77 (2.63 to 17.45)	< 0.001	5.06 (2.57 to 9.94)	< 0.001

Results are from a multivariable analysis; unadjusted: performed with one covariate at a time; adjusted: performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified as 0, 1, or 2+

Factors	Unadjusted hazard ratio	<i>P</i> value	Adjusted hazard ratio	<i>P</i> value
IV	45.17 (17.80 to 114)	< 0.001	3.67 (1.76 to 7.64)	0.001
Unknown	22.47 (7.76 to 65.04)	< 0.001	1.48 (0.63 to 3.46)	0.37
Treatment intent categories				
<i>Curative intent</i>				
Major resection	1 (reference)		1 (reference)	
Polypectomy	1.65 (0.72 to 3.80)	0.24	4.13 (1.64 to 10.42)	0.003
<i>Non-curative intent</i>				
Major resection	16.21 (12.55 to 20.94)	< 0.001	9.45 (5.87 to 15.21)	< 0.001
Bypass/stoma	25.20 (16.28 to 39.02)	< 0.001	19.04 (10.54 to 34.40)	< 0.001
<i>Best supportive care</i>	20.11 (14.78 to 27.35)	< 0.001	21.15 (12.49 to 35.81)	< 0.001
Results are from a multivariable analysis; unadjusted: performed with one covariate at a time; adjusted: performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified as 0, 1, or 2+				

Patients with stage I-III disease that underwent a major resection with curative intent

Table 4 presents the characteristics of all 1021 patients with colon cancer, stages I-III, that were treated with a major resection with curative intent (R0 and R1). Of these patients, 487 (48%) were males and 534 were females, with mean ages of 71.7 (range: 32.9–91.2, SD: 10.6) and 72.8 (range: 20.3–99.6, SD: 11.1) years, respectively. The mean number of patients per calendar year increased from 17.5 patients/y in 1980–1989 to 38.7 patients/y in 2010–2016. The mean number of octogenarian patients per year increased from 3.6 to 9.3 patients, respectively.

Table 4

Characteristics of patients with colon cancer stages I-III that underwent major resections with curative intent

Characteristic	Age group (years)							P value
	Total, n = 1021	< 65, n = 233	65–74, n = 327	75–79, n = 201	80–84, n = 171	85–89, n = 66	≥ 90, n = 23	
Sex								0.043 ^a
Females	534 (52)	121 (52)	156 (48)	105 (52)	99 (58)	38 (58)	15 (65)	
Males	487 (48)	112 (48)	171 (52)	96 (48)	72 (42)	28 (42)	8 (35)	
Calendar year								0.011 ^b
1980–1989	175 (17)	50 (29)	61 (35)	28 (16)	27 (15)	9 (5)	0 (0)	
1990–1999	241 (24)	55 (23)	79 (33)	49 (20)	35 (15)	16 (7)	7 (3)	
2000–2009	334 (33)	82 (25)	92 (28)	64 (19)	67 (20)	18 (5)	11 (3)	
2010–2016	271 (27)	46 (17)	95 (35)	60 (22)	42 (16)	23 (9)	5 (2)	
ASA score								< 0.001 ^b
1–2	591 (58)	190 (82)	215 (66)	94 (47)	69 (40)	19 (29)	4 (17)	
3	395 (39)	41 (18)	101 (31)	101 (50)	93 (54)	44 (67)	15 (65)	
4	35 (3)	2 (1)	11 (3)	6 (3)	9 (5)	3 (5)	4 (17)	
Localization								< 0.001 ^a
Right colon	573 (56)	103 (44)	179 (55)	130 (65)	105 (61)	43 (65)	13 (57)	
Left colon	448 (44)	130 (56)	148 (45)	71 (35)	66 (39)	23 (35)	10 (43)	

Values are the number of patients (%), unless otherwise indicated. ^aCochran-Armitage exact trend test; ^bOrdinal logistic regression with age group as covariate; ^cNominal logistic regression with age group as covariate

Age group (years)								
Stage (TNM)								0.050 ^b
I	154 (15)	33 (14)	54 (17)	25 (12)	27 (16)	12 (18)	3 (13)	
II	548 (54)	125 (54)	157 (48)	116 (58)	91 (53)	43 (65)	16 (70)	
III	319 (31)	75 (32)	116 (35)	60 (30)	53 (31)	11 (17)	4 (17)	
R-status								0.44 ^c
R0 - resection	983 (96.3)	223 (96)	317 (97)	196 (98)	160 (94)	64 (97)	23 (100)	
R0 - resection with perforation	20 (2)	7 (3)	5 (2)	3 (2)	4 (2)	1 (2)	0 (0)	
R1 - resection	18 (2)	2 (1)	4 (1)	2 (1)	7 (4)	1 (2)	0 (0)	
Type of resection								0.003 ^c
Right hemicolectomy	504 (49)	87 (37)	156 (48)	113 (56)	99 (58)	37 (56)	12 (52)	
Transverse resection	24 (2)	5 (2)	9 (3)	4 (2)	2 (1)	4 (6)	0 (0)	
Left hemicolectomy	131 (13)	39 (17)	44 (14)	27 (13)	12 (7)	4 (6)	5 (22)	
Sigmoid and high anterior resections	267 (26)	75 (32)	89 (27)	41 (20)	43 (25)	16 (24)	3 (13)	
Hartmann's operation	35 (3)	10 (4)	12 (4)	5 (2)	3 (2)	3 (4)	3 (13)	
Subtotal resection	55 (5)	15 (6)	17 (5)	10 (5)	11 (6)	2 (3)	0 (0)	
Other resections	5 (1)	2 (1)	0 (0)	1 (1)	1 (1)	1 (2)	0 (0)	
Emergency surgery								0.13 ^a
Yes	141 (14)	30 (13)	47 (14)	20 (10)	27 (16)	7 (11)	10 (43)	
No	880 (86)	203 (87)	280 (86)	181 (90)	144 (84)	59 (89)	13 (57)	
Values are the number of patients (%), unless otherwise indicated. ^a Cochran-Armitage exact trend test; ^b Ordinal logistic regression with age group as covariate; ^c Nominal logistic regression with age group as covariate								

Postoperative complications and 100-day mortality

In 9.6% of cases, the Clavien-Dindo score was 3 or more. Anastomotic leakage was diagnosed in 2.5% (26/1021), and wound dehiscence in 1.7% (17/1021) of patients. A reoperation was required after 12.1% (17/141) of emergency resections, compared to 5.6% (49/880) of elective resections ($p = 0.004$). Table 5 presents the risk factors we identified that were associated with postoperative complications.

Table 5

Factors associated with postoperative complications^a after major resections with curative intent (R0 and R1); n = 1021^b

Factors	Unadjusted odds ratio (95% CI)	P value	Adjusted odds ratio (95% CI)	P value
Age (years)				
<65	1 (reference)		1 (reference)	
65–74	1.51 (1.08 to 2.11)	0.016	1.35 (0.94 to 1.96)	0.11
75–79	2.13 (1.47 to 3.08)	< 0.001	1.51 (0.99 to 2.28)	0.053
80–84	2.83 (1.93 to 4.17)	< 0.001	1.96 (1.27 to 3.03)	0.002
85–89	2.96 (1.73 to 5.03)	< 0.001	2.14 (1.18 to 3.87)	0.013
≥90	7.60 (3.11 to 18.58)	< 0.001	5.36 (2.11 to 13.61)	< 0.001
Female sex	0.99 (0.78 to 1.25)	0.95	1.15 (0.89 to 1.49)	0.28
Calendar year				
1980–1989	1 (reference)		1 (reference)	
1990–1999	0.46 (0.32 to 0.67)	< 0.001	0.44 (0.30 to 0.66)	< 0.001
2000–2009	0.43 (0.30 to 0.60)	< 0.001	0.50 (0.33 to 0.94)	0.001
2010–2016	0.47 (0.33 to 0.68)	< 0.001	0.61 (0.40 to 0.94)	0.025
ASA score				
1–2	1 (reference)		1 (reference)	
3	2.00 (1.56 to 2.56)	< 0.001	1.43 (1.08 to 1.90)	0.013
4–5	19.07 (9.56 to 38.05)	< 0.001	10.86 (5.19 to 22.73)	< 0.001
Emergency surgery	2.51 (1.75 to 3.60)	0.001	2.30 (1.57 to 3.39)	< 0.001
Anaemia (g/dL haemoglobin)				
Female ≥ 12.0, Male ≥ 13.0	1 (reference)		1 (reference)	
Female 10-11.9, Male 11-12.9	1.97 (1.47 to 2.64)	< 0.001	2.26 (1.64 to 3.11)	< 0.001
Female < 10, Male < 11	4.76 (3.53 to 6.43)	< 0.001	5.61 (4.01 to 7.84)	< 0.001
Surgery duration (minutes)				

^aComplications were classified according to Clavien-Dindo grades; ^bPatients included those with stages I-III colon cancer during 1980–2016

Factors	Unadjusted odds ratio (95% CI)	P value	Adjusted odds ratio (95% CI)	P value
<90	1 (reference)		1 (reference)	
90–179	1.01 (0.73 to 1.38)	0.96	0.96 (0.67 to 1.39)	0.84
≥180	2.24 (1.48 to 3.38)	0.001	1.52 (0.89 to 2.58)	0.13
Blood loss (mL)				
0-200	1 (reference)		1 (reference)	
201–400	1.21 (0.88 to 1.65)	0.24	1.48 (1.04 to 3.11)	0.029
401–800	1.67 (1.20 to 2.32)	0.002	2.16 (1.45 to 3.21)	< 0.001
>800	3.69 (2.41 to 5.65)	< 0.001	4.13 (2.44 to 7.01)	< 0.001
^a Complications were classified according to Clavien-Dindo grades; ^b Patients included those with stages I-III colon cancer during 1980–2016				

Among patients with colon cancer stages I-III, mortality within 100 days after admission was 4.7% (48/1021). The 100-day mortality rates increased successively with increasing age. The rates were 0.4% among patients aged < 65 years, 10.1% in octogenarian patients, and 34.8% in patients above 90 years old ($p < 0.001$). Table 6 presents the factors we identified that were associated with death within 100 days.

Table 6

Factors associated with 100-day mortality after major resections with curative intent (R0 or R1); n = 1021^a

Factor	Dead within 100 days (%)	Unadjusted odds ratio (95% CI)	P value	Adjusted odds ratio (95% CI)	P value
Age (years)					
< 65	1/233 (0.4)				
65–74	8/327 (2.4)	1 (reference)		1 (reference)	
75–79	7/201 (3.5)				
80–84	15/171 (8.8)	4.48 (2.17 to 9.25)	< 0.001	3.78 (1.67 to 8.56)	0.001
85–89	9/66 (13.6)	7.35 (3.11 to 17.37)	< 0.001	7.15 (2.68 to 19.1)	< 0.001
≥ 90	8/23 (34.8)	24.83 (9.22 to 66.88)		11.3 (3.50 to 36.6)	< 0.001
Female sex		0.53 (0.29 to 0.97)	0.038	0.47 (0.23 to 0.95)	0.036
Calendar year				0.96 (0.93 to 0.99)	0.013
1980–1989	13/175 (7.4)	1 (reference)		1 (reference)	
1990–1999	12/241 (5.0)	0.65 (0.29 to 1.47)	0.30	0.45 (0.16 to 1.23)	0.12
2000–2009	12/334 (3.6)	0.46 (0.21 to 1.04)	0.062	0.33 (0.12 to 0.90)	0.030
2010–2016	11/271 (4.1)	0.53 (0.23 to 1.21)	0.13	0.40 (0.15 to 1.05)	0.063
ASA score				3.77 (2.12 to 6.69)	< 0.001
1–2	10/591 (1.7%)	1 (reference)		1 (reference)	
3	22/395 (5.6%)	3.42 (1.61 to 7.32)	0.001	1.77 (0.676 to 4.12)	0.18
4	16/35 (45.7%)	48.93 (19.64 to 122)	< 0.001	15.43 (5.27 to 45.2)	< 0.001
Emergency surgery		9.69 (5.30 to 17.70)	< 0.001	6.28 (3.06 to 12.90)	< 0.001

^aPatients included those with stages I-III colon cancer during 1980–2016; results are from a logistic regression analysis, with death as the dependent variable; unadjusted: performed with one covariate at a time; adjusted: performed with all listed covariates included simultaneously

Long-term relative survival, local recurrence, and metastasis

Overall, the five-year relative survival rate was 83.2% (95% CI: 79.4. to 86.8) for all patients with stages I-III disease that underwent major resections with a curative intent. Patients aged 65–74 years had the lowest five-year relative survival rate: 79.0% (95% CI: 72.9 to 84.4), compared to 88.4% (95% CI: 77.0 to 99.1) in octogenarian patients.

When we excluded patients that died within the first 100 days, the overall five-year relative survival rate was 87.2% (95% CI: 83.3 to 90.7). In this group, patients aged 65–74 years had the lowest five-year relative survival rate: 81.0% (95% CI: 74.9 to 86.4), compared to 98.2% (95% CI: 86.1 to 109.3) in octogenarian patients.

Factors associated with relative long-term survival are presented in Table 7. Long-term relative survival rates did not differ significantly between the different age groups. A similar multivariable analysis performed in a selected group of patients with stage III colon cancer revealed that patients with left-sided colon cancer had better survival than those with right-sided colon cancer (OR = 0.55, 95% CI: 0.33 to 0.92; $p = 0.022$). This effect was not found in separate analyses of patients with stage I or stage II colon cancer.

Table 7

Factors associated with relative long-term relative survival, among patients that survived 100 days; n = 973^a

Factor	Unadjusted hazard ratio (95% CI)	<i>P</i> value	Adjusted hazard ratio (95% CI)	<i>P</i> value
Age (years)				
< 65	1 (reference)		1 (reference)	
65–79	1.29 (0.83 to 2.01)	0.26	0.99 (0.61 to 1.59)	0.97
≥ 80	0.49 (0.14 to 1.69)	0.26	0.65 (0.31 to 1.36)	0.26
Female sex	1.58 (0.998 to 2.43)	0.051	1.60 (1.05 to 2.44)	0.029
Calendar year				
1980–1989	1 (reference)		1 (reference)	
1990–1999	0.82 (0.48 to 1.38)	0.45	0.76 (0.45 to 1.28)	0.30
2000–2009	0.47 (0.27 to 0.84)	0.010	0.54 (0.31 to 0.95)	0.031
2010–2016	0.37 (0.19 to 0.72)	0.004	0.49 (0.26 to 0.90)	0.022
CCI	1.39 (1.14 to 1.68)	0.001	1.40 (1.21 to 1.75)	0.003
ASA score				
1–2	1 (reference)		1 (reference)	
3	1.81 (1.19 to 2.74)	0.005	1.63 (1.02 to 2.59)	0.040
4	4.89 (2.10 to 11.38)	< 0.001	4.37 (1.84 to 10.37)	0.001
Emergency surgery	3.05 (1.97 to 4.72)	< 0.001	2.12 (1.34 to 3.36)	0.001
Left vs. right colon	0.95 (0.62 to 1.44)	0.80	0.74 (0.49 to 1.14)	0.17
TNM-stage				
I	1 (reference)		1 (reference)	
II	3.79 (0.47 to 30.54)	0.21	1.90 (0.53 to 6.73)	0.32
III	16.67 (2.14 to 129)	0.007	8.03 (2.36 to 27.3)	0.001

^aPatients included those with stages I-III colon cancer during 1980–2016, treated with a major resection with curative intent (R0 and R1). Results are from a multivariable analysis; unadjusted: performed with one covariate at a time; adjusted: performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified as 0, 1, 2, or 3+

Factor	Unadjusted hazard ratio (95% CI)	<i>P</i> value	Adjusted hazard ratio (95% CI)	<i>P</i> value
R-status				
R0 - resection	1 (reference)		1 (reference)	
R0 - resection with perforation	9.96 (5.51 to 17.99)	< 0.001	4.97 (2.55 to 9.71)	< 0.001
R1 - resection	3.61 (1.26 to 10.33)	0.017	3.26 (1.18 to 9.04)	0.023
^a Patients included those with stages I-III colon cancer during 1980–2016, treated with a major resection with curative intent (R0 and R1). Results are from a multivariable analysis; unadjusted: performed with one covariate at a time; adjusted: performed with all listed covariates included simultaneously. CCI: Charlson Comorbidity Index, classified as 0, 1, 2, or 3+				

Local recurrence was diagnosed in 4.4% (43/973) of patients. The overall estimated five-year local recurrence rate was 4.5% (95% CI: 3.7 to 5.3). The estimated five-year local recurrence rates after an R0 resection, an R1 resection, or a resection with tumour perforation were 4.3% (95% CI: 3.6 to 5.0), 43.2% (95% CI: 10.2 to 76.2), and 57.5% (95% CI: 19.1 to 95.9), respectively. The estimated five-year local recurrence rates were not affected by age.

Metastatic disease was diagnosed in 20% (195/973) of patients. The overall estimated five-year metastasis rate was 22.5% (95% CI: 19.5 to 25.5). The estimated five-year local metastasis rates after an R0 resection, an R1 resection, or a resection with tumour perforation were 21.2% (95% CI: 18.2 to 24.2), 45.5% (95% CI: 17.9 to 73.1), and 70.4% (95% CI: 47.2 to 93.6), respectively. The estimated five-year metastasis rates were not affected by age.

Chemotherapy

Starting in 1993, adjuvant chemotherapy was given to 53% (72/137) of patients under 75 years of age that underwent a major resection with curative intent for stage III disease. Among these patients, 28% (16/58) received adjuvant chemotherapy in 1993–2004, and 71% (56/79) received adjuvant chemotherapy in 2005–2016. Among patients aged 75–84 years, a selected group of 13% (11/85) received adjuvant chemotherapy. Among patients treated with a major resection with curative intent for stage II disease, 7% (15/214) received adjuvant chemotherapy.

Among patients that underwent palliative surgery or best supportive care, 34.5% (158/458) received palliative chemotherapy. This rate remained stable throughout the study period. The percentage of patients given palliative chemotherapy decreased as age increased. Palliative chemotherapy was given to 76% (79/104) of patients < 65 years, compared to 2.7% (4/148) of octogenarian patients.

Discussion

In this series, the rate of patients selected for surgical treatment decreased as patient age increased. Nevertheless, postoperative morbidity and 100-day mortality rates increased as patient age increased. During the study period, the percentage of octogenarian patients that underwent a major resection with curative intent increased, and the 100-day mortality was reduced. However, among patients that survived the first 100 days, long-term relative survival was independent of age.

All patients

Previous studies have pointed out age-related disparities in multimodal cancer treatments [9, 20]. In patients with colon cancer, individual treatment plans are based on accurate disease staging. In the first period (1980–1989) of the present study, we observed a transient trend towards a higher proportion of older patients with unknown disease stages. During the study period, we found significant progress in staging availability and precision, and focus was placed on the importance of preoperative staging, irrespective of patient age. Nevertheless, the proportion of patients with unknown stages among octogenarian patients in this series was low, compared to the proportions based on national data from several European countries [21]. Moreover, the disease stages at admission were equally distributed across the age groups, and the proportion of patients that presented with stage IV disease (25%) was comparable to proportions reported previously [22, 23].

Surgery is the cornerstone of colon cancer treatment. The primary objective of surgery is either radical resection or endoscopic resection, for early-stage tumours. Palliative surgery may be indicated as part of a multimodal treatment in patients with advanced disease or in cases with obstruction. Overall, the percentage of patients that underwent surgical treatments in this series was 89%. This percentage decreased as age increased. Surgery was performed in 93% of patients younger than 80 years and 82% of octogenarian patients. These findings were comparable to national data from European countries, where surgical treatment rates ranged between 59% and 79% among patients 80 years and older [21]. Variations in the overall rates of patients that undergo surgical treatment for colon cancer among different series are likely to depend on demographic, socioeconomic, and clinically related factors. The availability of healthcare services in our catchment area was high, and the threshold for referring patients to the hospital, irrespective of age, was low. However, because comorbidity increased with age, the rate of patients considered unsuitable for surgical treatment was relatively high among older patients.

The overall rate of patients that underwent emergency surgery in this series was 16%, and the rate increased with increasing age. Previous studies have shown significant variability (8–34%) in the rates of emergency surgery; these differences might be due to differences in the definition of emergency surgery and the selection of patient cohorts [24–26]. The rate of emergency surgery in this series was lower than the 25% reported previously, in a comparable population-based study from Sweden [27]. We observed that the rate of emergency surgery declined throughout the 37 years of the study. This finding might be related to a continuous increase in the availability of health care services, including the implementation of fast-track examinations, when alarm symptoms indicated colorectal cancer, and a higher societal awareness of this disease.

In parallel with the increases in population aging and the number of older patients admitted to hospital with colon cancer, the rate of octogenarian patients that underwent surgery increased. Hence, the proportion of octogenarian patients considered eligible for surgery has increased. A comparison of general health between the current and previous generations is difficult to assess objectively, and we lack evidence that older people in the current generation are healthier than those in previous generations [28, 29]. However, comorbid disease treatments and perioperative care have improved during the last few decades, and these advances have lowered the threshold for surgery [30–32].

The literature has shown variability in the rates of short-term mortality among patients with colon cancer. Clearly, differences in patient populations and differences in patient selection procedures for different treatment options, primarily surgical treatments, have major impacts on the outcome. In the present study, the overall 100-day mortality was 14.3%, and it increased, with increasing age, to 23.7% among octogenarian patients. These rates were comparable to rates reported in other unselected population-based series [26, 33]. We found that comorbidity, advanced TNM-stages, and emergency surgery had profound negative effects on the 100-day mortality. These associations were consistent with those demonstrated in previous reports [34, 35]. We noted a 50% reduction in the overall 100-day mortality rate, between the first and last decades of the observational period. The basis for this improvement was multifactorial, but it was driven by the general, continuous progress in medical treatments during the study period. Although we observed a significant increase in short-term mortality with increasing age, the long-term relative survival rates of young and old patient groups converged over time, and after five years, survival was independent of age. The five-year relative survival among all patients was 58.5%, comparable to rates reported in previous studies on unselected series of patients with colon cancer [36].

Patients with stages I-III disease that underwent a major resection with curative intent

Among patients with stages I-III disease at diagnosis, 92.6% (1021/1102) were treated with a major resection with curative intent, comparable to the proportions reported previously in studies on colon cancer [37]. Although the rate was lower among octogenarian patients (90.1%, 237/263), it was similar to the overall rate, which indicated that the approach to surgical treatment remained consistent, irrespective of age. During the first part of this study, the selection of patients for a major resection with curative intent was performed by a traditional interdisciplinary team, which included the surgeon and the anaesthesiologist. This selection was primarily based on a clinical evaluation combined with the ASA-score. Later, the focus changed, and treatment decisions were increasingly performed by multidisciplinary teams, which also included oncologists, radiologists, and pathologists [5].

The overall rate of postoperative morbidity, defined as a Clavien-Dindo score of 3 or more, was 9.6%, and the overall 100-day mortality was 4.7%. We observed a significant reduction in both postoperative morbidity and mortality during the study, and as in other series, we confirmed that high ASA scores and the need for emergency surgery had negative impacts on both endpoints. Moreover, high peri-operative blood loss increased the postoperative morbidity, which highlighted the importance of the surgical

technique [38]. Finally, preoperative anaemia was significantly associated with an increased risk of postoperative complications. In a previous meta-analysis by Fowler et al, preoperative anaemia was also associated with a poor postoperative outcome [39]. Accordingly, methods for detecting and treating preoperative anaemia would be beneficial.

The major challenge in treating colon cancer, which was noted in this series and confirmed by others, is the significant increase in postoperative morbidity and mortality with increasing age, even after a thorough patient selection process. In this series, octogenarian patients selected to undergo major curative surgery had a significantly increased risk of postoperative morbidity and mortality compared to younger patients. The mortality rate was 0.4% among patients aged < 65 years, and it increased by 25-fold, to 10.1%, in octogenarian patients.

Nevertheless, the five-year relative survival rate in this series was equivalent across age groups, consistent with findings in previous series [40, 41, 36, 42]. Among patients that survived 100 days after surgery, long-term survival was most significantly negatively impacted by the TNM stage, the R-status, and the presence of a tumour perforation [40, 41, 36, 42]. As observed previously [40, 41, 36, 42], the negative effect of emergency surgery persisted past the postoperative period. This finding highlighted the need to enhance the focus and follow-up for this group of patients.

As the population ages, octogenarian patients will become the most common group with colon cancer. Consequently, measures are needed to reduce the excess rates of postoperative morbidity and mortality among older patients. Increasing the focus on the process of selecting patients to different levels of treatment will be highly important, both for the individual patient and for the healthcare system. It is essential to perform geriatric assessments systematically in the preoperative work-up [43–45], pay attention to the concept of prehabilitation [46], and increase focus on patient preferences [47]. Recent reports have demonstrated the value of a geriatric assessment in summarizing the patient's degree of frailty and predicting postoperative morbidity and mortality for older patients with colon cancer [48]. The Society for Geriatric Oncology has recommended these assessments for all patients with cancer that are over 70 years of age [49]. In a systematic review, more than half of older patients with cancer were considered to be in a pre-frailty or frailty condition [50], and both these conditions were associated with adverse postoperative outcomes.

Most efforts to reduce postoperative morbidity and mortality rates have focused on the peri-operative and immediate postoperative statuses. Thus, the concept of prehabilitation prior to surgery has not gained sufficient attention. As part of this concept, the geriatric assessment evaluates several individual modifiable factors relevant to status optimization prior to surgery [51]. Moreover, a multidisciplinary team approach was shown to improve the postoperative outcome in frail patients [52]. Currently, an ongoing prospective multicentre study is examining multimodal prehabilitation for patients with colorectal cancer. Hopefully, those results will provide valuable information regarding the role of prehabilitation in the future management of older patients with cancer [46].

Numerous factors contribute to heterogeneity in the group of older patients with cancer. It is important to consider that personal patient preferences regarding treatment decisions might vary substantially among older patients. In the late stages of life, some needs, like preserving the remaining quality of life, may outweigh the need for radical treatment [47, 53]. It has been shown that the physician's recommendation was the most decisive factor in influencing the patient's decision [54]. That finding emphasized the importance of a thorough, and preferably evidence-based, foundation for the physician's advice.

Strengths and weaknesses

The main strength of this study was the transparent presentation of a consecutive, population-based cohort of patients with colon cancer that were treated in accordance with current evidence-based guidelines over a period of 37 years. Our institution was the primary hospital for a stable population throughout this extensive observational period, and thus, the cohort was suitable for evaluating trends over time. We believe that octogenarian patients with colon cancer will emerge as an important entity; thus, the results from this series provide important contributions to the current state of the field.

The main limitation of the study was its design. Due to its observational nature, we could not investigate causality. Moreover, the results may not be applicable to the older population, in general. Frail and unfit patients might not have been referred to our hospital, due to their clinical status. Finally, unknown or unrecorded confounders might have affected decisions regarding patient selection and treatment.

Conclusion

This study showed that octogenarian patients treated for colon cancer had adverse 100-day mortality rates, but among those that survived 100 days postoperatively, the long-term survival rate was equivalent to that of younger patients. The increasing fraction of older patients in years to come will become a major challenge in treating colon cancer. In addressing that challenge, early disease detection, followed by prehabilitation, a multidisciplinary approach with a geriatric assessment, and a meticulous post-operative follow up will be essential factors for improving treatment results and surmounting current standards.

Abbreviations

ASA: American Society of Anaesthesiology

CCI: Charlson Comorbidity Index

CI: Confidence interval

SD: Standard deviation

Declarations

Acknowledgments

We declare no conflicts of interest in connection with the current study.

Ethics approval and consent to participate

The Regional Committee for Medical and Health Research Ethics (REC) approved the study (2016/2172/REK midt).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Funding

This study was funded by the Department of Surgery, Levanger Hospital, Nord-Trøndelag Hospital Trust, Norway.

Availability of data and materials

The dataset used for this study is located on a secure server in the Levanger Hospital data system. The database was confirmed by comparing data with corresponding data in the Norwegian Cancer Registry 1980-2016.

Authors' contributions

ØH: Made substantial contributions to the design of the work, the interpretation of data, and drafting and revising the manuscript.

THE: Made substantial contributions to the design of the work, the analysis and interpretation of data, and drafting and revising the manuscript.

AX: Made substantial contributions to the design of the work and revising the manuscript.

SL: Made substantial contributions to the design of the work, the analysis and interpretation of data, and revising the manuscript.

BHE: Made substantial contributions to the design of the work, the analysis and interpretation of data, and drafting and revising the manuscript.

All authors read and approved the final manuscript.

References

1. Danckert B FJ EG, Hansen HL, Johannesen TB, Khan S, Køtlum JE, Ólafsdóttir E, Schmidt LKH, Virtanen A, Storm HH. NORDCAN: Cancer Incidence, Mortality, Prevalence and Survival in the Nordic Countries, Version 82 (26032019) Association of the Nordic Cancer Registries Danish Cancer Society <http://www-depiarcfr/NORDCAN/NO/frameasp>.
2. Norwegian Institute of Public Health (NIPH); Public Health Report - Health Status in Norway 2018 (2018).
3. Kunitake H, Zingmond DS, Ryoo J, Ko CY (2010) Caring for octogenarian and nonagenarian patients with colorectal cancer: what should our standards and expectations be? *Dis Colon Rectum* 53 (5):735–743. doi:10.1007/DCR.0b013e3181cdd658
4. Hoydahl O, Edna TH, Xanthoulis A, Lydersen S, Endreseth BH (2020) Long-term trends in colorectal cancer: incidence, localization, and presentation. *BMC Cancer* 20 (1):1077. doi:10.1186/s12885-020-07582-x
5. Helsedirektoratet (2019) Nasjonalt handlingsprogram med retningslinjer for diagnostikk, behandling og oppfølging av kreft i tykktarm og endetarm..
6. Pallis AG, Papamichael D, Audisio R, Peeters M, Folprecht G, Lacombe D, Van Cutsem E (2010) EORTC Elderly Task Force experts' opinion for the treatment of colon cancer in older patients. *Cancer Treat Rev* 36 (1):83–90. doi:10.1016/j.ctrv.2009.10.008
7. Millan M, Merino S, Caro A, Feliu F, Escuder J, Francesch T (2015) Treatment of colorectal cancer in the elderly. *World J Gastrointest Oncol* 7 (10):204–220. doi:10.4251/wjgo.v7.i10.204
8. Orimo H, Kamiya N (2008) [Redefining the concept of elderly—for "successful aging" society]. *Nihon Rinsho* 66 (8):1605–1614
9. Hardiman KM, Cone M, Sheppard BC, Herzig DO (2009) Disparities in the treatment of colon cancer in octogenarians. *Am J Surg* 197 (5):624–628. doi:10.1016/j.amjsurg.2008.12.018
10. Stornes T, Wibe A, Endreseth BH (2016) Complications and risk prediction in treatment of elderly patients with rectal cancer. *Int J Colorectal Dis* 31 (1):87–93. doi:10.1007/s00384-015-2372-x
11. Kang T, Kim HO, Kim H, Chun HK, Han WK, Jung KU (2015) Age Over 80 is a Possible Risk Factor for Postoperative Morbidity After a Laparoscopic Resection of Colorectal Cancer. *Ann Coloproctol* 31 (6):228–234. doi:10.3393/ac.2015.31.6.228

12. Charlson ME, Pompei P, Ales KL, MacKenzie CR (1987) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 40 (5):373–383. doi:10.1016/0021-9681(87)90171-8
13. Anesthesiologists ASo (2014) ASA Physical Status Classification System.
14. Nutritional anaemias. Report of a WHO scientific group (1968). *World Health Organ Tech Rep Ser* 405:5–37
15. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240 (2):205–213. doi:10.1097/01.sla.0000133083.54934.ae
16. Sobin LH, Wittekind C (2002) *TNM Classification of Malignant Tumours*, 6th ed. New York: Wiley
17. Wittekind C, Compton CC, Greene FL, Sobin LH (2002) TNM residual tumor classification revisited. *Cancer* 94 (9):2511–2516. doi:10.1002/cncr.10492
18. Coviello PWDE (2015) Estimating and modeling relative survival. *The Stata Journal* 15 (1):186–215
19. The Human Mortality Database. (2020). <http://www.mortality.org>.
20. Lee SM, Shin JS (2020) Colorectal Cancer in Octogenarian and Nonagenarian Patients: Clinicopathological Features and Survivals. *Ann Coloproctol* 36 (5):323–329. doi:10.3393/ac.2020.01.19.2
21. Vermeer NCA, Claassen YHM, Derks MGM, Iversen LH, van Eycken E, Guren MG, Mroczkowski P, Martling A, Johansson R, Vandendael T, Wibe A, Moller B, Lippert H, Portielje JEA, Liefers GJ, Peeters K, van de Velde CJH, Bastiaannet E (2018) Treatment and Survival of Patients with Colon Cancer Aged 80 Years and Older: A EURECCA International Comparison. *Oncologist* 23 (8):982–990. doi:10.1634/theoncologist.2017-0551
22. Benitez Majano S, Di Girolamo C, Rachet B, Maringe C, Guren MG, Glimelius B, Iversen LH, Schnell EA, Lundqvist K, Christensen J, Morris M, Coleman MP, Walters S (2019) Surgical treatment and survival from colorectal cancer in Denmark, England, Norway, and Sweden: a population-based study. *Lancet Oncol* 20 (1):74–87. doi:10.1016/S1470-2045(18)30646-6
23. Maringe C, Walters S, Rachet B, Butler J, Fields T, Finan P, Maxwell R, Nedrebo B, Pahlman L, Sjøvall A, Spigelman A, Engholm G, Gavin A, Gjerstorff ML, Hatcher J, Johannesen TB, Morris E, McGahan CE, Tracey E, Turner D, Richards MA, Coleman MP, Group IMW (2013) Stage at diagnosis and colorectal cancer survival in six high-income countries: a population-based study of patients diagnosed during 2000–2007. *Acta Oncol* 52 (5):919–932. doi:10.3109/0284186X.2013.764008
24. Ellis L, Coleman MP, Rachet B (2012) How many deaths would be avoidable if socioeconomic inequalities in cancer survival in England were eliminated? A national population-based study, 1996–2006. *Eur J Cancer* 48 (2):270–278. doi:10.1016/j.ejca.2011.10.008
25. Rametta S, Grosso G, Galvano F, Mistretta A, Marventano S, Nolfo F, Buscemi S, Gangi S, Basile F, Biondi A (2013) Social disparities, health risk behaviors, and cancer. *BMC Surg* 13 Suppl 2:S17. doi:10.1186/1471-2482-13-S2-S17

26. Sjo OH, Larsen S, Lunde OC, Nesbakken A (2009) Short term outcome after emergency and elective surgery for colon cancer. *Colorectal Dis* 11 (7):733–739. doi:10.1111/j.1463-1318.2008.01613.x
27. Jestin P, Nilsson J, Heurgren M, Pahlman L, Glimelius B, Gunnarsson U (2005) Emergency surgery for colonic cancer in a defined population. *Br J Surg* 92 (1):94–100. doi:10.1002/bjs.4780
28. WHO (2018) Ageing and health. WHO. <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
29. (NIPH) NloPH (2016) Health among the elderly in Norway.
30. Kerstjens HA, Groen HJ, van Der Bij W (2001) Recent advances: Respiratory medicine. *BMJ* 323 (7325):1349–1353. doi:10.1136/bmj.323.7325.1349
31. Grover FL, Cleveland JC, Jr., Shroyer LW (2002) Quality improvement in cardiac care. *Arch Surg* 137 (1):28–36. doi:10.1001/archsurg.137.1.28
32. Fowler AJ (2013) A Review of Recent Advances in Perioperative Patient Safety. *Ann Med Surg (Lond)* 2 (1):10–14. doi:10.1016/S2049-0801(13)70020-7
33. Niemelainen S, Huhtala H, Ehrlich A, Kossi J, Jamsen E, Hyoty M (2020) Risk factors of short-term survival in the aged in elective colon cancer surgery: a population-based study. *Int J Colorectal Dis* 35 (2):307–315. doi:10.1007/s00384-019-03488-8
34. Howard R, Yin YS, McCandless L, Wang S, Englesbe M, Machado-Aranda D (2019) Taking Control of Your Surgery: Impact of a Prehabilitation Program on Major Abdominal Surgery. *J Am Coll Surg* 228 (1):72–80. doi:10.1016/j.jamcollsurg.2018.09.018
35. Soyalp C, Yuzkat N, Kilic M, Akyol ME, Demir CY, Gulhas N (2019) Operative and prognostic parameters associated with elective versus emergency surgery in a retrospective cohort of elderly patients. *Aging Clin Exp Res* 31 (3):403–410. doi:10.1007/s40520-018-0976-z
36. Niemelainen S, Huhtala H, Ehrlich A, Kossi J, Jamsen E, Hyoty M (2020) Long-term survival following elective colon cancer surgery in the aged. A population-based cohort study. *Colorectal Dis* 22 (11):1585–1596. doi:10.1111/codi.15242
37. Brouwer NPM, Bos A, Lemmens V, Tanis PJ, Huguen N, Nagtegaal ID, de Wilt JHW, Verhoeven RHA (2018) An overview of 25 years of incidence, treatment and outcome of colorectal cancer patients. *Int J Cancer* 143 (11):2758–2766. doi:10.1002/ijc.31785
38. Morner M, Gunnarsson U, Jestin P, Egenvall M (2015) Volume of blood loss during surgery for colon cancer is a risk determinant for future small bowel obstruction caused by recurrence—a population-based epidemiological study. *Langenbecks Arch Surg* 400 (5):599–607. doi:10.1007/s00423-015-1317-8
39. Fowler AJ, Ahmad T, Phull MK, Allard S, Gillies MA, Pearse RM (2015) Meta-analysis of the association between preoperative anaemia and mortality after surgery. *Br J Surg* 102 (11):1314–1324. doi:10.1002/bjs.9861
40. Weerink LBM, Gant CM, van Leeuwen BL, de Bock GH, Kouwenhoven EA, Faneyte IF (2018) Long-Term Survival in Octogenarians After Surgical Treatment for Colorectal Cancer: Prevention of

- Postoperative Complications is Key. *Ann Surg Oncol* 25 (13):3874–3882. doi:10.1245/s10434-018-6766-1
41. Mulcahy HE, Patchett SE, Daly L, O'Donoghue DP (1994) Prognosis of elderly patients with large bowel cancer. *Br J Surg* 81 (5):736–738. doi:10.1002/bjs.1800810540
 42. Mothes H, Bauschke A, Schuele S, Eigendorff E, Altendorf-Hofmann A, Settmacher U (2017) Surgery for colorectal cancer in elderly patients: how can we improve outcome? *J Cancer Res Clin Oncol* 143 (9):1879–1889. doi:10.1007/s00432-017-2438-y
 43. Ellis G, Gardner M, Tsiachristas A, Langhorne P, Burke O, Harwood RH, Conroy SP, Kircher T, Somme D, Saltvedt I, Wald H, O'Neill D, Robinson D, Shepperd S (2017) Comprehensive geriatric assessment for older adults admitted to hospital. *Cochrane Database Syst Rev* 9:CD006211. doi:10.1002/14651858.CD006211.pub3
 44. Dogrul RT, Dogrul AB, Konan A, Caglar O, Sumer F, Caliskan H, Kizilarlanoglu MC, Kilic MK, Balci C, Arik G, Aycicek GS, Ozsurekci C, Halil M, Cankurtaran M, Yavuz BB (2020) Does Preoperative Comprehensive Geriatric Assessment and Frailty Predict Postoperative Complications? *World J Surg* 44 (11):3729–3736. doi:10.1007/s00268-020-05715-8
 45. participants P, Audisio RA, Pope D, Ramesh HS, Gennari R, van Leeuwen BL, West C, Corsini G, Maffezzini M, Hoekstra HJ, Mobarak D, Bozzetti F, Colledan M, Wildiers H, Stotter A, Capewell A, Marshall E (2008) Shall we operate? Preoperative assessment in elderly cancer patients (PACE) can help. A SIOG surgical task force prospective study. *Crit Rev Oncol Hematol* 65 (2):156–163. doi:10.1016/j.critrevonc.2007.11.001
 46. van Rooijen S, Carli F, Dalton S, Thomas G, Bojesen R, Le Guen M, Barizien N, Awasthi R, Minnella E, Beijer S, Martinez-Palli G, van Lieshout R, Gogenur I, Feo C, Johansen C, Scheede-Bergdahl C, Roumen R, Schep G, Slooter G (2019) Multimodal prehabilitation in colorectal cancer patients to improve functional capacity and reduce postoperative complications: the first international randomized controlled trial for multimodal prehabilitation. *BMC Cancer* 19 (1):98. doi:10.1186/s12885-018-5232-6
 47. Niemeyer-Guimaraes M, Schramm FR (2017) The Exercise of Autonomy by Older Cancer Patients in Palliative Care: The Biotechnoscientific and Biopolitical Paradigms and the Bioethics of Protection. *Palliat Care* 9:1178224216684831. doi:10.1177/1178224216684831
 48. Rostoft S (2017) Integration of Geriatric Assessment in the Care of Patients with Gastrointestinal Malignancies. *Visc Med* 33 (4):275–280. doi:10.1159/000475452
 49. Wildiers H, Heeren P, Puts M, Topinkova E, Janssen-Heijnen ML, Extermann M, Falandry C, Artz A, Brain E, Colloca G, Flamaing J, Karnakis T, Kenis C, Audisio RA, Mohile S, Repetto L, Van Leeuwen B, Milisen K, Hurria A (2014) International Society of Geriatric Oncology consensus on geriatric assessment in older patients with cancer. *J Clin Oncol* 32 (24):2595–2603. doi:10.1200/JCO.2013.54.8347
 50. Handforth C, Clegg A, Young C, Simpkins S, Seymour MT, Selby PJ, Young J (2015) The prevalence and outcomes of frailty in older cancer patients: a systematic review. *Ann Oncol* 26 (6):1091–1101.

doi:10.1093/annonc/mdu540

51. Ronning B, Wyller TB, Jordhoy MS, Nesbakken A, Bakka A, Seljeflot I, Kristjansson SR (2014) Frailty indicators and functional status in older patients after colorectal cancer surgery. *J Geriatr Oncol* 5 (1):26–32. doi:10.1016/j.jgo.2013.08.001
52. van der Vlies E, Smits AB, Los M, van Hengel M, Bos WJW, Dijksman LM, van Dongen EPA, Noordzij PG (2020) Implementation of a preoperative multidisciplinary team approach for frail colorectal cancer patients: Influence on patient selection, prehabilitation and outcome. *J Geriatr Oncol* 11 (8):1237–1243. doi:10.1016/j.jgo.2020.04.011
53. Rostoft S, van den Bos F, Pedersen R, Hamaker ME (2021) Shared decision-making in older patients with cancer - What does the patient want? *J Geriatr Oncol* 12 (3):339–342. doi:10.1016/j.jgo.2020.08.001
54. Puts MT, Tapscott B, Fitch M, Howell D, Monette J, Wan-Chow-Wah D, Krzyzanowska M, Leighl NB, Springall E, Alibhai SM (2015) A systematic review of factors influencing older adults' decision to accept or decline cancer treatment. *Cancer Treat Rev* 41 (2):197–215. doi:10.1016/j.ctrv.2014.12.010

Figures





Figure 1

Survival of patients with colon cancer during 1980-2016. (a) 5-year relative survival for all patients in each age group; table columns represent the number of patients at risk at surgery (time = 0) and every 1

year thereafter. (b) Long term relative survival for all patients, classified by treatment intent; table columns represent the number of patients at risk at surgery (time = 0) and every 2.5 years thereafter.



< 65 years	232	216	186	166	139	121	104	78	54	44	31	23	14
65 - 74 years	319	277	206	157	119	78	54	36	26	15	10	5	1
75 - 79 years	194	162	122	83	54	36	18	11	3	1	1		
80 - 84 years	156	130	90	60	32	15	5						
85 - 89 years	57	47	26	10	6	2							
90 years +	15	13	8	2	0								

Figure 2

Relative survival after colon cancer resection with curative intent in patients that survived 100 days. Relative survival is stratified by age group. Table columns represent the number of patients at risk at surgery (time = 0) and every 2.5 years thereafter.