Effectiveness of Allied Health Clinical Supervision Following the Implementation of an Organisational Framework

Marcus Gardner (✉ mgardner@bendigohealth.org.au)

La Trobe University

Carol McKinstry

La Trobe University

Byron Perrin

La Trobe University

Research Article

Keywords: Clinical supervision, allied health workforce, MCSS-26, ANOVA

Posted Date: September 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-914459/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on February 26th, 2022. See the published version at <https://doi.org/10.1186/s12913-022-07636-9>.

Abstract

Background: Clinical supervision makes an important contribution to high quality patient care and professional wellbeing for the allied health workforce. However, there is limited research examining the longitudinal implementation of clinical supervision for allied health. The aim of this study was to determine the effectiveness of clinical supervision for allied health at a regional health service and clinicians' perceptions of the implementation of an organisational clinical supervision framework.

Methods: A cross-sectional study was conducted as a phase of an overarching participatory action research study. The MCSS-26 tool was used to measure clinical supervision effectiveness with additional open-ended questions included to explore the implementation of the clinical supervision framework. MCSS-26 findings were compared with an initial administration of the MCSS-26 5 years earlier. MCSS-26 data (total scores, summed domain and sub-scale scores) were analysed descriptively and reported as mean and standard deviation values. Differences between groups were analysed with independent-samples t-test (t) and one-way between groups ANOVA.

Results: There were 125 responses to the survey (response rate 50%). The total MCSS-26 score was 78.5 (S.D. 14.5). The total MCSS-26 score was unchanged compared with the initial administration. There was a statistically significant difference in clinical supervision effectiveness between speech pathology and physiotherapy ($F = 2.9$, $p = 0.03$) and higher MCSS-26 scores for participants whose clinical supervisor was a senior clinician and those who chose their clinical supervisor. Seventy percent of participants perceived that the organisation's clinical supervision framework was useful and provided structure and consistent expectations for clinical supervision.

Conclusions: Clinical supervision was effective for allied health in this regional setting and clinical supervision effectiveness was maintained over a 5 year period. The implementation of an organisational clinical supervision framework may have positively impacted on the effectiveness of clinical supervision for some professions.

Background

Allied health clinicians play a vital role in the provision of high quality patient care across the continuum of care in health care systems (1, 2). The allied health workforce are a significant proportion of the health workforce, ranking as the largest workforce behind nursing, and also medicine in some settings (3). Professions included as allied health disciplines differs across settings and countries. The professionals commonly described as allied health include physiotherapy, occupational therapy, speech pathology, podiatry, dietetics, exercise physiology (4, 5). Additionally, in some contexts social work, psychology and allied health assistants are also defined as allied health (6). Across the majority of allied health professions clinical supervision is practised to enhance the professional development of clinicians and high quality patient care (7).

Clinical supervision has been defined as “the formal provision, by approved supervisors, of a relationship-based education and training that is work-focused and which manages, supports, develops and evaluates the work of colleague/s” (8^{p. 440}). Clinical supervision supports effective clinical governance and the professional wellbeing of allied health clinicians (7). In conjunction with other forms of professional support, clinical supervision positively impacts on allied health workforce recruitment and retention, particularly in non-metropolitan settings (9, 10). While clinical supervision is widely used by allied health professionals, there can be difficulties accessing clinical supervisors, inconsistent approaches to clinical supervision across different professions and settings and variable effectiveness of clinical supervision across allied health professions (11–14). Other challenges include poor translation of best practice clinical supervision approaches into practice, such as provision for supervisees to choose their clinical supervisor, and inadequate training of clinical supervisors (15, 16).

Clinical supervision frameworks have been recommended as a policy strategy to improve the effectiveness of clinical supervision for allied health (17–21). Allied health clinical supervision frameworks can facilitate a common understanding of clinical supervision, agreed roles and responsibilities, and promote consistent approaches (17). Structured frameworks could also assist health care organisations to evaluate the implementation of clinical supervision and identify areas for improvement (17). Multidisciplinary allied health clinical supervision guidelines within health service organisations can influence the quality of clinical supervision by promoting the formation of safe and effective supervisory relationships that focus on the professional development of the supervisee (21). However, internationally, few allied health professions and jurisdictions have developed clinical supervision frameworks and most have been reported to be of low quality (22). Additionally, there is limited longitudinal research measuring the impact of implemented clinical supervision frameworks (7).

In 2015, a regional health service in Victoria, Australia introduced an organisational clinical supervision framework for allied health. An action research group, comprising of allied health managers at the health service, was established to oversee the implementation and evaluation of the framework. The action research group informed the cycles of planning, action and reflection as a part of the framework implementation. The framework included a multidisciplinary clinical supervision protocol which outlined expectations for clinical supervision and roles and responsibilities. Interprofessional training was conducted to support the implementation of the framework.

When the framework was implemented in 2015, an initial survey was conducted. The findings revealed that while clinical supervision was perceived to be effective overall, there were significant differences in perceived effectiveness between some disciplines (14). To measure the longer-term perceived effectiveness of clinical supervision and the clinical supervision framework, a repeat survey was planned and undertaken.

The aim of this study was to determine the effectiveness of clinical supervision for allied health at an Australian regional health service and to contrast the findings with those of an earlier survey. A secondary aim was to explore clinicians’ perceptions of the organisational clinical supervision framework.

Methods

Research design

A cross-sectional study was conducted as a phase of an overarching participatory action research study. The findings of the survey reported in this paper were compared with the findings of an initial administration of the MCSS-26 survey at the health service 5 years prior (14). Additional questions were included in the later survey to explore the impact of the organisation's introduction of the clinical supervision framework and the COVID-19 pandemic on clinical supervision, identifying opportunities for future improvement of clinical supervision practice. Consistent with the action research design, the action research group assisted with the interpretation of findings (23). Ethics approval was provided by Bendigo Health and La Trobe University Ethics Committees (Approval number LNR/15/BHCG/26). All methods were carried out in accordance with the relevant guidelines and regulations and all participants provided informed consent.

Participants

Participants were allied health professionals and allied health assistants employed at a large regional Australian health service. The allied health professions included were physiotherapy, occupational therapy, social work, dietetics, speech pathology, podiatry, exercise physiology, psychology, audiology and allied health assistants. Participants included a range of staff experience levels and were employed across hospital, community and mental health settings. The survey was distributed electronically by the Director of Allied Health in July 2020 and was open for 5 weeks. To be eligible, participants had to be receiving clinical supervision at the health service for at least 12 months. Participation in the survey was voluntary and data were non-identifiable. Data from professions with less than ten responses were aggregated to reduce the risk of participant identification.

Outcome measures

To measure perceived effectiveness of clinical supervision, the MCSS-26 tool was used (24). The MCSS-26 has been validated for use by allied health professionals, has been found to have high test-retest reliability and was the same tool used in the 2015 study (24). The MCSS-26 is a self-completion tool, measuring the perceived effectiveness of clinical supervision from the supervisee's perspective (25). A summary of the MCSS-26 domains and sub-scales is provided in Table 1. The MCSS-26 tool contains 26 questions which are rated on a five point scale from strongly disagree (0) to strongly agree (4). The possible range of scores is between 0 and 104, with higher scores aligning with a higher level of perceived clinical supervision effectiveness. The developers of the MCSS-26 have recommended that a threshold score of ≥ 73 is indicative of effective clinical supervision (26). The MCSS-26 also contains questions relating to the supervisee, their clinical supervision and the clinical supervision session characteristics. Additional closed and open-ended survey questions were included to explore factors potentially influencing the effectiveness of clinical supervision, the perceived impact of the

implementation of the health service’s clinical supervision framework and suggestions for improving clinical supervision practice.

Table 1
Description of MCSS-26 domains and sub-scales (25)

Normative domain <ul style="list-style-type: none">• <i>Importance/value of clinical supervision</i>: the importance of receiving clinical supervision to improve patient care• <i>Finding time</i>: the time available for supervisees to participate in clinical supervision
Formative domain <ul style="list-style-type: none">• <i>Trust/rapport</i>: trust with the supervisor and the ability to raise sensitive or confidential issues• <i>Supervisor advice/support</i>: the level of support and guidance provided by the clinical supervisor
Restorative domain <ul style="list-style-type: none">• <i>Improved care/skills</i>: the extent to which clinical supervision improves skills and quality of care• <i>Reflection</i>: support for the supervisee to reflect on complex issues

Data analysis

All available survey data were analysed. Frequency data for valid responses were reported as raw numbers and percentages. MCSS-26 data (total scores, summed domain and sub-scale scores) were analysed descriptively and reported as mean and standard deviation values. Differences between groups were analysed with independent-samples t-test (t) and one-way between groups ANOVA. Due to small numbers, some profession categories were collapsed for comparison of groups. Sub-group analysis included variables of interest informed by the action research group, including supervisor grade level, supervisor allocation, clinical supervision frequency and use of clinical supervision agreements. Total MCSS-26 scores for allied health and each profession in this study were compared descriptively with data from the 2015 study. All quantitative analyses were performed using SPSS version 26. Open ended survey question data were thematically coded using content analysis by two researchers independently and then discussed together to reach consensus (26).

Results

Responses

One hundred and twenty-five allied health clinicians responded to the survey, with a response rate of 50.0% of eligible participants.

Participant characteristics

The responding participants were predominantly female (n = 102; 81.6 %) and the mean age was 38.2 years (S.D. 11.5). Over half of the participants were in grade 2 positions (n = 67; 53.6%), early to mid-career clinicians with generally 2 to 7 years of experience. Nearly a quarter were at grade 3 or 4 levels (n = 30; 24.4%), mainly senior clinicians with more than 7 years of experience. Approximately one sixth were in grade 1 positions (n = 19; 15.2%), graduates and early career clinicians generally with up to 2 years' experience. There was a similar number of participants working in hospital (n = 48; 38.4 %) and community settings (n = 51; 40.8 %), with the remainder working across both. Approximately half of the participants had attended clinical supervision training (n = 61; 48.8 %). Participant characteristics are summarised in Table 2.

Table 2
Supervisee characteristics (n = 125)

	No. of participants	%
Sex	18	14.4
Male	102	81.6
Female	5	4.0
Missing		
Staff Grade[^]	19	15.2
Grade 1	67	53.6
Grade 2	30	24.0
Grade 3 and 4	3	2.4
Management	1	0.8
Other	5	4.0
Missing		
Discipline	36	28.8
Occupational therapy	33	26.4
Physiotherapy	22	17.6
Social work	7	5.6
Dietetics	10	8.0
Speech pathology	2	1.6
Psychology	3	2.4
Podiatry	4	3.2
Exercise physiology	1	0.7
Other	2	1.5
allied health Assistants	5	4.0
Missing		

	No. of participants	%
Work setting	48	38.4
Hospital	51	40.8
Community	21	16.8
Both	5	4.0
Missing		
Supervisor allocated or chosen	104	83.2
Allocated	12	9.6
Chosen	3	2.4
Other	6	4.8
Missing		
Attended training	61	48.8
Yes	56	44.8
No	8	6.4
Missing		

Clinical supervision session characteristics

Over half of the participants' clinical supervisors were senior clinicians (grade 3 or 4) (n = 64; 52.5%) while approximately a quarter were in management (n = 29; 23.8%) or grade 2 positions (n = 26; 21.3%). Most participants had clinical supervisors who were allocated to them (n = 104; 83.2 %). Clinical supervision sessions mostly occurred monthly, within the workplace and for under one hour. Most participants received individual clinical supervision (n = 115; 83.3 %), with a smaller number participating in group clinical supervision (n = 14; 10.7%) and 75% had a clinical supervision agreement in place (n = 94). Characteristics of supervision sessions are provided in Table 3.

Table 3
Characteristics of supervision sessions (n = 125)

	No. of participants	%
Frequency of supervision sessions	1	0.8
Weekly	12	9.6
Fortnightly	82	65.6
Monthly	18	14.4
2–3 months	6	4.8
>3 months	6	4.8
Missing		
Location of supervision session	99	79.2
Within the workplace	6	4.8
Away from the workplace	14	11.2
Both	6	4.8
Missing		
Type of supervision	115	83.3
One to one	3	2.2
Triad	14	10.7
Group	6	4.3
Other		
Duration of supervision sessions	1	0.8
<15 mins	8	6.4
15–30 mins	12	8.0
31–45 mins	84	67.2
46–60 mins	18	12.0
>60 mins	2	1.6
Missing		

	No. of participants	%
Clinical supervision agreement in place	94	75.2
Yes	23	18.4
No	8	6.4
Missing		

MCSS-26 results

Clinical supervision was perceived by participants to be effective overall (Table 4). The average total MCSS-26 score across all professions was 78.5 (S.D. 14.5). Scores in the *formative* domain were highest of the three domains, with the sub-scale relating to *reflection* scoring highest, indicating that participants perceived that clinical supervision assisted them to reflect and learn from practice issues. The lowest domain score was for the *normative* domain. Of the two sub-scales in the *normative* domain, there was a relatively high score for the *importance/value* sub-scale and a lower score for *finding time* sub-scale indicating that while participants valued clinical supervision, they found it difficult to make time for it.

Table 4
MCSS-26 total, domain and sub-scale scores for all allied health professions

MCSS-26 scores	No. of items	Possible range	Mean (S.D.)	Mean (out of 100) (+/- S.D.) ¹
Normative domain	9	0–36	26.1 (5.4)	72.5 (5.2)
Importance/value of clinical supervision	5	0–20	16.3 (2.9)	81.5 (2.8)
Finding time	4	0–16	9.8 (3.3)	61.2 (3.2)
Formative domain	7	0–28	21.8 (4.5)	77.9 (4.3)
Improved care/skills	4	0–16	12.2 (2.9)	76.2 (2.8)
Reflection	3	0–12	9.6 (2.1)	80.0 (2.0)
Restorative domain	10	0–40	30.6 (6.9)	76.5 (6.6)
Trust/support	5	0–20	15.7 (3.2)	78.5 (3.1)
Supervisor support/advice	5	0–20	14.8 (4.2)	74.0 (4.0)
Total	26	0-104	78.5 (14.5)	75.5 (13.9)

¹ Mean expressed out of 100 to enable comparison across domains and subscales

Participants whose clinical supervisor was at a grade 3 or 4 level (82.0; S.D. 12.4) had higher MCSS-26 scores than those whose clinical supervisor was a manager (73.7; S.D. 17.2, $F = 2.8$, $p = 0.04$). There were higher MCSS-26 scores for those who had input into choosing their clinical supervisors (86.6; S.D. 11.58) than those who did not (78.2; S.D. 13.8, $t = -2.3$, $p = 0.047$). Those who had clinical supervision monthly or more frequently (80.7; S.D. 14.2) had higher scores than those who had clinical supervision every 2 months or more (71.5; S.D. 9.7, $F = 4.7$, $p = 0.01$). Attendance at training and use of clinical supervision agreements did not influence MCSS-26 scores.

Scores were higher for speech pathologists (88.5; S.D. 10.2), social workers (81.1; S.D. 16.7) and occupational therapists (80.4, S.D. 10.2) compared to physiotherapists (74.9; S.D. 8.9) and clinicians in the “other” group of professions (73.9; S.D. 13.9). ANOVA analysis for the impact of the supervisee’s profession on MCSS-26 scores demonstrated a statistically significant difference between speech pathology and physiotherapy ($F = 2.9$, $p = 0.03$). There were no other significant differences between professional groups.

Impact of organisational clinical supervision framework

Seventy percent of participants who responded agreed or strongly agreed that the organisation's clinical supervision framework had been useful. Twenty-two percent were neutral while 8% disagreed. Themes from open-ended responses relating to perceived impact of the framework included the framework *provides structure and consistent expectations*, *lack of awareness* of the framework, and the need for *framework modifications*. Some participants indicated that the framework *provides structure and consistent expectations* for clinical supervision practice, while in contrast others had a *lack of awareness* of the framework or perceived that it had made no change. Some responses indicated the need for *framework modifications*, such as redesigning clinical supervision documentation templates.

Considering future improvements to the organisational clinical supervision framework, themes associated with responses included *no changes*, *choice of clinical supervisor*, *group or peer models*, *access to clinical supervision education and training*, and *clarity around the purpose of clinical supervision*. The themes arising from those who proposed improvements were: having input into the *choice of clinical supervisor*, increased opportunities for *group or peer models* of clinical supervision; increased *access to clinical supervision education and training*, and improved *clarity around the purpose of clinical supervision*, particularly for senior clinicians and those in specialist roles.

Comparison of 2020 results with 2015 results

The total MCSS-26 mean score for all allied health professions did not change between 2015 (78.5, S.D. 13.9) and 2020 (78.5, S.D. 14.5) results (see Table 5). The restorative domain score increased from 2015 (30.0, S.D. 7.3) to 2020 (30.6, S.D. 6.9) and the normative domain score decreased from 2015 (26.6, S.D. 4.7) to 2020 (26.1, S.D. 5.4). The formative domain mean score was unchanged. The total MCSS-26 score for physiotherapy increased from 2015 (70.9, S.D. 11.3) to 2020 (74.9, S.D. 8.9).

Table 5
Comparison of MCSS-26 total, domain and sub-scale scores for all disciplines, 2015 and 2020

	2015 survey, mean (SD)	2020 survey, mean (SD)
Normative domain	26.6 (4.7)	26.1 (5.4)
<i>Importance/value of clinical supervision</i>	16.9 (2.4)	16.3 (2.9)
<i>Finding time</i>	9.7 (3.2)	9.8 (3.3)
Formative domain	21.8 (4.6)	21.8 (4.5)
<i>Improved care/skills</i>	12.4 (2.8)	12.2 (2.9)
<i>Reflection</i>	9.4 (2.3)	9.6 (2.1)
Restorative domain	30.0 (7.3)	30.6 (6.9)
<i>Trust/support</i>	15.1 (3.6)	15.7 (3.2)
<i>Supervisor support/advice</i>	14.9 (4.3)	14.8 (4.2)
Total	78.5 (13.9)	78.5 (14.5)

Remaining professional group scores were similar between 2015 and 2020, apart from the “other profession” group which decreased from 2015 (83.6, S.D. 12.2) to 2020 (73.9, S.D. 13.9), however, there were different professions included in the “other profession” group in 2015 and 2020. Table 6 provides a comparison MCSS-26 scores for discipline groups between the two administrations of the survey.

Table 6
Total MCSS-26 scores for individual allied health professions, 2015 and 2020

Discipline	Mean (SD), 2015	Mean (SD), 2020
Social work	81.3 (13.6)	81.1 (16.7)
Physiotherapy	70.9 (11.3)	74.9 (8.9)
Occupational therapy	82.8 (14.4)	80.4 (10.2)
Speech pathology ¹	-	88.5 (10.2)
Dietetics ²	70.4 (12.0)	-
Other professions ³	83.6 (12.2)	73.9 (13.9)
Allied health	78.5 (13.9)	78.5 (14.5)
¹ insufficient number of responses in 2015 to analyse speech pathology separately (included in “other professions” group in 2015)		
² insufficient number of responses in 2020 to analyse dietetics separately (included in “other professions” group in 2020)		
³ other profession group in 2015 included: speech pathology, psychology, podiatry, exercise physiology and allied health assistance; other profession group in 2020 included: dietetics, psychology, podiatry, exercise physiology and allied health assistance		

Discussion

Clinical supervision was perceived by supervisees to be effective overall. All professions with more than ten responses had MCSS-26 scores above the threshold score for effective clinical supervision of 73. There was variation in the effectiveness of clinical supervision across allied health professions. While the overall results were similar to those reported in other allied health studies using the MCSS-26, the score for speech pathology was higher than in previous studies (13, 28, 29). This novel finding, contrasts with previous studies, where the professions with the most effective clinical supervision were those counselling-based professions with more established traditions of clinical supervision, such as social work and psychology (13, 28). The lower score for physiotherapy is consistent with several other studies (13, 17, 18). This may be explained by other research findings where physiotherapists reported greater satisfaction when direct models of clinical supervision are used rather than a mix of direct and reflective clinical supervision (30, 31) which was the predominant model used by physiotherapists in this study.

The effectiveness of clinical supervision in this study was influenced by the level of experience and seniority of the clinical supervisor. Clinical supervision was most effective when provided by a senior clinician (grade 3) and least effective when provided by a manager. This finding builds on existing evidence that clinical supervision should be separated from line management to prevent blurring the boundaries of these functions and clinical supervision becoming overly focused on administrative functions (15, 21, 32–34). Additionally, there is a power differential when a manager is the clinical supervisor which may result in “supervisee guarding” where supervisees avoid discussing issues around work skills and performance for fear of being viewed as incompetent (35).

Managers should also participate in clinical supervision that is separated from operational accountability to enable them to access professional support and maintain and develop their own clinical supervision skills. Strategies suggested by participants in this study to address this issue included implementing peer group models of clinical supervision for senior clinicians which could ensure that clinical supervision is separated from operational issues. Another potential strategy was exploring cross-organisational models of clinical supervision for senior clinicians and clinical managers. This could enable experienced clinicians to source clinical supervision that is relevant to their clinical speciality or role and promote sharing of evidence-based practice between organisations. Facilitating cross-organisational approaches would require the development of coordinated state or jurisdictional clinical supervision policies to address issues such as establishing registers of clinical supervisors and cost-sharing between organisations.

In this study, another factor associated with increased effectiveness of clinical supervision was where supervisees had chosen their clinical supervisor. The theme of clinicians wanting to have input into choosing their supervisor was present in responses from participants suggesting improvements for clinical supervision at the health service. This finding is consistent with the recommendations made by other researchers as best practice clinical supervision (36, 37). Aside from social workers who had a process for enabling supervisees to choose supervisors, the health service’s model of supervision was based on hierarchical allocation of supervisors within the same profession and speciality. This is a similar model of supervisor allocation reported within health services in other allied health clinical supervision studies (13, 28). When discussing this finding, the action research group determined that it was not practical to enable all staff to choose their supervisor within the existing allied health workforce structure in this health service. This perspective is supported by a study involving allied health in community health settings that reported that workforce structures containing insufficient numbers of senior clinicians was a barrier to accessing and allocating clinical supervisors (11). However, it may be possible for some clinicians, such as experienced clinicians or those who have had the same supervisor for an extended period, to have input into the choice of their supervisor.

The overall MCSS-26 score in this study for the health service was the same as the score reported in the 2015 study, indicating that the overall effectiveness of clinical supervision was maintained during the 5 year period between surveys. There were minimal changes in the domain and sub-scale scores with the sub-scales relating to the importance of clinical supervision and its role in assisting reflection scoring

higher in both surveys and finding time for clinical supervision scoring the lowest in both. The lack of change in overall effectiveness of clinical supervision, despite the increased focus on clinical supervision, could be associated with clinicians having a greater awareness of the benefits of clinical supervision and therefore having higher expectations for clinical supervision.

For some individual professions there were changes in scores between surveys. While the 2015 study did not include enough responses from speech pathology to enable this profession to be analysed individually, the high score for speech pathology in 2020 was notable. There was also a modest increase in the score for physiotherapy. Neither of these professions had structured frameworks for clinical supervision in place prior to the implementation of the organisational framework in 2015, which may have contributed to the increased effectiveness of clinical supervision for speech pathology and physiotherapy.

Participants perceived that the organisational framework specified expectations for clinical supervision and provided a structured approach. In contrast, social work and occupational therapy, who did have existing frameworks in place prior to the introduction of the organisational framework, had no change or a slight decrease in their MCSS-26 scores. Previous studies have recommended that structured frameworks for clinical supervision are required to improve the quality of clinical supervision (17, 18). However, this study is the first that the authors are aware of to provide a longitudinal comparison of the impact of implementing a clinical supervision framework for allied health, demonstrating that such initiatives can have a positive impact on clinical supervision. This finding strengthens the need for future policy direction to focus on the development of common allied health clinical supervision frameworks to facilitate consistent approaches to clinical supervision implementation and education.

Whilst the response rate in this study was 50%, it was within the range of response rates reported in other studies using the MCSS-26 survey (16, 28, 29). However, the low number of responses in some professions restricted the analysis of comparison between groups and there may have been a positivity bias amongst those who responded. Additionally, the ability to draw conclusions when comparing the 2015 and 2020 survey findings was limited as a formal pre-post design was not used, consistent with recommendations for the use of MCSS-26 and an action research approach.

Conclusion

Clinical supervision was effective for allied health in this regional setting and effectiveness been maintained over a 5 year period. Strategies to improve clinical supervision practice in the future include providing opportunities for supervisees to choose their clinical supervisor and introducing peer group supervision for experienced clinicians. The implementation of an organisational clinical supervision framework was perceived to be useful and may have positively impacted on the effectiveness of clinical supervision for some professions. There is a need for future policies to focus on the development of common, structured frameworks to support quality clinical supervision across allied health.

Declarations

Ethics approval and consent to participate

Ethics approval was provided by the Bendigo Health and La Trobe University Ethics Committees (Approval number LNR/15/BHCG/26). All participants provided consent to participate in the survey.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and analysed during the current study are not publicly available due to conditions of ethical approval. However, a de-identified version of the data is available from the corresponding author on reasonable request.

Competing interests

Nil to declare.

Funding

This research did not use any funding.

Author's contributions

All authors contributed to conceptualisation, methodology, visualisation, draft preparation, writing review and editing of the manuscript. BP and MG performed quantitative analysis and MG and CM performed qualitative analysis.

Acknowledgements

The authors wish to acknowledge and thank the members of the action research reference group for their contribution and continued involvement and guidance in the research project and participants who provided their data.

Author details

¹ La Trobe Rural Health School, P.O. Box 199, Bendigo, Vic. 3552, Australia. ² Bendigo Health, P.O. Box 126, Bendigo, Victoria. 3552. Australia.

References

1. Philip K. Allied health: untapped potential in the Australian health system. *Australian Health Review*. 2015 Jul 6;39(3):244-7.
2. Davis SF, Enderby P, Harrop D, Hindle L. Mapping the contribution of Allied Health Professions to the wider public health workforce: a rapid review of evidence-based interventions. *Journal of Public Health*. 2017 Mar 1;39(1):177-83.
3. Nancarrow SA, Young G, O'Callaghan K, Jenkins M, Philip K, Barlow K. Shape of allied health: an environmental scan of 27 allied health professions in Victoria. *Australian Health Review*. 2017 Jun 20;41(3):327-35.
4. Association of Schools Advancing Allied Health Professions. What is Allied Health? 2020. Available from: <http://www.asahp.org/what-is/>
5. National Health Service. Health Education England Allied Health Professions. 2020. Available from: <https://www.hee.nhs.uk/our-work/allied-health-professions>
6. Allied Health Therapy and Science disciplines: Position paper. Department of Health, Victoria; 2015. Available from: <https://www2.health.vic.gov.au/about/publications/policiesandguidelines/allied-health-categories-position-paper>
7. Dawson M, Phillips B, Leggat S. Clinical supervision for allied health professionals: a systematic review. *Journal of Allied Health*. 2013;42(2):65-73.
8. Milne D. An empirical definition of clinical supervision. *British Journal of Clinical Psychology*. 2007 Nov;46(4):437-47.
9. Ducat WH, Kumar S. A systematic review of professional supervision experiences and effects for allied health practitioners working in non-metropolitan health care settings. *Journal of Multidisciplinary Healthcare*. 2015;8:397.
10. Keane S, Lincoln M, Rolfe M, Smith T. Retention of the rural allied health workforce in New South Wales: a comparison of public and private practitioners. *BMC health services research*. 2013 Dec;13(1):1-9.
11. Wilson E, Taylor NF. Clinical Supervision for Allied Health Professionals Working in Community Health Care Settings: Barriers to Participation. *Journal of Allied Health*. 2019;48(4):270-9.
12. Martin P, Kumar S, Lizarondo L, Baldock K. Debriefing about the challenges of working in a remote area: A qualitative study of Australian allied health professionals' perspectives on clinical supervision. *PloS one*. 2019;14(3):e0213613.
13. Snowdon DA, Millard G, Taylor NF. Effectiveness of clinical supervision of allied health professionals: a survey. *Journal of Allied Health*. 2016;45(2):113-21.
14. Gardner M, McKinstry C, Perrin B. Effectiveness of Allied Health Clinical Supervision: A Cross-Sectional Survey of Supervisees. *Journal of Allied Health*. 2018;47(2):126-32.
15. Martin P, Kumar S, Lizarondo L, VanErp A. Enablers of and barriers to high quality clinical supervision among occupational therapists across Queensland in Australia: findings from a qualitative study. *BMC Health Services Research*. 2015;15(1):1-8.

16. Saxby C, Wilson J, Newcombe P. Can clinical supervision sustain our workforce in the current healthcare landscape? Findings from a Queensland study of allied health professionals. *Australian Health Review*. 2015;39(4):476-82.
17. Fitzpatrick S, Smith M, Wilding C. Quality allied health clinical supervision policy in Australia: a literature review. *Australian Health Review*. 2012;36(4):461-5.
18. Snowdon DA, Millard G, Taylor NF. Effectiveness of clinical supervision of physiotherapists: a survey. *Australian Health Review*. 2015;39(2):190-6.
19. Nancarrow SA, Wade R, Moran A, Coyle J, Young J, Boxall D. Connecting practice: a practitioner centred model of supervision. *Clinical Governance: An International Journal*. 2014; 19(3): 235-252.
20. Bell KE, Hall F, Pager S, Kuipers P, Farry H. Developing allied health professional support policy in Queensland: a case study. *Human Resources for Health*. 2014 Dec;12(1):1-8.
21. Snowdon DA, Sargent M, Williams CM, Maloney S, Caspers K, Taylor NF. Effective clinical supervision of allied health professionals: a mixed methods study. *BMC health services research*. 2020 Dec;20(1):1-1.
22. Gardner M, McKinstry C, Perrin B. Clinical Supervision Frameworks for Allied Health Professionals: A Systematic and Critical Review. *Internet Journal of Allied Health Sciences and Practice*. 2021;19(3) [in press].
23. McNiff J. *Action research: Principles and practice*: Routledge; 2013.
24. Winstanley J, White E. The MCSS-26©: Revision of the Manchester Clinical Supervision Scale© Using the Rasch Measurement Model. *Journal of Nursing Measurement*. 2011;19(3):160-78.
25. Winstanley J. Manchester clinical supervision scale. *Nursing Standard* (through 2013). 2000;14(19):31.
26. Winstanley J. *The Manchester Clinical Supervision Scale user guide*. Osman Consulting. 2000.
27. Krippendorff K. *Content analysis: An introduction to its methodology*: Sage publications; 2018.
28. Dawson M, Phillips B, Leggat S. Effective clinical supervision for regional allied health professionals - the supervisee's perspective. *Australian Health Review*. 2012;36(1):92-7.
29. Kumar S, Osborne K, Lehmann T. Clinical supervision of allied health professionals in country South Australia: A mixed methods pilot study. *Australian Journal of Rural Health*. 2015;23(5):265-71.
30. Snowdon DA, Cooke S, Lawler K, Scroggie G, Williams K, Taylor NF. Physiotherapists prefer clinical supervision to focus on professional skill development: a qualitative study. *Physiotherapy Canada*. 2020;72(3):249-57.
31. Snowdon DA, Sargent M, Williams CM, Maloney S, Caspers K, Taylor NF. Effective clinical supervision of allied health professionals: a mixed methods study. *BMC Health Services Research*. 2020;20(1):1-11.
32. Dawson M, Phillips B, Leggat SG. Effective clinical supervision for regional allied health professionals: the supervisor's perspective. *Australian Health Review*. 2013;37(2).

33. Martin P, Baldock K, Kumar S, Lizarondo L. Factors that contribute to high-quality clinical supervision of the rural allied health workforce: lessons from the coalface. *Australian Health Review*. 2019;43(6):682-8.
34. Martin P, Lizarondo L, Kumar S. A systematic review of the factors that influence the quality and effectiveness of telesupervision for health professionals. *Journal of Telemedicine and Telecare*. 2018;24(4):271-81.
35. Martin P, Copley J, Tyack Z. Twelve tips for effective clinical supervision based on a narrative literature review and expert opinion. *Medical Teacher*. 2014;36(3):201-7.
36. Saxby C, Wilson J, Newcombe P. Does best practice clinical supervision lead to better outcomes? Findings from a Queensland study of community allied health professionals. *Advances in Clinical Supervision Innovation & Practice*. 2013:23.
37. Ducat W, Martin P, Kumar S, Burge V, Abernathy L. Oceans apart, yet connected: Findings from a qualitative study on professional supervision in rural and remote allied health services. *Australian Journal of Rural Health*. 2016;24(1):29-35.