

# Sense of Coherence and Strategies for Coping with Stress Among Nurses

Katarzyna Betke (✉ [katarzyna.betke@cm.umk.pl](mailto:katarzyna.betke@cm.umk.pl))

Nicolaus Copernicus University Faculty of Medicine: Uniwersytet Mikołaja Kopernika w Toruniu  
Collegium Medicum im Ludwika Rydygiera w Bydgoszczy <https://orcid.org/0000-0001-9379-1099>

**Małgorzata Anna Basińska**

Kazimierz Wielki University in Bydgoszcz Institute of Psychology: Uniwersytet Kazimierza Wielkiego  
Instytut Psychologii

**Anna Andruszkiewicz**

Nicolaus Copernicus University in Toruń Ludwik Rydygier Collegium Medicum in Bydgoszcz:  
Uniwersytet Mikołaja Kopernika w Toruniu Collegium Medicum im Ludwika Rydygiera w Bydgoszczy

---

## Research article

**Keywords:** stress, nurses, coping strategies, work environment, stress coping strategies, the sense of coherence

**Posted Date:** October 19th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-91555/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

The nursing profession is associated with constant presence of difficult situations and stress, which arise from responsibility for the highest values – human life and health. With demographic changes in the society, the demand for nursing care increases. Looking after the health of nursing staff becomes a non-material investment in employees. One of the most important health potentials described in literature is the sense of coherence. It can significantly contribute to maintaining good health, modify one's functioning in a stressful working environment and influence the choice of strategies for coping with stress.

## Aim

The aim of the study was to describe the specific relationship between the sense of coherence and strategies for coping with stress in a group of professionally active nurses.

## Methodology and methods

The study involved 90 female nurses and 1 male nurse aged 22-52, the group was diversified in terms of: education, work system, marital status and place of residence. The study was conducted with the use of: Sense of Coherence Questionnaire SOC-29 and Inventory to Measure Coping Strategies with Stress - Mini-COPE. The study was conducted in accordance with the principles of scientific research set out in the Helsinki Declaration.

## Results

The sense of coherence value in the test group was  $M = 134.24$  ( $SD = 19.55$ ). In stressful situations nurses most often used active strategies to cope with stress: Planning  $M = 2.10$  ( $SD = 0.54$ ), Seeking Emotional Social Support  $M = 1.95$  ( $SD = 0.68$ ) and Seeking Instrumental Social Support  $M = 1.95$  ( $SD = 0.69$ ), and the least frequently: Alcohol/Drug Use  $M = 0.28$  ( $SD = 0.48$ ). The level of the sense of coherence and its components differentiated the strategies of coping with stress used in the examined group of nurses.

## Conclusions

The research confirmed that the sense of coherence serves as a health potential in a stressful working environment. Nurses with a stronger sense of coherence used more adaptive strategies to deal with stress than those with average or low levels.

# Background

Nursing is a profession commonly considered difficult and not always appreciated by the society, but there is a great need for it. Nowadays, it is a profession which creates high expectations for people who practise it. Nurses are the largest medical professional group. They are individuals with higher education degrees, who make their own decisions concerning the care administered to patients and who provide a wide range of health services. They carry out their everyday tasks in continuous contact with other people, which is linked to holding responsibility for the highest values: human life and health. At the same time, apart from the duties performed, more expectations are set by their superiors, other members of the therapeutic team, as well as patients and their relatives. The European Agency for Safety and Health at Work and the results of worldwide studies place nursing among the most stressful professions [1]. Numerous studies have shown that nurses very often consider their work stressful [2–5]. The sources of stress in the work of a nurse include: bad work organisation, irregular work, shift work disrupting the natural biological body rhythm, periodically causing high overloads, lack of satisfactory remuneration, lack of recognition by superiors, lack of prospects for professional development and a poor flow of information in the team. The profession is also associated with exposure to harmful biological, chemical, physical and psychological factors, as well as stressful cooperation with patients and their families [3–4, 6–8]. Long-term functioning in a stressful working environment has an undeniable impact on health. It also triggers the need for specific actions, which are defined as coping strategies. Coping with stress, according to the authors of the transactional stress theory - Richard Lazarus and Susan Folkman - is a constantly changing cognitive and behavioural effort, focused on specific external and internal struggles, in situations seen as burdensome or exceeding personal capabilities. In order to cope with difficulties, a person can choose active ways of solving problems, considered pro-health or evasive strategies. Dealing with stress can be considered a behaviour significant for human well-being. The use of personal resources has an effect on health, well-being and effective functioning in the work environment [9]. The choice of adaptive strategies for coping with stress can also be influenced by a generalised and relatively stable dispositional orientation, which is characteristic for a given individual and which determines the fundamental way of perceiving, feeling and understanding the world around and one's own life, i.e. the so-called Aaron Antonovsky's sense of coherence (SOC). There are three correlated components of the sense of coherence: the sense of comprehensibility, the sense of manageability and the sense of meaningfulness [10]. According to Aaron Antonovsky, "the key to health", i.e. the main factor thanks to which a person can cope with stress and not fall ill or can recover faster in case of illness is precisely the sense of coherence [10–13]. The functional aspect of the sense of coherence is most noticeable in situations of stress and its management. The role of this construct is to prevent transformation of a state of tension into a state of stress, as well as to promote the use of more adaptive strategies for coping with difficult situations [14]. The sense of coherence influences the way of perceiving stressful situations occurring during the performance of professional tasks, favours engaging in problem-oriented strategies and also plays a role in protecting employees from the negative effects of stress experienced. Studies conducted in a group of over a thousand nurses in Poland have shown that a strong sense of coherence reduces the level of experienced stress, irrespective of the strength of the stressors and weakens its negative effects [11, 15]. The current situation, i.e. a shortage of nursing staff, the ageing of this professional group and the fact that nursing graduates do not take up this kind of employment creates a

need to look for solutions to these problems. According to Małgorzata Anna Basińska and Anna Andruszkiewicz, it seems justified to take actions aimed at reducing stressors in everyday work, as well as to develop and strengthen personal resources and the ability to cope with difficult situations. Creating a friendly work environment, strengthening health meta-resources, as well as providing training aimed at developing an ability to choose constructive strategies for coping with stress should become a permanent component of investment in employees [16]. It seems necessary to implement a strategy to prevent stress and its consequences in the workplace. Today, stress should not be seen as a weakness, but rather as a phenomenon that can be managed by creating a culture of openness and understanding [17].

The aim of the study was to describe the specific relationship between the sense of coherence and strategies for coping with stress in a group of professionally active nurses.

## Methods

The study involved 90 female nurses and 1 male nurse aged 22–52 ( $M = 38$ ,  $SD = 7.46$ ), the group was diversified in terms of socio-demographic variables and variables connected to professional work. The majority of respondents declared their marital status to be “in a relationship”, the least of them were “separated” or “widowed”. The majority of the respondents had higher education vocational degrees, the average number of years in the profession was over 16 years ( $M = 16.597$ ;  $SD = 8.35$ ), and the majority of the respondents were unit nurses. The study was conducted in accordance with the principles of scientific research set out in the Helsinki Declaration. The research was approved by the Bioethics Committee of October 20, 2015 (KB 666/2015). The individuals participating in the study on a voluntary basis were healthy and did not have a professional or other relationship with the researcher. Sets of questionnaires were provided to nurses in post-graduate education.

To measure the sense of coherence, the Sense of Coherence Questionnaire (SOC-29) by Antonovsky was used, in an adapted Polish version (Koniarek et al., 1993). It helps estimate the intensity of the sense of coherence and its three components: the sense of comprehensibility, the sense of manageability and the sense of meaningfulness. An evaluation of the Polish version of the SOC-29 Questionnaire showed a very high reliability of the tool - Cronbach's alpha index = 0.78 [14]. To evaluate strategies for coping with stress, the Inventory to Measure Coping Strategies with Stress (Mini-COPE) by Carver, Scheier and Weintraub was used, in an adapted Polish version (Juczyński, Ogińska-Bulik, 2009). The questionnaire consists of 28 statements relating to 14 stress coping strategies. The respondent addresses each statement on a scale from 0 (I almost never do this) to 3 (I almost always do this). The Polish version is characterized by high reliability, internal consistency expressed as Cronbach's alpha = 0.78 [11].

## Results

The analyses presented below were performed on the basis of a correlation model. The following statistical tests were used:

1. to describe group variables the mean (M) and standard deviation (SD), and number (f) and percentage (%);
  
2. to determine the direction and strength of correlation between variables Pearson's linear correlation test (when there was a linear correlation between variables) or Spearman's nonparametric test (when variables were qualitative);
  
3. to assess the significance of differences between multiple means ANOVA variance analysis or ANOVA rank analysis by Kruskal-Wallis (in the absence of a normal distribution);
  
4. NIR test; to examine the significance of differences between individual variables.

The first step of the analysis was to calculate average values for all variables. The results obtained were presented in the form of raw average results. Individual tables differ in terms of the number of presented studies due to exclusion of incomplete questionnaires from the analyses. The studied group of nurses obtained an average value of the sense of coherence of M = 134.24. The values of the sense of coherence and its components were highly differentiated [insert Table 1.].

Table 1  
Mean raw scores (M), standard deviation (SD), in terms of sense of coherence and its components (N = 88) in the studied group of nurses

<b>Dimensions sense of coherence</b>	<b>M</b>	<b>SD</b>	<b>Minimum</b>	<b>Maximum</b>
<b>Sense of coherence</b>	134.24	19.551	79.00	196.00
<b>Sense of comprehensibility</b>	44.56	9.190	23.00	77.00
<b>Sense of manageability</b>	47.24	8.203	24.00	63.00
<b>Sense of meaningfulness</b>	42.44	5.952	26.00	56.00

The nurses from the study group most often declared the use of strategies classified as active ways of coping with stress, focused on the problem [insert Table 2.].

Table 2  
Average raw results (M), standard deviation (SD) in the scope of coping with stress strategies (N = 91) in the studied group of nurses

<b>Strategies for coping with stress</b>	<b>M</b>	<b>SD</b>	<b>Minimum</b>	<b>Maximum</b>
<b>Active Coping</b>	<b>2.19</b>	<b>0.475</b>	<b>1.00</b>	<b>3.00</b>
<b>Planning</b>	<b>2.10</b>	<b>0.540</b>	<b>0.50</b>	<b>3.00</b>
<b>Positive Reinterpretation and Growth</b>	<b>1.82</b>	<b>0.593</b>	<b>0.00</b>	<b>3.00</b>
<b>Acceptance</b>	1.71	0.558	0.50	3.00
<b>Humour</b>	0.65	0,575	0.00	3.00
<b>Religion</b>	1.30	0.937	0.00	3.00
<b>Seeking Emotional Social Support</b>	<b>1.95</b>	<b>0.681</b>	<b>0.00</b>	<b>3.00</b>
<b>Seeking Instrumental Social Support</b>	<b>1.95</b>	<b>0.693</b>	<b>0.00</b>	<b>3.00</b>
<b>Suppression of Competing Activities</b>	1.46	0.720	0.00	3.00
<b>Denial</b>	0.91	0.649	0.00	3.00
<b>Restraint Coping</b>	1.33	0.620	0.00	2.50
<b>Alcohol/Drug Use</b>	0.28	0.484	0.00	2.00
<b>Behavioural Disengagement</b>	0.68	0.569	0.00	2.00

The next stage of the analysis was to answer the question of: Does the level of the sense of coherence and its components differentiate the strategies of coping with stress used in the examined group? The nurses were divided into three subgroups based on the results obtained: individuals with a low sense of coherence, with an average sense of coherence and with a high sense of coherence according to the criterion (mean  $\pm$  ½ SD).

### **The sense of coherence and the strategies used to cope with stress**

The results of the analyses show that the sense of coherence significantly differentiates the studied group in terms of the applied coping strategies [insert Table 3].

Table 3  
Sense of coherence and strategies for coping with stress (N = 88) –  
results of analysis of variance

Strategies for coping with stress	Statistical test F / H	P
Active Coping	0.44	0.645
Planning	0.19	0.826
Positive Reinterpretation and Growth	2.59	0.081
Acceptance	0.99	0.375
Humour	1.95	0.148
Religion	1.72	0.185
Seeking Emotional Social Support	3.06	0.052
Seeking Instrumental Social Support	0.90	0.411
Suppression of Competing Activities	0.40	0.670
Denial	<b>7.58</b>	<b>0.001</b>
Restraint Coping	1.58	0.211
Alcohol/Drug Use	H (2, N = 88) = 5.150	0.076
Behavioural Disengagement	<b>5.32</b>	<b>0,007</b>
Focus on and Venting of Emotions	<b>7.46</b>	<b>0.001</b>

Nurses belonging to the subgroup with a high sense of coherence used Denial, Behavioural Disengagement and Focus on and Venting of Emotions strategies less frequently than the subgroup with low and average sense of coherence. The respondents from the subgroup with a high sense of coherence were significantly less likely to use the Alcohol/Drug Use strategy than those from the subgroup with a low sense of coherence, who most often declared their choice of this strategy. The picture of statistically significant differences in the application of the Alcohol/Drug Use strategy between individuals from the particular subgroups in terms of the sense of coherence is presented in Fig. 1.

### **The sense of comprehensibility and the strategies used to cope with stress**

The results show that the sense of comprehensibility significantly differentiates the studied group in terms of the applied coping strategies [insert Table 4.].

Table 4  
Sense of comprehensibility and strategies for coping with stress-  
analysis of variance

Strategies for coping with stress	Statistical test	P
	F / H	
Active Coping	0.193	0.825
Planning	0.027	0.974
Positive Reinterpretation and Growth	2.518	0.087
Acceptance	0.507	0.604
Humour	2.997	0.055
Religion	<b>H = 11.800</b>	<b>0.003</b>
Seeking Emotional Social Support	0.910	0.407
Seeking Instrumental Social Support	1.133	0.327
Suppression of Competing Activities	0.382	0.684
Denial	<b>6.822</b>	<b>0.002</b>
Restraint Coping	3.047	0.053
Alcohol/Drug Use	2.746	0.070
Behavioural Disengagement	1.490	0.231
Focus on and Venting of Emotions	<b>H = 13.635</b>	<b>0.001</b>

Nurses from the subgroup with high levels of comprehensibility were significantly less likely to use the Denial coping strategy than those belonging to the subgroups with low and average levels of comprehensibility. Kruskal-Wallis test was then used to examine the differentiating role of the sense of comprehensibility in the Religion strategy [Kruskal-Wallis test:  $H(2, N = 88) = 11.8002$   $p = .003$ ] and Focus on and Venting of Emotions [Kruskal-Wallis test:  $H(2, N = 88) = 13.635$   $p = .001$ ] [insert Figs. 2,3].

Nurses from the subgroup with high levels of comprehensibility were significantly statistically less likely to use the Religion and Focus on and Venting of Emotions coping strategy than those belonging to the subgroups with low and average levels of comprehensibility.

### **The sense of manageability and the strategies used to cope with stress**

The results show that the sense of manageability significantly differentiates the studied group in terms of the applied coping strategies [insert Table 5.].

Table 5

The sense of manageability and used to strategies for coping with stress

<b>Strategies for coping with stress</b>	<b>Statistical test</b>	<b>P</b>
	<b>F / H</b>	
<b>Active Coping</b>	H (2, N = 88) = .585	0.746
<b>Planning</b>	0.49	0.617
<b>Positive Reinterpretation and Growth</b>	0.17	0.843
<b>Acceptance</b>	0.46	0.635
<b>Humour</b>	0.18	0.833
<b>Religion</b>	H (2, N = 88) = 3.422	0.181
<b>Seeking Emotional Social Support</b>	<b>H (2, N = 88) = 6.902</b>	<b>0.032</b>
<b>Seeking Instrumental Social Support</b>	2.48	0.090
<b>Suppression of Competing Activities</b>	1.26	0.289
<b>Denial</b>	<b>3.71</b>	<b>0.029</b>
<b>Restraint Coping</b>	0.82	0.444
<b>Alcohol/Drug Use</b>	H (2, N = 88) = 5.653	0.059
<b>Behavioural Disengagement</b>	<b>3.13</b>	<b>0.049</b>
<b>Focus on and Venting of Emotions</b>	<b>8.66</b>	<b>0.000</b>

Nurses from the subgroup with high levels of manageability were significantly more likely to use the Seeking Emotional Social Support coping strategy than those belonging to the subgroups with low and average levels of manageability. Nurses from the subgroup with a low level of sense of manageability were the least likely to choose this strategy.

Nurses from the subgroup with high levels of manageability were significantly less likely to use the Denial and Behavioural Disengagement coping strategy than those belonging to the subgroup with a low level of manageability. Respondents from the subgroup with high levels of manageability were significantly less likely to choose the Focus on and Venting of Emotions coping strategy than those belonging to the subgroups with low and average levels of manageability.

### **The sense of meaningfulness and the strategies used to cope with stress**

The results of the analyses show that the sense of meaningfulness significantly differentiates the studied group in terms of the applied coping strategies [insert Table 6.].

Table 6  
Sense of meaningfulness and used strategies for coping with stress-  
analysis of variance

Strategies for coping with stress	Statistical test F / H	P
Active Coping	0.030	0.971
Planning	0.090	0.914
Positive Reinterpretation and Growth	H (2, N = 88) = 2.034	0.362
Acceptance	0.595	0.554
Humour	H (2, N = 88) = 4.019	0.134
Religion	0,007	0.993
Seeking Emotional Social Support	<b>3.838</b>	<b>0.025</b>
Seeking Instrumental Social Support	<b>H (2, N = 88) = 7.787</b>	<b>0.020</b>
Suppression of Competing Activities	1.606	0.207
Denial	1.122	0.331
Restraint Coping	1.032	0.361
Alcohol/Drug Use	<b>H (2, N = 88) = 3.005</b>	0.223
Behavioural Disengagement	2.432	0.094
Focus on and Venting of Emotions	<b>3.806</b>	<b>0.026</b>

Nurses from the subgroup with high levels of sense of meaningfulness were significantly statistically more likely to use the Seeking Emotional Social Support coping strategy than those belonging to the subgroups with low and average levels of meaningfulness. The respondents from the subgroup with a high level of sense of meaningfulness were less likely to use the Focus on and Venting of Emotions strategy than those with a low and average sense of meaningfulness.

The analysis showed that nurses from the subgroup with an average level of sense of meaningfulness significantly more often chose the strategy of Seeking Instrumental Social Support in difficult situations. Nurses from the subgroup with a low level of sense of meaningfulness were the least likely to choose this strategy [insert Fig. 4.].

The analyses showed that respondents from the subgroup with a low sense of meaningfulness were significantly more likely to use the Alcohol/Drug Use strategy than those belonging to the subgroups with a high and average sense of coherence [insert Fig. 5.].

## Discussion

This study confirmed a correlation between the sense of coherence and coping strategies in the examined group of nurses. These results correspond to conclusions presented by both Polish and foreign researchers [2, 13, 14, 18] concerning the pro-health and buffer role played by the sense of coherence in a stressful working environment.

The studied group presented a varied level of sense of coherence and its components, similarly to the results published in literature. The research on the sense of coherence showed that nurses had a sense of coherence at a level similar to that indicated by Polish researchers - the lowest level of the sense of coherence among nurses was reported by Koniarek (1992):  $M = 124.60$  [19], while the highest - by Kretowicz (2011):  $M = 145.02$  [19]. Polish nurses presented a similar level of the sense of coherence to paediatric nurses in Hong Kong:  $M = 135.75$  [20], in South Africa:  $M = 137.92$  ( $SD = 20.46$ ) [21], in Brazil:  $M = 144.60$  ( $SD = 22.60$ ) [22], as well as in Japan:  $M = 124.40$  ( $SD = 21.20$ ) [23].

The sense of coherence can protect an individual from the destructive effects of difficult and stressful situations. Many studies have shown that a high sense of coherence significantly reduces the effects of stress and helps cope with the difficulties and challenges of life [2, 12, 14, 16, 18, 19, 21, 24]. The components of the sense of coherence also have a significant preventive effect [10, 14]. The sense of coherence determines the way of perceiving stressful situations at work, favours engaging in problem-oriented strategies and also plays a role in protecting employees from the negative effects of stress experienced. Studies conducted in a group of over a thousand nurses in Poland have shown that a high sense of coherence reduces the level of experienced stress, irrespective of the strength of the stressors and weakens the negative effects of stress. The buffering role of the sense of coherence was confirmed primarily in relation with the sense of manageability and the sense of meaningfulness [11, 15].

This is important for nursing staff, whose work is performed in situations of emotional stress and tension. Antonovsky proved that in stressful situations a high sense of coherence encourages more adaptive strategies for coping with difficult situations, i.e. focusing on the task instead of on emotions, and seeking social support [14]. Medical Universities in Poland include Antonovsky's theory into practice, offering, in accordance with current teaching curricula and standards, study modules which help students strengthen their sense of coherence. These include, among others, psychology with elements of interpersonal communication, health promotion, basics of psychotherapy. The process of strengthening the sense of coherence should be continued in the working environment of nursing staff. According to Korcz and Korcz-Biernat, it seems reasonable to take measures that would allow nursing staff to increase their participation in decision-making, also through partnership-based relations with other members of the interdisciplinary team. The authors also emphasize the importance of developing an ability to cope with stress at work [25]. The objectives of salutogenesis are addressed at people who in their everyday work help others and perform duties in conditions of high emotional tension and stress. The author of the theory of salutogenesis stated that his book "Unraveling the Mystery of Health. How People Manage

Stress and Stay Well” was also written with nurses in mind, hoping for his theory to be empirically verified by that professional group [10].

The examined group of nurses more often reported choosing pro-health, problem-oriented stress management strategies (i.e. Active Coping, Planning, Seeking Instrumental Social Support and Seeking Emotional Social Support), which corresponds to the results of other authors' research in this area. Polish nurses most often reported choosing active and constructive actions as a strategy for coping with occupational stress [16]. According to Perek et al., this is a typical way of coping with stress in this occupational group [7]. Active stress management strategies are more commonly used by people with a higher level of coherence. This was confirmed by our own research. Basińska and Andruszkiewicz, using the AVEM Questionnaire and the Latack Scale, studied stress management strategies and work-related experiences among 150 nurses working at different wards. The authors proved that there were no differences between coping strategies chosen by nurses depending on the department they worked at. Those who coped with problems at work through the use of Suppression of Competing Activities/ Behavioural Disengagement strategy used offensive problem-solving strategies less often. Nurses who coped with problems through the Positive Reinterpretation and Growth strategy used more offensive ways of solving problems and were more satisfied with their lives. More often, offensive methods and a task-based mode were used by nurses using Active Coping, Seeking Emotional Social Support and Seeking Instrumental Social Support strategies [16]. The results of the Polish study correspond to the reports of other researchers. Australian nurses, as Lim indicates, choose problem-oriented strategies more often than emotion-oriented ones [26]. Studies on the use of stress management strategies by French nurses indicate that they most often choose Active Coping, Restraint Coping and Positive Reinterpretation and Growth strategies [27]. The study also found that nurses turned to strategies to reduce emotional tension, such as: Seeking Emotional Social Support, Positive Reinterpretation and Growth and Acceptance. The least respondents chose evasive strategies, which are primarily aimed at forgetting or distracting attention from the problem.

The main objective of the study was to assess the correlation between the sense of coherence and the strategies used to deal with stress. These study confirmed the existence of such a correlation. The examined nurses with a high level of coherence were less likely to use evasive strategies, i.e. Denial, Behavioural Disengagement and Focus on and Venting of Emotions, than those with a low and average level of sense of coherence, which has a positive impact on their health and professional functioning. According to Antonovsky, *“people with a strong sense of coherence choose a specific strategy of coping, which in their opinion is most adequate for a given stressor”*, therefore, they use more adaptive techniques to deal with problems that may contribute to better functioning and maintaining good health [28]. They are also able to make optimal use of the resources available to them, updating and flexibly adapting to changing circumstances those coping strategies that prove to be the most effective in a particular situation. The choice of stress management strategies that are beneficial to human health has an effect on maintaining health potential and functioning in a stressful work environment. Active strategies make it easier to stay healthy. Evasive strategies may reduce the positive impact of the sense of coherence and make it difficult to maintain good health.

In our study, we also observed an increased tendency for choosing an evasive stress management strategy - Alcohol/Drug Use - by nurses with low levels of coherence. This trend is in line with Antonovsky's theory. People with a weak sense of coherence more often experience negative emotions such as shame, discouragement, anxiety, which in effect inhibits action-taking. They perceive problems inaccurately, which leads to a failure to make the necessary efforts to cope effectively [11–13, 20].

Empirical research has shown that the sense of coherence and its components correlate with strategies used to deal with stressful situations. A high level of comprehensibility reduces the intensity of assessment of stressful situations, especially those related to threat and loss. The sense of manageability works in a similar way, which additionally influences the perception of the situation as a challenge. Whereas a strong sense of meaningfulness maximizes the perception of stressors as challenges and limits the perception of a situation as a loss. People with a strong sense of meaningfulness take on the challenges of life, search for meaning and make an effort to solve and deal with problems [12, 14].

The study found that nurses with a stronger sense of comprehensibility, manageability and meaningfulness were more likely to choose emotionally focused strategies, and least likely to use non-adaptive strategies, whose consequences may be detrimental to the one's functioning and health. The sense of meaningfulness, the component Antonovsky treats as a motivational-emotional one, and according to Antonovsky, the most important one in creating the sense of coherence, correlated with the more frequent choice of the Seeking Emotional Social Support strategy, and the least frequently with the choice of the Focus on and Venting of Emotions strategy. What is important, the weaker the sense of meaningfulness of the respondents, the more often they used the Alcohol/Drug Use strategy. The use of this and other evasive strategies by nurses may contribute to deterioration of health, ineffective functioning in the work environment and absenteeism, which in the current situation of shortages in nursing staff is an undesirable situation. In relation to the above, it is justified to implement psycho-educational measures developing an ability to choose adaptive stress management strategies, which should be addressed especially at people with a weak sense of coherence. Strengthening the potential of the sense of coherence is in line with both Antonovsky's idea and the contemporary trend of positive psychology [29].

Investing in employees means reaching to their internal resources and developing their ability to cope with difficult situations. This investment will contribute to effective operation at work and better quality of services provided to the society. Implementation of strategies aimed at preventing stress and its consequences in the workplace should take place on both a personal and organisational level. Stress management programmes including education and training on stress management are a personal-level strategy that provides support for nurses experiencing stress. An organisation-focused approach addresses work-related stressors by reducing or eliminating them. Long-term implementation of this strategy will allow for better management of nursing staff and for ensuring adequate resources [30].

The presented research results may constitute a source of knowledge for designing psycho-educational programmes addressed both at students - future nurses and active professionals in order to minimize the occurrence of risky behaviours and to help develop skills to cope with them.

## Conclusions

1. The level of the sense of coherence and its components in the examined group of nurses was similar to results reported for this group to date.
2. The sense of coherence serves as a health potential in a stressful working environment.
3. Nurses with a stronger sense of coherence more often chose adaptive coping strategies, considered pro-health strategies.
4. It is recognised that strengthening health potential i.e. the sense of coherence is justified, as it can significantly modify an individual's functioning in a stressful working environment.

## Declarations

### Authors' contributions

All authors contributed to the conception and the design of the study, K.B. collected, analysed and interpreted the data, and drafted the manuscript. M.A.B. and A.A. commented on the manuscript and edited the text.

### Competing interests

The authors declare no competing interests.

### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Ethics approval and consent to participate

The study also included nurses who participated in postgraduate vocational training programs. The study was conducted according to the principles laid down in the Declaration of Helsinki. It was approved by the Ethics Committee of the Nicolaus Copernicus University in Toruń, Ludwik Rydygier Collegium Medicum in Bydgoszcz 2015/10/20 (KB 666/2015). The individuals participating in the study on a voluntary basis were healthy and did not have a professional or other relationship with the researcher. All nurses were informed verbally and/or in text about topic, goals, expected benefits and possible risks of the study (informed consent) before participating in the study. Filling out the questionnaire after receiving information was interpreted as consent to participate.

## Consent for publication

Not applicable

## Availability of data and materials

The data generated and analysed during the current study are not publicly available due to the sensitivity of the data.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

K.B. carried out the research and conception of the study in close cooperation with A.A.

K.B. contributed to the data collection. Interviews were transcribed, analysed and interpreted by M-A.B. and discussed with A.A.. All authors reviewed the drafts of the manuscript, gave comprehensive feedback and approved the final manuscript.

## Acknowledgements

We would like to thank all participants for their participation in this research,


## References

1. European study of working conditions. 2015, <https://www.eurofound.europa.eu/pl/data/european-working-conditions-survey> (2015, accessed 10 June 2019).
2. Khamisa N, Oldenburg B, Peltzer K, Ilic D. Work Related Stress, Burnout, Job Satisfaction and General Health of Nurses. *Int J Environ Res Public Health*. 2015;12(1):652–66.
3. Basińska MA, Andruszkiewicz A. Styles of coping with stress in life and applied techniques of coping with stress at work by nurses. In: Łepecka-Klusek C and Turowski K, editors *Mental health promotion: materials from the Nationwide Scientific Conference; May 28–29, 1999*. Lublin, 1999, pp. 21–28.
4. Holman D, Johnson S, O'Connor E. Stress management interventions: Improving subjective psychological well-being in the workplace. In: Diener R, Oishi S, Tay L, editors. *Handbook of well-being*. Salt Lake City: UT: DEF Publishers; 2018. DOI: nobascholar.com.
5. Kowalczyk K, Krajewska-Kułak E. Influence of selected sociodemographic factors on psychosocial workload of nurses and association of this burden with absenteeism at work. *Med Pr*. 2015;66(5):615–24.
6. Kamau C, Medisauskaite A, Lopes B. Inductions Buffer Nurses' Job Stress, Health, and Organizational Commitment. *Arch Occup Environ Health*. 2015;70(6):305–8.

7. Perek M, Kózka M, Twarduś K. Difficult situations in the work of pediatric nurses and ways of dealing with them. *Probl Piel.* 2007;15(4):223–8.
8. Beisert M. Symptoms, mechanisms and causes of nurses' burnout. In: Sęk H, editor. *Occupational burnout. Causes, prevention.* Warsaw: PWN Publisher; 2010. p. 183.
9. Lazarus RS, Folkman S. *Stress, appraisal and coping.* New York: Springer; 1984. pp. 18–23, 141.
10. Antonovsky A. *Health, Stress and Coping.* San Francisco: Jossey-Bass; 1979.
11. Jażdżewska N, Kretowicz K, Żuralska R. Hope for success and a sense of coherence in the nurses' professional group. *Probl Piel.* 2018;26(1):9–14.
12. Ando M, Kawano M. Relationships among moral distress, sense of coherence, and job satisfaction. *Nur Ethics.* 2018;25(5):571–9.
13. Wijesinghe S, Parshall MB. Impermanence and Sense of Coherence. *J Transcult Nurs.* 2016;27(2):157–65.
14. Antonovsky A. *Unraveling The Mystery of Health. How People Manage Stress and Stay Well.* San Francisco: Jossey-Bass Publishers; 1987.
15. Dudek B, Koniarek J. Sense of coherence and perception of the relationship between working conditions, stress and self-assessment of health. *Alkohol Narkom.* 1996;1(22):65–73.
16. Basińska MA, Andruszkiewicz A. Strategies of coping with occupational stress by nurses and their behavior and work-related experiences. *Pol For Psycholog.* 2010;15(2):169–92.
17. Michie S. Causes and management of stress at work. *Occup Environ Med.* 2002;59(1):67–72.
18. Basińska MA, Andruszkiewicz A. Types of behaviors and experiences at work in the group of nurses depending on the declared religiosity. *Pielęg XXI w.* 2005;4:21–5.
19. Mazzella Ebstein AM, Sanzero Eller L, See Tan K, et al. The relationships between coping, occupational stress, and emotional intelligence in newly hired oncology nurses. *Psycho-Oncology* 2019; 28(2): 278–283. DOI: <https://doi.org/10.1002/pon.4937> 2019, 28 (2): 278–283.
20. Shiu AT-Y. The significance of sense of coherence for the perceptions of task characteristics and stress during interruptions amongst a sample of public health nurses in Hong Kong: Implications for nursing management. *Public Health Nurs.* 1998;15(4):273–80.
21. Van der Colff J, Rothmann S. Occupational stress, sense of coherence, coping burnout and work engagement of registered nurses in South Africa. *SAJIP.* 2009;35:1–10.
22. Rodrigues Costa Schmidt D and Spadoti. Dantas RA. Analysis of Validity and Reliability of the Adapted Portuguese Version of Antonovsky's Sense of Coherence Questionnaire Among Nursing Professionals. *Rev Latino-Am Enfermagem* 2011; 19 (1): 42–9.
23. Kikuchi Y, Nakaya M, Ikeda M, et al. Relationship between depressive state, job stress, and sense of coherence among female nurses. *Breast Cancer Res Treat* 2014; 18: 32–35.
24. Manor-Binyamini I, Nator M. Parental coping with adolescent developmental disabilities in terms of stress, sense of coherence and hope within the Druze community of Israel. *Res Dev Disabil.* 2016;55:358–67.


25. Korcz I, Korcz-Biernat E. Sense at work and burnout syndrome. In: Pietrulenwicz B, editors. *Possibilities to improve the learning process - selected issues*. Zielona Gora, 2005, pp. 251–258.
26. Lim J, Bogossian F, Ahern K. Stress and coping in Australian nurses: a systematic review. *Int Nurs Rev*. 2010;57:22–31.
27. Lala AL, Sturzu LM, Picard JP. et al. Coping behavior and risk and resilience stress factors in French regional emergency medicine unit workers: a cross-sectional survey. *J Med Life*. 2016;9(4):363–8.
28. Andruszkiewicz A, Wrońska I. The sense of coherence of nurses and the type of work related behaviors. *Pielęg XXI w*. 2003;5:47–51.
29. O'Connor SA, Grant AM. Positive psychology meets real estate: The positive built workplace environment. *Corporate Real Estate Journal* 2018-19; 8 (2): 136–153.
30. Sargazi O, Foroughameri G, Miri S. et al. Improving the professional competency of psychiatric nurses: Results of a stress inoculation training program. *Psych Res*. 2018. DOI:<https://doi.org/10.1016/j.psychres.2018.10.057>.

## Figures




**Figure 1**

Graph results of using Alcohol/Drug Use strategy to the levels of the sense of coherence



**Figure 2**


Graph results of using Religion strategy to the levels of the sense of comprehensibility



1 – Low sense of comprehensibility 2 – Average sense of comprehensibility 3 – High sense of comprehensibility

**Figure 3**


Graph results of using Focus on and Venting of Emotions strategy to the levels of the sense of comprehensibility



1 – Low sense of meaningfulness 2 – Average sense of meaningfulness 3 – High sense of meaningfulness

**Figure 4**

Graph results of using Seeking Instrumental Social Support strategy to the levels of the sense of meaningfulness



**Figure 5**

Graph results of using Alcohol/Drug Use strategy to the levels of the sense of meaningfulness

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [5.10SenseofCoherenceQuestionare.docx](#)
- [5.10.MiniCope.docx](#)