Serum Levels of Vitamin D, Retinol, Zinc and CRP in Relation To Obesity Among Children and Adolescents

Yan Zou

Zhejiang Provincial Center for Disease Control and Prevention <https://orcid.org/0000-0001-9740-317X>

Ronghua Zhang (✉ yingyang901@yeah.net)

Zhejiang Provincial Center for Disease Control and Prevention

Lichun Huang

Zhejiang Provincial Center for Disease Control and Prevention

Dong Zhao

Zhejiang Provincial Center for Disease Control and Prevention

Danting Su

Zhejiang Provincial Center for Disease Control and Prevention

Jia Meng

Zhejiang Provincial Center for Disease Control and Prevention

Yueqiang Fang

Zhejiang Provincial Center for Disease Control and Prevention

Research

Keywords: Vitamin D, Retinol, C-reactive protein, Obesity

Posted Date: September 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-915638/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at European Journal of Medical Research on April 4th, 2022. See the published version at <https://doi.org/10.1186/s40001-022-00670-7>.

Abstract

Background: The aim of this study was to examine the possible association between serum micronutrients (vitamin D, retinol, Zinc), C-reactive protein (CRP) and obesity among children and adolescents.

Methods: Weight and height were measured and serum 25-hydroxy vitamin D, serum retinol, serum Zinc and CRP were measured in 2818 children and adolescents (6-17 years of age), using the data of 10 investigation sites in Zhejiang Province. The difference of micronutrients (vitamin D, retinol, zinc), CRP among different BMI status were explored by ANOVA and chi-square test. The associated factors of micronutrients and CRP of overweight and obesity were explored by multifactor analysis.

Results: There were significant differences between male students and female students on BMI, 25(OH)D3, CRP, and there were significant differences between children and adolescents and between students living in urban area and rural area on BMI, 25(OH)D3, retinol, zinc ($p < 0.05$). There were significant differences on 25(OH)D3 and CRP level, and on retinol level among children and adolescents with different nutritional conditions ($p < 0.05$). In multifactor analysis, we found that gender (OR=0.626, 95%CI:0.486-0.806), living area (OR=0.666, 95%CI:0.515-0.860), age group (OR=0.894, 95%CI:0.856-0.934), high CRP concentration (OR=1.084, 95%CI:1.046-1.122) and appropriate retinol level (OR=1.536, 95%CI:1.185-1.991) were associated with students with obesity.

Conclusion: Gender, living area, age group, CRP concentration and vitamin A status were associated with children and adolescents with overweight and obesity. More attention in the intervention of overweight and obesity should be paid to boys living in urban areas, and high serum CRP and retinol level should also be concerned.

Introduction

The lancet commented that the prevalence of overweight and obesity in children and adolescents (5–19 years) was increasing[1]. The finding are consistent with other published studies[2]. WHO estimates that worldwide obesity has nearly tripled since 1975 and over 340 million children and adolescents aged 5–19 were overweight or obese in 2016[3]. In China, the prevalence of obesity also increased rapidly[4–5].

Obesity can lead to the increase of non-communicable diseases[4–6], especially children who are obese also have more metabolic and cardiovascular risk factors[7]. Children with excess weight are likely to become adults with obesity[8]. Early prevention of overweight and obesity is of great significance to children and their life cycle.

Meanwhile, vitamin A, vitamin D and zinc deficiencies have been reported among children and adolescents[9–10]. Because micronutrients play an important role in energy metabolism, it is necessary to study the relationship between micronutrients and obesity. C-reactive protein (CRP), as an indicator of low-grade inflammation, has been considered a possible risk factor for cardiovascular diseases associated with obesity[11], and CRP also plays as a major determinant of central obesity[12]. This study intends to learn the possible association between serum micronutrients, CRP and obesity among children and adolescents.

Methods

Study design

The data of this study are from the China National Nutrition and Health Survey 2016-2017 (CHNNS2016-2017). Children and adolescents from 10 investigation sites in Zhejiang Province were selected as participants to form provincial representative sample of Zhejiang Province.

Field investigation, physical examination and [blood specimen collection](#) was conducted between September 2016 and November 2017 in the sampling site. The nutritional status of children and adolescents were evaluated, and their level of

micronutrients (vitamin D, Retinol, Zinc), and CRP were measured.

Sampling method and study population

The data of this study is a part of the China National Nutrition and Health Survey 2016-2017, and that was a cross-sectional survey designed to examine the health and nutritional status of children and adolescents. In the China National Nutrition and Health Survey 2016-2017, stratified multistage probability sampling method was used and there were 10 surveillance sites including urban and rural area of Zhejiang provincial coverage were selected. According the the plan of the survey , at least 280 children and adolescents between 6 and 17 years old in each surveillance site should be interviewed. In this analysis, 2818 children and adolescents were included.

Ethics approval was obtained from the Ethical Committee of Zhejiang Provincial Center for Disease Control and Prevention. All student guardian provided written informed consent after the research protocols were carefully explained to them. Thus, informed consents from the parents/guardians of all participants were received.

Measurements and data collection

Demographic information were collected by general information questionnaire. Anthropometrical measurements were conducted by well-trained health workers of local community health center who followed a reference protocol recommended by the WHO[13]. Height was measured without shoes to the nearest 0.2 cm using a portable SECA stadiometer, and weight was measured without shoes and in light clothing to the nearest 0.1 kg on a calibrated beam scale. Body mass index(BMI) was calculated as weight (kg)/height(m)². Overweight and obesity were defined by the BMI cut-off points recommended in "Screening for overweight and obesity among school-age children and adolescents" (WS/T 586-2018) and low weight were defined by the BMI cut-off points recommended in "Screening standard for malnutrition of school-age children and adolescents" (WS/T 456-2014). Blood sample were collected to detect the concentration of retinol, and we judged marginal deficiency and deficiency of vitamin A among children and adolescents by boundary values of serum retinol level in "Method for Vitamin A deficiency screening"(WS/T 553-2017). Blood sample were collected to detect the concentration of 25-hydroxyvitamin D (25(OH)D), and according to standards recommended by the American endocrine association, we judged deficiency and inadequacy of vitamin D among children and adolescents by boundary values of the concentration of 25-hydroxyvitamin D (25(OH)D)[14].

Statistical analysis

All the data were analyzed with SAS9.4(SAS Institute Inc, Cary, NC, USA). Continuous variables were present in the form of mean and standard. Category variables were present in the form of number and percentage. The difference of micronutrients (vitamin D, retinol, zinc), CRP among different BMI status were explored by ANOVA and chi-square test. The associated factors of micronutrients and CRP of overweight and obesity were explored by multifactor analysis. A two side P<0.05 was utilized to assess statistical significance.

Results

Subject characteristics

Totally, there were 2818 children and adolescents included in this study, and males accounted for 50.2%. The prevalence of low weight, eutrophic, overweight, obese were 6.7%, 59.2%, 22.9% and 11.2%, respectively. The serum concentration of 25(OH)D3, retinol, zinc and CRP were 2.8 ± 0.7 ng/ml, 2.7 ± 0.6 mg/dL, 110.9 ± 42.6 mg/dL and 1.0 ± 4.1 mg/dL respectively.

Serum Levels Of 25(Oh)d3, Retinol, Zinc And Crp Characteristics

There were significant differences between males and females on BMI, 25(OH)D3, CRP, and males had higher serum concentration of 25(OH)D3, CRP than that of females. There were significant differences between children and adolescents and between that living in urban areas and rural areas on BMI, 25(OH)D3, retinol, zinc ($p < 0.001$) (Table 1).

Table 1
BMI and serum levels of 25(OH)D3, retinol, zinc and CRP among children and adolescents stratified by gender, age group and living area

Variables	Total	Gender		p	Age group		P	Living area		
		Males	Females		Children	Adolescents		Urban	Rural	p
Height(cm)	145.7 ± 17.4	147.1 ± 18.8	144.3 ± 15.7	0.000	130.5 ± 7.9	156.6 ± 11.3	0.000	147.1 ± 17.5	145.4 ± 17.3	0.442
Weight(kg)	40.4 ± 15.1	41.8 ± 16.2	39.0 ± 13.9	0.005	28.6 ± 7.2	48.4 ± 12.8	0.000	41.2 ± 15.4	39.6 ± 14.9	0.005
BMI (kg/cm ²)	18.3 ± 3.8	18.6 ± 3.7	18.1 ± 3.9	0.001	16.6 ± 2.9	19.5 ± 3.5	0.000	18.7 ± 4.1	18.0 ± 3.4	0.000
25(OH)D3(ng/ml)	2.8 ± 0.7	2.9 ± 0.7	2.7 ± 0.7	0.000	3.0 ± 0.7	2.7 ± 0.7	0.000	2.7 ± 0.7	2.9 ± 0.7	0.000
Retinol(mg/dL)	2.7 ± 0.6	2.6 ± 0.6	2.7 ± 0.5	0.074	2.5 ± 0.6	2.7 ± 0.5	0.000	2.8 ± 0.5	2.6 ± 0.6	0.000
Zinc(mg/dL)	110.9 ± 42.6	112.1 ± 41.3	109.7 ± 44.0	0.148	114.7 ± 55.5	108.6 ± 34.1	0.000	107.0 ± 28.8	114.9 ± 52.8	0.001
CRP (mg/dL)	1.0 ± 4.1	1.2 ± 5.3	0.8 ± 2.3	0.042	0.9 ± 2.5	1.0 ± 4.4	0.707	1.1 ± 4.9	0.9 ± 3.0	0.339

There were significant differences on serum concentrations of 25(OH)D3, CRP and retinol among children and adolescents with different nutritional conditions ($p < 0.05$). Children and adolescents with obesity had higher concentration of 25(OH)D3 and CRP (Table 2). Children and adolescents with different nutritional conditions had different vitamin A status ($p < 0.05$), while their vitamin D status were similar (Table 3). Trend chi-square test showed that the higher the vitamin A status, the larger the BMI value ($p = 0.009$).

Table 2
Serum levels of 25(OH)D3, retinol, zinc and CRP among children and adolescents with different nutritional status

	Low weight (N = 177)	Euthophic (N = 1980)	Overweight (N = 304)	Obese (N = 297)	F	p
25(OH)D3(ng/ml)	2.8 ± 0.7	2.8 ± 0.7	2.8 ± 0.7	2.9 ± 0.7	2.612	0.050
Retinol(mg/dL)	2.6 ± 0.6	2.7 ± 0.6	2.7 ± 0.6	2.7 ± 0.5	12.309	0.069
Zinc(mg/dL)	108.0 ± 34.7	110.9 ± 46.9	110.5 ± 29.8	113.5 ± 27.1	0.671	0.570
CRP(mg/dL)	0.8 ± 2.7	0.8 ± 3.7	1.1 ± 2.6	2.4 ± 6.9	15.022	0.000

Table 3
Vitamins D and A status among children and adolescents with different nutritional status

	Vitamin D status*				Vitamin A status**		
	Deficiency	Inadequacy	Normal	Appropriate	Deficiency	Marginal deficiency	Appropriate
Low weight	3(1.7%)	61(34.5%)	83(46.9%)	30(16.9%)	11(6.2%)	49(27.7%)	117(66.1%)
Euthophic	37(2.0%)	656(35.7%)	850(46.2%)	297(16.1%)	78(4.2%)	478(26.0%)	1285(69.8%)
Overweight	7(2.3%)	110(36.2%)	138(45.4%)	49(16.1%)	14(4.6%)	65(21.4%)	225(74.0%)
Obese	1(0.3%)	86(29.0%)	144(48.5%)	66(22.2%)	14(4.7%)	54(18.2%)	229(77.1%)
* $\chi^2 = 15.443$, p = 0.079							
** $\chi^2 = 12.704$, p = 0.048; trend $\chi^2 = 6.886$, p = 0.009							

Multifactor analysis for the associated micronutrients and CPR of overweight and obesity

In multifactor analysis, we found that gender(OR = 0.687, 95%CI:0.569–0.829), living area (OR = 0.668, 95%CI:0.552–0.809), age(OR = 0.952, 95%CI:0.922–0.982), high CRP concentration (OR = 1.080,95%CI:1.043–1.119) and appropriate vitamin A status (OR = 1.318, 95%CI:1.094–1.589) were associated with children and adolescents with overweight. And gender(OR = 0.626, 95%CI:0.486–0.806), living area (OR = 0.666, 95%CI:0.515–0.860), age(OR = 0.894, 95%CI:0.856–0.934), high CRP concentration (OR = 1.084,95%CI:1.046–1.122) and appropriate vitamin A status (OR = 1.536, 95%CI:1.185–1.991) were associated with children and adolescents with obesity(Table 4).

Table 4
Multifactor analysis for the associated micronutrients and CRP of overweight and obesity

Factors	B	S.E.	Wald	P	OR	95% C.I.for OR	
						Lower	Upper
Multifactor analysis for children and adolescents with overweight							
Gender	-0.376	0.096	15.356	0.000	0.687	0.569	0.829
Living area	-0.403	0.098	17.073	0.000	0.668	0.552	0.809
Age	-0.049	0.016	9.359	0.002	0.952	0.922	0.982
CRP (mg/dL)	0.077	0.018	18.296	0.000	1.080	1.043	1.119
Vitamin A status	0.276	0.095	8.431	0.004	1.318	1.094	1.589
Vitamin D status	0.058	0.068	0.726	0.394	1.059	0.928	1.209
Multifactor analysis for children and adolescents with obesity							
Gender	-0.469	0.129	13.182	0.000	0.626	0.486	0.806
Living area	-0.407	0.131	9.708	0.002	0.666	0.515	0.86
Age	-0.112	0.022	24.908	0.000	0.894	0.856	0.934
CRP (mg/dL)	0.080	0.018	19.913	0.000	1.084	1.046	1.122
Vitamin A status	0.429	0.132	10.501	0.001	1.536	1.185	1.991
Vitamin D status	0.149	0.09	2.732	0.098	1.161	0.973	1.386

Discussion

In the present study, gender, living area, age group, CRP concentration and vitamin A status were associated with children and adolescents with overweight and obesity.

In line with our findings, some systematic reviews have reported on growing prevalence of obesity and overweight among children and adolescents. A meta-analysis concluded that the prevalence of obesity in children and adolescents aged 2 to 19 years in the United States in 2011–2014 was 17.0%, while the prevalence of extreme obesity was 5.8%[15]. A systematic review in Asia, reported that the overall prevalence by gender was 7.0% and 4.8% in boys and girls for obesity in children, and 11.7% and 10.9% in boys and girls respectively for overweight in children[16]. We found that the prevalence of overweight and obesity were 22.9% and 11.2%, respectively, higher than the previous literature. Similarly, we also found that the prevalence among males were higher than females. We found higher prevalence of obesity and overweight in urban areas. Conversely, a study in Bangladesh reported that

overweight and obesity was associated with rural participants[17]. A study of China base on 1995–2014, revealed that overweight and obesity increased in Chinese children and adolescents, particularly in rural areas[5]. Our study was carried out in 2016–2017, whether there is a long-term trend in the distribution of obesity in urban and rural areas still needs to be further studied.

CRP is produced in the liver, macrophages, and adipose tissues. Elevated serum CRP is associated with obesity among children. A study in Japan reported that CRP increased the risk of obesity in school children[18]. Nappo reported that CRP levels are associated to higher body mass and overweight/obesity risk in a large population of European children. Children with higher baseline levels of CRP were at higher risk of developing overweight/obesity during growth[19]. We confirmed that

CRP level was higher in obese or overweight than that in non-obese children, as reported in previous studies. The mechanism lies in that obese children present increased oxidative stress and impaired inflammation and insulin sensitivity, which in turn result in similar impaired endothelial dysfunction and early signs of atherosclerosis[20]. Obesity is a pro-inflammatory state that may predispose patients to acute coronary syndrome characterized by chronic low grade inflammation resulting in endothelial dysfunction[21]. Thus, CRP play an important role in the prevention of chronic inflammation associated with obesity in the early stages of life.

Our previous study reported that the prevalence of vitamin A deficiency was 4.5%, and the prevalence of marginal deficiency of vitamin A was 24.7%[22]. At present, the nutritional status of vitamin A is not optimistic. This study found that the retinol level of obese students is higher than that of non obese children, which may be related to the dietary structure of obese children[23]. Liang reported that vitamin A metabolism may be disordered in obese children, although children with obesity have higher vitamin levels than lean children[24]. Gajewska reported that BMI value may influence the vitamin A status in obese children after therapy[25]. Gajewska suggest there is an occurrence of relationships between vitamin A and oxidized LDL in prepubertal obese children. Vitamin A concentrations is associated with dyslipidemia[26]. The relationship and mechanism between obesity and vitamin A should be further explored in the future study.

Shaikh reported that children with high BMI showed extremely high prevalence of vitamin D deficiency[27]. Kang reported that during the coronavirus disease-2019 (COVID-19) pandemic, increased childhood obesity and vitamin D deficiencies were observed[28]. Meta-analysis reported that children and adolescents with obesity have higher risk of vitamin D deficiency[29–30]. Also, A positive association between obesity and lower 25(OH)D serum concentration was found among Chinese adults[31]. Our study found that children and adolescents with obesity had higher concentration of 25(OH)D3, while in the multifactor analysis, vitamin D status was not the influencing factors for children and adolescents with obesity or overweight. The next step of research will further explore the influencing factors of vitamin D status, in order to better explore the relationship between vitamin D and obesity. It is suggested to encourage physical exercise, the reduction of screen time, and healthy eating habits in order to reduce the prevalence of overweight and obesity in children and adolescents and the impact of associated comorbidities, to keep the appropriate micronutrient status[32].

In China, measurement of serum CRP levels is not included in school health checkups, and it is difficult to obtain informed consent for blood sampling from school children and their parents. Thus, few serological surveys have been performed in healthy children. We performed serological tests in this the China National Nutrition and Health Survey 2016–2017 and evaluated the relationship between the test results and obesity.

Limitations of the present study should be noted. Since it is a cross-sectional study, causality cannot be established. Obesity and micronutrient status may be related to dietary intake and physical activity, which needs to be further explored in future research.

Conclusions

In summary, the present study documented gender, living area, age group, CRP concentration and vitamin A status were associated with children and adolescents with overweight and obesity. More attention in the intervention of overweight and obesity should be paid to boys living in urban areas, and high serum CRP and retinol level should also be concerned.

Declarations

Acknowledgments

We are grateful to all the school health care staffs participating in this study.

Authors' contributions

Conceptualization, YZ and RZ; methodology, YZ and LH; software, DS and DZ; investigation, YF and JM; resources, YZ and RZ; writing, YZ; project administration, YZ. The authors read and approved the final manuscript.

Funding

This study was supported by the Basic Public Welfare Research Plan of Zhejiang Province (LGF19H260002) and National Key Research and Development Project (2018YFC1603103).

Availability of data and materials

The supporting data can be acquired via correspondence author.

Ethical approval and consent to participate

This study was approved by the ethics committee of Zhejiang Provincial center for disease control and prevention. A signed consent by each participant was asked to provide.

Consent for publication

All the authors were consent for publication.

Conflict of interests

The authors declare no conflict of interest.

References

1. Dietz WH. Double-duty solutions for the double burden of malnutrition. *Lancet*. 2017;390(10113):2607–8.
2. de Onis M, Blössner M, Borghi E. Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr*. 2010;92(5):1257–64. doi:10.3945/ajcn.2010.29786.
3. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.
4. Dong B, Zou Z, Song Y, et al. Adolescent Health and Healthy China 2030: A Review. *J Adolesc Health*. 2020;67(5S):24–31.
5. Dong Y, Jan C, Ma Y, et al. Economic development and the nutritional status of Chinese school-aged children and adolescents from 1995 to 2014: an analysis of five successive national surveys. *Lancet Diabetes Endocrinol*. 2019;7(4):288–99.
6. Di Cesare M, Soric M, Bovet P, et al. The epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. *BMC Med*. 2019;17(1):212.
7. Friedemann C, Heneghan C, Mahtani K, et al. Cardiovascular disease risk in healthy children and its association with body mass index: systematic review and meta-analysis. *BMJ*. 2012;345:e4759.
8. Freedman DS, Khan LK, Dietz WH, et al. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa Heart Study. *Pediatrics*. 2001;108(3):712–8.
9. Labadarios D, Steyn NP, Maunder E, et al. The National Food Consumption Survey (NFCS): South Africa, 1999. *Public Health Nutr*. 2005;8(5):533–43.
10. Senekal M, Nel J, Malczyk S, et al. Provincial Dietary Intake Study (PDIS): Micronutrient Intakes of Children in a Representative/Random Sample of 1- to < 10-Year-Old Children in Two Economically Active and Urbanized Provinces in South Africa. *Int J Environ Res Public Health*. 2020;17(16):5924.
11. Ridker PM. Inflammatory biomarkers and risks of myocardial infarction, stroke, diabetes, and total mortality: implications for longevity. *Nutr Rev*. 2007;65(12 Pt 2):253–9.
12. Santos AC, Lopes C, Guimarães JT, et al. Central obesity as a major determinant of increased high-sensitivity C-reactive protein in metabolic syndrome. *Int J Obes (Lond)*. 2005;29(12):1452–6.

13. WHO. Physical Status: The use and interpretation of anthropometry; Report of a WHO Expert Committee. World Health Organization Technical Report Series no. 854; WHO: Geneva, Switzerland. 1995;1-452.
14. American Endocrine Association. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. 2011.
15. Ogden CL, Carroll MD, Lawman HG, et al. Trends in Obesity Prevalence Among Children and Adolescents in the United States, 1988–1994 Through 2013–2014. *JAMA*. 2016;315(21):2292–9.
16. Mazidi M, Banach M, Kengne AP. Lipid and Blood Pressure Meta-analysis Collaboration Group. Prevalence of childhood and adolescent overweight and obesity in Asian countries: a systematic review and meta-analysis. *Arch Med Sci*. 2018;14(6):1185–203.
17. Mostafa I, Hasan M, Das S, et al. Changing trends in nutritional status of adolescent females: a cross-sectional study from urban and rural Bangladesh. *BMJ Open*. 2021;11(2):e044339.
18. Nishide R, Ando M, Funabashi H, Yoda Y, Nakano M, Shima M. Association of serum hs-CRP and lipids with obesity in school children in a 12-month follow-up study in Japan. *Environ Health Prev Med*. 2015;20:116–22.
19. Nappo A, Iacoviello L, Fraterman A, Gonzalez-Gil EM, Hadjigeorgiou C, Marild S, Molnar D, Moreno LA, Peplies J, Sioen I, Veidebaum T, Siani A, Russo P. High-sensitivity C-reactive protein is a predictive factor of adiposity in children: results of the identification and prevention of dietary- and lifestyle-induced health effects in children and infants (IDEFICS) study. *J Am Heart Assoc*. 2013;2:e000101.
20. Giannini C, de Giorgis T, Scarinci A, Cataldo I, Marcovecchio ML, Chiarelli F, Mohn A. Increased carotid intima-media thickness in pre-pubertal children with constitutional leanness and severe obesity: the speculative role of insulin sensitivity, oxidant status, and chronic inflammation. *Eur J Endocrinol*. 2009;161:73–80.
21. Singh R, Verma A, Aljabari S, Vasylyeva TL. Urinary biomarkers as indicator of chronic inflammation and endothelial dysfunction in obese adolescents. *BMC Obes*. 2017;4,11.
22. Zou Y, Zhang R, Huang L, Su D, He M, Fang Y, Zhao D, Wang W, Han D. Sociodemographic disparity in the nutritional status among children and adolescents in Zhejiang Province. *Eur J Clin Nutr*. 2021;75:307–13.
23. Zou Y, Zhang R, Xia S, Huang L, Meng J, Fang Y, Ding G. Dietary Patterns and Obesity among Chinese Adults: Results from a Household-Based Cross-Sectional Study. *Int J Environ Res Public Health*. 2017;14,487.
24. Liang X, He Y, Chen M, Ping Q, Chen J. The association of lecithin retinol acyltransferase and the 25(OH)D receptor with pediatric overweight and obesity. *Eur J Clin Nutr*. 2019;73:1626–9.
25. Gajewska J, Ambroszkiewicz J, Szamotulska K, Klemarczyk W, Weker H, Chowska M. Associations Between Antioxidant Vitamin Status, Dietary Intake, and Retinol-Binding Protein 4 Levels in Prepubertal Obese Children After 3-Month Weight Loss Therapy. *J Clin Res Pediatr Endocrinol*. 2020, doi: 10.4274/jcrpe.galenos.2020.2020.0207. Epub ahead of print. PMID: 33261247.
26. Gajewska J, Ambroszkiewicz J, Weker H, Chowska M. Relations between oxidized low-density lipoproteins and fat-soluble vitamin concentrations in obese children - preliminary study. *Dev Period Med*. 2017;21:266–71.
27. Al Shaikh A, Farahat F, Abaalkhail B, Kaddam I, Aseri K, Al Saleh Y, Al Qarni A, Al Shuaibi A, Tamimi W. Prevalence of Obesity and Overweight among School-Aged Children in Saudi Arabia and Its Association with Vitamin D Status. *Acta Biomed*. 2020;91,e2020133.
28. Kang HM, Jeong DC, Suh BK, Ahn MB. The Impact of the Coronavirus Disease-2019 Pandemic on Childhood Obesity and Vitamin D Status. *J Korean Med Sci*. 2021;36,e21.
29. Fiamenghi VI, Mello ED. Vitamin D deficiency in children and adolescents with obesity: a meta-analysis. *J Pediatr (Rio J)*. 2020;S0021-7557(20):30207–2.
30. Pereira-Santos M, Costa PR, Assis AM, Santos CA, Santos DB. Obesity and vitamin D deficiency: a systematic review and meta-analysis. *Obes Rev*. 2015;16:341–9.
31. Ren Z, Zhao A, Wang Y, Meng L, Man-Yau, Szeto I, Yang C, Wang M, Zhang J, Wu W, Wang P, Zhang Y. Association of serum 25-hydroxy vitamin D with obesity-related indices in Chinese adults: A cross-sectional study. *Food Sci Nutr*.

2021;9:2260–8.

32. Kumar AS, Kelly. Review of childhood obesity: from epidemiology, etiology, and comorbidities to clinical assessment and treatment. Mayo Clin Proc. 2017;92:251–65.