

# Reduction of Testosterone Levels in *Schistosoma Haematobium* or *Schistosoma Mansoni* Infected Men: A Cross-Sectional Study in Two Schistosomiasis-Endemic Areas of the Adamawa Region of Cameroon

Hermine BOUKENG JATSA (✉ [mjatsa@yahoo.fr](mailto:mjatsa@yahoo.fr))

University of Yaounde I: Universite de Yaounde I <https://orcid.org/0000-0002-9838-8570>

Ulrich MEMBE FEMOE

University of Yaounde I: Universite de Yaounde I

Calvine NOUMEDEM DONGMO

University of Yaounde I: Universite de Yaounde I

Romuald Issiaka NGASSAM KAMWA

University of Maroua: Universite de Maroua

Betrand NONO FESUH

University of Yaounde I: Universite de Yaounde I

Louis-Albert TCHUEM TCHUENTE

University of Yaounde I: Universite de Yaounde I

Pierre KAMTCHOUING

University of Yaounde I: Universite de Yaounde I

---

## Research Article

**Keywords:** male genital schistosomiasis, testosterone, testes, *Schistosoma haematobium*, *Schistosoma mansoni*, Cameroon

**Posted Date:** October 12th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-915844/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## Abstract

**Background:** The incidence of schistosomiasis-induced male reproductive dysfunction and infertility is probably underestimated comparing to female genital schistosomiasis. This study aimed to investigate the impact of *S. haematobium* or *S. mansoni* infection on the reproductive function of adult men in Tibati and Wouldé, two schistosomiasis endemic areas in the Adamawa region of Cameroon.

**Methods:** A total of 89 men in the reproductive age (range: 14 – 56 years) from two localities were enrolled in the study with 51 in Tibati and 38 in Wouldé. Each participant was submitted to a questionnaire to document data on sociodemographic and stream contact behaviors. A medical examination was performed to measure the circumference of the testes and to evaluate genital tract pathologies. Stool and urine samples were collected and screened for the presence of *S. haematobium* or *S. mansoni* ova. Blood serum was used to evaluate the level of transaminases and testosterone.

**Results:** *S. haematobium* was present only in Tibati with a prevalence of 31.37%. *S. mansoni* prevalence was 3.92% at Tibati and 44.71% in Wouldé. The intensity of infection was  $22.12 \pm 9.57$  eggs/10 mL for *S. haematobium* and  $128.10 \pm 3.76$  egg for *S. mansoni*. Serum transaminases activity and the mean testicular circumference of *Schistosoma*-positive individuals were close to those of negative ones. The testes size was however higher in *S. mansoni*-positive individuals than in *S. haematobium*-positive ones ( $P < 0.05$ ). The serum testosterone level of *S. haematobium* and *S. mansoni*-positive men was significantly reduced by 56.07% ( $P < 0.001$ ) and 51.94% ( $P < 0.01$ ) respectively in comparison to that of *Schistosoma*-negative ones. A significant and negative correlation was established between schistosomiasis and the low level of serum testosterone. Male genital tract pathologies such as scrotal abnormalities, varicocele, nodular epididymis, inguinal hernia and hydrocele were recorded in both *Schistosoma*-positive and *Schistosoma*-negative men. However, no significant link was established between schistosomiasis infection and these pathologies.

**Conclusion:** These results demonstrated that infection with *S. haematobium* or *S. mansoni* is associated with a low production of the reproductive hormone testosterone, and may be a major cause of male infertility.

## Introduction

Schistosomiasis is an acute and chronic disease caused by blood flukes of the genus *Schistosoma*. In Cameroon, three species of human schistosomes occur: *S. haematobium*, *S. mansoni* and *S. guineensis*. People are infected while exposed to infested water during routine agricultural, domestic, occupational, and recreational activities. In 2016, the global burden of schistosomiasis was estimated at 2.50 million disability-adjusted life years (DALYs). Schistosomiasis remains a major public health problem with at least 229 million people worldwide and almost 90% of them living in sub-Saharan Africa, requiring preventive chemotherapy in 2018 [1]. In Cameroon, it was estimated that 47% of the population (9 484 894 million) was at risk of schistosomiasis infection and needed preventive treatment in 2010 [2]. The three northern regions of the country, namely Far North, North and Adamawa are the most affected, and some health districts in the South-West, Littoral, West and Centre regions showed high prevalence of infection. Epidemiological data collected in some schools of the Adamawa region in 2006 revealed that 14.2% of the children were infected with at least one of the schistosomes species [3]. The strategy of choice to control schistosomiasis remains preventive chemotherapy with praziquantel. School-aged children are the primary target of these therapeutic interventions since they are the most vulnerable group and the most affected by the infection [4]. In Cameroon, fishermen, farmers and irrigation workers are also high-risk groups for schistosome infection; but they are not enrolled in deworming campaigns with praziquantel [3, 5]. *Schistosoma* – infected adults generally suffered from chronic schistosomiasis with severe consequences as liver fibrosis, portal hypertension, ascites, calcification of the bladder, kidney failure and genital dysfunction [2]. Genital schistosomiasis occurs when *Schistosoma* circulating eggs invade the reproductive tissues of the host. It has been associated with female and male reproductive health problems such as ectopic pregnancies, abortion, low sperm count, coitus pain, vaginal bleeding, genital organs enlargement and infertility [6–8]. Female genital schistosomiasis (FGS) is well documented and its prevalence is considerable in *S. haematobium* endemic areas (33 to 75%) [6, 9–12]. In contrast to FGS, male genital schistosomiasis (MGS) is under reported and its burden is underestimated. *S.*

*haematobium* and *S.mansoni* are the two species involved in MGS. The main findings reported in patients who have travelled or lived in schistosomiasis endemic areas are swelling of scrotum and other genital organs, dilatation and calcification of the prostate and seminal vesicles, haemospermia, hydrocele, changes in semen/ejaculate, infertility and urethral discharge [7, 13]. Case reports have mentioned the presence of *S. haematobium* or *S. mansoni* eggs in male genital organs and semen of some young and adult patients [14–24]. Variation in the levels of sex hormones was also reported in some male patients with *S. haematobium* and / or *S. mansoni* [25–29].

In Cameroon, investigation on the association of schistosomiasis and the male reproductive tract have drawn no attention from the researchers. To the best of our knowledge, this is the first report on male genital schistosomiasis in Cameroon. The aim of this study was to evaluate the impact of the infection with *S. haematobium* or *S. mansoni* on the reproductive tract of adult men in Tibati and Wouldé, two schistosomiasis endemic areas in the Adamawa region of Cameroon.

## Methods

### Study area

Cameroon is divided in 10 regions at the first level, 58 divisions at the second level, and 360 sub-districts at the third level. According to the United Nations, the population of Cameroon was estimated to 25 876 380 inhabitants in 2019, with 1 200 970 inhabitants in the Adamawa region [30]. The Adamawa region is divided in five divisions: Vina, Mbéré, Mayo Banyo, Djerem and Faro et Déo. Tibati, the capital of the “Djerem” division is situated between 6°47' N and 12°63' E. It is a semi – urban area with a population density of 7 inhabitants /km<sup>2</sup>. Situated in “Pagneré” neighborhood, Lake Pagneré is the biggest one in the city. Trading, fishing and fish mongering are the main activities in Tibati. Wouldé is a village in the Mayo-Baléo sub-district in “Faro et Déo” division. It is situated between 7°43' N and 12°47' E and surrounded by volcanic hills. At the entrance of this rural area flows the river Wouldé with its thermal source. The population density is 6.6 inhabitants /km<sup>2</sup> and the major occupations in this community is farming, livestock and trading (Fig. 1). The vegetation of the Adamawa region falls within the rainforest – savannah mosaic and the climate is a sudano – guinean type with an annual rainfall of 1500 mm. Transmission of schistosomiasis presumably takes place at the lake, river, ponds and marshy fields. Fulbe and Kutin ethnic groups are predominant in Tibati and Wouldé and they are mainly Muslims.

### Ethical considerations

This study was approved by the National Ethics Committee of Cameroon (N° 072/CNE/DNM/08). Surveys were conducted in the communities with the approval of administrative authorities. The objectives of the study were explained to the populations and an informed consent was signed by each participant. Results were sent to each participant at the end of the study and all infected persons were treated with praziquantel.

### Study design

People involved in the study were all men volunteers, residents of Lake Pagneré in Tibati and of “Chefferie” neighbourhood in Wouldé. Seriously ill individuals showing incapacitation were not included in the study. A questionnaire was administered to each participant for personal information, fertility status and history of sexually transmissible infections. Participants were submitted to physical examination by a male medical practitioner to assess by palpation genital tract pathologies as varicocele, hydrocele, nodular epididymis, scrotal abnormalities and inguinal hernia. The circumference of the two testes was also measured. Initially planned in the study, sperm collection to search for the presence of schistosome ova and to conduct semen analysis was not carried out because of cultural considerations (masturbation being a taboo in those localities). Urine and stool samples as well as blood were collected from each participant for parasitological and biochemical analyses.

### Samples collection

Urine and stool samples were collected in 60 mL plastic screw-cap vials and blood in 5 mL dry tubes, between 10.00 a.m and 14.00 p.m. The samples were transported to the laboratory of the Divisional Hospital of Tibati or to the Health district of

Tignère for processing. In the laboratory, each urine sample was agitated to ensure adequate dispersal of eggs and 10 mL of urine were filtered through a Nucleopore® filter. The filters were examined by microscopy for the presence of schistosome eggs. Stool samples were examined by a single thick smear technique using a 41.7 mg Kato-Katz template. Number of eggs for each schistosome specie was counted and intensity of infection was calculated and expressed as eggs per gram of feces (epg) for *S. mansoni* or eggs per 10 ml of urine (eggs/10 ml) for *S. haematobium*. Serum obtained from blood samples after coating were kept as -20°C and transported to the laboratory in a cold ice box for the assay.

## Evaluation of the level of transaminases

Serum activities of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) were evaluated by colorimetric method using the Fortress Commercial kits (Fortress Diagnostics, UK) according the protocol given by the manufacturer. Absorbances were read against the blank at 505 nm and the ALT and AST activities were determined from their respective standard curve.

## Testosterone assay

Concentrations of serum testosterone was evaluated by an enzyme immunoassay using the Human Testosterone ELISA kit (Invitrogen). A volume of 50 µL of standards and samples were dispensed in microtiter plates. After adding 100 µL of testosterone – HRP conjugate (testosterone labeled with horseradish peroxidase), plates were incubated for 60 min at room temperature. The competitive reaction was stopped by washing with the wash buffer. The chromogenic solution of 3, 3', 5, 5' tetramethylbenzidine (100 µL) was added to each well and after incubation for 30 min at room temperature, 100 µL of the stop reagent (HCl 2N) was added. Absorbance was read at 450 nm and testosterone concentrations were determined from the standard curve.

## Data analysis

The data for this study was entered into Excel and exported to the R version 3.5.0 software for statistical analysis. In univariate analysis, measures of central tendency and dispersion were reported for quantitative variables while the frequencies and percentages were reported for the modalities of qualitative variables. In bivariate analysis, frequencies and percentages were determined respectively for *Schistosoma*-infected and non-infected subjects to evaluate its prevalence across groups of other qualitative variables. Equality of prevalence rates were tested using the Pearson's Chi-squared test for equality of proportions. Differences in distributions for quantitative variables between SCH-positive and SCH-negative subjects were tested using the Kolmogorov-Smirnov test. Univariate logistic regression models were used to identify factors which are affected by *Schistosoma* infection. For the quantitative results of biochemical parameters, data were analysed using GraphPrism 8.0.1, by one-way analysis of variance (ANOVA) and differences between groups were assessed using the Tukey multiple comparison post-test. In all, level of significance was set at 5%. The intensity of individual infection for *S. mansoni* (light: 1–99 epg; moderate: 100–399 epg; and heavy  $\geq$  400 epg) and for *S. haematobium* (light: < 50 eggs/10mL and heavy:  $\geq$  50 eggs/10mL) was classified according to the WHO guidelines [31].

## Results

### Socio-demographic characteristics of the participants

A total number of 89 men in reproductive age (range: 14–56 years) were enrolled in this study with 51 in Tibati and 38 in Wouldé. The mean age of the population was  $32.21 \pm 1.16$  years. Regarding the marital status of the participants, 53 (58.43%) were married, 32 (35.96%) were single with girlfriends, 4 (4.49%) were divorced and 1 (1.12%) was widowed. The results revealed that 49.44 % of the population were permanently in contact with schistosomiasis transmission sites (STS). Considering the study population per locality, 76.32% of the participants in Wouldé, and 29.41% in Tibati were exposed to STS. (Table 1).

Table 1  
Sociodemographic characteristics of the population of Tibati and Wouldé

Sociodemographic indicators	Category	Study areas	SCH-negative	SCH-positive	Total	Mean ± SEM / Prevalence
			(n = 54)	(n = 35)		
Age (years)	Mean number		33.87	29.66	89	32.21 ± 1.16
Weight (kg)	Mean number		64.37	61.43	89	63.21 ± 0.89
Height (cm)	Mean number		169.22	166.17	80	167.89 ± 0.88
Profession	No frequent contact with water	Tibati	21	15	36	70.59%
		Wouldé	2	7	9	23.68%
		Total	23	22	45	51.68%
	Frequent contact with water	Tibati	12	3	15	29.41%
		Wouldé	19	10	29	76.32%
		Total	31	13	44	49.44%
Marital status	Single		16	16	32	35.96%
	Divorced		4	0	4	4.49%
	Married		34	18	52	58.43%
	Widowed		0	1	1	1.12%

Considering the indicator “profession”, the prevalence is the percentage of people by locality whose occupation exposes them or not to frequent contact with water.

SCH-negative: negative to *Schistosoma* infection

SCH-positive: positive to *Schistosoma* infection

## Prevalence and intensity of infection

Table 2 summarizes the prevalence and intensity of *S. haematobium* and *S. mansoni* infection in the male adult individuals involved in this study. *S. haematobium* was the most prevalent species in Tibati with 31.37 % while the prevalence of *S. mansoni* was 3.92%. In Wouldé, *S. haematobium* was absent among the population of study and the prevalence of *S. mansoni* was 44.74 %. The eggs count for urinary schistosomiasis ranged from 2 to 2620 eggs/10 mL and from 24 to 1224 epg for intestinal schistosomiasis. The geometric mean infection intensity was 22.12 ± 9.57 eggs/10 mL for *S. haematobium* and 128.10 ± 3.76 epg for *S. mansoni*. According to the degree of intensity of infection, 47.37% of *S. mansoni*-infected men presented light-intensity infection, 31.58% moderate-intensity infection and 21.05% heavy-intensity infection. Light-intensity infection with *S. haematobium* represented 68.75% while heavy-intensity infection was 31.25%. Hematuria was present in 80% of the men presenting *S. haematobium* heavy-intensity infection (Fig. 2).

Table 2  
Prevalence and intensity of *Schistosoma* infection in the male adult population of Tibati and Wouldé

Localities	<i>Schistosoma haematobium</i>				<i>Schistosoma mansoni</i>			
	Number of subjects	Number of positive	Prevalence (%)	Intensity of infection (eggs/10 mL)	Number of subjects	Number of positive	Prevalence (%)	Intensity of infection (eggs/g)
Tibati	51 (57.30%)	16	31.37	22.12 (6.64–73.72)	51	2	3.92	24
Wouldé	38 (42.70%)	0	0	/	38	17	44.74	128.10 (67.66–242.60)

Geometric mean with 95% CI is used to expressed the intensity of infection

## Serum activity of transaminases

The serum activity of transaminases was evaluated in the study population. The alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities of *S. haematobium* and *S. mansoni*-positive men were close to those of SCH-negative men. There was thus no significant difference of the serum activity of transaminases between SCH-positive men and SCH-negative ones (Table 3).

Table 3  
Transaminases activities of *Schistosoma haematobium* and *Schistosoma mansoni*-positive individuals of Tibati and Wouldé

Transaminases	SCH-negative	<i>S. haematobium</i> -positive	<i>S. mansoni</i> -positive
ALT (UI/L)	12.84 ± 1.26	12.19 ± 1.66	13.80 ± 1.52
AST (UI/L)	42.03 ± 4.26	43.40 ± 4.70	40.27 ± 2.44

## Testicular size and serum level of testosterone

The mean testicular circumference of SCH-negative men was 9.26 ± 0.38 cm and the one of *S. haematobium* or *S. mansoni*-positive men was 8.28 ± 0.49 cm and 10.71 ± 0.62 cm respectively. These results revealed a no significant difference between SCH-positive individuals and SCH-negative ones. The testicular circumference of *S. mansoni*-positive men was however higher than that of *S. haematobium*-positive men ( $P < 0.05$ ) (Fig. 3a).

The serum concentration of testosterone was 3.40 ± 0.48 ng/mL and 3.72 ± 0.58 ng/mL for *S. haematobium* and *S. mansoni*-positive men respectively versus 7.74 ± 1.07 ng/mL for SCH-negative men. In comparison to SCH-negative group, *Schistosoma* infection induced a significant reduction of the serum level of testosterone by 56.07% in *S. haematobium* positive ( $P < 0.001$ ) and by 51.94% in *S. mansoni* positive ( $P < 0.01$ ) (Fig. 3b). A strong correlation was observed between *Schistosoma* infection and the serum level of testosterone in the study population. The Spearman's rho correlation coefficient was -0.54 ( $P = 0.0101$ ) and -0.56 ( $P = 0.0006$ ) for the groups of *S. haematobium* and *S. mansoni* infected men respectively (Fig. 4).

## Prevalence of genital tract pathologies

Table 4 shows the prevalence of genital tract pathologies in SCH-negative and SCH-positive men. Physical examination of the population of study revealed that some men infected with *S. haematobium* or *S. mansoni* presented one or two of the

following pathologies: varicocele, hydrocele, nodular epididymis, scrotal abnormalities and inguinal hernia. Scrotal abnormalities were predominant with 25 and 52.63% of the cases in *S. haematobium* or *S. mansoni* infection versus 40.74% of cases in non-infected men. Varicocele and nodular epididymis were also detected in about 20.00% of SCH-positive men. These pathologies were also present in SCH-negative men. The results of the bivariate logistic regression model show that the genital tract pathologies are not related to SCH infection: no statistical difference was observed between SCH-negative and SCH-positive men.

Table 4

Prevalence of the genital tract pathologies in *Schistosoma haematobium* or *Schistosoma mansoni*-positive men of Tibati and Wouldé

Pathology	SCH-negative (n = 54)	<i>Schistosoma</i> -infected individuals							
		<i>S. haematobium</i> (n = 16)				<i>S. mansoni</i> (n = 19)			
		Proportion	Proportion	Odds ratio	CI (95%)	P-value	Proportion	Odds ratio	CI (95%)
Scrotal abnormalities	22 (40.74%)	4 (25.00%)	0.4848	0.1565–1.6790	0.3782	10 (52.63%)	1.6160	0.5820–4.6740	0.4267
Nodular epididymis	13 (24.07%)	2 (12.50%)	0.4734	0.0968–2.1130	0.4919	5 (26.32%)	1.3990	0.4457–4.568	0.7461
Varicocele	11 (20.37%)	3 (18.78)	0.9020	0.2407–3.3764	> 0.9999	4 (21.05%)	1.0424	0.3250–3.9692	> 0.9999
Inguinal hernia	7 (12.96%)	1 (6.25%)	0.4387	0.0365–3.0033	0.6694	4 (21.05%)	1.8901	0.5449–7.9074	0.4508
Hydrocele	4 (7.41%)	1 (6.25%)	0.7678	0.05893–5.4633	> 0.9999	2 (10.53%)	1.6538	0.2865–7.8212	0.6257

## Discussion

In schistosomiasis endemic areas, people are infected during their routine activities when they are exposed to contaminated water. In this study about half of the population was exposed to schistosomiasis transmission sites due their agricultural and fishing activities. Three quarters of the study population in Wouldé was made up of farmers working in marshy fields around Wouldé river. The study population in Wouldé was then a high-risk group for schistosomiasis infection [3]. The prevalence of *S. haematobium* or *S. mansoni* infection was within the interval 10–49%; which reflects a moderate occurrence in the study population in Tibati or Wouldé. Despite this moderate endemicity of schistosomiasis, 80% of men with *S. haematobium* heavy-intensity infection presented hematuria. It reflects the increase of egg population in the urothelium and is sometimes linked to bladder and ureteral calcification and renal failure [32].

Among other complications of urinary schistosomiasis, male reproductive function impairment and even infertility are quite common. *S. haematobium* is reported to be more involved than *S. mansoni* [33]. The symptoms include epididymitis, haemospermia, pain during urination, prostatitis, dilatation and calcification of seminal vesicles and testicular inflammation [7, 34, 35]. In the current study, the mean testicular circumference of *S. mansoni*-positive men was significantly higher than that of *S. haematobium*-positive men. The increase size of the testes could presumably be the consequence of numerous eggs trapped in that organ or / and the inflammation induces by their presence in the testes. The presence of *S. haematobium* or *S. mansoni* ova in the testes has effectively been reported in men living in SCH-endemic areas or tourists who have visited these areas [14, 19, 21–23, 36]. *Schistosoma* ova in the testes induces a direct damage of the testicular tissue by bilharzial inflammation and granuloma formation [37] that could impair with the testicular steroidogenesis. The testis is an exocrine and endocrine gland which synthesizes and produces testosterone through the process of steroidogenesis in the Leydig cells and sperm through spermatogenesis in the seminiferous tubules [38]. When *Schistosoma* eggs invade the testes,

inflammatory cells migrate around eggs and generate granulomatous lesions. Lenzi et al. [39] has described the bilharzial granuloma as a structure composed by macrophages, lymphocytes, eosinophils, neutrophils, giant cells and fibroblasts surrounding schistosome eggs. The modulation of the Leydig cells steroidogenesis by macrophages has been demonstrated [40–42]. Under normal physiological and non-inflammatory conditions, testicular interstitial macrophages play an important role in Leydig cell proliferation and differentiation, and also stimulate its steroidogenic function [40, 41]. When macrophages are activated, they produce inflammatory cytokines such as tumor necrosis factor-alpha (TNF- $\alpha$ ) and interleukin-1 (IL-1) that inhibit the Leydig cells steroidogenesis, and thus the testosterone production. They also act as transcriptional repressors of steroidogenic enzyme gene expression [40, 43]. In fact, the exposure of TM3 Leydig cells to inflammatory cytokines TNF- $\alpha$ , IL-1 $\beta$  and IL-6 resulted in a decrease of Leydig cells viability and testosterone concentrations [42]. Moreover, the endocrine function of the testis is perturbed by the reactive oxygen species (ROS) generated by macrophages. They impair the Leydig cell mitochondria by inhibiting the Steroidogenic Acute Regulatory (StAR) protein expression [40]. The significant decrease of testosterone concentration recorded during the current study in *S. haematobium*-positive men and in *S. mansoni*-positive men could be linked to the inhibition of the Leydig cells steroidogenesis by secretory products of the granulomas' inflammatory cells. In addition, a significant correlation between *S. haematobium* or *S. mansoni* infection and the reduced testosterone level was established in our study. Hormonal imbalance in SCH-infected individuals is less reported. Authors have recorded a decreased level of hydroxyprogesterone, testosterone and dihydrotestosterone in boys and men presenting hepatosplenic schistosomiasis [26–29]. This was associated to an increase concentration of estradiol, estrone and estriol [25, 29], suggesting an intense aromatization of androgens to estrogens in SCH-infected persons. Such sex steroids imbalance has led to delayed puberty in young boys and to hypogonadism in men [27, 29].

Authors have described hydrocele, testicular atrophy and nodular scrotum in patients presenting testicular schistosomiasis [21]. In this study we denoted genital tract pathologies such as scrotal abnormalities, nodular epididymis, varicocele, hydrocele and inguinal hernia in SCH-positive men. Since the prevalence of these pathologies was statistically comparable between SCH-positive and SCH-negative men, it would have become a challenge to establish a correlation between schistosomiasis or reduced testosterone level and these pathologies. The testosterone level of SCH-negative men in this study was however similar to that of a control group of men of reproductive age involved in a study on the effect of agropesticides on the male reproductive function of farmers in a village in the West region of Cameroon [44].

The consequence of the testosterone depletion in SCH-positive men in Tibati and Wouldé would be a poor semen quality since spermatogenesis relies on the ability of Leydig cells to produce testosterone [38]. Due to traditional and cultural beliefs, it wasn't possible to conduct semen analysis in this study; the population of our study areas considering masturbation (method of sperm collection) as a taboo. Some authors have however reported *Schistosoma* egg excretion in the semen, low volume of ejaculate, decreasing of sperm viability, oligoasthenoteratozoospermia, azoospermia and sperm apoptosis [15–18, 20, 36, 45–47]. Leukocytospermia, mainly eosinophilia, and secretion of inflammatory cytokines in the semen were also reported in SCH-infected men [15, 17, 20, 36, 45]. A clinical scheme presenting a low testosterone level or/and a poor semen quality could presumably conducted to subfertility or infertility.

## Conclusion

The results from this study demonstrated that *S. haematobium* and *S. mansoni* infections are associated with the reduction of the male reproductive hormone testosterone of adult men living in schistosomiasis endemic areas in Cameroon. The recommendations from this study are that the national control programme for schistosomiasis and intestinal helminthiasis in Cameroon should accelerate the extension of schistosomiasis treatment to reach all individuals at risk in communities. In addition, a program of sensitization and surveillance of young people and adults suffering from male genital schistosomiasis should be set up in endemic areas.

## Declarations

*Ethics approval and consent to participate*

The protocol of the current study was approved by the National Ethics Committee of Cameroon (N° 072/CNE/DNM/08). All participants signed an informed consent.

### ***Consent for publication***

All presentations of data have consent of publication.

### ***Availability of data and materials***

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

### ***Competing interests***

The authors declare that they have no competing interests

### ***Funding***

Financial support was provided by the “Société d’Andrologie de Langue Française » (SALF) for the realization of biochemical study.

### ***Authors' contributions***

HBJ, LATT and PK conceived and design this study. HBJ and RIKN conducted field study. HBJ, UMF and CND computerized the raw data from the surveys and performed statistical analysis with BNF. HBJ realized biochemical assays and wrote the first draft of the manuscript with UMF. All authors read and approved the final manuscript.

### ***Acknowledgements***

Authors are grateful to M. Jean Marie Kagho, the medical practitioner who conduct medical examination of the participants

### ***Authors' information***

<sup>1</sup>Laboratory of Animal Physiology, Department of Animal Biology and Physiology, Faculty of Science, University of Yaoundé I, P.O. Box 812, Yaoundé, Cameroon

<sup>2</sup>Centre for Schistosomiasis and Parasitology, P.O. Box 7244, Yaoundé, Cameroon

<sup>3</sup>Laboratory of Parasitology and Ecology, Department of Animal Biology and Physiology, Faculty of Science, University of Yaoundé I, P.O. Box 812, Yaoundé, Cameroon

<sup>4</sup>Department of Biological Science, Faculty of Science, University of Maroua, P.O. Box 46, Maroua, Cameroon

<sup>5</sup>Laboratory of Mathematical engineering and information system, Department of Mathematics, National Advances School of Engineering of Yaoundé, University of Yaoundé I, P.O. Box 8390, Yaoundé, Cameroon

## **References**

1. World Health Organization. Schistosomiasis [Internet]. 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/schistosomiasis>.
2. World Health Organization. Progress report 2001–2011 and strategic plan 2012–2020. 2013.
3. Tchuem Tchuente L-A. National Programme for the Control of Schistosomiasis and Intestinal Helminthiasis 2003–2019 Progress report Cameroon. 2020.

4. World Health Organization. Preventive chemotherapy in human helminthiasis. 2006.
5. Tchuente LT, Rollinson D, Stothard JR, Molyneux D. Moving from control to elimination of schistosomiasis in sub-Saharan Africa: time to change and adapt strategies. *Infect Dis Poverty*. 2017;6:1–14.
6. Kjetland EF, Leutscher PDC, Ndhlovu PD. A review of female genital schistosomiasis. *Trends Parasitol*. 2012;28:58–65.
7. Kayuni S, Lampiao F, Makaula P, Juziwelo L, Lacourse EJ, Reinhard-Rupp J, et al. A systematic review with epidemiological update of male genital schistosomiasis (MGS): A call for integrated case management across the health system in sub-Saharan Africa. *Parasite Epidemiol Control*. 2018;3:e00077.
8. Sturt AS, Webb EL, Francis SC, Hayes RJ, Bustinduy AL. Beyond the barrier: Female Genital Schistosomiasis as a potential risk factor for HIV-1 acquisition. *Acta Trop*. 2020;209:105524.
9. Catteau X, Fakhri A, Albert V, Doukoure B, Noël J-C. Genital Schistosomiasis in European Women. *ISRN Obstet Gynecol*. 2011;2011:1–4.
10. World Health Organization. Female genital schistosomiasis: A pocket atlas for clinical health-care professionals. 2015.
11. Helling-Giese G, Kjetland EF, Gundersen SG, Poggensee G, Richter J, Krantz I, et al. Schistosomiasis in women: Manifestations in the upper reproductive tract. *Acta Trop*. 1996;62:225–38.
12. Kjetland EF, Ndhlovu PD, Gomo E, Mduluza T, Midzi N, Gwanzura L, et al. Association between genital schistosomiasis and HIV in rural Zimbabwean women. *AIDS*. 2006;20:593–600.
13. Stecher CW, Kallestrup P, Kjetland EF, Vennervald B, Petersen E. Considering treatment of male genital schistosomiasis as a tool for future HIV prevention: a systematic review. *Int J Public Health*. 2015;60:839–48.
14. Lopes RI, Leite KR, Prando D, Lopes RN. Testicular schistosomiasis caused by *Schistosoma mansoni*: A case report from Brazil. *Brazilian J Infect Dis*. 2007;11:523–4.
15. Pérignon A, Pelicot M, Consigny PH. Genital schistosomiasis in a traveler coming back from Mali. *J Travel Med*. 2007;14:197–9.
16. Chohan KR, Kling CA, Byler TK. *Schistosoma haematobium* ova in human semen: a case report. *Fertil Steril Reports*. 2020;in press:1–3.
17. van Delft F, Visser L, Polderman A, van Lieshout L. Cough and alterations in semen after a tropical swim. *Neth J Med*. 2007;65:304–6.
18. Al-Qahtani SM, Droupy SJ. Testicular schistosomiasis. *Saudi Med J*. 2010;31:325–7.
19. Mutengo M, Mudenda V, Mwansa J, Kaonga K, Sianongo S, Wamulume H, et al. Presence of Schistosomiasis in genital biopsies from patients at the University Teaching Hospital in Lusaka, Zambia. *Med J Zambia*. 2010;36:114–8.
20. Tayrab E, Ashmarg AK, Shareef H, Bedawi SB, Aradaib IE. Association of *Schistosoma mansoni* with infertility in a sudanese patient from schistosomiasis area of endemicity: A case report. *Res J Med Sci*. 2010;4:125–7.
21. Rambau PF, Chandika A, Chalya PL, Jackson K. Scrotal swelling and testicular atrophy due to schistosomiasis in a 9-year-old boy: A case report. *Case Rep Infect Dis*. 2011;2011:1–3.
22. Adisa J, Egbujo EM, Yahaya BA, Echejoh G. Primary infertility associated with *Schistosoma mansoni*: A case report from the Jos Plateau, north central Nigeria. *Afr Health Sci*. 2012;12:563–5.
23. Ondo CZ, Sarr A, Sow Y, Thiam I, Fall B, Sow D, et al. Bilharziome testiculaire à *Schistosoma haematobium* : à propos de deux observations. *Progrès en Urol*. 2014;24:67–9.
24. Ekenze SO, Modekwe VO, Nzegwu MA, Ekpemo SC, Ezomike UO. Testicular schistosomiasis mimicking malignancy in a child: A case report. *J Trop Pediatr*. 2015;61:304–9.
25. Rizk A, Kader MA, Hashmat H, El Agouz W, Abdalla M. Sex steroids in bilharzial liver affection. 1. estrogens. *Acta Biol Med Ger*. 1980;39:991–3.
26. Heshmat H, Rizk A, Abdel Kader M, Abdalla M. Sex steroids in bilharzial liver infections. *Radioimmunoassay Relat Proced Med* 1982. Vienna: International Atomic Energy Agency; 1982. p. 847.

27. Cavaliere H, Leite Z, Medeiros-Neto G. Serum immunoreactive somatomedin-C levels in growth failure and delayed puberty associated with chronic hepatosplenic schistosomiasis. *Clin Endocrinol (Oxf)*. 1986;24:617–26.
28. el-Ridi AM, Aboul-Magd LA, Edris A, Rashed SM, Hamadto HA, Abdalla KF. Serum testosterone level variation in bilharzial male patients. *J Egypt Soc Parasitol*. 1987;17:87–92.
29. Saad AH, Abdelbaky A, Osman AM, Abdallah KF, Salem D. Possible role of *Schistosoma mansoni* infection in male hypogonadism. *J Egypt Soc Parasitol*. 1999;29:307–23.
30. United Nations, Department of Economic and Social Affairs, Population Division. 2019. World population prospects 2019, Online Edition. Rev. 1.
31. WHO. Prevention and control of schistosomiasis and STH. WHO Technical Report. 2002. p. 912: 1–57.
32. Regis L, Lozano F, Lopez-Pacios AM, Morote J. Urogenital schistosomiasis: A diagnosis to consider in patients with hematuria in Europe. *UroToday Int J*. 2013;6.
33. Ribeiro AR, Luis C, Fernandes R, Botelho MC. Schistosomiasis and infertility: what do we know? *Trends Parasitol*. 2019. p. 964–71.
34. Vilana R, Corachán M, Gascón J, Valls E, Bru C. Schistosomiasis of the male genital tract: transrectal sonographic findings. *J Urol*. 1997;158:1491–3.
35. Shebel HM, Elsayes KM, Atta AE, Elguindy HM, El-Diasty YM. TA. Genitourinary schistosomiasis: Life cycle and radiologic-pathologic findings. *Radiographics*. 2012;32:1031–46.
36. Hassan A, El-Mogy S, Zalata K, Mostafa T. Testicular schistosomiasis: A case study. *Fertil Steril*. 2011;95:2124.e1-2124.e4.
37. Abdel-Naser MB, Altenburg A, Zouboulis CC, Wollina U. Schistosomiasis (bilharziasis) and male infertility. *Andrologia*. 2019;51:1–6.
38. O'Donnell L, Stanton P, de Kretser DM. Endocrinology of the male reproductive system and spermatogenesis. In: Feingold K, Anawalt B, A B, editors. *Endotext*. MDText.com, Inc.; 2017.
39. Lenzi HL, Kimmel E, Schechtman H, Pelajo-Machado M, Romanha WS, Pacheco RG, et al. Histoarchitecture of schistosomal granuloma development and involution: morphogenetic and biomechanical approaches. *Mem Inst Oswaldo Cruz*. 1998;93 SUPPL.:141–51.
40. Hales DB. Testicular macrophage modulation of Leydig cell steroidogenesis. *J Reprod Immunol*. 2002. p. 3–18.
41. Hutson JC. Physiologic interactions between macrophages and Leydig cells. *Exp Biol Med*. 2006;231:1–7.
42. Leisegang K, Henkel R. The *in vitro* modulation of steroidogenesis by inflammatory cytokines and insulin in TM3 Leydig cells. *Reprod Biol Endocrinol*. 2018;16:1–11.
43. Hales DB, Diemer T, Hales KH. Role of cytokines in testicular function. *Endocrine*. 1999;10:201–17.
44. Manfo FPT, Moundipa PF, Déchaud H, Tchana AN, Nantia EA, Zobot MT, et al. Effect of agropesticides use on male reproductive function: A study on farmers in Djutitsa (Cameroon). *Environ Toxicol*. 2012;27:423–32.
45. Leutscher PDC, Pedersen M, Raharisolo C, Jensen JS, Hoffmann S, Lisse I, et al. Increased prevalence of leukocytes and elevated cytokine levels in semen from *Schistosoma haematobium*-infected individuals. *J Infect Dis*. 2005;191:1639–47.
46. Kini S, Dayoub N, Raja A, Pickering S, Thong J. Schistosomiasis-induced male infertility. *BMJ Case Rep*. 2009;2009.
47. Leutscher PDC, Høst E, Reimert CM. Semen quality in *Schistosoma haematobium* infected men in Madagascar. *Acta Trop*. 2009;109:41–4.

## Figures





Figure 1

The vegetation of the Adamawa region falls within the rainforest – savannah mosaic and the climate is a sudano – guinean type with an annual rainfall of 1500 mm. Transmission of schistosomiasis presumably takes place at the lake, river, ponds and marshy fields. Fulbe and Kutin ethnic groups are predominant in Tibati and Woulde and they are mainly Muslims.

***S. mansoni* intensity of infection**



- ⋯ Light-intensity infection
- Moderate-intensity infection
- ▒ Heavy-intensity infection

***S. haematobium* intensity of infection**





Figure 2


Degree of the intensity of *Schistosoma* infection in the male adult population of Tibati and Wouldé Light-infection intensity: 1 - 99 epg; moderate-infection intensity: 100 - 399 epg; and heavy-infection intensity  $\geq 400$  epg for *S. mansoni* and for *S. haematobium*, light-infection intensity:  $< 50$  eggs/10mL and heavy-infection intensity:  $\geq 50$  eggs/10mL




**Figure 3**

Mean testes circumferences (A) and serum testosterone levels (B) in *Schistosoma haematobium* and *Schistosoma mansoni*-positive men in Tibati and Wouldé Data are expressed as mean  $\pm$  SEM. ANOVA followed by Tukey multiple comparison post-test was used for statistical analysis.  $\alpha P < 0.05$ : values are significantly different from those of *S. haematobium*-positive men. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ : values are significantly different from those of SCH-negative men.

**A. Correlation between *S. haematobium* infection and serum testosterone level**



**B. Correlation between *S. mansoni* infection and serum testosterone level**



**Figure 4**

Correlation curves between Schistosoma haematobium (A) or Schistosoma mansoni (B) intensity of infection and serum testosterone levels