Non-invasive Myocardial Work Index Contributes to Early Identification of Impaired Left Ventricular Myocardial Function in Uremic Patients With Preserved Left Ventricular Ejection Fraction

Huizhen Zhu

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University
<https://orcid.org/0000-0001-7494-1629>

Yanan Li

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Cunying Cui

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Danqing Huang

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Yuanyuan Liu

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Ruijie Liu

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Qingqing Zhao

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Ying Wang

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Lin Liu (✉ liulin@zzu.edu.cn)

Fuwai Central China Cardiovascular Hospital: Central China Fuwai Hospital of Zhengzhou University

Research

Keywords: Echocardiography, Uremia, Hemodialysis, Myocardial work, Ventricular function, Left

Posted Date: September 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-917401/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Cardiac damage is the leading cause of death in uremic patients. This study aimed to evaluate the application of non-invasive myocardial work index (NIMWI) by echocardiography in assessing the left ventricular (LV) systolic function in uremic patients.

Methods

Twenty-six uremic patients and twenty-seven age- and sex-matched healthy volunteers were enrolled in the study. Except for the conventional echocardiographic parameters, the LV myocardial work (MW) parameters including GWI (myocardial global work index), GCW (global constructive work), GWW (global wasted work), and GWE (global work efficiency) were calculated in study participants. Differences in MW parameters between the uremic and normal groups were compared by independent-sample t-test. Receiver operating characteristic (ROC) curves were constructed for MW parameters to detect abnormal LV systolic function in uremic patients.

Results

Compared with the normal group, GWW was significantly increased and GWE decreased in the uremic group ($P < 0.05$). Area under the curve (AUC) for GWE by the ROC analysis was 0.966. The best threshold, sensitivity and specificity values of GWE to detect abnormality of LV systolic function in uremic patients were 92.5%, 0.89 and 0.96 respectively.

Conclusions

NIMWI may be applied to assess the global MW of uremic patients. The presence of reduced GWE can help identify impaired left ventricular myocardial function in uremic patients with preserved LV ejection fraction with a high sensitivity and specificity.

Background

Chronic kidney disease (CKD) is a common disease which seriously threatens patients' health and life. The disease causes damage to multiple systems and organs, especially in its end-stage (i.e. uremia). Indeed, cardiac damage is the leading cause of death in uremic patients [1]. Earlier studies have demonstrated that myocardial contraction function of the uremic patients has been impaired when the left ventricular ejection fraction (LVEF) is normal [2]. Therefore, objective and accurate method to evaluate left ventricular (LV) systolic function in uremic patients is urgently needed.

The pressure-volume loop method measured by cardiac catheterization can assess the LV systolic function and myocardial oxygen consumption accurately [3]. However, this method is fairly invasive and has limited application in clinical practice. LVEF, measured by conventional transthoracic

echocardiography, is the indicator most commonly used to predict mortality or morbidity of cardiovascular disease; Nonetheless it is often influenced by pre- and post-loads and the cardiac structure [4]. Previous studies reported that the global longitudinal strain (GLS) obtained by two dimensional (2D) speckle tracking technology is more sensitive for diagnosing abnormalities of LV systolic function than conventional LVEF [5]. However, myocardial strain is susceptible to the cardiac afterload, which can affect the accuracy of myocardial function evaluation. The non-invasive myocardial work index (NIMWI) is a new method based on 2D speckle tracking technology that combines myocardial deformation and LV pressure to evaluate LV myocardial work (MW), which is more objective than GLS alone [6]. Russell et al. demonstrated a good correlation between NIMWI assessment and invasive and direct measurements of MW [7].

Previous studies have indicated that the NIMWI can evaluate the LV MW in patients with CKD without dialysis [8], but the MW in CKD patients undergoing haemodialysis treatment has not been studied. Thus, this study aimed to evaluate the feasibility of the NIMWI for assessing LV global MW in uremic patients with preserved ejection fraction undergoing haemodialysis, and provided reference values for the clinical evaluation of LV systolic function in these patients.

Results

1. Clinical characteristics

Compared with the normal group, the uremic group showed higher systolic blood pressures (SBP) and diastolic blood pressures (DBP). The differences between groups were statistically significant (both $P < 0.05$). There were no differences in gender, age, heart rate (HR) and body surface area (BSA) between the two groups ($P > 0.05$) (Table 1).

2. Traditional echocardiographic measurement

Compared with the normal group, the uremic group showed increased interventricular septum end-diastolic thickness (IVSTd), LV posterior wall end-diastolic thickness (LVPWTd), LV end-diastolic diameter (LVEDd), LV mass index (LVMI) and decreased GLS. The differences between groups were statistically significant ($P < 0.05$). There were no significant differences in LVEF between the two groups ($P > 0.05$) (Table 2).

3. Comparison of MW parameters

Compared with the normal group, the uremic group showed increased global wasted work (GWW) and decreased global work efficiency (GWE). The differences observed between groups were statistically significant ($P < 0.05$). However, differences in global work index (GWI) and global constructive work (GCW) between the two groups were not statistically significant ($P > 0.05$) (Table 3, Figure 1).

4. Receiver operating characteristic (ROC) curve analysis

Based on ROC curve analysis of main MW parameter, area under the curve (AUC) of GWE and GWW were 0.966 and 0.925 respectively. GWE was the best diagnostic index for identifying impaired LV myocardial function in uremic patients with preserved LV ejection fraction. The sensitivity and specificity were 89% and 96%, [respectively](#). The best threshold of GWE was 92.5% (Figure 2).

5. Intra- and inter-observer variability

Intra-class correlation coefficients indicated that GWI, GCW, GWW and GWE between the intra- and inter-observer measurements were consistent (Table 4).

Discussion

Uremia is the final stage of CKD progression. Serious damage to the kidney functions promotes the production and accumulation of toxins in the body, resulting in multi-organ dysfunction and poor health. Cardiovascular complications are the main cause of death in patients with uremia [9]. Therefore, early detection of changes in myocardial contraction in uremic patients is important to enable the early control of cardiovascular complications. The results of our study showed that compared with the control group, there were no change in LVEF in uremic patients ($P > 0.05$), but the uremic group showed increased GWW and decreased GWE ($P < 0.05$), indicating that the NIMWI can identify LV systolic dysfunction in uremic patients with preserved ejection fraction. GWE was the best diagnostic index for identifying impaired LV myocardial function in uremic patients with preserved LV ejection fraction, and its sensitivity, specificity and optimal threshold were 89%, 96% and 92.5%, [respectively](#).

Traditional evaluations of MW mainly rely on invasive pressure measurement, the feasibility of which is limited in conventional clinical practice [10,11]. The results of MW index assessment of MW have a strong correlation with the measurement results of invasive cardiac catheterization [12]. This finding confirms the feasibility of the NIMWI for assessing MW. The NIMWI based on 2D speckle tracking technology, which combines myocardial deformation and non-invasive LV pressure to assess MW [7], is more comprehensive than GLS for the assessment of cardiac function in patients with altered afterload, and the technique can detect impaired myocardial function early, and its feasibility has been established in hypertension, hypertrophic cardiomyopathy, dilated cardiomyopathy and cardiac resynchronization [13-16].

Compared with the normal group, there were no difference in GWI and GCW, but GWW increased and GWE decreased in the uremic group, indicating that a decreased GWE of the left ventricle in uremic patients. GWE is the ratio of GCW to the sum of GCW and GWW, and there is no obvious change in the GCW. Thus, it can be seen that the decrease of GWE is mainly related to the increase of LV GWW. GWW refers to the work done by the left ventricle that is not conducive to ejection, including the total work done by cardinal elongation at systole and cardinal shortening at isovolumic diastole. Any factor affecting myocardial elongation at systole and myocardial shortening at isovolumic diastole may increase GWW. An earlier study [17] demonstrated that several factors in the body of with uremic patients, such as sodium and water retention and renin-angiotensin-aldosterone system

activation, participate in the development of hypertension. The presence of long-term high loads leads to increased LV myocardial stiffness, ventricular remodeling, and myocardial oxygen consumption [18,19]. Luca et al. [20] found that secondary nephritic anaemia in uremic patients aggravates cardinal ischaemia, hypoxia, damages cardinal cells by calcium and phosphorus imbalance, and causes microvascular ischemia, interstitial fibrosis and cardiac myocyte hypertrophy by long-term haemodialysis-related myocardial stunning. These phenomena negatively affect myocardial contractions, resulting in an increase in global wasted work (GWW), and a decrease in the global work efficiency of the left ventricle.

However, this study had some limitations that should be noted. It was a single centre study with a relatively small sample size and no long-term follow-up of uremic patients, which will be encompassed in the next phase of the study.

Conclusions

NIMWI may be applied to assess the global MW of uremic patients. AUC, sensitivity, and specificity for GWE parameter by ROC analysis were high, revealing the accuracy of GWE for the detection of the LV systolic dysfunction in the uremic patients. The reproducibility of MW parameters can help to improve its application in clinical practice.

Methods

1. Study population

A total of 26 uremic patients who were treated in the nephrology department of our hospital from November 2020 to June 2021 were consecutively identified. All uremic patients were treated with regular haemodialysis three times a week and with LVEF $\geq$ 50% according to the 2016 European Society of Cardiology Guidelines [21]. Patients were excluded if there were valvular disease, LV outflow tract obstruction, arrhythmia, essential hypertension, diabetes, previous myocardial infarction, medium and large pericardial effusion, LV systolic dysfunction with LVEF<50%, and patients with a poor acoustic window of transthoracic echocardiography.

Twenty-seven healthy volunteers matched for sex, age and body surface area (BSA) were selected as the normal group. All volunteers had no history of heart disease, hypertension, diabetes or medication. Their physical examination, electrocardiogram, chest radiograph, renal function and echocardiography results were normal.

All subjects provided informed consent prior to participating in this study, and the study protocols were approved by the Hospital Ethics Committee. The participant selection process was illustrated in the flowchart in Figure 3.

2. Echocardiography

GE Vivid E95 (GE Vingmed Ultrasound AS, Horten, Norway) colour Doppler ultrasound system equipped with M5Sc-D 1.4-4.6 MHz probe was used for echocardiographic image acquisition.

The subjects were instructed to assume the left-lateral position and an electrocardiogram was attached. Transthoracic echocardiography was used to measure the LVEDd, LVPWTd and IVSTd of the participants. The modified biplane Simpson method was used to calculate LVEF from the apical four- and two-chamber views. The LVMI was calculated as LV mass divided by BSA. We obtained the dynamic two-dimensional images from the LV apical four-, two- and three-chamber views at frame rate of 58 - 69 frames/sec (average of 67.4 ± 6 frames/sec) for at least three cardiac cycles in the resting state. The images were copied, saved and exported to the workstation in Digital Imaging and Communication in Medicine (DICOM) format on a mobile hard disk for offline analysis.

All two-dimensional images and measurements were performed according to American Society of Echocardiography guideline [22], and all parameters were averaged over three consecutive cardiac cycles. The SBP and DBP of the participants were measured using a brachial artery cuff.

3. Two-dimensional speckle tracking image analysis

The stored images were analysed offline by an Echo PAC BT203 (GE Vingmed Ultrasound, Horten, Norway) workstation. The durations of the LV isovolumic systole and ejection period were first determined according to the opening and closing of the aortic and mitral valves, and then the automatic functional imaging mode was selected. The endocardium of the LV long axis, four-chamber and two-chamber view were depicted from the level of the mitral valve annulus to the LV apex in sequence. The software automatically tracked the endocardium and epicardium of the left ventricle. The intracardiac and epicardial wrap were manually adjusted to ensure satisfactory tracking. Finally, Approve was clicked, and the blood pressure measured by the brachial artery cuff was entered into the MW menu to obtain the LV pressure-strain loops (PSL), GLS and NIMWI. The NIMWI parameters included GWI, GCW, GWW and GWE, as shown in Figure 4.

4. Statistical analysis

All statistical analyses were performed using SPSS version 25.0 (SPSS Inc., IBM, Chicago, USA). Continuous variables were expressed as mean $\pm$ standard deviation if normally distributed. Independent-sample *t*-test was used to compare continuous variables between the uremic group and normal group. The chi-squared test was used to compare count data.

ROC curves were performed to determine the optimal sensitivity and specificity of MW parameter. The AUC was calculated to assess the performance of MW parameter for the detection impaired myocardium in uremic patients. According to the Youden index, the best threshold of each tested MW parameter was estimated. Ten subjects were randomly selected, and two observers used intra-class correlation coefficients to carry out repeatability tests for the myocardial work parameters GWI, GCW, GWW and GWE. A *P*-value < 0.05 was considered statistically significant.

Abbreviations

NIMWI: Non-invasive myocardial work index

LV: Left ventricular

MW: Myocardial work

GWI: Global work index

GCW: Global constructive work

GWW: Global wasted work

GWE: Global work efficiency

ROC: Receiver operating characteristic

AUC: Area under the curve

CKD: Chronic kidney disease

LVEF: Left ventricular ejection fraction

GLS: Global longitudinal strain

2D: Two dimensional

SBP: Systolic blood pressures

DBP: Diastolic blood pressures

HR: Heart rate

BSA: Body surface area

IVSTd: Interventricular septum end-diastolic thickness

LVPWTd: Left ventricular posterior wall end-diastolic thickness

LVEDd: Left ventricular end-diastolic diameter

LVMI: Left ventricular mass index

ROC: Receiver operating characteristic

AUC: Area under the curve

DICOM: Digital Imaging and Communication in Medicine

PSL: Pressure-strain loops

Declarations

Acknowledgments

Not applicable.

Funding

This study was supported by National Natural Science Foundation of China (82071950), National Natural Science Foundation of Henan for Excellent Young Scientists (202300410364), Medical Science and Technology Research Project of Henan Province (SB201901099, LHGJ20200084).

Contributions

Performed the literature review: HZZ, YNL; Carried out echocardiography measurements: HZZ, YNL, QQZ, YW; Selected participants and collected the clinical data: HZZ, YNL, DQH; Checked the validity of data: HZZ, YNL, LL; Data analysis: HZZ, YNL, YYL; Wrote the manuscript: HZZ; Prepared the manuscript: YW, RJL; Supported the experiments financially: LL; Checked the paper: LL, YNL; All authors have checked and confirmed the paper. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The study was approved by Fuwai Central China Cardiovascular Hospital, Henan Provincial People's Hospital, Central China Fuwai Hospital of Zhengzhou University, People's Hospital of Zhengzhou University.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Department of Ultrasound, Fuwai Central China Cardiovascular Hospital, Henan Provincial People's Hospital, Central China Fuwai Hospital of Zhengzhou University, People's Hospital of Zhengzhou University, Zhengzhou, Henan451464, China.

References

1. Yan P, Li H, Hao C, Shi H, Gu Y, Huang G et al. 2D-Speckle Tracking Echocardiography Contributes to Early Identification of Impaired Left Ventricular Myocardial Function in Patients with Chronic Kidney Disease. *Nephron Clinical Practice*. 2011;118(3):232-40. 'doi:'10.1159/000321383.
2. Ma Y, Zhang B, Zhang Y, Dong Y, Zhang R. Ultrasonic image analysis of longitudinal strain in uremic patients with preserved left ventricular ejection fraction. *BIOMED ENG ONLINE*. 2018;17(1):112. 'doi:'10.1186/s12938-018-0536-y.
3. Tulner SA, Klautz RJ, van Rijk-Zwikker GL, Engbers FH, Bax JJ, Baan J et al. Perioperative assessment of left ventricular function by pressure-volume loops using the conductance catheter method. *ANESTH ANALG*. 2003;97(4):950-7. 'doi:'10.1213/01.ane.0000078814.81889.b3.
4. Konstam MA, Abboud FM. Ejection Fraction. *CIRCULATION*. 2017;135(8):717-9. 'doi:'10.1161/CIRCULATIONAHA.116.025795.
5. Tops LF, Delgado V, Marsan NA, Bax JJ. Myocardial strain to detect subtle left ventricular systolic dysfunction. *EUR J HEART FAIL*. 2017;19(3):307-13. 'doi:'10.1002/ejhf.694.
6. Boe E, Skulstad H, Smiseth OA. Myocardial work by echocardiography: a novel method ready for clinical testing. *European Heart Journal - Cardiovascular Imaging*. 2019;20(1):18-20. 'doi:'10.1093/ehjci/jey156.
7. Russell K, Eriksen M, Aaberge L, Wilhelmsen N, Skulstad H, Remme EW et al. A novel clinical method for quantification of regional left ventricular pressure-strain loop area: a non-invasive index of myocardial work. *EUR HEART J*. 2012;33(6):724-33. 'doi:'10.1093/eurheartj/ehs016.
8. Ke QQ, Xu HB, Bai J, Xiong L, Li MM. Evaluation of global and regional left ventricular myocardial work by echocardiography in patients with chronic kidney disease. *Echocardiography*. 2020;37(11):1784-91. 'doi:'10.1111/echo.14864.
9. Sun M, Kang Y, Cheng L, Pan C, Cao X, Yao H et al. Global longitudinal strain is an independent predictor of cardiovascular events in patients with maintenance hemodialysis: a prospective study using three-dimensional speckle tracking echocardiography. *The International Journal of Cardiovascular Imaging*. 2016;32(5):757-66. 'doi:'10.1007/s10554-016-0836-x.

10. P. F, J. LB. Pressure-volume loops: A dynamic approach to the assessment of ventricular function. W.B. Saunders. 1994;8(1):84.
11. Mclaurin LP, Grossman W, Stefadouros MA, Rolett EL, Young DT. A New Technique for the Study of Left Ventricular Pressure-Volume Relations in Man. CIRCULATION. 1973;48(1):56-64.
12. Hubert A, Le Rolle V, Leclercq C, Galli E, Samset E, Casset C et al. Estimation of myocardial work from pressure-strain loops analysis: an experimental evaluation. European Heart Journal-Cardiovascular Imaging. 2018;19(12):1372-9. 'doi':10.1093/ehjci/jey024.
13. Chan J, Edwards NFA, Khandheria BK, Shiino K, Sabapathy S, Anderson B et al. A new approach to assess myocardial work by non-invasive left ventricular pressure - strain relations in hypertension and dilated cardiomyopathy. European Heart Journal - Cardiovascular Imaging. 2019;20(1):31-9. 'doi':10.1093/ehjci/jey131.
14. Galli E, Vitel E, Schnell F, Le Rolle V, Hubert A, Lederlin M et al. Myocardial constructive work is impaired in hypertrophic cardiomyopathy and predicts left ventricular fibrosis. Echocardiography. 2019;36(1):74-82. 'doi':10.1111/echo.14210.
15. Cui C, Liu L, Li Y, Liu Y, Huang D, Hu Y et al. Left Ventricular Pressure - Strain Loop Based Quantitative Examination of the Global and Regional Myocardial Work of Patients with Dilated Cardiomyopathy. Ultrasound in Medicine & Biology. 2020;46(10):2834-45. 'doi':10.1016/j.ultrasmedbio.2020.06.008.
16. Galli E, Leclercq C, Fournet M, Hubert A, Bernard A, Smiseth OA et al. Value of Myocardial Work Estimation in the Prediction of Response to Cardiac Resynchronization Therapy. J AM SOC ECHOCARDIOG. 2018;31(2):220-30. 'doi':10.1016/j.echo.2017.10.009.
17. Harrison DG. The Mosaic Theory revisited: common molecular mechanisms coordinating diverse organ and cellular events in hypertension. J AM SOC HYPERTENS. 2013;7(1):68-74. 'doi':10.1016/j.jash.2012.11.007.
18. Brilla CG, Funck RC, Rupp H. Lisinopril-Mediated Regression of Myocardial Fibrosis in Patients With Hypertensive Heart Disease. Circulation: Journal of the American Heart Association. 2000;102(12):1388-93.
19. Kuznetsova T, D Hooge J, Kloch-Badelek M, Sakiewicz W, Thijs L, Staessen JA. Impact of Hypertension on Ventricular-Arterial Coupling and Regional Myocardial Work at Rest and during Isometric Exercise. J AM SOC ECHOCARDIOG. 2012;25(8):882-90. 'doi':10.1016/j.echo.2012.04.018.
20. Di Lullo L, Gorini A, Russo D, Santoboni A, Ronco C. Left Ventricular Hypertrophy in Chronic Kidney Disease Patients: From Pathophysiology to Treatment. CARDIORENAL MED. 2015;5(4):254-66. 'doi':10.1159/000435838.

21. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. EUR J HEART FAIL. 2016;18(8):891-975. 'doi:'10.1002/ehf.592.

22. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L et al. Recommendations for Cardiac Chamber Quantification by Echocardiography in Adults: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. European Heart Journal-Cardiovascular Imaging. 2015;16(3):233-71. 'doi:'10.1093/ehjci/jev014.

Tables

Table 1 Comparison of general clinical data in the study groups

Variable	Normal group <i>n</i> =27	Uremic group <i>n</i> =26		
			χ^2	<i>P</i> -value
Male/Female	13/14	13/13	3.000	0.083
			<i>t</i>	<i>P</i> -value
Age (years)	44.07±9.34	49.11±10.56	1.842	0.071
BSA (m ²)	1.72±0.16	1.75±0.18	0.732	0.468
HR (times.min ⁻¹)	69.00±6.87	72.84±10.76	0.732	0.130
SBP (mmHg)	120.37±8.49	157.50±16.30 ^a	10.456	<0.001
DBP (mmHg)	78.59±6.16	98.19±12.07 ^a	7.484	<0.001

Values are mean ± standard deviation.

BSA body surface area, **HR** heart rate, **SBP** systolic blood pressure, **DBP** diastolic blood pressure

^a*p*< 0.05 vs. normal group, 1mmHg=0.133kPa

Table 2 Conventional echocardiography parameters in the study groups

Variable	Normal group <i>n</i> =27	Uremic group <i>n</i> =26	<i>t</i>	<i>P</i> -value
IVSTd (mm)	8.94±0.81	11.96±1.24 ^a	10.461	<0.001
LVPWTd (mm)	8.96±0.75	12.00±1.29 ^a	10.360	<0.001
LVEDd (mm)	43.33±3.92	50.69±6.71 ^a	4.848	<0.001
LVEF (%)	64.29±2.05	62.61±4.38	-1.777	0.084
LVMl (g/m ²)	73.02±13.86	138.10±34.40 ^a	8.965	<0.001
GLS (%)	-18.77±0.93	-14.11±2.26 ^a	9.717	<0.001

Values are mean ± standard deviation.

IVSTd interventricular septum end-diastolic thickness, *LVPWTd* left ventricular posterior wall end-diastolic thickness, *LVEDd* left ventricular end-diastolic diameter, *LVEF* left ventricular ejection fraction, *LVMl* left ventricular mass index, *GLS* global longitudinal strain

^a*p* < 0.05 vs. normal group.

Table 3 Comparisons of MW parameters in the study groups

Variable	Normal group <i>n</i> =27	Uremic group <i>n</i> =26	<i>t</i>	<i>P</i> -value
GWI (mmHg%)	1927.74±164.01	1863.23±284.71	-1.006	0.321
GCW (mmHg%)	2086.55±199.59	2163.26±324.98	1.040	0.309
GWW (mmHg%)	83.77±37.42	238.34±114.89 ^a	6.637	<0.001
GWE (%)	95.48±1.80	88.80±3.08 ^a	-9.655	<0.001

Values are mean ± standard deviation.

GWI global work index, *GCW* global constructive work, *GWW* global wasted work, *GWE* global work efficiency

^a*P* < 0.05 vs. normal group.

Table 4 Intra-and inter-observer variability

Variable	Intra-observer ICC		Inter-observer ICC	
	ICC	95%CI	ICC	95%CI
GWl	0.995	0.979-0.999	0.992	0.970-0.998
GCW	0.961	0.843-0.990	0.993	0.970-0.998
GWW	0.961	0.842-0.990	0.972	0.889-0.993
GWE	0.983	0.932-0.996	0.971	0.883-0.993

GWl global work index, GCW global constructive work, GWW global wasted work, GWE global work efficiency, ICC intraclass correlation coefficients, CI confidence interval

Figures

Figure 1

A: Left ventricular MW parameters of the normal group. B: Left ventricular MW parameters of uremic patients. In each group, the LV PSL, 17-segment bull's-eye representation of GWI, and the MW parameters were given, and the MW parameters included GWI, GCW, GWW, and GWE. MW: myocardial work; PSL: pressure-strain loops; GLS: global longitudinal strain; GWl: global work index; GCW: global constructive work; GWW: global waste work; GWE: global work efficiency

Figure 2

ROC curve analysis of the main MW parameters for identifying impaired LV myocardial function in uremic patients with preserved LV ejection fraction. On the left: ROC curve of GWW parameter (AUC=0.925). On the right: ROC curve of GWE parameter (AUC=0.966). GWE had better AUC than GWW with the sensitivity, specificity and optimal threshold of 89%, 96% and 92.5%, respectively. ROC: receiver operating characteristic; GWE: global work efficiency

Figure 3

Flow chart of study populations

Figure 4

Schematic diagram of the NIMWI measurement results. (a) Changes in systolic blood pressure measured by the cuff as a function of the global longitudinal strain of the left ventricle. Red curve: Overall pressure and strain of the left ventricle. The area under curve represents the global myocardial work index of the left ventricle. (b) Column chart of the global constructive work and waste work of the left ventricle. (c) Bull's-eye diagram of the 17-segment myocardial work index of the left ventricle. (d) Results of the NIMWI parameters. NIMWI: non-invasive myocardial work index