

Role of Exogenous-Applied Salicylic Acid, Zinc And Glycine Betaine To Improve Drought-Tolerance In Wheat During Reproductive Growth Stages

Ramadan Shemi (✉ ramadangamil@agr.cu.edu.eg)

Southwest University

Rui Wang

Southwest University

El-Sayed M.S. Gheith

Cairo University

Hafiz Athar Hussain

Chinese Academy of Agricultural Sciences

Linna Cholidah

Southwest University

Kangping Zhang

Southwest University

Sai Zhang

Southwest University

Longchang Wang

Southwest University

Research Article

Keywords: Drought stress, Glycine betaine, Salicylic acid, Zinc, Antioxidant defense system, Wheat (*Triticum aestivum* L.).

Posted Date: October 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-923487/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Plant Biology on December 1st, 2021. See the published version at <https://doi.org/10.1186/s12870-021-03367-x>.

Abstract

Background: Drought has become a dangerous threat to reduce crop productivity throughout the world. Exogenous application of regulators, micronutrients, or osmoprotectants for inducing drought-tolerance in field crops has been effectively adopted. A controlled pot study was performed to investigate the relative efficacy of salicylic acid (SA), zinc (Zn), and glycine betaine (GB) as foliar applications on the growth, tissues pigments content, relative water content (RWC), leaf gas exchange, antioxidant enzymes activity, reactive oxygen species (ROS) accumulation, osmolytes contents and the yield parameters of wheat plants subjected to two soil water conditions (85% field capacity: well-watered, 50% field capacity: water-deficient) during reproductive growth stages.

Results: Water deficient conditions significantly decreased the growth, yield parameters, RWC, photosynthesis pigment, and gas exchange attributes except for intercellular CO₂ concentration. However, foliar applications remarkably improved the growth parameters under water deficit conditions. Drought stress statistically increased the contents of hydrogen peroxide (H₂O₂), superoxide anion radical (O₂^{•-}), and malonaldehyde (MDA), and elevated the harmful oxidation to cell lipids in plants, however, they were considerably reduced by foliar applications. Activities of all antioxidant enzymes, proline content, and soluble sugar were increased in response to foliar applications under water deficit conditions.

Conclusions: Overall, foliar application of GB, SA, and Zn compounds improved the drought-tolerance in wheat by decreasing the ROS accumulation, promoting enzymatic antioxidants, and increasing osmolytes accumulation. Finally, GB treatment was most effective in thoroughly assessed parameters of wheat followed by SA and Zn applications to alleviate the adverse effects of drought stress.

Background

Wheat (*Triticum aestivum* L.) is one of the most important cereal crops which is highly susceptible to drought stress [1]. Universally, the wheat crop is the primary vegetable protein source in the diet of humans, containing a higher protein content than other main grain crops such as maize and rice. The increasing population and food demands further necessitate the increase in wheat production to ensure social stability and future food security [2, 3]. In China, wheat is the second-most important food crop after rice and the third-most important crop in total production overall, after maize and rice. The cultivated area of wheat in China was estimated to be 23.73 million ha and it yielded about 133.60 million tonnes year⁻¹ [4]. Wheat is sensitive cereal crop to drought stress after maize, especially during the critical growth periods (e.g., flowering stage). However, drought led to the yield reduction among different species, in which maize had higher yield reduction (39.3%) compared to wheat (20.6%) by 40% water deficiency [5]. Drought is the most important one of abiotic stress factors that negatively influence the growth and production of various field crops [6–10]. Drought causes various changes in crop plants through different morphological, physiological, and biochemical responses [11–13]. Many studies have reported that the defense system of antioxidant was increased through the synthesis of antioxidants such as APX and GR and enzymatic antioxidants such as POD, SOD, and CAT, which were stimulated to

quench the ROS production in plants during drought stress [3, 6, 14–16]. Moreover, Datir et al. [17] have reported that the antioxidant enzymes, free proline contents, GB accumulation, MDA, and H₂O₂ contents were increased under drought induced by polyethylene glycol (PEG) in wheat cultivars. In addition, Chen et al. [18] remarked that the drought stress slowed down wheat growth and reduced grain yield by impacting the anthesis and grain-filling process, and decreased leaf water potential, stomatal conductance, and the photosynthesis.

Foliar application of various growth regulators, micronutrients, and osmoprotectants can play the key role in inducing drought-tolerance in plants at various plant growth stages. SA is a multifunctional plant hormone that can effectively ameliorate the adverse effects caused by biotic and abiotic stresses [19] by modulating different growth responses, and physiological and biochemical characteristics in plants [10, 20–23]. A useful role of the salicylic acid has been noticed in improving drought stress tolerance in wheat by improving the relative water contents, leaf gas exchange soluble carbohydrate, proteins, proline contents, the activity of antioxidant enzymes and yield attributes [24–26]. Furthermore, Zinc is an essential micronutrient that participates in many physiological functions and structure of the regulatory cofactor of many enzymes, carbohydrate, chlorophyll production, pollen function, fertilization, metabolism of RNA, protein synthesis, and the DNA functions [27, 28]. Water deficit condition at grain filling phase statistically reduced the plant height, yield parameters, RWC, chlorophyll contents, and decreased activity of SOD, POD, and CAT, while exogenous applications of zinc and salicylic acid had positive impact on all these parameters and mitigated the harmful effects of water deficit on wheat plants [24, 29, 30]. Rahmani et al. [31] reported that supplemental Zn improved the drought-tolerance in safflower by increasing proline, relative water content, and chlorophyll contents, and the yield and its components. Nevertheless, GB has an important role as an osmoprotectants, which help the plants to resist drought through improved leaf gas exchange and chlorophyll content in maize plants [32]. Wheat genotypes prevent the damage of PEG induced osmotic stress via various mechanisms such as osmolytes accumulation [33]. In addition, net photosynthesis, transpiration rate and yield attributes were decreased under drought stress at the tillering, flowering, and milking stages in wheat plants, while exogenous application of GB enhanced them under drought treatment [34, 35]. Similarly, Hasanuzzaman et al. [36] recorded that GB played an important role in reducing aggregation and detoxification of reactive oxygen species, hence recovering photosynthesis and decreasing oxidative stress. Our previous study, Shemi et al. [37] have illustrated that exogenous application of GB, Zn and SA substantially improved the activity of CAT, SOD, and APX enzymes, and decreased the contents of MDA and H₂O₂, and these changes were beneficial to protect maize leaf tissues from oxidative harm in cell membranes.

However, there is no investigation has been carried out to assess the relative effects of SA, Zn, or GB compounds in inducing drought-tolerance of the wheat crop during reproductive growth stages. The current study was aimed to test the hypothesis that the foliar applications of salicylic acid, zinc, and/or glycine betaine can ameliorate the damaging effects of drought stress in wheat during reproductive growth stages by observing the changes in the scavenging ability of antioxidant defense system, which constitutes the executive for the protective response system. The objectives of the study were to (1)

Investigate the response of wheat growth, yield, and its attributes to the exogenous applications of SA, Zn, and GB under different soil water conditions during reproductive growth stages; (2) Determine the effect of respective exogenous applications on chlorophyll contents, RWC, leaf gas exchange, antioxidant enzymes and MDA content, ROS and osmolytes accumulations under different soil water conditions; and (3) Compare the efficacy of concerned exogenous applications to alleviate the harmful effects of drought stress under different soil water conditions.

Materials And Methods

Experimental design and plant growth conditions

The controlled pot experiment was conducted during the winter cultivating season of 2018–2019 at the glasshouse of the College of Agronomy and Biotechnology, Southwest University, Chongqing, China. The experimental area lies at longitude 106° 26' 02" E, latitude 29° 49' 32" N, and altitude 220 m. During the cultivation season, the average minimum and maximum temperatures were 10°C to 26°C, and the relative humidity was between 75–87%. The experiment was implemented in a completely randomized design (CRD) with two factors: two soil water conditions (well-watered condition at 85 % of field capacity, and water-deficient condition at 50 % of field capacity), and four spraying treatments [0.00 double distilled water (CK), 140 mg l⁻¹ salicylic acid (SA), 4 g l⁻¹ zinc (Zn), and 11.5 g l⁻¹ glycine betaine (GB)]. The effective concentrations of these spraying treatments were based on enhancements in the growth and yield of various crops under water-deficient [10, 25, 30, 35, 37]. The experiment included eight treatments, and each treatment involved twelve pots. A winter wheat cultivar Yumai-13, which was bred by Sichuan Agricultural University and Chongqing Academy of Agricultural Sciences in China, was used as plant material, and the seeds were obtained from Wheat Research Institute, College of Agronomy and Biotechnology, Southwest University, Chongqing, China.

Each plastic pot (25 cm diameter, 30 cm depth) was filled with 8 kg air-dried and sieved (0.5 mm) soil, which was collected from the experimental station at the College of Agronomy and Biotechnology, Southwest University. Experimental soil was clay loam, and had the following physical and chemical properties: pH, organic matter, electrical conductivity (EC), bulk density, soil water content at field capacity (FC), total N, available phosphorous, and available potassium were 6.25, 12.58 g kg⁻¹, 0.45 ds/m, 1.44 g cm⁻³, 24.35%, 0.98 g kg⁻¹, 15.53 mg kg⁻¹, and 86.11 mg kg⁻¹, respectively. At the time of soil filling, 2.7 g controlled-release urea (44.6% N), 3.5 g calcium superphosphate (12% P₂O₅), 1.5 g potassium chloride (60% K₂O) were applied for each pot. Fifteen uniform grains were manually sown in the third week of November in pots at depth of 4–5 cm. Thinning was conducted 10 days after germination, and seven uniform seedlings per pot were selected for the subsequent studies. Each pot was irrigated to 85% FC by tap water till the start of drought stress treatments.

Soil Water Conditions

The plants were subjected to two soil water conditions for 30 days, from booting (Feekes 10 stage) until milk (Feekes 11 stage) stages of wheat: well-watered condition (85% of field capacity; WW) and water-deficient condition (50% of field capacity; WD). During the drought period, the pots were weighed daily to keep the required soil water levels by adding proper water volumes. Soil water contents for 85% and 50% field capacity were 22.5 % and 11.25 %, respectively [37]. Soil water content (SWC) was determined using the following equation: $SWC \% = [(FW - DW) / DW] \times 100$, where FW was the fresh weight of soil sample from the inner area of each pot, and DW was the dry weight of soil sample after oven drying at 85° C for 3 days [54].

Spraying Treatments

After 7 and 15 days of drought treatment, the wheat plants under each water condition were sprayed with 0.00 (double distilled water; CK), 140 mg l⁻¹ SA (hydroxybenzoic acid-2 C₇H₆O₃, MW = 138.12 g mol⁻¹), 4 g l⁻¹ Zn (zinc sulfate heptahydrate ZnSO₄ · 7H₂O, MW = 287.54 g mol⁻¹) and 11.5 g l⁻¹ GB (betaine C₅H₁₁NO₂, MW = 117.14 g mol⁻¹) [37]. Tween-20 (0.05%) was added with foliar applications as a surfactant at the time of treatment.

Measurements And Analysis

Wheat plants were sampled after 15 days of foliar application treatments to measure growth, photosynthetic pigments, RWC, photosynthesis gas exchange, and biochemical assays. Completely expanded, undamaged, and healthy wheat plant leaves (2nd leaf from the top) from all replicates were sampled. After washing, wheat leaves were frozen with liquid N₂ immediately and stored at -80°C for biochemical analyses, and analysis of yield attributes were recorded at harvesting time.

Growth, Yield, And Its Attributes

Four pots were randomly selected and plants of each pot were taken to measure plant height, fresh weight pot⁻¹, dry weight pot⁻¹, and leaf area pot⁻¹. Total leaf area pot⁻¹ was measured with LI-3100 leaf area meter (Li-COR, CID, Inc., USA). The dry weight of plants pot⁻¹ was estimated following oven drying at 85°C for 48 hours. At full maturity (plants at 160-days old), five pots were randomly selected, and plants of each pot were harvested to measure the number of tillers pot⁻¹, number of spikes pot⁻¹, number of grains spike⁻¹, grain weight spike⁻¹ (g), 1000-grain weight (g), biological yield pot⁻¹ (g), grain yield pot⁻¹ (g) and harvest index (HI). The HI was computed as the percent ratio of grain yield and biological yield according to Donald [55].

Photosynthetic pigments and RWC

Chlorophyll (Chl. a, Chl. b and total Chl.) contents were determined in the 2nd leaf from the top according to Peng and Liu [56]. The extraction of a 200 mg leaf blade sample was done with 10 ml ethanol-acetone (1:2, v/v), and the extract was moved to a 15 ml centrifuge tube. The tubes were put in the dark to avoid light for 24 hours until the sample changed into a white color. The chlorophyll content was calculated by the following equation: Chlorophyll a content (mg/g tissue) = $(12.7D_{663} - 2.69D_{645}) \times V / (1000 \times W)$, Chlorophyll b content (mg/g tissue) = $(22.7 D_{645} - 4.68D_{663}) \times V / (1000 \times W)$, and total chlorophyll (mg/g tissue) = $D_{652} \times V / (34.5 \times W)$ / Chl a + Chl b, where, D_{663} , D_{645} and D_{652} , respectively are the corresponding wavelengths of the light density value, V is the volume of extracting liquid and W is the weight of fresh leaf. The RWC of wheat leaves was measured according to Barrs and Weatherley [57]. Fresh leaves were cut into small segments (1.5 cm length), weighed fresh weight (FW), then floated in distilled water for 4 h under low light to register saturated weight (SW), and then dried in an oven until constant weight at 80 °C for 24 hours to record dry weight (DW). RWC was computed as: $RWC = (FW - DW) / (SW - DW) \times 100 \%$.

Photosynthesis gas-exchange

Net photosynthesis rate (Pn), transpiration rate (Tr), stomatal conductance (Gs), and intercellular CO₂ concentration (Ci) were registered using a portable infrared gas analyzer-based photosynthesis system (LI-6400; LiCor, Inc., Lincoln, NE, USA) at 09:30-11:30 am from the fully expanded leaf (2nd leaf from top). Air relative humidity and ambient CO₂ concentrations were about 78 % and 370 $\mu\text{mol CO}_2 \text{ mol}^{-1}$, respectively during the collection of data.

Assay of enzymatic antioxidants and lipid peroxidation

Antioxidant enzyme activity was reported using commercial kits for glutathione reductase (GR, A111), superoxide dismutase (SOD, A500), catalase (CAT, A501), ascorbate peroxidase (APX, A304), and peroxidase (POD, A502), by following the manufacturer's instructions (Sino Best Biological Co., Ltd., China). The absorbance readings of GR, SOD, CAT, APX, and POD were detected at 340 nm, 560 nm, 240 nm, 290 nm, and 470 nm, respectively using an ultraviolet (UV)-visible spectrophotometer, and their activities were expressed as units per fresh weight ($\text{U g}^{-1} \text{FW}$). One unit of GR activity was expressed as the amount of enzyme depleting 1 μmol NADPH in 1 min, one unit of SOD activity was defined as the amount of enzyme needed to reduce the reference rate to 50% of maximum inhibition, one unit of CAT activity was measured as the amount of enzyme that decomposes 1nmol H₂O₂ at 240 nm min^{-1} in 1 g fresh weight, one unit of APX was estimated as the amount of enzyme required for catalyzing 1 μmol ASA at 290 nm 2 min^{-1} of 1 g fresh weight in 1 ml of a reaction mixture, and one unit of POD activity was demonstrated as the absorbance change of 0.01 at 470 nm min^{-1} for 1 g fresh weight in 1 ml of a reaction mixture [37]. Lipid peroxidation in wheat leaves was assayed as MDA content, and was measured by thiobarbituric (TBA) method using MDA Detection Kit (A401). Lipid hydroperoxide degradation products could condense with thiobarbituric acid (TBA) to yield red compounds [37]. The absorbance for MDA content was recorded at 532 and 600 nm and expressed as nmol g^{-1} fresh weight.

Determination of reactive oxygen species accumulation

Hydrogen peroxide (H_2O_2) and superoxide anion radical ($\text{O}_2^{\cdot-}$) contents in the wheat leaves were noted using the commercial 'H₂O₂ Detection Kit (A400)' and 'O₂^{·-} Detection kit (A407)', respectively, according to the manufacturer's instructions. H₂O₂ content was estimated at 415 nm and represented as $\mu\text{mol g}^{-1}$ fresh weight. Super oxygen anion serotonin reacted with hydrochloride to produce NO₂⁻. The NO₂⁻ interacted with amino benzene and alpha-pyridoxine to produce red compounds at 530 nm which had a characteristic absorption peak [37]. The content of O₂^{·-} was measured at 530 nm and expressed as $\mu\text{mol g}^{-1}$ fresh weight.

Estimation of osmolytes accumulation

Proline and soluble sugar contents in wheat leaves were determined using commercial kits for proline (PRO, A605) and soluble sugar contents (SSC, B602), according to the manufacturer's instructions (Sino Best Biological Co., Ltd., China). The absorbance reading of the toluene layer was estimated at 520 nm, on a spectrophotometer, and proline (Sigma, St Louis, MO, USA) was used for the standard curve [37]. Proline content was expressed as $\mu\text{g g}^{-1}$ fresh weight. The absorbance reading of SSC was detected at 620 nm using an ultraviolet (UV)-visible spectrophotometer [37]. Soluble sugar content was articulated as mg g^{-1} fresh weight.

Statistical analysis

The collected data were analyzed following the analysis of variance (ANOVA) according to the Two-way Factorial Design using Statistical Software Package MSTAT-C [58]. The significant differences among mean values were estimated according to Least significant difference test (L.S.D) at a 95% confidence level [59]. Sigma Plot 10.0 (Systat Software Inc., San Jose, CA, USA) was used for the graphical presentation of the data.

Results

Growth and yield attributes

Water deficient condition statistically ($p < 0.05$) hampered the wheat growth and yield parameters in terms of plant height, fresh weight of plants pot^{-1} , dry weight of plants pot^{-1} , leaf area of plants pot^{-1} , number of tillers pot^{-1} , number of spikes pot^{-1} , number of grains spike⁻¹, grain weight spike⁻¹, 1000-grain weight, biological yield pot^{-1} , grain yield pot^{-1} and harvest index in comparison to well-watered condition. However, results showed that growth and yield parameters were improved by SA, Zn, and GB spraying treatments in both stressed and non-stressed plants (Tables 1,2, and 3). As expected, one or more spraying treatments significantly enhanced all parameters in both soil water conditions except for

the number of grains spike⁻¹ as compared to the control. Respective spraying treatments increased plant height by 6.94%, 6.92% and 8.05%, fresh weight of plants pot⁻¹ by 18.88%, 11.30% and 23.16%, dry weight of plants pot⁻¹ by 15.82%, 9.34% and 28.30%, leaf area of plants pot⁻¹ by 39.34%, 33.22% and 47.63%, number of tillers pot⁻¹ by 13.07%, 8.97% and 15.29%, number of spikes pot⁻¹ by 13.78%, 9.72% and 16.33%, number of grains spike⁻¹ by 3.47%, 1.23 and 4.95%, grain weight spike⁻¹ by 13.15%, 2.26% and 18.42%, 1000-grain weight by 9.75%, 4.29% and 13.42%, biological yield pot⁻¹ by 17.63%, 12.95% and 25.16%, grain yield pot⁻¹ by 27.99%, 15.23% and 37.36% and harvest index by 8.98%, 1.96% and 10.00%, under water-deficient condition, respectively, as compared to the values of the control treatment. Overall, maximum growth and yield parameters were recorded from the plants treated with GB followed by SA and Zn spraying treatments as compared with control under both soil water conditions.

Table 1
Influences of soil water conditions and foliar treatments on wheat growth parameters.

Soil water conditions	Foliar treatments	Plant height (cm)	Fresh weight of plants pot ⁻¹ (g)	Dry weight of plants pot ⁻¹ (g)	Leaf area of plants pot ⁻¹ (cm ²)	No. of tillers pot ⁻¹	No. of spikes pot ⁻¹
WW	CK	63.39 ^{bc} ± 1.67	73.13 ^{bc} ± 2.92	17.79 ^d ± 0.73	962.74 ^c ± 54.55	18.62 ^c ± 0.56	16.67 ^b ± 0.33
	SA	70.58 ^a ± 1.40	81.20 ^a ± 3.31	21.95 ^b ± 0.61	1067.01 ^b ± 61.24	20.59 ^{ab} ± 0.55	18.34 ^a ± 0.61
	Zn	68.67 ^{ab} ± 2.64	79.48 ^{ab} ± 3.62	20.73 ^{bc} ± 0.73	1015.75 ^{bc} ± 53.79	19.83 ^b ± 0.60	17.83 ^b ± 0.33
	GB	71.86 ^a ± 2.71	82.38 ^a ± 4.11	23.57 ^a ± 0.94	1148.27 ^a ± 65.15	21.27 ^a ± 0.79	19.00 ^a ± 0.34
	Means	68.62	79.05	21.00	1048.44	20.07	17.96
WD	CK	57.73 ^d ± 0.70	43.16 ^f ± 2.38	14.66 ^f ± 0.15	428.14 ^f ± 21.73	14.38 ^f ± 0.61	12.55 ^d ± 0.65
	SA	61.74 ^c ± 0.59	51.31 ^d ± 2.04	16.98 ^e ± 0.30	596.60 ^{de} ± 23.37	16.26 ^d ± 0.53	14.28 ^c ± 0.72
	Zn	61.73 ^c ± 0.33	48.04 ^{de} ± 2.57	16.03 ^e ± 0.72	570.37 ^e ± 27.85	15.67 ^{de} ± 0.41	13.77 ^c ± 0.63
	GB	62.38 ^{bc} ± 1.70	53.16 ^d ± 2.10	18.81 ^d ± 0.35	632.10 ^d ± 16.41	16.58 ^d ± 0.44	14.60 ^c ± 0.47
	Means	60.89	48.92	16.61	556.80	15.72	13.79

Values are means (± SE) of three replicates. For L.S.D.'s results, means with various letters indicate the significant differences ($p < 0.05$). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.

Table 2

Influences of soil water conditions and foliar treatments on wheat yield and its components.

Soil water conditions	Foliar treatments	No. of grains spike ⁻¹	Grain weight spike ⁻¹ (g)	1000-grain weight (g)	Biological yield pot ⁻¹ (g)	Grain yield pot ⁻¹ (g)	Harvest index (%)
WW	CK	38.12 ^a ± 0.97	1.48 ^{bc} ± 0.03	38.97 ^{bc} ± 0.93	58.93 ^{cd} ± 0.73	22.77 ^{cd} ± 0.73	38.58 ^b ± 2.33
	SA	41.40 ^a ± 0.99	1.73 ^{ab} ± 0.07	41.64 ^{ab} ± 0.76	67.80 ^{ab} ± 2.67	27.28 ^{ab} ± 0.83	40.46 ^{ab} ± 2.82
	Zn	39.42 ^a ± 1.41	1.57 ^b ± 0.03	39.89 ^b ± 0.85	66.32 ^{bc} ± 3.41	25.42 ^{bc} ± 0.24	38.62 ^b ± 0.80
	GB	42.02 ^a ± 1.71	1.78 ^a ± 0.03	42.52 ^a ± 1.65	70.33 ^a ± 2.61	29.24 ^a ± 0.77	41.70 ^a ± 2.01
	Means	40.23	1.64	40.75	65.84	26.18	39.83
WD	CK	35.69 ^a ± 1.32	1.14 ^e ± 0.04	31.87 ^f ± 0.23	43.16 ^g ± 0.88	15.36 ^f ± 0.29	35.60 ^{cd} ± 0.74
	SA	36.93 ^a ± 1.05	1.29 ^{cd} ± 0.06	34.98 ^d ± 1.08	50.77 ^e ± 1.71	19.66 ^{de} ± 0.34	38.80 ^b ± 1.20
	Zn	36.13 ^a ± 0.54	1.20 ^{de} ± 0.04	33.24 ^{ef} ± 1.39	48.75 ^{ef} ± 1.39	17.70 ^e ± 0.69	36.30 ^c ± 1.07
	GB	37.46 ^a ± 1.12	1.35 ^c ± 0.03	36.15 ^{cd} ± 0.39	54.02 ^{de} ± 2.22	21.10 ^d ± 0.53	39.16 ^{ab} ± 1.50
	Means	36.55	1.24	34.05	49.17	18.45	37.46
Values are means (± SE) of three replicates. For L.S.D.'s results, means with various letters indicate the significant differences ($p < 0.05$). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.							

Table 3

p-values of the two-way factorial analysis of growth, yield, and physiological and biochemical parameters of wheat as affected by various foliar treatments under both soil water conditions.

Parameters	Main factors effects		Interaction effects
	S	F	S × F
Plant height	< 0.0001	< 0.0065	< 0.0392
Fresh weight of plants	< 0.0001	< 0.0249	< 0.0486
Dry weight of plants	< 0.0001	< 0.0001	< 0.0372
Leaf area of plants	< 0.0001	< 0.0038	< 0.0435
Number of tillers	< 0.0001	< 0.0037	< 0.0467
Number of spikes	< 0.0001	< 0.0051	< 0.0245
Number of grains	< 0.0004	< 0.1066	< 0.4751
Grain weight	< 0.0001	< 0.0001	< 0.0371
1000-grain weight	< 0.0001	< 0.0061	< 0.0497
Biological yield	< 0.0001	< 0.0007	< 0.0378
Grain yield	< 0.0001	< 0.0001	< 0.0212
Harvest index	< 0.0073	< 0.0054	< 0.0485
Chlorophyll a content	< 0.0087	< 0.0019	< 0.0347
Chlorophyll b content	< 0.0180	< 0.0097	< 0.0492
Total chlorophyll content	< 0.0163	< 0.0010	< 0.0345
RWC	< 0.0001	< 0.0273	< 0.0479
Net photosynthesis rate	< 0.0001	< 0.0003	< 0.0198
Transpiration rate	< 0.0001	< 0.0008	< 0.0245
Stomatal conductance	< 0.0001	< 0.0001	< 0.0359
Intercellular CO ₂ concentration	< 0.0100	< 0.0204	< 0.0457
APX activity	< 0.0001	< 0.0002	< 0.0021
GR activity	< 0.0001	< 0.0002	< 0.0102
POD activity	< 0.0001	< 0.0001	< 0.0260

p-values are considered as significant ($p < 0.05$, $n = 3$) and highly significant ($p < 0.01$, $n = 3$). 'S': effect of soil water conditions; 'F': effect of foliar treatments; 'S × F': effect of the interaction between two variables.

Parameters	Main factors effects		Interaction effects
	S	F	S × F
CAT activity	< 0.0001	< 0.0001	< 0.0002
SOD activity	< 0.0001	< 0.0001	< 0.0302
MDA content	< 0.0001	< 0.0001	< 0.0430
H ₂ O ₂ content	< 0.0001	< 0.0001	< 0.0469
O ₂ ⁻ content	< 0.0001	< 0.0001	< 0.0041
Free proline content	< 0.0001	< 0.0001	< 0.0006
Total soluble sugar	< 0.0001	< 0.0001	< 0.0019

p-values are considered as significant ($p < 0.05$, $n = 3$) and highly significant ($p < 0.01$, $n = 3$). 'S': effect of soil water conditions; 'F': effect of foliar treatments; 'S × F': effect of the interaction between two variables.

Photosynthesis Gas-exchange

Net photosynthesis rate (Pn), transpiration rate (Tr), stomatal conductance (Gs) and intercellular CO₂ concentration (Ci) were statistically ($p < 0.05$) influenced by soil water conditions. However, results noticed that photosynthesis gas exchange parameters were increased by SA, Zn, and GB spraying treatments except for Ci (Fig. 2 and Table 3). Pn and Tr were significantly affected by SA and GB treatments under water-deficient conditions and GB under well-watered conditions, while Gs was considerably impacted by concerned spraying treatments under both soil water conditions except for Zn under the well-watered condition, and Ci was substantially reduced by SA and GB treatments under water-deficient condition. Under the water-deficient condition, respective spraying treatments promoted the Pn by 34.74%, 30.11%, and 49.03%, Tr by 21.84%, 13.15%, and 33.42%, and Gs by 58.97%, 35.89%, and 69.23%, while they decreased the Ci by 13.01%, 9.94%, and 16.22%, respectively, as compared to the values of the control treatment. In general, the highest Pn, Tr and Gs, and the lowest Ci were registered from the plants treated with GB followed by SA and Zn spraying treatments as compared with control treatment under both soil water conditions.

Enzymatic antioxidants and MDA content

Ascorbate peroxidase (APX), glutathione reductase (GR), peroxidase (POD), catalase (CAT), and superoxide dismutase (SOD) activities and malonaldehyde (MDA) content were significantly ($p < 0.05$) affected by soil water conditions. Results indicated that APX, GR, POD, CAT, and SOD activities were

increased and MDA content was decreased by SA, Zn, and GB spraying treatments in both stressed and non-stressed plants (Fig. 3 and Table 3). GR and CAT activities and MDA contents were statistically impacted by concerned spraying treatments in both soil water conditions except for Zn under well-watered condition; while APX, POD, and SOD activities were significantly influenced by concerned spraying treatments under water-deficient condition, and POD by GB application and SOD by SA application under well-watered condition. Under the water-deficient condition, respective spraying treatments improved the activity of APX by 47.82%, 29.89% and 42.39%, GR by 59.62%, 42.85% and 48.44%, POD by 26.08%, 18.84% and 35.30%, CAT by 62.16%, 50.95% and 120.11%, SOD by 61.42%, 34.64% and 42.20%, but reduced MDA content by 29.09%, 16.64% and 26.51%, respectively, as compared to the values of the control treatment. Overall, the highest APX, GR, and SOD activities and the lowest MDA contents were recorded from the plants treated with SA followed by GB and Zn spraying treatments, while the highest POD and CAT activities were recorded from the plants treated with GB followed by SA and Zn spraying treatments as compared with control treatment under both soil water conditions.

Reactive Oxygen Species Accumulation

The level of reactive oxygen species (ROS) accumulation was statistically ($p < 0.05$) increased under water-deficient conditions. Results showed that hydrogen peroxide (H_2O_2) and superoxide anion radical ($O_2^{\cdot-}$) contents were decreased by SA, Zn, and GB spraying treatments in both stressed and non-stressed plants (Fig. 4 and Table 3). H_2O_2 and $O_2^{\cdot-}$ contents were significantly impacted by concerned spraying treatments under both soil water conditions except for Zn under well-watered conditions. Under the water-deficient condition, respective spraying treatments decreased the H_2O_2 by 29.33%, 23.99%, and 26.45%, and $O_2^{\cdot-}$ by 36.10%, 25.26%, and 28.97%, respectively, as compared to the values of the control treatment. Overall, results indicated that the lowest H_2O_2 and $O_2^{\cdot-}$ contents were recorded from the plants treated with SA followed by GB and Zn spraying treatments as compared with control treatment under both soil water conditions.

Osmolytes Accumulation

Free proline contents and total soluble sugar were significantly ($p < 0.05$) enhanced by the water-deficient condition. Results indicated that free proline contents and total soluble sugar were increased with SA, Zn, and GB spraying treatments in both stressed and non-stressed plants (Fig. 5 and Table 3). Free proline content and total soluble sugar were statistically affected by concerned spraying treatments under water-deficient conditions and GB under well-watered conditions. Under the water-deficient condition, respective spraying treatments enhanced free proline content by 53.57%, 33.41%, and 67.07%, and total soluble sugar by 36.01%, 22.76%, and 56.25%, respectively, as compared to the values of the control treatment. Overall, results showed that the highest free proline and total soluble sugar contents were recorded from

the plants treated with GB followed by SA and Zn spraying treatments as compared with control treatment under both soil water conditions.

Discussion

Drought is a critical agricultural hazard that negatively influences the crop production and may cause a disequilibrium between the defense systems of antioxidant and ROS accumulation, resulting in oxidative damages [6, 8–10, 38]. Bayoumi et al. [39] reported that drought led to a decrease in plant height, the number of tillers, 1000-grain weight, spike length, biological and grain yields, and harvest index. Drought stress may hinder the total growth and development of many field crops, and the reproductive growth stages are oversensitive under drought stress [8, 37, 38, 40]. The presented data (Tables 1, 2, and 3) indicated that water-deficient conditions statistically ($p < 0.05$) hampered wheat growth and yield parameters. However, one or more spraying treatments significantly improved growth and yield parameters except for the number of grains spike⁻¹ compared to control treatment in both stressed and non-stressed plants. Previously, plant height and seedling fresh and dry weights were considerably decreased by water deficit, while an exogenous application with GB, Zn and SA treatments markedly promoted maize growth [37]. In this regard, Anjum et al. [32] and Raza et al. [35] revealed that growth parameters, yield, and yield attributes were statistically decreased by drought stress, while they were significantly increased by exogenous application of GB treatment in maize and wheat plants. Drought stress significantly disrupted the plant height and yield parameters in wheat [8]. Many studies indicated that, salicylic acid, zinc, and glycine betaine played a role to regulate protective responses under diverse stresses in many plant species [10, 24, 29, 37, 41]. In this study, the impairment in wheat growth performance and yield parameters under water-deficient conditions could be ascribed to excessive production of ROS (H_2O_2 and $O_2^{\cdot-}$) which led to oxidative damage to lipids membrane and raised MDA content (Figs. 3 and 4). In agreement with our results, Mittler [42] and Miller et al. [14] revealed that the increase of ROS accumulation could destroy the cell membrane and lead to immediate damage to lipids, proteins, photosynthetic pigments, nucleic acids, as well as cell structure, and finally caused the death of cells and loss of plant biomass. In contrast to the adverse impacts of the water-deficient condition, drought-tolerance induced by spraying treatments was associated with enhanced contents of Chl. a, Chl. b and total Chl., and RWC (Fig. 1), improved photosynthesis gas-exchange (Fig. 2), activities of APX, GR, POD, CAT, SOD and osmolytes accumulation (Figs. 3 and 5), and decreased the MDA, H_2O_2 , and $O_2^{\cdot-}$ contents (Figs. 3 and 4). Growth and yield increase by these exogenous applications under water-deficient conditions is an outside indicator of metabolism alteration in the plant cells. In the past, many studies have indicated the harmful effects of drought stress on the growth performance and yields of various grain crops [10, 11, 37, 40, 43, 44]. However, the damage range under drought stress were differed with the strength of stresses and the crop growth stages [9].

Our results indicated that Chl. a, Chl. b, and total Chl. contents, and RWC were severely decreased in wheat plants by drought stress (Fig. 1). However, SA, Zn, and GB spraying treatments enhanced these parameters in both stressed and non-stressed plants. The decreases in photosynthetic pigments as one

of the most substantial restrict factors for plant photosynthetic activity under abiotic stress were documented by numerous other studies [8, 45, 46]. Hussain et al. [11] found that drought stress enhanced the deterioration of chlorophyll in maize plants. In this study, photosynthetic pigments were decreased under water-deficient conditions. Previously, Anjum et al. [32] indicated that the Chl. a, Chl. b and total Chl. contents in maize were reduced under progressive water deficit, while these pigments were substantially increased by GB spray application. However, RWC is a beneficial variable to assess the physiological water status of plant leaves [47]. In this regard, our previous study, Shemi et al. [37] stated that drought stress significantly reduced RWC, whilst exogenous applications with GB, Zn and SA treatments improved the RWC under drought stress. Bayoumi et al. [39] reported that the RWC was decreased under drought stress in wheat genotypes. Likewise, drought stress at the grain filling phase substantially reduced RWC and chlorophyll content, while exogenous applications of zinc and salicylic acid had a positive impact on these parameters and alleviated the harmful effects of water deficit on wheat plants [30].

In the present study, net photosynthesis rate, transpiration rate, stomatal conductance, and intercellular CO_2 concentration were considerably affected by soil water condition, while they were increased by SA, Zn, and GB spraying treatments except for intercellular CO_2 concentration (Fig. 2). Anjum et al. [32] found that the gas exchange parameters were considerably reduced in maize cultivars under water-stressed conditions, while GB application considerably improved gas exchange rate under water deficit as compared with control treatment. Shemi et al. [37] found that the net photosynthesis rate, transpiration, and stomatal conductance were decreased by water deficit, while exogenous application of GB, Zn and SA treatments enhanced the CO_2 assimilation and improved physiological water status. Under water-deficit conditions, plants tend to close the stomata, which causes reduction in CO_2 availability and decrease the photosynthetic rate and finally reduces plant yield [48, 49]. In our study, the water-deficient condition significantly increased the APX, GR, POD, CAT, and SOD activities (Fig. 3). However, the increases in activity of antioxidant enzymes were not adequate to protect against ROS accumulations and were not enough to repair the injuries of the oxidative stress triggered by the water-deficient condition. Exogenous applications of salicylic acid, zinc, and glycine betaine to wheat plants exposed to the water-deficient condition increased the activity of these antioxidant enzymes (Fig. 3). Exogenous applications modified the enzymatic and non-enzymatic antioxidants in wheat plants under stress conditions and efficiently quench the harmful ROS (Figs. 3 and 4). In the past, several studies have noticed that antioxidant enzymes and non-enzymes activities were increased, but the contents of H_2O_2 , $\text{O}_2^{\cdot-}$ and MDA were decreased by exogenous applications of Zn, SA, or GB under drought stress in various plants [10, 21, 24, 29, 35, 37, 41, 50]. In this regard, Miller et al. [14] indicated that diverse enzymatic antioxidants could maintain the equilibrium between production and scavenging of ROS, whereas activities of these enzymes may protect the plants under stress conditions. Furthermore, the foliar application of GB, Zn and SA treatments substantially enhanced the activity of SOD, CAT, and APX enzymes and scavenged ROS accumulation in maize [37]. The SOD enzyme is considered as the first line for protecting against ROS accumulation, which stimulates the transformation of superoxide anion radical ($\text{O}_2^{\cdot-}$) to O_2 and H_2O_2 [51]. The CAT enzyme stimulates the conversion of H_2O_2 to H_2O and

molecular oxygen, where H_2O_2 is considered a powerful and harmful oxidizing agent [52]. Malondialdehyde (MDA) is regarded as an appropriate indicator for lipid peroxidation in cell membranes [42]. The present study showed that MDA content in wheat leaves was statistically increased under the water-deficient condition and it was well associated with H_2O_2 and $O_2^{\cdot-}$ contents, while respective foliar applications decreased MDA, H_2O_2 , and $O_2^{\cdot-}$ contents (Figs. 3 and 4). This finding indicated that exogenous applications raised the ability of plants to cope with oxidative stresses, and therefore improved the drought-tolerance.

The presented results revealed that ROS accumulation (such as H_2O_2 and $O_2^{\cdot-}$ contents) was statistically increased under the water-deficient condition and decreased by SA, Zn, and GB spraying treatments (Fig. 4). This was also proved by Wang and Jin [29], Sofy [42], Talaat et al. [6], Abdel-Motagally and El-Zohri [8], Datir et al. [17], Hasanuzzaman et al. [36] and Shemi et al. [37] who indicated ROS accumulation was considerably increased under drought stress and decreased by exogenous applications of SA, Zn, and GB in various field crops. In addition to the enzymatic defense system, some compatible solutes vigorously participate in the amelioration of drought stress. Proline and soluble sugar are very important for the osmoregulation process in plants under drought stress. In this study, proline and soluble sugar in wheat leaves were enhanced under the water-deficient condition and they were also significantly increased by SA, Zn, and GB spraying applications (Fig. 5). This phenomenon can be considered as a portion of the mechanism to inhibit water loss in plants by adjusting the osmotic condition. In our previous study, free proline and soluble sugar in maize were improved by SA, Zn and GB treatments under drought stress [37]. Furthermore, Hussain et al. [11], and Abdel-Motagally and El-Zohri [8] indicated that proline content and soluble sugar were noticeably enhanced in tested plant leaves under drought stress compared to control treatments. Nasrin et al. [41] revealed that proline and soluble sugar content were significantly increased by the application of salicylic acid under different irrigation schedules. Moreover, El Tayeb and Ahmed [53] remarked that the accumulation of sugars content in shoots and roots of wheat cultivars was substantially promoted under water stress and increased by exogenous application of SA. Proline content was substantially increased under water deficit as compared to well-watered conditions in wheat genotypes [39].

As noticed in this study, all precedent results proposed that foliar applications could be regarded as protection compounds against oxidative damage caused by drought stress, which could improve the photosynthesis pigments and RWC, leaf gas exchange, the capacity of the antioxidant defense system, decrease the MDA, H_2O_2 , and $O_2^{\cdot-}$ contents, and increase the osmolyte accumulation in water-stressed plants, thus they might be considered as an important strategy to improve plant growth and yield attributes under drought stress (Fig. 6).

Conclusions

The present study pointed out that the exogenous application of SA, Zn, and GB treatments could partially mitigate the harmful effects of water stress on the growth of wheat through increasing

chlorophyll and RWC contents, modulating gas exchange traits, enhancing expression of antioxidant enzymes, scavenging ROS and increasing accumulation of osmolytes. These mechanisms are very important to sustained wheat production in comparatively water-meager regions. Among the foliar application compounds, GB was most effective followed by SA and Zn in promoting the growth and yield attributes of wheat. It could be concluded that exogenous GB, SA, and Zn applications could minimize the loss of wheat yield triggered by drought stress.

Abbreviations

SA: Salicylic acid; Zn: Zinc; GB: Glycine betaine; CK: double distilled water; RWC: Relative water content; ROS: Reactive oxygen species; H₂O₂: Hydrogen peroxide; O₂^{•-}: superoxide anion radical; MDA: Malonaldehyde; GR: Glutathione reductase; SOD: Superoxide dismutase; CAT: Catalase; APX: Ascorbate peroxidase; POD: Peroxidase; PEG: Polyethylene glycol; Chl.: Chlorophyll; Pn: Net photosynthesis rate; Tr: Transpiration rate; Gs: Stomatal conductance; Ci: Intercellular CO₂ concentration; WW: Well-watered condition; WD: Water-deficient condition; SWC: Soil water content.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

All authors agreed to publish.

Availability of data and material

All the data in this work available in this manuscript and supplementary files

Competing interests

The authors have no competing interests to declare. This work is original and has never been published before. We have no conflict with any other scientist.

Funding

This work was supported by the China Scholarship Council (CSC No. 2017GBJ001939)/ Special Fund for Agro-Scientific Research in the Public Interest (No. 201503127) and the Natural Science Foundation Project of China (No. 31871583).

Authors' contributions

R.S, L.W, and EL. Gh conceived and designed the research. R.S, R.W, H.H, and L.C performed the experiment. R.S wrote the initial draft of the manuscript. R.S, and R.W contributed to reagents, materials, and analysis tools. R.S, L.W, EL. Gh, K.Z, and S.Z revised subsequent versions of the manuscript, prepared the final version, and provided technical guidance. All authors have read and approved the manuscript.

Acknowledgements

The present work is part of the Ph.D. dissertation of the first author. The first author gives his gratitude to the teams of Key Laboratory of Eco-environments in Three Gorges Reservoir Region, and Engineering Research Center of South Upland Agriculture, Ministry of Education, China for their help and support in the experiment. We thank Mrs. Esraa Hussein for helping us to make the figures.

References

1. Ibrahim MU, Khaliq A, Hussain S, Murtaza G. Sorghum Water Extract Application Mediates Antioxidant Defense and Confers Drought Stress Tolerance in Wheat. *J. Plant Growth Regul.* 2021; 1–12.
2. Hussain S, Khaliq A, Tanveer M, Matloob A, Hussain HA. Aspirin priming circumvents the salinity-induced effects on wheat emergence and seedling growth by regulating starch metabolism and antioxidant enzyme activities. *Acta Physiol. Plant.* 2018; 4: 1–12.
3. Thind S, Hussain I, Ali S, Hussain S, Rasheed R, Ali B, Hussain H A. Physiological and Biochemical Bases of Foliar Silicon-Induced Alleviation of Cadmium Toxicity in Wheat. *J. Soil Sci. Plant Nut.* 2020; 4: 2714–2730.
4. FAOSTAT. Population data. Food and Agricultural Organization of United Nation, Roma. 2019; On line at <http://Faostat.fao.org/download/O/OA/E>.
5. Daryanto S, Wang L, Jacinthe PA. Global synthesis of drought effects on maize and wheat production. *Plos One.* 2016; 5: e0156362.
6. Talaat NB, Shawky BT, Ibrahim AS. Alleviation of drought-induced oxidative stress in maize (*Zea mays L.*) plants by dual application of 24-epibrassinolide and spermine. *Environ Exp Bot.* 2015; 47–58.
7. Kosová K, Urban MO, Vítámvás P, Prasil IT. Drought stress response in common wheat, durum wheat, and barley: transcriptomics, proteomics, metabolomics, physiology, and breeding for an enhanced drought tolerance. In: Hossain MA, Wani SH, Bhattacharjee S, Burritt DJ, Tran LSP (Eds.), *Drought Stress Tolerance in Plants: Molecular and Genetic Perspectives*, vol. 2. Springer International Publishing. 2016; 277–314.
8. Abdel-Motagally FMF, El-Zohri M. Improvement of wheat yield grown under drought stress by boron foliar application at different growth stages. *J Saudi Soc Agric Sci.* 2018; 2: 178–185.

9. Hussain HA, Hussain S, Khaliq A, Ashraf U, Anjum S A, Men S, Wang L. Chilling and drought stresses in crop plants: implications, cross talk, and potential management opportunities. *Front. plant sci.* 2018; 393.
10. Sohag AAM, Tahjib-Ul-Arif M, Brestič M, Afrin S, Sakil MA, Hossain MT, Hossain MA, Hossain MA. Exogenous salicylic acid and hydrogen peroxide attenuates drought stress in rice. *Plant Soil Environ.* 2020; 7–13.
11. Hussain HA, Men S, Hussain S, Chen Y, Ali S, Zhang S, Zhang K, Li Y, Xu Q, Liao C, Wang L. Interactive effects of drought and heat stresses on morpho-physiological attributes, yield, nutrient uptake and oxidative status in maize hybrids. *Sci. Rep.* 2019; 3890.
12. Nisa W, Nisa V, Nagoo SA, Dar ZA. Drought tolerance mechanism in wheat: A review. *Pharma Innovation.* 2019; 2: 714–724.
13. Xu Q, Fu H, Zhu B, Hussain HA, Zhang K, Tian X, ... Wang L. Potassium Improves Drought Stress Tolerance in Plants by Affecting Root Morphology, Root Exudates and Microbial Diversity. *Metabolites.* 2021; 3: 131.
14. Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R. Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant Cell Environ.* 2010; 453–467.
15. Scheibe R, Beck E. Drought, Desiccation, and Oxidative Stress. In: Lüttge U, Beck E, Bartels D (eds) *Plant Desiccation Tolerance. Ecological Studies (Analysis and Synthesis)*, Vol. 215. Springer, Berlin, Heidelberg. 2011; 209–231.
16. Hussain HA, Men S, Hussain S, Zhang Q, Ashraf U, Anjum SA ... Wang L. Maize tolerance against drought and chilling stresses varied with root morphology and antioxidative defense system. *Plants.* 2020; 6: 720.
17. Datir SS, Inamdar A. Biochemical responses of wheat cultivars to peg-induced drought stress. *Russ Agric Sci.* 2019; 5–12.
18. Chen X, Min D, Yasir TA, Hu YG. Evaluation of 14 morphological, yield-related and physiological traits as indicators of drought tolerance in Chinese winter bread wheat revealed by analysis of the membership function value of drought tolerance (MFVD). *Field Crops Res.* 2012; 195–201.
19. Maruri-López I, Aviles-Baltazar NY, Buchala A, Serrano M. Intra and extracellular journey of the phytohormone salicylic acid. *Front Plant Sci.* 2019; 10.
20. Hayat Q, Hayat S, Irfan M, Ahmad A. Effect of exogenous salicylic acid under changing environment: A review. *Environ Exp Bot.* 2010; 14–25.
21. Chen YE, Cui JM, Li GX, Yuan M, Zhang ZW, Yuan S. Effect of salicylic acid on the antioxidant system and photosystem II in wheat seedlings. *Biol Plant.* 2016; 139–147.
22. Noreen S, Fatima K, Athar HUR, Ahmad S, Hussain K. Enhancement of physio-biochemical parameters of wheat through exogenous application of salicylic acid under drought stress. *J Anim Plant Sci.* 2017; 153–163.
23. Fardus J, Matin MA, Hasanuzzaman M, Hossain MA, Hasanuzzaman M. Salicylic acid-induced improvement in germination and growth parameters of wheat under salinity stress. *J Anim Plant Sci.*

- 2018; 1: 197–207.
24. MR, Sofy. Application of salicylic acid and zinc improves wheat yield through physiological processes under different levels of irrigation intervals. *Int J Plant Res.* 2015; 5: 136–156.
 25. Sedaghat M, Sarvestani ZT, Emam Y, Bidgoli AM. Physiological and antioxidant responses of winter wheat cultivars to strigolactone and salicylic acid in drought. *Plant Physiol Bioch.* 2017; 59–69.
 26. Sharma M, Gupta SK, Majumder B, Maurya VK, Deeba F, Alam A, Pandey V. Salicylic acid mediated growth, physiological and proteomic responses in two wheat varieties under drought stress. *J Proteom.* 2017; 28–51.
 27. Khan HR, McDonald GK, Rengel Z. Zinc fertilization and water stress affects plant water relations, stomatal conductance and osmotic adjustment in chickpea (*Cicer arietinum* L.). *Plant and Soil.* 2004; 271–284.
 28. Nasiri Y, Najafi N. Effect of soil and foliar application of zinc and iron on flowering and essential oil of chamomile at green house conditions. *Acta Agric Slov.* 2015; 1–4.
 29. Wang H, Jin J. Effects of zinc deficiency and drought on plant growth and metabolism of reactive oxygen species in maize (*Zea mays* L.). *Agric Sci China.* 2007; 8: 988–995.
 30. Yavas I, Unay A. Effects of zinc and salicylic acid on wheat under drought stress. *J Anim Plant Sci.* 2016; 4: 1012–1018.
 31. Rahmani F, Sayfzadeh S, Jabbari H, Valadabadi SA, Masouleh EH. Alleviation of drought stress effects on safflower yield by foliar application of zinc. *Int J Plant Prod.* 2019; 4: 297–308.
 32. Anjum SA, Farooq M, Wang LC, Xue LL, Wang SG, Wang L, Zhang S, Chen M. Gas exchange and chlorophyll synthesis of maize cultivars are enhanced by exogenously-applied glycinebetaine under drought conditions. *Plant Soil Environ.* 2011; 7: 326–331.
 33. Marcinska I, Czyczylo-Mysza I, Skrzypek E, Filek M, Grzesiak S, Grzesiak MT, Janowiak F, Hura T, Dziurka M, Dziurka K, Nowakowska A, Quarrie SA. Impact of osmotic stress on physiological and biochemical characteristics in drought susceptible and drought resistant wheat genotypes. *Acta Physiol Plant.* 2013; 451–461.
 34. Raza MAS, Saleem MF, Ashraf MY, Ali A, Asghar HN. Glycinebetaine applied under drought improved the physiological efficiency of wheat (*Triticum aestivum* L.) plant. *Soil Environ.* 2012; 1: 67–71.
 35. Raza MAS, Saleem MF, Shah GM, Khan IH, Raza A. Exogenous application of glycinebetaine and potassium for improving water relations and grain yield of wheat under drought. *J Soil Sci Plant Nut.* 2014; 2: 348–364.
 36. Hasanuzzaman M, Banerje A, Bhuyan MHMB, Roychoudhury A, Al Mahmud J, Fujita M. Targeting glycinebetaine for abiotic stress tolerance in crop plants: physiological mechanism, molecular interaction and signaling. *Phyton-Int J Exp Bot.* 2019; 3: 185–221.
 37. Shemi R, Wang R, Gheith ESM, Hussain H.A, Hussain S, Irfan M. ... Wang, L. Effects of salicylic acid, zinc and glycine betaine on morpho-physiological growth and yield of maize under drought stress. *Sci. Report.* 2021; 1: 1–14.

38. Anjum SA, Ashraf U, Tanveer M, Khan I, Hussain S, Shahzad B, Zohaib A, Abbas F, Saleem MF, Ali I, Wang LC. Drought induced changes in growth, osmolyte accumulation and antioxidant metabolism of three maize hybrids. *Front Plant Sci.* 2017; 69.
39. Bayoumi TY, Eid MH, Metwali EM. Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *Afr J Biotechnol.* 2008; 2341–2352.
40. Zhang B, Li W, Chang X, Li R, Jing R. Effects of favorable alleles for water-soluble carbohydrates at grain filling on grain weight under drought and heat stresses in wheat. *Plos One.* 2014; 7: e102917.
41. Nasrin R, Ali E, Jahanfar D, Soodabeh J. Salicylic acid induced changes on antioxidant capacity, pigments and grain yield of soybean genotypes in water deficit condition. *J Plant Interact.* 2017; 1: 457–464.
42. R, Mittler. Oxidative stress, antioxidants and stress tolerance.. *Trends Plant Sci.* 2002; 405–410.
43. Kamara AY, Menkir A, Badu-Apraku B, Ibikunle O. The influence of drought stress on growth, yield and yield components of selected maize genotypes. *J Agric Sci.* 2003; 1: 43–50.
44. Rollins JA, Habte E, Templer SE, Colby T, Schmidt J, Korff MV. Leaf proteome alterations in the context of physiological and morphological responses to drought and heat stress in barley (*Hordeum vulgare* L.). *J Exp Bot.* 2013; 11: 3201–3212.
45. Abdelkader AF, Aronsson H, Sundqvist C. High salt-stress in wheat leaves causes retardation of chlorophyll accumulation due to a limited rate of protochlorophyllide formation. *Physiol Plant.* 2007; 157–166.
46. Wang GP, Hui Z, Li F, Zhao MR, Zhang J, Wang W. Improvement of heat and drought photosynthetic tolerance in wheat by overaccumulation of glycinebetaine. *Plant Biotechnol Rep.* 2010; 213–222.
47. Kadioglu A, Saruhan N, Saglam A, Terzi R, Acet T. Exogenous salicylic acid alleviates effects of long term drought stress and delays leaf rolling by inducing antioxidant system. *Plant Growth Regul.* 2011; 1: 27–37.
48. Chaves MM, Maroco JP, Pereira JS. Understanding plant responses to drought from genes to the whole plant. *Funct Plant Biol.* 2003; 239–264.
49. Oneto CD, Otegui ME, Baroli I, Beznec A, Faccio P, Bossio E, Blumwald E, Lewi D. Water deficit stress tolerance in maize conferred by expression of an isopentenyltransferase (IPT) gene driven by a stress-and maturation-induced promoter. *J Biotechnol.* 2016; 66–77.
50. Suriyan C, Thapanee S, Chalernpol K. Glycinebetaine alleviates water deficit stress in indica rice using proline accumulation, photosynthetic efficiencies, growth performances and yield attributes. *Aust. J Crop Sci.* 2013; 2: 213–218.
51. Alscher PG, Erturk N, Heath LS. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot.* 2002; 1331–1341.
52. K, Asada. Ascorbate peroxidase-a hydrogen peroxide-scavenging enzyme in plants. *Physiol Plant.* 1992; 2: 235–241.

53. El Tayeb MA, Ahmed NL. Response of wheat cultivars to drought and salicylic acid. *Am-Euras J Agron.* 2010; 1: 1–7.
54. Coombs J, Hall DO, Long SP, Scurlock JMO. *Techniques in bioproductivity and photosynthesis.* Pergamon, Oxford. 1987.
55. CM, Donald. In search of yield. *J Aust Inst Agric Sci.* 1962; 171–178.
56. Peng YS, Liu E. A comparative study of methods of extracting chlorophyll. *Acta Agriculturae Universitatis Pekinensis.* 1992; 247–250.
57. Barrs HD, Weatherley PE. A re-examination of the relative turgidity technique for estimating water deficits in leaves. *Aust J Biol Sci.* 1962; 413–428.
58. University, Michigan State. *MSTAT-C: Micro-computer Statistical Program, Version 2.* Michigan State University, East Lansing, USA. 1990.
59. Steel RC, Torrie SH. *Principles and Procedures of Statistics.* Mc Grauc Hill Book Company, Inc., New York, London. 1997.

Figures




Figure 1

Influences of soil water conditions and foliar treatments on chlorophyll a content, chlorophyll b content, total chlorophyll content and relative water content. Every column in each graph represents the mean (\pm SE) of three replicates. Various letters above columns indicate the significant differences among means ($p < 0.05$). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.




Figure 2

Influences of soil water conditions and foliar treatments on net photosynthesis rate (Pn), transpiration rate (Tr), stomatal conductance (Gs) and intercellular CO₂ concentration (Ci). Every column in each graph represents the mean (\pm SE) of three replicates. Various letters above columns indicate the significant differences among means ($p < 0.05$). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.




Figure 3

Influences of soil water conditions and foliar treatments on the activities of ascorbate peroxidase (APX), glutathione reductase (GR), peroxidase (POD), catalase (CAT), superoxide dismutase (SOD) and malonaldehyde (MDA) content. Every column in each graph represents the mean ($\pm SE$) of three replicates. Various letters above columns indicate the significant differences among means ($p < 0.05$). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.




Figure 4

Influences of soil water conditions and foliar treatments on hydrogen peroxide (H₂O₂) and superoxide anion (O₂⁻) contents. Every column in each graph represents the mean (±SE) of three replicates. Various letters above columns indicate the significant differences among means (p < 0.05). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.




Figure 5

Influences of soil water conditions and foliar treatments on the accumulations of free proline content and total soluble sugar. Every column in each graph represents the mean (±SE) of three replicates. Various letters above columns indicate the significant differences among means (p < 0.05). WW, well-watered; WD, water-deficient; CK, control (double distilled water); SA, salicylic acid; Zn, zinc; GB, glycine betaine.




Figure 6

The various mechanisms of foliar salicylic acid (SA), zinc (Zn) and glycine betaine (GB) treatments induced drought-tolerance in wheat plants. RWC, relative water content; APX, ascorbate peroxidase; GR, glutathione reductase; POD, peroxidase; CAT, catalase; SOD, superoxide dismutase; ROS, reactive oxygen species; MDA, malonaldehyde.