

Prevalence of Healthy Aging Among Community Dwelling Adults Age 70 and Older From Five European Countries

Simeon Schietzel (✉ simeon.schietzel@usz.ch)

Department of Geriatric Medicine and Aging Research, University Hospital Zurich and University of Zurich, Zurich, Switzerland.

Patricia O. Chocano-Bedoya

Center on Aging and Mobility, University Hospital Zurich, City Hospital Waid & Triemli and University of Zurich, Zurich, Switzerland

Angelique Sadlon

Center on Aging and Mobility, University Hospital Zurich, City Hospital Waid & Triemli and University of Zurich, Zurich, Switzerland

Michael Gagesch

Department of Geriatric Medicine and Aging Research, University Hospital Zurich and University of Zurich, Zurich, Switzerland.

Walter C. Willett

Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, Massachusetts.

Endel J. Orav

Department of Biostatistics, Harvard T. H. Chan School of Public Health, Boston, Massachusetts.

Reto W. Kressig

University Department of Geriatric Medicine Felix Platter and University of Basel, Basel, Switzerland.

Bruno Vellas

Gérontopôle de Toulouse, Institut du Vieillissement, Center Hospitalo-Universitaire de Toulouse, Toulouse, France.

René Rizzoli

Division of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland.

José A. P. da Silva

Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal.

Michael Blauth

Department for Trauma Surgery, Medical University of Innsbruck, Innsbruck, Austria.

John A Kanis

Centre for Metabolic Bone Diseases, University of Sheffield Medical School, UK

Andreas Egli

Center on Aging and Mobility, University Hospital Zurich, City Hospital Waid & Triemli and University of Zurich, Zurich, Switzerland

Heike A. Bischoff-Ferrari

Center on Aging and Mobility, University Hospital Zurich, City Hospital Waid & Triemli and University of Zurich, Zurich, Switzerland

Research Article

Keywords: Healthy Aging, DO-HEALTH, older, senior

Posted Date: October 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-923536/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Geriatrics on March 2nd, 2022. See the published version at <https://doi.org/10.1186/s12877-022-02755-8>.

Abstract

Background: To compare the prevalence of healthy aging among adults age 70 and older from 5 European countries recruited for the DO-HEALTH clinical trial. Participants were selected for absence of prior major health events.

Methods: Cross-sectional analysis of DO-HEALTH baseline data. All 2,157 participants (mean age 74.9, SD 4.4; 61.7% women) were included and 2,123 had data for all domains of the healthy aging status (HA) definition. HA was assessed based on the Nurses' Health Study (NHS) definition requiring four domains: no major chronic diseases, no disabilities, no cognitive impairment (Montreal Cognitive Assessment, MoCA ≥ 25), no mental health limitation (GDS-5 < 2 , and no diagnosis of depression). Association between HA and age, BMI, gender, and physical function (sit-to-stand, gait speed, grip strength) was assessed by multivariate logistic regression analyses adjusting for center.

Results: Overall, 41.8% of DO-HEALTH participants were healthy agers with significant variability by country: Austria (Innsbruck) 58.3%, Switzerland (Zurich, Basel, Geneva) 51.2%, Germany (Berlin) 37.6%, France (Toulouse) 36.7% and Portugal (Coimbra) 8.8% ($p < 0.0001$). Differences in prevalence by country persisted after adjustment for age. In the multivariate model, younger age (OR = 0.95, 95% CI 0.93 to 0.98), female gender (OR = 1.36, 95% CI 1.03 to 1.81), lower BMI (OR = 0.94, 95% CI 0.91 to 0.96), faster gait speed (OR = 4.70, 95% CI 2.68 to 8.25) and faster performance in sit-to-stand test (OR = 0.90, 95% CI 0.87 to 0.93) were independently and significantly associated with HA.

Conclusions: Despite the same inclusion and exclusion criteria preselecting relatively healthy adults age 70 years and older, HA prevalence varied significantly between countries and was highest in Austria and Switzerland, lowest in Portugal. Independent of country, younger age, female gender, lower BMI and better physical function were associated with HA.

Trial registration: DO-HEALTH was registered under the protocol NCT01745263 at the International Trials Registry (clinicaltrials.gov), and under the protocol number 2012-001249-41 at the Registration at the European Community Clinical Trial System (EudraCT).

Background

For the five DO-HEALTH countries, the UN Population Division estimated in 2019 that the proportion of older adults will grow considerably. Between 2015 and 2050 the population age 65 and older was predicted to grow from 18–28.7% in Switzerland, from 18.8–29.4% in Austria, from 21.2–30% in Germany, from 18.9–27.8% in France and from 20.8–34.8% in Portugal.(1) The number of older adults with age-related chronic diseases will increase accordingly.(2, 3) Therefore, enabling more older adults to stay healthy and active longer is of major public health importance, and directly linked to reducing the burden of frailty and age-related chronic diseases.(4)

Defining a healthy aging status (HA) is however challenging due to the inherent multidimensionality of human health, including physical and mental functionality to actively participate in life.(5) To date, there is no consensus on a definition. Numerous definitions exist, aiming to capture one or multiple dimensions of human health in older age. Approaches range from simple concepts like absence of diseases and disabilities,(6) up to extensive evaluations of physical and mental diseases as well as cognitive and social function.(7–10)

Based on the requirements of a given HA definition, the prevalence of HA varies from 0.4–41% even in the same population, (11) and reaches up to 92% in one-dimensional definitions of HA.(12, 13) With regard to important European studies on HA (6, 9, 10, 14–21) that focused on adults not younger than 60 and not older than 85 years of age, (9, 19–21) reported prevalence ranges from 1.6–49.3%. A well-validated concept is the Nurses' Health Study (NHS) HA definition,(6, 17, 22–26) which has been linked to healthy survival after the age of 70.(22–24)

In our study, we applied the NHS definition of HA among all participants of DO-HEALTH, a multicentre randomized trial including five countries to explore differences in HA in Europe among pre-selected relatively healthy adults age 70 years and older, living in the community, overall and by subgroups of age and by country. Further, we investigated cross-sectional correlates of HA in this sample of relatively healthy older adults.

Methods

Participants and study design

This is a cross-sectional analysis of baseline data collected in DO-HEALTH, a multi-center randomized controlled trial designed to test the effects of vitamin D, omega-3 and a home-exercise program among community-dwelling adults age 70 years and older (ClinicalTrials.gov Identifier: NCT01745263). The total population included 2157 community-dwelling older adults from Switzerland (Zurich, Basel, Geneva), Austria (Innsbruck), Germany (Berlin), France (Toulouse) and Portugal (Coimbra). By targeting 40% of participants who experienced a fall in the year prior to enrollment, the study aimed to also include pre-frail older adults. For inclusion in DO-HEALTH, participants had to be without major health events (i.e. cancer, angina pectoris, myocardial infarction, stroke, severe kidney or liver disease) in the five years prior to enrollment, a Mini Mental Status Examination (MMSE) score of at least 24 (the MMSE as the cognitive measure for inclusion in DO-HEALTH correlated with outcome measure MOCA in DO-HEALTH at the baseline exam by 0.495), and sufficient mobility to be able to come to the study center. The study was approved by the ethical and regulatory agencies of all 5 countries. Written informed consent was obtained from all participants prior to screening procedures.

Definition of healthy aging status and operationalization in DO-HEALTH

We defined HA according to the definition used in the NHS,(22–24) which includes four domains: no major chronic diseases, no disabilities, no impairment in cognitive function and no mental health limitations. For an in-depth description of items and how they were operationalized in DO-HEALTH see Table 1 and **Appendix 1**.

Table 1

Operationalization of the NHS definition of the healthy aging status phenotype in DO-HEALTH

NHS	DO-HEALTH
No major chronic diseases	
NHS questionnaire 1. No type 2 diabetes 2. No congestive heart failure 3. No kidney failure 4. No chronic obstructive pulmonary disease 5. No cancer other than non-melanoma skin cancer 6. No myocardial infarction 7. No coronary artery bypass surgery, or angioplasty 8. No stroke 9. No Parkinson's disease 10. No multiple sclerosis 11. No amyotrophic lateral sclerosis	Sangha 1. No type 2 diabetes 2. No heart disease 3. No kidney disease 4. No lung disease 5. No cancer other than non-melanoma skin cancer * 6. No myocardial infarction * 7. No coronary intervention * 8. No stroke * 9. No Parkinson`s disease † 10. No multiple sclerosis † 11. No amyotrophic lateral sclerosis †
No disabilities	

NHS	DO-HEALTH
<p>SF-36</p> <p>No limitation in:</p> <ol style="list-style-type: none"> 1. Bathing 2. Dressing 3. Climbing one flight of stairs 4. Walking >1 mile or walking several blocks 5. Moving a table 6. Bowling or Golf 7. Pushing a vacuum cleaner <p>No more than moderate limitations in:</p> <ol style="list-style-type: none"> 8. Lifting or carrying groceries 9. Bending, kneeling, or stooping 10. Lifting heavy objects 11. Climbing several flights of stairs 	<p>PROMIS</p> <p>No limitation in:</p> <ol style="list-style-type: none"> 1. Wash and dry your body 2. Dress yourself, including shoelaces and buttons 3. Climb up five steps 4. Walk a block on flat ground 5. Stand up from an armless straight chair ± 6. Get in and out of bed ± 7. Get on and off the toilet ± <p>No more than moderate limitations in:</p> <ol style="list-style-type: none"> 8. Run errands and shop 9. Do chores such as vacuuming or yard work 10. Reach and get down a 2-Kg object from above head 11. Bend down and pick up items from the floor ±
<p>No impairment in cognitive function</p>	
<p>TICS score >31</p>	<p>MoCA score >=25 §</p>
<p>No mental health limitation</p>	
<p>SF-36, MHI score >84/100</p> <p>How much of the time during the last month have you:</p> <ol style="list-style-type: none"> 1. Felt so down in the dumps that nothing could cheer you up? 2. Been a happy person? 3. Felt downhearted and blue? 4. Been a very nervous person? 5. Felt calm and peaceful? 	<p>GDS-5 score < 2/5</p> <ol style="list-style-type: none"> 1. Do you feel pretty worthless the way you are now? 2. Are you basically satisfied with your life? 3. Do you often feel helpless? 4. Do you prefer to stay at home rather than going out and doing new things? 5. Do you often get bored? <p>Plus: No diagnosis of depression (Sangha)</p>

NHS	DO-HEALTH
<p>NHS = NursesHea < hStudy. DO - HEALTH = Vita min D3 - Ω s self-administered comorbidity questionnaire. TICS= Telephone Interview for Cognitive Status. MoCA = Montreal Cognitive Assessment. SF-36 = Short Form 36 Health Survey Questionnaire. MHI = Mental Health Index. GDS-5 = Geriatric Depression Scale. * Exclusion criteria of DO-HEALTH. † Not assessed, but severe cases unlikely to be included as seniors with advanced mobility limitations were not included in DO-HEALTH. ± Substitutes, not matching corresponding items from the NHS definition but picturing elements of the NHS requirements regarding disabilities and physical functions. § In analogy to the TICS cutoff, separating seniors with normal cognition from those with mild cognitive impairment (Brand et al. Ciesielska et al.).</p>	

(1) no major chronic diseases

Based on the Sangha self-administered comorbidity questionnaire which included diabetes, lung, heart and kidney disease,(27) and considering exclusion criteria from the trial (no cancer, no mobility impairments).

(2) no impairment in cognitive function

We used the Montreal Cognitive Assessment questionnaire (MoCA),(28) with a cut off score of ≥ 25 as suggested by a recent meta-analysis of 20 studies among community-dwelling seniors.(29)

(3) no mental health limitations

We required healthy agers to score < 2 points in the Geriatric Depression Scale short form, comprising five questions (GDS-5),(30) and no self-reported diagnosis of depression, ascertained by the Sangha.(31)

(4) no disabilities

We used the Patient Reported Outcome Measurement Information Questionnaire (PROMIS),(32) to evaluate basic and instrumental activities of daily living as well as higher physical function. We required that healthy agers had no limitation in simple activities (e.g. dressing, climbing 5 steps) and no more than moderate limitations in more complicated activities (e.g. running errands and shopping).

Ascertainment of covariates

Body mass index (BMI) was ascertained dividing participants` weight in Kg (assessed wearing not more than underwear) by their height squared in meters (assessed without shoes). Physical function was ascertained with the timed repeated sit-to-stand task and with a test of grip strength.(33) In the timed repeated sit-to-stand test, participants had to rise up 5 times from a chair without the help of their arms. Grip strength was measured in Kilo Pascal (KPa) with the Martin Vigorimeter (KLS Martin Group, Tuttlingen Germany). We used mean values from three attempts performed with the dominant hand. Formal education was assessed in total years of formal education completed. All measurements were performed by trained study nurses following standardized procedures in each site.

Statistical analysis

Baseline characteristics of DO-HEALTH participants are presented as means (standard deviations) for continuous variables and frequencies (percentages) for categorical variables and stratified by country. We conducted ANOVA and chi square tests for comparisons of all characteristics by center,

The prevalence of the HA definition and its components was evaluated as percentages and stratified by age groups (70-74, 75-79 and 80 years and older) and countries. To evaluate that the differences in prevalence between countries were not influenced by age, we also present for each country the age-adjusted prevalence using a logistic regression model with HA as the outcome and age and country as predictors. In addition, we compared the characteristics of participants with HA vs. those without (age, BMI, education, faller status, timed sit-to-stand performance, grip strength, gait speed) using logistic regression adjusting each variable for each other and adjusting for center.

Results

Baseline characteristics of participants

The total number of participants with complete data on all domains required to define HA by the NHS tool was 2,123. Among them, 1,306 (61.5%) were women. Participants` mean age was 74.9 (4.4) years. The mean number of comorbidities assessed by Sangha questionnaire was 1.7 (1.4) and mean MMSE and MoCA scores were 28.5 (1.5) and 25.7 (3.3) respectively. Women had worse physical function with regard to both repeated sit-to-stand and gait speed findings (Table 2).

Table 2
Characteristics of participants by country

	Total	Austria	France	Germany	Portugal	Switzerland
	(n=2123)	(n=199)	(n=281)	(n=348)	(n=297)	(n=998)
Age, years	74.9 (4.4)	74.1 (4.1)	75.2 (4.3)	73.3 (2.7)	76 (5)	75.1 (4.6)
Women, N (%)	1306 (61.5%)	102 (51.3)	165 (58.7)	246 (70.7)	189 (63.6)	604 (60.5)
BMI, kg/m ²						
Men	26.6 (3.5)	25.5 (3.3)	26.8 (3.2)	26.7 (3)	28 (3.5)	26.4 (3.6)
Women	26.2 (4.7)	25 (4.4)	25.1 (4.3)	26.9 (4.7)	29.1 (4.4)	25.5 (4.5)
Mean number of comorbidities	1.7 (1.4)	1.5 (1.3)	2.0 (1.4)	1.7 (1.3)	2.6 (1.7)	1.4 (1.2)
MoCA score	25.7 (3.3)	26.9 (2.5)	27 (2.3)	25.3 (2.3)	21.9 (4.3)	26.3 (2.8)
GDS-5 score	1.8 (2.3)	1.3 (1.4)	2.4 (2.4)	1.2 (1.5)	4.1 (3.6)	1.2 (1.6)
Education, years	12.7 (4.3)	12 (3.7)	13.3 (3.9)	14.5 (3.3)	7.9 (5.4)	13.4 (3.4)
Prior fall, N (%)	884 (41.6)	98 (49.3)	118 (42)	124 (35.6)	120 (40.4)	424 (42.5)
Sit-to-stand, s						
Men	11.2 (4)	8.8 (2.9)	13.5 (3.3)	9.5 (2.1)	15.6 (5.7)	10.4 (2.9)
Women	11.9 (4.3)	10.2 (3.0)	14.4 (4.2)	9.3 (2.3)	17 (5.6)	11.0 (3.0)
Gait speed, m/s						
Men	1.15 (0.22)	1.17 (0.22)	1.14 (0.22)	1.31 (0.18)	1.10 (0.23)	1.12 (0.22)
Women	1.10 (0.23)	1.12 (0.22)	1.06 (0.19)	1.26 (0.2)	0.93 (0.23)	1.09 (0.21)

Values are means and standard deviations unless otherwise noted. Y = years, BMI = Body Mass Index, MoCA = Montreal cognitive Assessment, GDS-5 = Geriatric Depression Scale short form, s = seconds, m/s = meters per second.

Components of the HA definition

For the total study population and based on the four components of the NHS HA definition, 77.6% of participants reported absence of major chronic diseases, 73.3% reported no disabilities in simple tasks and no more than moderate disabilities in complex tasks, 69.9% reported no cognitive impairment and 84.7% had no mental health problems. Healthy agers had a mean MoCA score of 27.5 (SD=1.6) and mean GDS-5 of 0.84 (SD=1.06) whereas the non-healthy agers had a mean MoCA score of 24.4 (SD=3.6) and GDS-5 of 2.44 (SD=2.72).

Prevalence of HA overall, and by age and country

In total, 41.8% of participants (887/2123) met all four requirements and thereby qualified as healthy agers. By country, participants from Austria had the highest prevalence of HA with 58.3%, followed by participants from Switzerland 51.2%, Germany 37.6% and France 36.7% whereas the prevalence of HA in participants from Portugal was the lowest with 8.8% ($p < .0001$). (Figure 1a and 1b). After adjustment by age, Austria and Switzerland remained the countries with highest prevalence of HA and Portugal was the lowest (Figure 1a).

The prevalence was significantly higher in younger participants (47.6% among those 70 to <75 years, 38.7% among 75 to <80 years and 24.6% among those 80 years and older, $p < 0.0001$) (Figure 2a).

Performance in the four components of the HA definition by country

With regard to the four components of the NHS HA definition each country showed a differential performance profile. (Figure 2b).

Characteristics of healthy agers

In the unadjusted analyses, healthy agers were on average younger (74.0 vs. 75.5 years, $P < .0001$), had more years of education, (13.4 vs 12.1, $P < .0001$), had a lower BMI (25.3 vs. 27.1 kg/m², $p < .0001$) and had better physical function (faster gait speed, higher grip strength, shorter time in the sit-to-stand test, and less likely to have a prior fall). (Table 3).

Table 3
Characteristics associated with healthy aging status

	Healthy Agers*	Non-Healthy Agers*	p-value†	Multivariate model‡	
	n=887	n=1236		OR (95%CI)	p-value
Age, years	74.0 (3.8)	75.5 (4.7)	< .0001	0.95 (0.93,0.98)	.0001
Women [%]	61.3%	61.6%	0.88	1.36 (1.03,1.81)	0.0319
Education, years	13.4 (3.5)	12.1 (4.7)	< .0001	1.02 (0.99,1.05)	0.233
Prior fall [%]	38.9%	42.6%	0.03	0.87 (0.72,1.07)	0.1813
BMI, Kg/m ²	25.3 (3.8)	27.1 (4.4)	< .0001	0.94 (0.91,0.96)	< .0001
Gait speed, m/s					
Men	1.20 (0.20)	1.12 (0.23)	< .0001	4.70 (2.68,8.25)	< .0001
Women	1.18 (0.20)	1.04 (0.24)	< .0001		
Grip strength, KPa					
Men	79.0 (15.7)	73.4 (17.5)	< .0001	1.01 (1.00,1.01)	0.2655
Women	53.2 (10.5)	48.9 (12.3)	< .0001		
Sit-to-stand, s					
Men	10.0 (2.9)	12.1 (4.5)	< .0001	0.90 (0.87,0.93)	< .0001
Women	10.3 (2.7)	13.1 (4.9)	< .0001		
*Values are means and SE unless noted. † p-values are from T-tests or X ² tests					
‡Variables in the multivariate model are adjusted for each other and for center.					

In the multivariate adjusted model, younger age, female gender, lower BMI, faster gait speed and a shorter time in sit-to-stand test were independently associated with a prevalent HA. For every additional year of age, participants had 5% lower odds of being healthy agers (OR = 0.95, 95% CI 0.93 to 0.98, P = .0001) and for every additional BMI point, participants had 6% lower odds of being healthy agers (OR = 0.94, 95% CI 0.91 to 0.96, P = < .0001). Women had 36% higher odds of being healthy agers (OR = 1.36, 95% CI 1.03 to 1.81, P = .0319). For every meter per second increase in gait speed participants had 4-times higher odds of being healthy agers (OR = 4.70, 95% CI 2.68 to 8.25, P = < .0001) but for every second increase in the sit-to-stand test, participants had 10% lower odds of being healthy agers (OR = 0.90, 95% CI 0.87 to 0.93, P = < .0001). Years of education, prior falls and grip strength were not independently associated with HA (Table 3).

Discussion

In this large cross-sectional study, we examined 2,123 DO-HEALTH trial participants recruited from 5 countries pre-selected to be relatively healthy older adults age 70 years and older. On average, 41.8% of participants met the HA criteria, but there were significant differences between countries. Prevalence of HA was highest in Switzerland (51.2%) and Austria (58.3%) and lowest in Portugal (8.8%). At the cross-sectional level, HA was independently and significantly associated with younger age, female gender, lower BMI and better physical function regarding gait speed and sit-to-stand test. Notably, however, education, prior falls and grip strength were not independently associated with HA.

Similar to our findings, results from a population-based study, the Survey of Health, Aging and Retirement in Europe (SHARE) revealed substantial variability in HA prevalence between countries ranging from 21.1% in Denmark to 1.6% in Poland among adults with a mean age of 72 (6.7) years.⁽⁹⁾ These differences support variations in health states of European older adults by country also at the population-based level. Given the pre-selection of relatively healthy adults with good mobility and cognitive function and no major health events in the 5 years prior to recruitment, DO-HEALTH compared with SHARE, shows a higher prevalence of HA in community-dwelling European adults age 70 and older. Also, SHARE used a different definition of HA,⁽⁸⁾ that included a domain of “social engagement”.

In our study, based on the NHS definition, Portugal had the lowest prevalence of HA in comparison to all other countries, despite the same inclusion and exclusion criteria and even after age adjustment. Consistent with our results, SHARE reported a lower prevalence of HA in southern countries of Europe (Italy 5.3%; Greece 7.7%; Spain 3.1%) compared to countries like Austria 10.2%, Switzerland 16.1% or Germany 11.6%.⁽⁹⁾ The observed difference in the prevalence of HA between Portugal and the five other European countries in DO-HEALTH could potentially be explained by differences in socioeconomic status.⁽³⁴⁾ The available median equivalent purchasing power per capita in Portugal in 2016 was around half as much as in France, Germany and Austria, and only round one third as much as in Switzerland.⁽³⁴⁾ Consistently, in DO-HEALTH, years of education as a surrogate to income, differed between countries and were lowest in Portugal (mean = 8.0, SD = 5.4 years) and highest in Germany (mean = 14.5, SD = 3.3 years). However, education was not independently associated with the total score of the HA definition in DO-HEALTH.

With regard to age, DO-HEALTH reflects a decline in HA with age even among this relatively healthy selection of older adults from 47.6–38.7% to 24.6 % in adults 70-74, 75-79 and more than 80 years of age, respectively. While this is best explained by a higher incidence of chronic diseases, disabilities, cognitive impairment and mental health limitations with advanced age, our data also supports the potential of being a healthy ager even at age 80 and older in one out of four cases.

With regard to gender, we found an independent association between HA and female gender. Reports from the Organisation for Economic Co-operation and Development (OECD) as well as results from

European studies of gender differences have repeatedly shown advantages concerning life expectancy, ischemic heart diseases, cancer and general health status for women compared with men.(35), (36), (37)

In support of the important role of physical activity in overall health and aging, HA in DO-HEALTH was independently associated with faster gait speed and better performance in the sit-to-stand.

Further, DO-HEALTH suggests that for every additional BMI point, older adults may have 6% lower odds of being healthy agers. In fact, healthy agers showed, on average, BMI levels around the upper limit of normal (25 [3.8] Kg/m²), and non-healthy agers were mildly overweight (27.1 [4.4] Kg/m²). These findings are consistent with the literature, where a normal BMI (18.5 - 24.9 Kg/m²) is associated with reduced mortality compared to the BMI values in the overweight and obese spectrum.(38–41) Alternatively, in unselected older adults, BMI level above normal have been associated with reduced mortality.(42–45) The optimal BMI for older adults is not known, optimal BMI level between 20 and 29.9 Kg/m² have been described.(44) DO-HEALTH might suggest that among relatively healthy older adults, the upper normal range may be most advantageous for HA, however, prospective investigations are needed to determine an optimal BMI in this population.

Our study has several strengths. We used a well-validated NHS definition of HA, extracted from standardized clinical health assessments derived from the baseline examination of a large clinical trial. Further, our study reflects extremely well phenotyped adults age 70 and older from 5 European countries including both southern and central Europe, hereby including possible differences also in terms of socioeconomic status, educational level and healthcare access. The observed differences in HA are conservative as we targeted relatively healthy older adults, and the observed pattern of a lower HA prevalence in southern Europe and specifically Portugal is supported by the literature.(9, 46) However, it is noteworthy that the cities and samples included in each study do not necessarily represent the entire country.

Our study also has limitations. The cross-sectional nature of our analysis does not allow us to draw a causal relationship between the covariates explored for their association with HA. Further, DO-HEALTH is not a population-based study but reflects a sample of relatively healthy older adults with the same inclusion and exclusion criteria applied in all 5 countries applied in a rigorous clinical trial setting. However, even though inclusion and exclusion criteria were identical between centers, bias due to different priorities in defined recruitment strategies between centers cannot be completely excluded.

Conclusions

In conclusion, in this sample of pre-selected relatively healthy European adults age 70 and older, the prevalence of HA differed across five countries with the lowest prevalence in Portugal compared with Austria, Switzerland, Germany and France. Independent of country, prevalence of HA was associated with higher age, lower BMI, female and better physical function. Further studies are needed to examine differences in HA between European countries at the population-based and prospective level.

Nonetheless, the cross-sectional assessment of HA in DO-HEALTH among relatively healthy older adults may reflect a conservative look at health disparities between countries and trigger efforts at the public health sphere.

Abbreviations

HA = Healthy Aging, NHS = Nurses` Health Study, MoCA = Montreal Cognitive Assessment, MMSE = Mini Mental Status Examination, PROMIS = Patient Reported Outcome Measurement Information Questionnaire, SHARE = Survey of Health, Aging and Retirement in Europe

Declarations

Ethical approval and consent to participate:

In our study we investigate baseline data of the DO-HEALTH trial. DO-HEALTH was approved by local/national ethics committee and regulatory authorities of all 5 countries (Switzerland, Germany, Austria, France, Portugal). For the coordinating centre of Do-HEALTH, the accredited ethics committee Cantonal Ethics Commission (CEC) Zurich, Switzerland, approved the trial. A project ethic committee provided ethical guidance to the consortium and ensured that the clinical trial was carried out within national and European regulations. DO-HEALTH was a Phase III trial (therapeutic confirmatory) and was conducted according to the ICH GCP Guidelines (CPMP/ICH/135/95). All participants gave their unpersuaded and full consent to participate in the trial.

Consent for publication:

Not applicable.

Availability of data and materials:

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

The study was funded by the Seventh Framework Program of the European Commission (grant agreement 278588), the University of Zurich (Chair for Geriatric Medicine and Aging Research), DSM Nutritional Products, Roche, NESTEC, Pfizer, and Streuli. The funding/ supporting organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the

data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Authors` contributions:

SS, PC, HBF, WW, EO analysed and interpreted the patient data. PC performed the statistical analyses supported by HBF, WW and EO. SS, PC and HBF were major contributor in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgments:

The authors explicitly express their gratitude to all participants and teams of the 7 recruitment sites for their dedication in collecting all DO-HEALTH data in the yearly clinical visits and in-person phone calls every 3 months.

References



1. United Nations, Department of Economic and Social Affairs, Population Division (2019). World Population Prospects 2019: Volume II: Demographic Profiles.
2. Active Ageing: A Policy Framework.: World Health Organization; 2002.
3. Jackson RD, Shidham S. The role of hormone therapy and calcium plus vitamin D for reduction of bone loss and risk for fractures: lessons learned from the Women's Health Initiative. *Curr Osteoporos Rep.* 2007;5(4):153–9.
4. European Union 2012. COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT AND THE COUNCIL. Taking forward the Strategic Implementation Plan of the European Innovation Partnership on Active and Healthy Ageing, Brussels 29.02.2012.
5. Global strategy and action plan on ageing and health. Geneva: World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.
6. Britton A, Shipley M, Singh-Manoux A, Marmot MG. Successful aging: the contribution of early-life and midlife risk factors. *Journal of the American Geriatrics Society.* 2008;56(6):1098–105.
7. Strawbridge WJ, Wallhagen MI, Cohen RD. Successful aging and well-being: self-rated compared with Rowe and Kahn. *The Gerontologist.* 2002;42(6):727–33.
8. McLaughlin SJ, Connell CM, Heeringa SG, Li LW, Roberts JS. Successful Aging in the United States: Prevalence Estimates From a National Sample of Older Adults. *The journals of gerontology Series B, Psychological sciences and social sciences.* 2010;65b(2):216–26.
9. Hank K. How "successful" do older Europeans age? Findings from SHARE. *The journals of gerontology Series B, Psychological sciences and social sciences.* 2011;66(2):230–6.
10. Kok AA, Aartsen MJ, Deeg DJ, Huisman M. Capturing the Diversity of Successful Aging: An Operational Definition Based on 16-Year Trajectories of Functioning. *The Gerontologist.* 2017;57(2):240–51.

11. Garfein AJ, Herzog AR. Robust Aging among the Young-Old, Old-Old, and Oldest-Old. *The Journals of Gerontology: Series B*. 1995;50B(2):S77-S87.
12. Montross L, Depp C, Daly J, Reichstadt J, Golshan S, Moore D, et al. Correlates of Self-Rated Successful Aging Among Community-Dwelling Older Adults. *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry*. 2006;14:43–51.
13. Depp CA, Jeste DV. Definitions and predictors of successful aging: a comprehensive review of larger quantitative studies. *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry*. 2006;14(1):6–20.
14. de Moraes JF, de Azevedo e Souza VB. Factors associated with the successful aging of the socially-active elderly in the metropolitan region of Porto Alegre. *Revista brasileira de psiquiatria (Sao Paulo, Brazil: 1999)*. 2005;27(4):302-8.
15. Bowling EGA. Enhancing the quality of extended life years. Identification of the oldest old with a very good and very poor quality of life. *Aging & mental health*. 1999;3(3):199–212.
16. Nosraty L, Sarkeala T, Hervonen A, Jylha M. Is there successful aging for nonagenarians? The vitality 90+ study. *Journal of aging research*. 2012;2012:868797.
17. Sabia S, Singh-Manoux A, Hagger-Johnson G, Cambois E, Brunner EJ, Kivimaki M. Influence of individual and combined healthy behaviours on successful aging. *CMAJ: Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2012;184(18):1985–92.
18. von Faber M, Bootsma-van der Wiel A, van Exel E, Gussekloo J, Lagaay AM, van Dongen E, et al. Successful aging in the oldest old: Who can be characterized as successfully aged? *Arch Intern Med*. 2001;161(22):2694–700.
19. Formiga F, Ferrer A, Megido MJ, Chivite D, Badia T, Pujol R. Low co-morbidity, low levels of malnutrition, and low risk of falls in a community-dwelling sample of 85-year-olds are associated with successful aging: the Octabaix study. *Rejuvenation research*. 2011;14(3):309–14.
20. Habib R, Nyberg L, Nilsson LG. Cognitive and non-cognitive factors contributing to the longitudinal identification of successful older adults in the betula study. *Neuropsychology, development, and cognition Section B, Aging, neuropsychology and cognition*. 2007;14(3):257–73.
21. Puig-Domingo M, Serra-Prat M, Merino MJ, Pubill M, Burdoy E, Papiol M. Muscle strength in the Mataro aging study participants and its relationship to successful aging. *Aging clinical and experimental research*. 2008;20(5):439–46.
22. Sun Q, Townsend MK, Okereke OI, Franco OH, Hu FB, Grodstein F. Adiposity and weight change in mid-life in relation to healthy survival after age 70 in women: prospective cohort study. *BMJ (Clinical research ed)*. 2009;339:b3796.
23. Sun Q, Townsend MK, Okereke OI, Franco OH, Hu FB, Grodstein F. Physical activity at midlife in relation to successful survival in women at age 70 years or older. *Arch Intern Med*. 2010;170(2):194–201.
24. Sun Q, Townsend MK, Okereke OI, Rimm EB, Hu FB, Stampfer MJ, et al. Alcohol Consumption at Midlife and Successful Ageing in Women: A Prospective Cohort Analysis in the Nurses' Health Study.

- PLoS Medicine. 2011;8(9).
25. Uotinen V, Suutama T, Ruoppila I. Age identification in the framework of successful aging. A study of older Finnish people. *International journal of aging & human development*. 2003;56(3):173–95.
 26. McLaughlin SJ, Jette AM, Connell CM. An examination of healthy aging across a conceptual continuum: prevalence estimates, demographic patterns, and validity. *J Gerontol A Biol Sci Med Sci*. 2012;67(7):783–9.
 27. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The self-administered comorbidity questionnaire: A new method to assess comorbidity for clinical and health services research. *Arthritis Rheum-Arthr*. 2003;49(2):156–63.
 28. Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*. 2005;53(4):695–9.
 29. Ciesielska N, Sokolowski R, Mazur E, Podhorecka M, Polak-Szabela A, Kedziora-Kornatowska K. Is the Montreal Cognitive Assessment (MoCA) test better suited than the Mini-Mental State Examination (MMSE) in mild cognitive impairment (MCI) detection among people aged over 60? Meta-analysis. *Psychiatria polska*. 2016;50(5):1039–52.
 30. Hoyl MT, Alessi CA, Harker JO, Josephson KR, Pietruszka FM, Koelfgen M, et al. Development and testing of a five-item version of the Geriatric Depression Scale. *Journal of the American Geriatrics Society*. 1999;47(7):873–8.
 31. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum*. 2003;49(2):156–63.
 32. Maska L, Anderson J, Michaud K. Measures of functional status and quality of life in rheumatoid arthritis: Health Assessment Questionnaire Disability Index (HAQ), Modified Health Assessment Questionnaire (MHAQ), Multidimensional Health Assessment Questionnaire (MDHAQ), Health Assessment Questionnaire II (HAQ-II), Improved Health Assessment Questionnaire (Improved HAQ), and Rheumatoid Arthritis Quality of Life (RAQoL). *Arthritis care & research*. 2011;63 Suppl 11:S4-13.
 33. Sipers WM, Verdijk LB, Sipers SJ, Schols JM, van Loon LJ. The Martin Vigorimeter Represents a Reliable and More Practical Tool Than the Jamar Dynamometer to Assess Handgrip Strength in the Geriatric Patient. *Journal of the American Medical Directors Association*. 2016;17(5):466.e1-7.
 34. Bundesamt für Statistik. Median equivalent purchasing power per capita in Europe. Eurostat - EU-SILC2016 (Version 15.02.2018).
 35. OECD. Health at a Glance 2019. 2019.
 36. Cash AWaK. The state of men's health in Western Europe. *The Journal of Men's Health and Gender* 2004 1:1, 60–66. 2004(1):60–6.
 37. Wang H, Dwyer-Lindgren L, Lofgren KT, Rajaratnam JK, Marcus JR, Levin-Rector A, et al. Age-specific and sex-specific mortality in 187 countries, 1970–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2012;380(9859):2071–94.

38. Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* (London, England). 2009;373(9669):1083–96.
39. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *Jama*. 2013;309(1):71–82.
40. Global BMIMC, Di Angelantonio E, Bhupathiraju Sh N, Wormser D, Gao P, Kaptoge S, et al. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet* (London, England). 2016;388(10046):776–86.
41. Aune D, Sen A, Prasad M, Norat T, Janszky I, Tonstad S, et al. BMI and all cause mortality: systematic review and non-linear dose-response meta-analysis of 230 cohort studies with 3.74 million deaths among 30.3 million participants. *BMJ* (Clinical research ed). 2016;353:i2156.
42. Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *The American journal of clinical nutrition*. 2014;99(4):875–90.
43. Chung WS, Ho FM, Cheng NC, Lee MC, Yeh CJ. BMI and all-cause mortality among middle-aged and older adults in Taiwan: a population-based cohort study. *Public health nutrition*. 2015;18(10):1839–46.
44. Tamakoshi A, Yatsuya H, Lin Y, Tamakoshi K, Kondo T, Suzuki S, et al. BMI and all-cause mortality among Japanese older adults: findings from the Japan collaborative cohort study. *Obesity* (Silver Spring). 2010;18(2):362–9.
45. Guallar-Castillón P, Balboa-Castillo T, López-García E, León-Muñoz LM, Gutiérrez-Fisac JL, Banegas JR, et al. BMI, Waist Circumference, and Mortality According to Health Status in the Older Adult Population of Spain. 2009;17(12):2232–8.
46. WHO. European health report 2018: More than numbers - evidence for all. 2018.

Figures




1a


1b

Figure 1

1a. Crude and adjusted* prevalence of healthy aging status by country *Adjusted for by age, gender, education, faller status, timed sit-to-stand performance, grip strength, gait speed. Bars show 95% confidence intervals. 1b. Prevalence of healthy agers by country



2a



2b

Figure 2

2a. Healthy aging status by age groups 2b. Participants (%) meeting requirements of single domains of the healthy aging status definition

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterial.docx](#)