

Serum Neurofilament Measurement Improves Clinical Risk Score for Outcome Prediction After Cardiac Arrest: Results of A Prospective Study

Sabina Hunziker (✉ sabina.hunziker@usb.ch)

Universitätsspital Basel <https://orcid.org/0000-0003-3648-3609>

Adrian Quinto

University Hospital Basel: Universitätsspital Basel

Maja Ramin-Wright

University Hospital Basel: Universitätsspital Basel

Christoph Becker

University Hospital Basel: Universitätsspital Basel

Katharina Beck

University Hospital Basel: Universitätsspital Basel

Alessia Vincent

University Hospital Basel: Universitätsspital Basel

Kai Tisljar

university hospital basel

Giulio Disanto

University hospital basel

Pascal Benkert

University Hospital Basel: Universitätsspital Basel

David Leppert

university hospital basel

Hans Pargger

university hospital basel

Stephan Marsch

university hospital basel

Raoul Sutter

University Hospital Basel: Universitätsspital Basel

Nils Peters

university hospital basel

Jens Kuhle

University Hospital Basel: Universitätsspital Basel

Research

Keywords: serum neurofilament, cardiac arrest, prognosis, CAHP, OHCA, cardiopulmonary resuscitation

Posted Date: October 19th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-92711/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on January 20th, 2021. See the published version at <https://doi.org/10.1186/s13054-021-03459-y>.

Abstract

Background: A recent study found serum neurofilament light chain (NfL) levels to be strongly associated with poor neurological outcome in patients after cardiac arrest. Our aim was to confirm these findings in an independent validation study and to investigate whether NfL improves the prognostic value of two cardiac arrest risk scores.

Methods: This prospective, single-center study included 164 consecutive adult cardiac arrest patients upon intensive care unit admission. We calculated two clinical risk scores (OHCA, CAHP) and measured NfL on admission using the single molecule array NF-light® assay. The primary endpoint was neurological outcome at hospital discharge assessed with the cerebral performance category (CPC) score.

Results: Poor neurological outcome (CPC \geq 3) was found in 60% (98/164) of patients, and 55% (91/164) died. Compared to patients with favorable outcome, NfL was 14-times higher in patients with poor neurological outcome (685 \pm 1787 vs. 49 \pm 111pg/mL), with an adjusted odds ratio of 3.4 (95%CI 2.1 to 5.6, p<0.001) and an area under the curve (AUC) of 0.82. Adding NfL to the clinical risk scores significantly improved discrimination of both the OHCA score (from AUC 0.82 to 0.89, p<0.001) and CAHP score (from AUC 0.89 to 0.92, p<0.05). Admission NfL showed better outcome prediction compared to neuron-specific enolase (NSE) (AUC 0.84 vs.0.69, p=0.01).

Conclusions: This study confirms the high performance of admission NfL alone and in combination with two clinical risk scores to prognosticate clinical outcome in patients after cardiac arrest. NfL should be considered as a standard laboratory measures in the evaluation of cardiac arrest patients.

Introduction

Despite the increased survival rates thanks to medical treatments, the mortality and risk for neurological deficits remains high for cardiac arrest patients[1–3]. Prognostication of outcome upon admission is difficult in these patients, yet, early identification of predictors for a poor outcome after out-of-hospital cardiac arrest (OHCA) would facilitate the therapeutic management, decision-making, and communication with relatives [4]. The associated high economic burden of comatose critically ill patients further increases the impact not just on an individual level, but also on society at large. For risk prediction of adverse clinical outcome, several clinical tools have been developed. Two scores were specifically developed for cardiac arrest patients including the Out-of-Hospital Cardiac Arrest score (OHCA) and the Cardiac Arrest Hospital Prognosis score (CAHP) [5–7]. These scores, however, still have limited accuracy and, thus, bear the risk of misclassifying patients. The improvement of these clinical scores by the addition of biomarkers reflecting pathophysiologic mechanisms of neural damage is an unmet medical need. Herein, different biomarkers indicating brain injuries after OHCA have been studied[8, 9], yet neuron-specific enolase (NSE) is currently the only biomarker recommended by guidelines as a prognostic blood marker for patients after cardiac arrest [9–12].

Recently, elevated levels of neurofilament light chain (NfL) in blood samples have been established as marker of neuronal damage in traumatic brain injury and many acute and chronic neurologic diseases[13–19]. NfL thus has potential to further improve assessment of neurological damage in OHCA patients. To date, one small pilot study investigated this biomarker in cerebrospinal fluid[20] and three studies in the blood of patients after cardiac arrest[21, 22]. Two studies analyzed neurofilament levels with standard immunoassays and reported associations of NfL with poor neurological outcome in patients after cardiac arrest [21, 22]. One of these studies used an ultrasensitive single molecule array (Simoa) assay to test prognostication of neurological outcome after OHCA in a large cohort of patients[23]. Serum NfL levels at 24 hours after cardiac arrest showed high discrimination regarding long-term poor neurological outcome with an area under the curve (AUC) of 0.94–0.95, which was better compared to other biomarkers (i.e. tau, neuron-specific enolase (NSE), and S100). Before wide-spread implementation of NfL, independent validation in a larger sample is mandatory.

Herein, the aim of this study was to externally validate the prognostic accuracy of NfL in a well characterized sample of patients after OHCA and to study whether NfL improves current OHCA specific risk scores, namely, the OHCA and CHAP scores.

Methods

Study setting

This is a prospective observational study including consecutive patients from November 2012 until February 2016 with data obtained in the COMMUNICATE trial at the University Hospital Basel, Switzerland. The main purpose of this study is to investigate novel biomarkers for risk stratification of OHCA patients. The methods used for this study have been published previously[7, 24, 25]. The study was approved by the local ethics committee. Patients or their relatives provided informed consent for study participation.

This manuscript adheres to the STROBE guidelines.

Study population

We included consecutive patients after cardiac arrest who were admitted to the ICU of the University Hospital Basel, a Swiss tertiary academic medical center, into the study. We did not use any exclusion criteria regarding the patients' characteristics and type or duration of the cardiac arrest except for patients having to be adults.

Data collection

We calculated all scores as suggested[5, 6, 26, 27] and as described in more detail in a previous publication[7]. In brief, we used data collected on the first day of admission. Resuscitation data were collected from clinical records, including no-flow time (time from cardiac arrest to start of basic life support (BLS)), low-flow time (time from start of BLS to return of spontaneous circulation (ROSC)), initial rhythm, setting and location of arrest, epinephrine dose given, as well as information on whether bystanders observed the cardiac arrest and started cardiopulmonary resuscitation (CPR). Clinical parameters (i.e., heart rate, blood pressure, Glasgow Coma Scale (GCS), respiratory rate, temperature, urine output), socio-demographics (i.e., age, gender, family situation, socio-economic status), comorbidities (i.e., coronary disease, congestive heart failure, hypertension, chronic obstructive pulmonary disease (COPD), malignant disease, diabetes, renal failure, liver failure) and cardiovascular risk factors (i.e., smoking status, diabetes, high blood lipids, family history) were recorded from medical records or during an interview with patients' relatives. We also collected initial levels of routine blood markers, such as sodium, potassium, pH, bicarbonate, lactate, haematocrit, white blood cell count, creatinine, urea, bilirubin.

We also calculated two risk scores that were specifically developed for outcome prediction in the cardiac arrest patient population including the out-of-hospital cardiac arrest (OHCA) score and the Cardiac Arrest Hospital Prognosis (CAHP) score[5, 6]. The OHCA score was developed in 2006 in France and later validated in the US and Switzerland[5, 7, 28]. This score, which incorporates initial rhythm, no-flow and low-flow intervals, and admission levels of serum creatinine and lactate into a risk model, showed a high performance for outcome prediction. The CAHP score was developed later and includes additional resuscitation specific variables, such as location of cardiac arrest, epinephrine dose as well as initial blood pH values and age [29].

Measurement of fluid biomarkers

We measured NfL in admission blood samples. After blood draw, all samples were pre-analytically processed, aliquoted, and frozen at -80°C . Serum NfL levels were measured in duplicates (intra-assay coefficients of variation $< 20\%$) by Simoa NfL assay as described previously[30]. Inter-assay CVs were 11.8% and 14.1% for two native serum samples (104 pg/ml and 31 pg/ml), respectively. We also measured NSE levels using an Electro-Chemi-Luminescent-Immuno-Assay (ECLIA) kit (Roche Diagnostics, Rotkreuz, Switzerland).

Outcome

Similar to previous studies, the primary endpoint was neurological outcome at hospital discharge assessed with the cerebral performance category (CPC) Score[31]. In line with previous studies, levels 1 (good recovery) and 2 (moderate disability) were considered as favourable neurological outcome, whereas levels 3 (severe disability), 4 (vegetative state) and 5 (death) were defined as unfavourable outcome[5, 6]. We also measured the modified Rankin Scale (mRS) at discharge. We defined levels 0 (no symptoms), 1 (no significant disability despite symptoms; able to carry out all usual duties and activities), 2 (slight disability; unable to carry out all previous activities, but able to look after own affairs without assistance), and 3 (moderate disability; requiring some help, but able to walk without assistance) as favourable outcome, and levels 4 (moderately severe disability, unable to walk and attend to bodily needs without assistance), 5 (severe disability; bedridden, incontinent and requiring constant nursing care and attention) and 6 (dead) as unfavourable outcome[32, 33].

Secondary endpoints were all-cause mortality at 30 days after discharge.

Statistical analysis

To characterise the patient cohort, descriptive statistics including means (\pm SD), medians and inter-quartile ranges were used for continuous variables as appropriate, whereas frequencies were reported for binary or categorical variables. We calculated univariable and multivariable logistic regression models to evaluate the association of NfL levels with the primary and secondary endpoints. To achieve a normal distribution, data of NfL levels were log transformed with a base of 10. Odds ratios (OR) and 95% confidence intervals (CI) were reported as a measure of association. Covariates used in the multivariate analysis were selected based on prior evidence of an association with unfavourable neurological outcome for patients with OHCA. Multivariate models were adjusted for age, gender and comorbidities. In a further step, we additionally included OHCA and CAHP scores into the models and also calculated AUCs of combined models to see whether NfL would improve discrimination of the scores. We also investigated the prognostic performance of NfL regarding sensitivity, specificity, positive and negative predictive values and likelihood ratios at three cut-offs (25, 50, 75) corresponding to the interquartile range and median.

STATA 15.0 was used for all statistical analyses and a two-sided p-value of < 0.05 was considered significant.

Results

Baseline characteristics

From the 164 included patients, 98 (60%) had poor neurological outcome and 91 (55%) patients died. The baseline characteristics of the cohort overall and stratified based on neurological outcome are shown in Table 1. Overall, the mean age was 63 years and 28.7% were female. Patients showed relevant comorbidities and cardiovascular risk factors. Patients with poor neurological outcome had less pre-existing coronary heart disease and a higher incidence of diabetes mellitus and malignant disease than patients with good neurological outcome. In patients with poor neurological outcome the cardiac arrest was more often unwitnessed, and they less frequently received bystander CPR and further had longer no-flow and low flow time than patients with a good neurological outcome.

Table 1
Baseline characteristics

Factor	CPC All	Good neurological outcome, CPC 1–2	Poor neurological outcome, CPC 3–5	p-value
N, n (%)	164 (100%)	66 (40%)	98 (60%)	
Sociodemographics				
Age, mean (SD)	63 (15)	60 (17)	65 (14)	0.027
Male, n (%)	117 (71.3%)	55 (83%)	62 (63%)	0.008
Comorbidities				
Coronary heart disease, n (%)	109 (66.5%)	50 (76%)	59 (60%)	0.044
Congestive heart failure, n (%)	20 (12.2%)	7 (11%)	13 (13%)	0.81
COPD, n (%)	15 (9.1%)	3 (5%)	12 (12%)	0.11
Liver disease, n (%)	3 (1.8%)	0 (0%)	3 (3%)	0.27
Hypertension, n (%)	86 (52.4%)	37 (56%)	49 (50%)	0.52
Diabetes, n (%)	39 (23.8%)	9 (14%)	30 (31%)	0.015
Chronic kidney disease, n (%)	24 (14.6%)	8 (12%)	16 (16%)	0.51
Malignant disease, n (%)	14 (8.5%)	2 (3%)	12 (12%)	0.047
Neurological disease, n (%)	13 (7.9%)	2 (3%)	11 (11%)	0.077
Resuscitation measures				
No-flow time, min, mean (SD)	4.22 (5.93)	1.70 (2.96)	6.02 (6.83)	< 0.001
Low-flow time, min, mean (SD)	20.42 (15.13)	16.53 (14.30)	23.12 (15.17)	0.006
Cardiac arrest setting				
At home	63 (38.4%)	16 (24%)	47 (48%)	< 0.001
In public	78 (47.6%)	43 (65%)	35 (36%)	
In hospital	23 (14.0%)	7 (11%)	16 (16%)	
Witnessed	137 (83.5%)	63 (95%)	74 (76%)	< 0.001
Bystander CPR	103 (62.8%)	51 (77%)	52 (53%)	0.002
Initial heart rhythm				
Ventricular fibrillation	6 (3.7%)	2 (3%)	4 (4%)	< 0.001
Ventricular tachycardia	88 (53.7%)	53 (80%)	35 (36%)	
Asystole	35 (21.3%)	2 (3%)	33 (34%)	
Pulseless electrical activity	29 (17.7%)	6 (9%)	23 (23%)	
Unknown	6 (3.7%)	3 (5%)	3 (3%)	
Epinephrine during resuscitation (mg), mean (SD)	2.18 (2.55)	1.08 (1.76)	2.96 (2.73)	< 0.001
Targeted Temperature Management (TTM), n (%)	109 (66.5%)	40 (61%)	69 (70%)	0.19
Blood markers				

Data presented as n (%) or mean (standard deviation) and median (Inter Quartile Range, IQR). COPD, chronic obstructive pulmonary disease; CPR, cardiopulmonary resuscitation; NfL, Neurofilament Light Chain.

Factor	CPC All	Good neurological outcome, CPC 1–2	Poor neurological outcome, CPC 3–5	p-value
NfL (pg/ml), mean (SD)	429 (1415)	49 (111)	685 (1787)	0.004
NfL (pg/ml), median (IQR)	51 (21, 173)	27 (13, 46)	116 (41, 330)	< 0.001

Data presented as n (%) or mean (standard deviation) and median (Inter Quartile Range, IQR). COPD, chronic obstructive pulmonary disease; CPR, cardiopulmonary resuscitation; NfL, Neurofilament Light Chain.

Primary endpoint: Poor neurological outcome measured with the cerebral performance category (CPC) score

NfL levels were higher in patients with poor neurological outcome compared to patients with good neurological outcome (mean/median 685/116 [SD ± 1787] vs mean/median 49/27 [SD ± 111], $p = 0.004$) with an univariate OR (of log-transformed NfL) of 2.9 (95%CI 2.0 to 4.3, $p < 0.001$) and a multivariate OR of 3.4 (95%CI 2.1 to 5.6, $p < 0.001$) after adjusting for age, gender and comorbidities (Table 2). Further, the risk for poor neurological outcome increased more than 40-fold for patients in the highest quartile compared to the lowest quartile of NfL concentrations (adjusted OR 47.1 [95%CI 9.8 to 227.0], $p < 0.001$).

Table 2
Comparison between scores to predict primary and secondary endpoints

Primary endpoint: Neurological outcome (CPC)										
	Good neurological outcome (CPC 1–2)	Poor neurological outcome (CPC 3–5)	p-value	Univariate		Multivariate* adjusted	NFL and OHCA score: multivariate* adjusted		NFL and CAHP score: multivariate* adjusted	
	n = 66	n = 98		OR (95%CI), p	AUC	OR (95%CI), p	OR (95%CI), p	AUC combined with NFL	OR (95%CI), p	AUC combined with NFL
NfL, mean (SD)	49 (111)	685 (1787)	0.004	2.93 (2.01, 4.27), p < 0.001	0.82	3.44 (2.13, 5.55), p < 0.001	2.81 (1.69, 4.67), p < 0.001	0.89	3.45 (1.89, 6.29), p < 0.001	0.92
NfL, median (IQR)	27 (13, 46)	116 (41, 330)	<0.001							
Quartiles of NFL:			< 0.001	<i>reference group</i>		<i>reference group</i>	<i>reference group</i>		<i>reference group</i>	
1st quartile	29 (44%)	12 (12%)								
2nd quartile	25 (38%)	16 (16%)		1.55 (0.62, 3.88), p = 0.35		1.35 (0.46, 4.03), p = 0.59	1.02 (0.28, 3.7), p = 0.97		1.01 (0.24, 4.23), p = 0.98	
3rd quartile	10 (15%)	31 (32%)		7.49 (2.81, 19.96), p < 0.001		5.59 (1.65, 18.94), p = 0.006	3.33 (0.82, 13.41), p = 0.091		3.99 (0.84, 18.96), p = 0.08	
4th quartile	2 (3%)	39 (40%)		47.12 (9.78, 227.01), p < 0.001		127.26 (16.53, 979.47), p < 0.001	92.39 (8.47, 1007.63), p < 0.001		140.48 (10.5, 1880.26), p < 0.001	
OHCA, mean (SD)	11 (16)	32 (17)	< 0.001	1.08 (1.05, 1.1), p < 0.001	0.82	1.08 (1.05, 1.11), p < 0.001	1.07 (1.04, 1.11), p < 0.001			
CAHP, mean (SD)	118 (41)	182 (38)	< 0.001	1.04 (1.03, 1.05), p < 0.001	0.89	1.04 (1.03, 1.05), p < 0.001			1.04 (1.02, 1.06), p < 0.001	
Secondary endpoint: Neurological outcome (mRs)										
	Good neurological outcome (mRs 0–2)	Poor neurological outcome (mRs 3–6)	p-value	Univariate		Multivariate* adjusted	NFL and OHCA score: multivariate* adjusted		NFL and CAHP score: multivariate* adjusted	
	n = 53	n = 111		OR (95%CI), p	AUC	OR (95%CI), p	OR (95%CI), p	AUC combined with NFL	OR (95%CI), p	AUC combined with NFL
NFL, mean (SD)	51 (123)	609 (1691)	0.018	2.72 (1.85, 3.98), p < 0.001	0.8	3.01 (1.88, 4.83), p < 0.001	2.47 (1.52, 4.04), p < 0.001	0.87	2.78 (1.64, 4.72), p < 0.001	0.88
NfL, median (IQR)	21 (12, 40)	92 (37, 284)	< 0.001							

* adjusted for age, gender, comorbidities;

Data presented as mean (SD) and median (Inter Quartile Range, IQR).; NfL = Neurofilament Light Chain (pg/ml); AUC = area under the curve; CAHP = Cardiac Arrest Hospital prognosis (-score); CPC = cerebral performance category; OHCA = Out-of-Hospital Cardiac Arrest (-score); OR = odds ratio; ROC = receiver operating characteristics curve.

Primary endpoint: Neurological outcome (CPC)										
Quartiles of NFL:	26 (49.1%)	15 (13.5%)	< 0.001	<i>reference group</i>		<i>reference group</i>	<i>reference group</i>		<i>reference group</i>	
1st quartile										
2nd quartile	18 (34.0%)	23 (20.7%)		2.21 (0.91, 5.37), p = 0.078		2.11 (0.73, 6.1), p = 0.17	1.8 (0.56, 5.83), p = 0.326		1.89 (0.55, 6.46), p = 0.309	
3rd quartile	7 (13.2%)	34 (30.6%)		8.42 (3, 23.64), p < 0.001		6.14 (1.73, 21.8), p = 0.005	3.6 (0.9, 14.41), p = 0.07		3.97 (0.94, 16.7), p = 0.06	
4th quartile	2 (3.8%)	39 (35.1%)		33.8 (7.13, 160.31), p < 0.001		66.08 (9.66, 452.17), p < 0.001	42.17 (4.72, 376.98), p = 0.001		53.32 (5.52, 514.78), p = 0.001	
OHCA, mean (SD)	11 (16)	30 (18)	< 0.001	1.06 (1.04, 1.09), p < 0.001	0.79	1.07 (1.04, 1.1), p < 0.001	1.06 (1.03, 1.08), p < 0.001			
CAHP, mean (SD)	117 (42)	175 (42)	< 0.001	1.03 (1.02, 1.04), p < 0.001	0.84	1.03 (1.02, 1.04), p < 0.001			1.03 (1.02, 1.04), p < 0.001	
Secondary endpoint: 30-day mortality										
	Survivors	Non-Survivors	p-value	Univariate		Age, gender, comorbidities adjusted	NFL and OHCA score: multivariate* adjusted		NFL and CAHP score: multivariate* adjusted	
	n = 131	n = 127		OR (95%CI), p	AUC	OR (95%CI), p	OR (95%CI), p	AUC combined with NFL	OR (95%CI), p	AUC combined with NFL
NfL, mean (SD)	136 (701)	664 (1763)	0.017	2.77 (1.94, 3.95), p < 0.001	0.83	2.97 (1.91, 4.62), p < 0.001	2.42 (1.54, 3.8), p < 0.001	0.87	2.59 (1.6, 4.2), p < 0.001	0.89
NfL, median (IQR)	27 (13, 47)	128 (45, 330)	<0.001							
Quartiles of NFL:	33 (45%)	8 (9%)	< 0.001	<i>reference group</i>		<i>reference group</i>	<i>reference group</i>		<i>reference group</i>	
1st quartile										
2nd quartile	25 (34%)	16 (18%)		2.64 (0.98, 7.14), p = 0.056		2.47 (0.75, 8.1), p = 0.135	2.29 (0.62, 8.42), p = 0.211		2.32 (0.58, 9.3), p = 0.235	
3rd quartile	11 (15%)	30 (33%)		11.25 (3.99, 31.71), p < 0.001		7.85 (2.19, 28.09), p = 0.002	5.49 (1.39, 21.76), p = 0.015		5.87 (1.35, 25.57), p = 0.019	
4th quartile	4 (5%)	37 (41%)		38.16 (10.52, 138.44), p < 0.001		92.29 (15.16, 561.71), p < 0.001	49.33 (7.29, 333.79), p < 0.001		65.12 (8.32, 509.95), p < 0.001	

* adjusted for age, gender, comorbidities;

Data presented as mean (SD) and median (Inter Quartile Range, IQR).; NfL = Neurofilament Light Chain (pg/ml); AUC = area under the curve; CAHP = Cardiac Arrest Hospital prognosis (-score); CPC = cerebral performance category; OHCA = Out-of-Hospital Cardiac Arrest (-score); OR = odds ratio; ROC = receiver operating characteristics curve.

Primary endpoint: Neurological outcome (CPC)										
OHCA, mean (SD)	14 (18)	32 (17)	< 0.001	1.06 (1.04, 1.09), p < 0.001	0.77	1.07 (1.04, 1.1), p < 0.001	1.05 (1.03, 1.08), p < 0.001			
CAHP, mean (SD)	123 (43)	182 (38)	< 0.001	1.03 (1.02, 1.04), p < 0.001	0.85	1.03 (1.02, 1.04), p < 0.001			1.03 (1.02, 1.04), p < 0.001	
* adjusted for age, gender, comorbidities;										
Data presented as mean (SD) and median (Inter Quartile Range, IQR).; NfL = Neurofilament Light Chain (pg/ml); AUC = area under the curve; CAHP = Cardiac Arrest Hospital prognosis (-score); CPC = cerebral performance category; OHCA = Out-of-Hospital Cardiac Arrest (-score); OR = odds ratio; ROC = receiver operating characteristics curve.										

NfL showed a comparable prognostic discrimination with an area under the curve (AUC) of 0.82 to both established clinical risk scores. Adding NfL to the risk scores further improved discrimination between good and poor neurological outcome (from AUC 0.82 with OHCA score alone to 0.89 in combination with NfL, p < 0.001, from AUC 0.89 with CAHP score alone to 0.92 in combination, p < 0.05).

Table 3 shows sensitivity and specificity, positive and negative predictive value as well as positive and negative likelihood ratios for NfL at three different cut-offs. At the calculated optimal cut-off (Youden -Index) of 50 pg/ml, we found a corresponding sensitivity of 72.4% and specificity of 81.8%, with 85.5% positive predictive value and 66.7% negative predictive value for poor neurological outcome. Further, we calculated prognostic measures based on receiver operating characteristic (ROC) curve analysis for the cut-offs at a NfL level of 25 pg/ml and 75 pg/ml. For the cut-off of 25 pg/ml, the sensitivity was 87.8% and specificity 47%, with 71.1% positive predictive value and 72.1% negative predictive value. Finally, a cut-off of 75 pg/ml revealed a sensitivity of 61.2%, specificity of 90.9%, a 90.9% positive predictive value, and a negative predictive value of 61.2%.

Table 3
Performance of NfL at different cut-off points to predict neurological Outcome

	CPC score			mRS			Mortality 30 days		
	NfL cut-off 25	NfL cut-off 50	NfL cut-off 75	NfL cut-off 25	NfL cut-off 50	NfL cut-off 75	NfL cut-off 25	NfL cut-off 50	NfL cut-off 75
Prevalence Pr(A)	59.80%	59.80%	59.80%	67.70%	67.70%	67.70%	55.50%	55.50%	55.50%
Sensitivity Pr(+ A)	87.80%	72.40%	61.20%	85.60%	66.70%	55.00%	91.20%	74.70%	63.70%
Specificity Pr(-N)	47.00%	81.80%	90.90%	50.90%	83.00%	90.60%	47.90%	79.50%	89.00%
ROC area (Sens. + Spec.)/2	0.67	0.77	0.76	0.68	0.75	0.73	0.7	0.77	0.76
Likelihood ratio (+) Pr(+ A)/Pr(+ N)	1.65	3.98	6.73	1.74	3.93	5.83	1.75	3.64	5.82
Likelihood ratio (-) Pr(-A)/Pr(-N)	0.26	0.34	0.43	0.28	0.4	0.5	0.18	0.32	0.41
Odds ratio LR(+)/LR(-)	6.35	11.83	15.79	6.17	9.78	11.71	9.56	11.43	14.28
Positive predictive value Pr(A+)	71.10%	85.50%	90.90%	78.50%	89.20%	92.40%	68.60%	81.90%	87.90%
Negative predictive value Pr(N-)	72.10%	66.70%	61.20%	62.80%	54.30%	49.00%	81.40%	71.60%	66.30%
Data presented as mean (SD); NfL = Neurofilament Light Chain (pg/ml); CAHP = Cardiac Arrest Hospital prognosis (-score); CPC = cerebral performance category; mRS = modified Rankin Scale; LLR + = positive likelihood ratio; LLR- = negative likelihood ratio; NPV = negative predictive value; PPV = positive predictive value.									

Secondary endpoints

Secondary endpoints were the modified Rankin Scale (mRS) and all-cause mortality at hospital discharge (Table 2).

In terms of the mRS, the mean/median NfL levels were also significantly higher in patients with poor neurological outcome (609 pg/ml/92 [SD ± 1691] vs 51/21 pg/ml [SD ± 123], $p = 0.018$) with a corresponding univariate OR of 2.72 (95%CI 1.85 to 3.98). These results stayed robust after adjusting in the full model (OR of 3.01 [95%CI 1.88 to 4.83]).

NfL showed a good prognostic discrimination for poor neurological outcome assessed with the mRS with an AUC of 0.80 which further improved in combination with the two risk scores to an AUC of 0.87 with OHCA and to an AUC of 0.88 with CAHP.

In Table 3 at the calculated optimal cut-off of 50 pg/ml, sensitivity was 66.7% and specificity was 83%, with 89.2% positive predictive value and 54.3% negative predictive value for poor neurological outcome. Further, we calculated the prognostic measures based on ROC curve analysis for the cut-offs at a NfL level of 25 pg/ml and 75 pg/ml. For the cut-off of 25 pg/ml, the calculated sensitivity was 85.6% and the specificity 50.9%, with 78.5% positive predictive value and 62.8% negative predictive value. Further, a cut-off of 75 pg/ml showed a sensitivity of 55%, specificity 90.6%, a 92.4% positive predictive value and a negative predictive value of 49%.

Similarly, NfL was significantly associated with mortality, also after adjusting in the full model (mean/median 664/128 [SD 1763] vs mean/median 136/27 [SD 701], $p = 0.017$) with a good prognostic discrimination (AUC 0.83). Also, we found an improvement of the prognostic value by combination of NfL with the OHCA and the CAHP risk scores (AUC from 0.77 to 0.87 and AUC 0.85 to 0.89 respectively).

At the calculated optimal cut-off of 50 pg/ml, we found a sensitivity of 74.7% and specificity of 79.5%, with 81.9% positive predictive value and 71.6% negative predictive value for poor neurological outcome. For the cut-off of 25 pg/ml, the sensitivity was 91.2% and specificity 47.9%, with 68.6% positive predictive value and 81.4% negative predictive value. Finally, a cut-off of 75 pg/ml revealed a sensitivity of 63.7%, specificity of 89%, an 87.9% positive predictive value, and a negative predictive value of 66.3%.

Figure 1 displays Kaplan–Meier curves for 30-day all-cause mortality based on the three different NfL cut-offs.

We additionally performed subgroup analyses to evaluate differences in associations of NfL with neurological outcome in specific patient groups. Results were also robust in these subgroups for different variables (Fig. 2).

We also compared NfL with NSE in patients who had available measurement of NSE on admission ($n = 131$). In this group, we found a significantly better prognostic performance of NfL compared to NSE (AUC NfL 0.84 vs. NSE 0.69, $p = 0.01$). Admission NfL also had a significantly higher AUC compared to admission NSE for neurological outcome measured with the mRS (AUC 0.82 vs 0.66, $p = 0.008$) and for mortality (AUC 0.83 vs 0.68, $p = 0.014$) (Fig. 3).

Discussion

In this prospective cohort of cardiac arrest patients, we externally validated previous research and found that serum NfL level is a reliable predictor for poor neurological outcome and in-hospital mortality. Associations of NfL and outcome also remained significant in models adjusted for age, gender, comorbidities and cardiac-arrest specific risk scores. In addition, our data show that NfL significantly improved the prognostic value of two established cardiac arrest specific scoring systems (OHCA and CHAP Score) to predict outcome after cardiac arrest. Results of the previous derivation studies and our prospective validation suggest that serum NfL has high potential to improve patient care regarding early risk stratification in the population of cardiac arrest patients.

Neurofilaments have emerged in recent years as biofluid markers of neuronal damage which causes the release of these intracellular cytoskeleton proteins into cerebrospinal fluid and blood. Due to the development of ultrasensitive methods such as Simoa, levels are now quantifiable in serum or plasma, allowing longitudinal measurements for monitoring purposes. NfL is currently the most widely used marker among the three sizes of neurofilaments, mainly due to superior assay performance. NfL levels closely reflect the rate and degree of disease acuity in many chronic and acute illnesses of the central nervous system, as well as in traumatic brain injury[13]. Further, in many of these conditions NfL levels are now established on the group level as predictors for later functional neurological and neuropsychological outcome. Importantly, despite these promising results in other neurological illnesses, there has been relatively little clinical data regarding NfL in cardiac arrest patients.

Our results are in line with two previous studies indicating that NfL is a good novel marker for prognosticating neurological outcome after cardiac arrest[21, 23]. While these studies focused on long-term outcomes after six months, our data suggest that NfL also helps to predict short-term clinical outcomes and thus may help navigate the therapeutic management in an early stage after cardiac arrest. We also investigated the potential of NfL to improve the cardiac specific OHCA and the CAHP scores[5, 24, 28]. Importantly, our data show that NfL measured early in the course after cardiac arrest further improves the discrimination of these clinical risk scores. An important advantage of these scores is that their

calculation is based on initial ICU parameters that are readily available upon patient admission, allowing to support clinical decision-making regarding the initial management of cardiac arrest patients. At this time, prognostication is particularly challenging as patients are often unconscious, limiting the clinical assessment. Early prognostic information, however, may help to inform relatives about expected risks and thus influence discussions about potential withdrawal of therapy. Clearly, interventional research is needed to understand how prognostic information influences these decisions in more detail.

In patients after cardiac arrest, NSE is currently the only biomarker recommended for prognostication by European and American guidelines[34, 35]. In a previous study, we found that NSE is highly predictive for adverse outcome when measured at day 3 of admission, while initial NSE levels measured on day 1 provided only little prognostic information[36]. The current analysis as well as another recent study found that the prognostic value of NSE on admission was inferior to NfL indicating that NfL may be a more accurate outcome marker early in the course of disease[23]. This may be explained by differences in marker kinetics and other factors such as influence of hemolysis on measurement characteristics. Importantly, admission NfL had a better prognostic performance to predict outcome compared to NSE. Thus, future guidelines should consider including NfL for early prognostication in patients after cardiac arrest as NSE shows only suboptimal results in the early course of disease.

In our study, NfL was measured at baseline within 24 hours after onset of cardiac arrest. Literature regarding the kinetics of serum NfL after cardiac arrest is still sparse, especially within the early course before 24 hours or after 72 hours. In previous stroke and cardiac arrest populations, NfL levels in patients with poor outcome were nearly doubled from 24 to 48 hours after cardiac arrest, reaching a steady state between 48 and 72 hours [19, 23]. As a limitation, we did not measure NfL at other time points than admission, but possibly looking at kinetics could further improve its prognostic value.

Strength And Limitations

Strengths of this study are the representative sample size with prospective and consecutive inclusion of patients and blinded analysis of blood markers. There are, however, some limitations: First, this is an observational study and is thus only hypothesis generating. Second, we did not measure NfL during follow-up and thus cannot make any statements about its kinetics. Third, external validation of the NfL cutoff levels proposed by our analysis is necessary before wide-spread use in clinical practice.

Conclusions

This study confirms the high performance of admission NfL alone and in combination with two clinical risk scores to prognosticate clinical outcome in patients after cardiac arrest. NfL should be considered as a standard laboratory measure in the evaluation of cardiac arrest patients.

Abbreviations

AUC = area under the curve; BLS = basic life support; CAHP = Cardiac Arrest Hospital Prognosis; CI = confidence interval; COPD = chronic obstructive pulmonary disease; CPC = cerebral performance category; CPR = cardiopulmonary resuscitation; GCS = Glasgow Coma Scale; ICU = intensive care; IQR = Inter Quartile Range;; mRS = modified Rankin Scale; NfL = Neurofilament light chain; NPV = negative predictive value; NSE = neuron-specific enolase; OHCA = out-of-hospital cardiac arrest; OR = odds ratio; OR = Odds ratio; PPV = positive predictive value; ROC = receiver-operating characteristic; ROSC = return of spontaneous circulation; SD = Standard deviation

Declarations

Ethics approval and consent to participate

Our study complies with the Declaration of Helsinki. The Ethics Committee of Northwest and Central Switzerland (Ethikkommission Nordwest- und Zentralschweiz, EKNZ) approved this study.

Consent for publication

Not applicable.

Availability of supporting data

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Sabina Hunziker and her research team is supported by the Swiss National Foundation (SNF) (Ref 10001C_192850/1 and 10531C_182422) and the Gottfried Julia Bangerter-Rhyner Foundation (8472/HEG-DSV) and the Swiss Society of General Internal Medicine (SSGIM). Jens Kuhle is supported by the Swiss National Research Foundation (320030_189140/1).

Author's contributions

SH conceived the design of the study and und wrote the study proposal. KB, AV, CB, KT, R.S., H.P., S.M. and SH collected the data and contributed to the evaluation and interpretation of data. S.H., A.Q., and M.R.W. performed statistical analysis of all data and wrote the first draft of the manuscript. J.K., G.D., P.B., D.L., N.P., and their team performed the laboratory analyses, and contributed to interpretation of data and writing the manuscript. All authors critically reviewed and approved the final manuscript.

Acknowledgements


We would like to acknowledge the medical ICU and laboratory staff of the University Hospital of Basel for making this study possible. We thank all patients and their family members for participating in this study.

References

1. Monsieurs KG, Nolan JP, Bossaert LL, Greif R, Maconochie IK, Nikolaou NI, Perkins GD, Soar J, Truhlar A, Wyllie J *et al*: **European Resuscitation Council Guidelines for Resuscitation 2015: Section 1. Executive summary.** *Resuscitation* 2015, **95**:1-80.
2. Young GB: **Clinical practice. Neurologic prognosis after cardiac arrest.** *N Engl J Med* 2009, **361**(6):605-611.
3. Blom MT, Beesems SG, Homma PC, Zijlstra JA, Hulleman M, van Hoeijen DA, Bardai A, Tijssen JG, Tan HL, Koster RW: **Improved survival after out-of-hospital cardiac arrest and use of automated external defibrillators.** *Circulation* 2014, **130**(21):1868-1875.
4. Nolan JP, Soar J, Cariou A, Cronberg T, Moulart VR, Deakin CD, Bottiger BW, Friberg H, Sunde K, Sandroni C *et al*: **European Resuscitation Council and European Society of Intensive Care Medicine 2015 guidelines for post-resuscitation care.** *Intensive care medicine* 2015, **41**(12):2039-2056.
5. Adrie C, Cariou A, Mourvillier B, Laurent I, Dabbane H, Hantala F, Rhaoui A, Thuong M, Monchi M: **Predicting survival with good neurological recovery at hospital admission after successful resuscitation of out-of-hospital cardiac arrest: the OHCA score.** *European heart journal* 2006, **27**(23):2840-2845.
6. Maupain C, Bougouin W, Lamhaut L, Deye N, Diehl JL, Geri G, Perier MC, Beganton F, Marijon E, Jouven X *et al*: **The CAHP (Cardiac Arrest Hospital Prognosis) score: a tool for risk stratification after out-of-hospital cardiac arrest.** *European heart journal* 2016, **37**(42):3222-3228.
7. Isenschmid C, Luescher T, Rasiyah R, Kalt J, Tondorf T, Gamp M, Becker C, Tisljar K, Sutter R, Schuetz P *et al*: **Performance of clinical risk scores to predict mortality and neurological outcome in cardiac arrest patients.** *Resuscitation* 2019, **136**:21-29.
8. Rossetti AO, Rabinstein AA, Oddo M: **Neurological prognostication of outcome in patients in coma after cardiac arrest.** *The Lancet Neurology* 2016, **15**(6):597-609.
9. Stammer P, Collignon O, Hassager C, Wise MP, Hovdenes J, Aneman A, Horn J, Devaux Y, Erlinge D, Kjaergaard J *et al*: **Neuron-Specific Enolase as a Predictor of Death or Poor Neurological Outcome After Out-of-Hospital Cardiac Arrest and Targeted Temperature Management at 33 degrees C and 36 degrees C.** *Journal of the American College of Cardiology* 2015, **65**(19):2104-2114.
10. Steffen IG, Hasper D, Ploner CJ, Schefold JC, Dietz E, Martens F, Nee J, Krueger A, Jorres A, Storm C: **Mild therapeutic hypothermia alters neuron specific enolase as an outcome predictor after resuscitation: 97 prospective hypothermia patients compared to 133 historical non-hypothermia patients.** *Crit Care* 2010, **14**(2):R69.
11. Oksanen T, Tiainen M, Skrifvars MB, Varpula T, Kuitunen A, Castren M, Pettila V: **Predictive power of serum NSE and OHCA score regarding 6-month neurologic outcome after out-of-hospital ventricular fibrillation and therapeutic hypothermia.** *Resuscitation* 2009, **80**(2):165-170.
12. Cronberg T, Rundgren M, Westhall E, Englund E, Siemund R, Rosen I, Widner H, Friberg H: **Neuron-specific enolase correlates with other prognostic markers after cardiac arrest.** *Neurology* 2011, **77**(7):623-630.
13. Khalil M, Pirpamer L, Hofer E, Voortman MM, Barro C, Leppert D, Benkert P, Ropele S, Enzinger C, Fazekas F *et al*: **Serum neurofilament light levels in normal aging and their association with morphologic brain changes.** *Nat Commun* 2020, **11**(1):812.
14. Shahim P, Gren M, Liman V, Andreasson U, Norgren N, Tegner Y, Mattsson N, Andreasen N, Ost M, Zetterberg H *et al*: **Serum neurofilament light protein predicts clinical outcome in traumatic brain injury.** *Scientific reports* 2016, **6**:36791.
15. Hansson O, Janelidze S, Hall S, Magdalinou N, Lees AJ, Andreasson U, Norgren N, Linder J, Forsgren L, Constantinescu R *et al*: **Blood-based NFL: A biomarker for differential diagnosis of parkinsonian disorder.** *Neurology* 2017, **88**(10):930-937.
16. Mattsson N, Cullen NC, Andreasson U, Zetterberg H, Blennow K: **Association Between Longitudinal Plasma Neurofilament Light and Neurodegeneration in Patients With Alzheimer Disease.** *JAMA neurology* 2019.

17. De Marchis GM, Katan M, Barro C, Fladt J, Traenka C, Seiffge DJ, Hert L, Gensicke H, Disanto G, Sutter R *et al*: **Serum neurofilament light chain in patients with acute cerebrovascular events.** *European journal of neurology* 2018, **25**(3):562-568.
18. Feneberg E, Oeckl P, Steinacker P, Verde F, Barro C, Van Damme P, Gray E, Grosskreutz J, Jardel C, Kuhle J *et al*: **Multicenter evaluation of neurofilaments in early symptom onset amyotrophic lateral sclerosis.** *Neurology* 2018, **90**(1):e22-e30.
19. Tiedt S, Duering M, Barro C, Kaya AG, Boeck J, Bode FJ, Klein M, Dorn F, Gesierich B, Kellert L *et al*: **Serum neurofilament light: A biomarker of neuroaxonal injury after ischemic stroke.** *Neurology* 2018, **91**(14):e1338-e1347.
20. Rosen C, Rosen H, Andreasson U, Bremell D, Bremner R, Hagberg L, Rosengren L, Blennow K, Zetterberg H: **Cerebrospinal fluid biomarkers in cardiac arrest survivors.** *Resuscitation* 2014, **85**(2):227-232.
21. Rana OR, Schroder JW, Baukloh JK, Saygili E, Mischke K, Schiefer J, Weis J, Marx N, Rassaf T, Kelm M *et al*: **Neurofilament light chain as an early and sensitive predictor of long-term neurological outcome in patients after cardiac arrest.** *International journal of cardiology* 2013, **168**(2):1322-1327.
22. Rundgren M, Friberg H, Cronberg T, Romner B, Petzold A: **Serial soluble neurofilament heavy chain in plasma as a marker of brain injury after cardiac arrest.** *Crit Care* 2012, **16**(2):R45.
23. Moseby-Knappe M, Mattsson N, Nielsen N, Zetterberg H, Blennow K, Dankiewicz J, Dragancea I, Friberg H, Lilja G, Insel PS *et al*: **Serum Neurofilament Light Chain for Prognosis of Outcome After Cardiac Arrest.** *JAMA neurology* 2019, **76**(1):64-71.
24. Isenschmid C, Kalt J, Gamp M, Tondorf T, Becker C, Tisljar K, Locher S, Schuetz P, Marsch S, Hunziker S: **Routine blood markers from different biological pathways improve early risk stratification in cardiac arrest patients: Results from the prospective, observational COMMUNICATE study.** *Resuscitation* 2018, **130**:138-145.
25. Metzger K, Gamp M, Tondorf T, Hochstrasser S, Becker C, Luescher T, Rasiah R, Boerlin A, Tisljar K, Emsden C *et al*: **Depression and anxiety in relatives of out-of-hospital cardiac arrest patients: Results of a prospective observational study.** *Journal of critical care* 2019, **51**:57-63.
26. Knaus WA, Draper EA, Wagner DP, Zimmerman JE: **APACHE II: a severity of disease classification system.** *Crit Care Med* 1985, **13**(10):818-829.
27. Le Gall JR, Lemeshow S, Saulnier F: **A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study.** *JAMA* 1993, **270**(24):2957-2963.
28. Hunziker S, Bivens MJ, Cocchi MN, Miller J, Saliccioli J, Howell MD, Donnino MW: **International validation of the out-of-hospital cardiac arrest score in the United States.** *Crit Care Med* 2011, **39**(7):1670-1674.
29. Martinell L, Nielsen N, Herlitz J, Karlsson T, Horn J, Wise MP, Uden J, Rylander C: **Early predictors of poor outcome after out-of-hospital cardiac arrest.** *Crit Care* 2017, **21**(1):96.
30. Disanto G, Barro C, Benkert P, Naegelin Y, Schadelin S, Giardiello A, Zecca C, Blennow K, Zetterberg H, Leppert D *et al*: **Serum Neurofilament light: A biomarker of neuronal damage in multiple sclerosis.** *Annals of neurology* 2017, **81**(6):857-870.
31. Jennett B, Bond M: **Assessment of outcome after severe brain damage.** *Lancet* 1975, **1**(7905):480-484.
32. Quinn TJ, Dawson J, Walters MR, Lees KR: **Reliability of the modified Rankin Scale: a systematic review.** *Stroke* 2009, **40**(10):3393-3395.
33. Rittenberger JC, Raina K, Holm MB, Kim YJ, Callaway CW: **Association between Cerebral Performance Category, Modified Rankin Scale, and discharge disposition after cardiac arrest.** *Resuscitation* 2011, **82**(8):1036-1040.
34. Callaway CW, Donnino MW, Fink EL, Geocadin RG, Golan E, Kern KB, Leary M, Meurer WJ, Peberdy MA, Thompson TM *et al*: **Part 8: Post-Cardiac Arrest Care: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care.** *Circulation* 2015, **132**(18 Suppl 2):S465-482.
35. Nolan JP, Cariou A: **Post-resuscitation care: ERC-ESICM guidelines 2015.** *Intensive care medicine* 2015, **41**(12):2204-2206.
36. Luescher T, Mueller J, Isenschmid C, Kalt J, Rasiah R, Tondorf T, Gamp M, Becker C, Sutter R, Tisljar K *et al*: **Neuron-specific enolase (NSE) improves clinical risk scores for prediction of neurological outcome and death in cardiac arrest patients: Results from a prospective trial.** *Resuscitation* 2019, **142**:50-60.

Figures



Number at risk					
—————	NfL (pg/ml) <25	43	36	35	35
.....	NfL (pg/ml) 25-50	38	24	23	23
—————	NfL (pg/ml) 50-75	17	8	7	7
-----	NfL (pg/ml) >75	66	13	12	9

Figure 1

Kaplan Meier curves for 30d all-cause mortality based on the three different NfL cut-offs Data is presented as number (n) percentage (%); NfL: Neurofilament Light Chain.




Figure 2

Subgroup analysis of NfL for primary endpoint (neurological outcome) Data is presented as univariable odds ratio (OR) and 95% confidence interval (95% CI). NfL: Neurofilament Light Chain; TTM: Targeted Temperature Management, COPD: Chronic obstructive pulmonary disease.




Figure 3

Areas under the curve for combination of NfL with NSE in a subgroup of 131 patients Area under the curve (AUC) for NSE compared with the AUC NfL for 3.1) the primary endpoint neurological outcome measured with cerebral performance category (CPC), 3.2) the endpoint neurological outcome measured with the modified ranking scale (mRS), 3.3) endpoint neurological mortality; ROC, receiver operating characteristic; NfL: Neurofilament Light Chain; NSE: Neuron specific enolase.