Psychological and Emotional Impact Of COVID-19 Pandemic on People Living with Chronic Disease: HIV and Cancer

Emanuele Focà (✉ emanuele.foca@unibs.it)

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Chiara Fornari

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Stefania Arsuffi

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Maria Chiara Vetrano

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Stefano Calza

Unit of Biostatistics and Bioinformatics, Department of Molecular and Translational Medicine, University of Brescia, Brescia, Italy

Stefano Renzetti

Unit of Biostatistics and Bioinformatics, Department of Molecular and Translational Medicine, University of Brescia, Brescia, Italy

Silvia Copeta

Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, Medical Oncology, University of Brescia at ASST Spedali Civili, Brescia, Italy.

Alfredo Berruti

Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, Medical Oncology, University of Brescia at ASST Spedali Civili, Brescia, Italy.

Francesco Castelli

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Silvia Compostella

Unit of Infectious Diseases of ASST Spedali Civili Hospital, Brescia, Italy

Eugenia Quiros-Roldan

University Department of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili Hospital, Brescia, Italy

Research Article

Keywords: COVID-19, Psychological effects, Resilience, Chronic diseases

Posted Date: October 29th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-927188/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at AIDS and Behavior on March 6th, 2022. See the published version at <https://doi.org/10.1007/s10461-022-03638-0>.

Abstract

People living with chronic disease (PLWCD) are the frailest category, both for the risk of severe COVID-19 illness and for the impact on the care continuum. Aim of this study was to analyze coping strategies and resilience in people living with HIV (PLWH) compared to people living with oncological diseases (PLWOD) during COVID-19 pandemic. We administrated an anonymous questionnaire, which explored the emotional experience, the demographic factors linked to a COVID-19-related stress syndrome, the patient's perception about the adequacy of clinical undertaking from the hospital and the resilience. We analyzed 324 questionnaires. There were no significant differences in prevalence of psychological distress among the whole cohort; however, PLWOD were calmer, less troubled, and more serene than PLWH. Moreover, PLWH smoked more, ate more and gained more weight than PLWOD. Most patients didn't feel lonely and continued to take pleasure from their activities. No differences in resilience were found between the groups. In the whole cohort lower levels of resilience were found in patients that were unemployed, with history of psychological disorders and in those who experienced more feelings of anger, anxiety and concern. PLWCD had high level of resilience, preserved their well-being, and activated adaptive coping during the pandemic.

Key Points

People living with chronic disease preserved their psychological well-being and activated adaptive coping strategies during the period of the quarantine. Furthermore, high level of resilience among the patients were detected.

Introduction

SARS-Cov-2 was discovered in Wuhan, China, in December 2019 and in short time caused an international health emergency [1–3]. The World Health Organization (WHO) declared the outbreak of a pandemic on March 11, 2020 [1]. COVID-19 caused the health systems distress around the world [4].

Italy and in particular Lombardy region suffered from an increasing number of cases with a high rate of intra-hospital mortality [5]. In the first phase in March-April 2020 the Brescia Hospital, a large tertiary hospital with 15709 beds, was one of the referral Hub-Hospital according to the high number of hospitalized patients.

The COVID-19 pandemic changed the certainty, activated fear, and inducted feelings like loneliness and isolation [3, 6]. The containment measures limited the individual freedom [6] and the burden on mental health conditions increased [7–10]. People living with chronic disease (PLWCD) are the most frail social category, not only for the risk for severe COVID-19 illness but also for the impact of the changes in the self-management [6, 11, 12].

Furthermore, COVID-19 pandemic had the potential to destroy the continuum of care [13–16] and people's lives [6, 17]. People living with HIV (PLWH) and people living with oncological disease (PLWOD) represent a

vulnerable population that may potentially be at higher risk to have severe COVID-19 compared to general population [2, 16, 18, 19]. This risk in PLWCD is predicted on potential interactions between COVID-19, presence of immunodeficiency and the co-existence of comorbidities [2, 4, 18]. Moreover, the stress due to the above-mentioned factors and the containments measures, like lockdown, have had an impact on mental health conditions [8] both in general population and in chronic patients [6].

The resilience-related factors, such as the perceived social support, are protective factors for the management of the chronicity [20] and for mitigating health challenges and stressful situations [4]. Several studies conducted in PLWH showed that depression is associated with treatment failure [25, 26], lower CD4+ t-cell count [27, 28]; while anxiety is linked with disengagement from care [29, 30] and non-adherence to antiretroviral therapy [31, 32]. On the other hand, the resilience and coping strategies developed by PLWH may potentially be a protective factor compared to general population [16, 33]. Indeed, considering the past, they could have a deep knowledge of pandemics and their effects and know how managing life in this specific context and how to cohabit with the stigma [16, 34]. Moreover, such similarities between the COVID-19 and HIV pandemics are present: the absence of vaccination at the time of the study and the perception of isolation and fear [16, 33].

According to the syndemic framework, we considered the biopsychosocial perspective of health and disease to understand the effects of the pandemic in PLWCD [19]. A syndemic is defined as two or more epidemic events that interact synergically to produce an increased burden in a specific population [19, 35]. This perspective was mostly applied in PLWH [36]: the personal experience of COVID-19 pandemic is assumed to overlap with others health challenges due to HIV, including the disease itself, mental burden, and other infections [37].

Given that higher rates of mental health problems have been reported in general population in the setting of the pandemic [4], collecting psycho-social health data among PLWCD is important. PLWH may have higher score of resilience and that COVID-19 may generate a weaker psychological impact in the face of reduced social support [38, 39] and disadvantaged socio-economic conditions [19] than PLWOD.

The aim of the study is to describe the emotional experiences and the psychological characteristics (including resilience) of PLWCD, such as PLWH compared to PLWOD, during COVID-19 pandemic in order to gain clues for the medical staff and to make more effective the psychological, clinical and therapeutic interventions.

Results

Participant Characteristics and Socio-economic Aspects

A total of 600 adults (≥ 18 years old. 315 PLWH vs 285 PLWOD) were recruited for the study, after being admitted to the Hospital for the routine visits. 211 patients (35,2% of the total) refused to take part to the study: 143 patients with HIV (32,17% women) and 68 patients with oncological disease (42,65% women). Lack of time and will were the main causes of refusal. Furthermore, 65 (10,8%) questionnaires were

excluded as were incomplete. Overall, 324 (54% of the total) were eligible for analyses: 167 PLWH and 157 PLWOD. PLWH were significant younger compared to PLWOD and PLWH had a significant longest history of disease compared to PLWOD (Table 1).

Additionally, from the socio-economic point of view, 153 patients (47,2%) were employed while 103 patients (31,8%) were retired. A greater number of PLWH than PLWOD were employed, while a significant greater number of PLWOD were retired ($p < 0.001$). At last, PLWH were significant more worried than PLWOD about economics and financial problems due to the pandemic (32,9% PLWH *vs* 9,56% PLWOD, $p < 0.001$). (Table 2).

Almost 25% of total patients referred to have a history of psychological problems without significant differences between groups. Depression was the most widespread psychological disorder among the groups. Furthermore, the majority of the patients (97,2%) reported to respect the care continuum during the pandemic. (Table S1 in supplementary materials).

Table 1
Participant Characteristics

Recruitment: n (%)	PLWH (N=315)	PLWOD (N=285)	Total (N=600)	
Participants	167 (51.5%)	157 (48.5%)	324(54%)	
Refusal	143 (67.8%)	68 (32.2%)	211(35.2%)	
Interrupted	5 (7.7%)	60 (92.3%)	65(10.8%)	
Characteristics	Observed		Total (N=324)	p- value
	PLWH (N=167)	PLWOD (N=157)		
Age:				<.001*
Year: mean (sd)	51.0 (11.2)	61.6 (12.5)	56.1 (13.0)	
Sex: n (%)				< .001*
Male	114 (68.3%)	73 (46.5%)	187 (57.7%)	
Female	52 (31.1%)	84 (53.5%)	136 (42.0%)	
Others	1 (0.6%)	0 (0%)	1 (0.3%)	
Education:				
Year: mean (sd)	10.6 (3.9)	10.7 (4)	10.6 (3.9%)	.880
Level of education: n (%)				.048*
Primary School diploma	17 (10.2%)	26 (16.6%)	43 (13.3%)	
Middle School diploma	74 (44.3%)	49 (31.2%)	123 (38.0%)	
High School diploma	55 (32.9%)	65 (41.4%)	120 (37.0%)	
Degree	21 (12.6%)	17 (10.8%)	38 (11.7%)	
Nationality: n (%)				.084
Italian	156 (93.4%)	153 (97.5%)	309 (95.4%)	
Foreigner	11 (6.6%)	4 (2.5%)	15 (4.6%)	
Diagnosis: n (%)				< .001*
< 5 years	17 (10.2%)	116 (73.9%)	133 (41.0%)	
≥ 5 years	150 (89.8%)	41 (26.1%)	191 (59.0%)	
Cohabitation during the interview: n (%)				< .001*
Single	48 (28.7%)	20 (12.7%)	68 (21.0%)	
With someone	119 (71.3%)	137 (87.3%)	256 (79.0%)	

Recruitment: n (%)	PLWH (N=315)	PLWOD (N=285)	Total (N=600)
Cohabitation during the quarantine (first wave): n (%)			< .001*
Single	42 (25.2%)	17 (10.8%)	59 (18.2%)
With someone	125 (74.9%)	140 (89.2%)	265 (81.8%)
Note: * Significant values.			

Table 2
Socio-economic aspects

Characteristics	Observed		Total (N=324)	p- value
	PLWH (N=167)	PLWOD (N=157)		
Employment: n (%)				< .001*
Employed	103 (61.7%)	50 (31.8%)	153 (47.2%)	
Pensioners	24 (14.4%)	79 (50.3%)	103 (31.8%)	
Housewife	11 (6.6%)	17 (10.8%)	28 (8.6%)	
Unemployed	21 (12.6%)	4 (2.6%)	25 (7.7%)	
Other	8 (4.8%)	6 (3.8%)	14 (4.3%)	
Student	0 (0.0%)	1 (0.6%)	1 (0.3%)	
Habitation: n (%)				< .001*
Spacious house with garden	123 (73.7%)	146 (93.0%)	269 (83.0%)	
Three-room apartment	20 (12.0%)	9 (5.7%)	29 (9.0%)	
Two-room apartment	17 (10.2%)	2 (1.3%)	19 (5.9%)	
Studio apartment	7 (4.2%)	0 (0.0%)	7 (2.2%)	
Concern for economics problems: n (%)				< .001*
Not at all	83 (49.7%)	121 (77.1%)	204 (63.0%)	
A little	29 (17.4%)	21 (13.4%)	50 (15.4%)	
A lot	55 (32.9%)	15 (9.6%)	70 (21.6%)	
Note: * Significant values.				

Information about COVID-19

Patients' general knowledge about important aspects of COVID-19 pandemic was analyzed (Table 3). The majority of patients (87%) reported to be alert to the COVID-19-related symptoms, such as anosmia,

ageusia, fever, cough, fatigue, and dyspnoea (86,2% of PLWH vs 87,9% of PLWOD). Participants were aware about the recommended prevention methods. In particular, 150 patients (46,9%) sustained that facial masks are enough to protect from the virus while 139 patients (43,4%) reported that not only facial masks but also physical distancing and hand washing are necessary to protect from the infection.

Furthermore, the association between living with chronic disease and living during the pandemic was analyzed. When asked about living with chronic diseases while facing the risk of contracting COVID-19, most of the participants (47,7%) reported that they no longer consider themselves at higher risk than other people (53% of PLWH vs 42% of PLWOD). However, a significant greater number of PLWOD than PLWH thought to be at risk (25,9% of PLWH vs 40,8% of PLWOD, $p = 0.018$) and were worried about their clinical outcome (35,5% of PLWH and 52,3% of PLWOD, $p = 0.007$). Lastly, most of the participants (62,7%) would like to get the vaccination as soon as it was available (58,7% of PLWH vs 66,9% of PLWOD).

In sum, most of the patients were careful about the symptoms of COVID-19 and they understood the importance of protecting themselves and reducing the diffusion of the virus: PLWOD were in general more aware and worried about the outcome of the COVID-19 infection than PLWH.

The 97,2% of patients were aware of the progress of the pandemic (Table S2 in supplementary materials). Patients consulted the COVID-19-related information from different sources, the most used ones were television, internet, and social media. Most of the patients (71,9%) reported that information obtained by the doctors were useful, while information provided by politicians were confused and catastrophic. 161 patients (49,7%) checked the news every 3-5 hours and PLWOD consulted the COVID-19-related information significant more frequently than PLWH (42,5% of PLWH vs 57,3% of PLWOD, $p = 0.013$).

Table 3
General knowledge

Characteristics	Observed		Total (N=324)	p- value
	PLWH (N=167)	PLWOD (N=157)		
Pay attention to Covid-19 symptoms: n (%)				.052**
Not at all	5 (3.0%)	11 (7.0%)	16 (4.9%)	
A little	18 (10.8%)	8 (5.1%)	26 (8.1%)	
A lot	144 (86.2%)	138 (87.9%)	282 (87.0%)	
The facial masks are enough to protect yourself: n (%)				.421
N-Miss	2	2	4	
Not at all	19 (11.5%)	12 (7.7%)	31 (9.7%)	
A little	73 (44.2%)	66 (42.6%)	139 (43.4%)	
A lot	73 (44.2%)	77 (49.7%)	150 (46.9%)	
Concern of being at risk: n (%)				.018*
N-Miss	1	0	1	
Not at all	88 (53.0%)	66 (42.0%)	154 (47.7%)	
A little	35 (21.1%)	27 (17.2%)	62 (19.2%)	
A lot	43 (25.9%)	64 (40.8%)	107 (33.1%)	
Worsening of the chronic disease: n (%)				.007*
N-Miss	1	2	3	
Not at all	83 (50.0%)	53 (34.2%)	136 (42.4%)	
A little	24 (14.5%)	21 (13.5%)	45 (14.0%)	
A lot	59 (35.5%)	81 (52.3%)	140 (43.6%)	
Vaccination: n (%)				.253
As soon as possible	98 (58.7%)	105 (66.9%)	203 (62.7%)	
Wait to see the effects	48 (28.7%)	39 (24.8%)	87 (26.9%)	
Never	21 (12.6%)	13 (8.28%)	34 (10.5%)	
Note: * significant values. ** tendency to significant values.				

Social support and coping strategies

Social and family networks were analyzed to identify the support received (Table S3 in supplementary materials). The 81,2% of participants reported to be very helped by the family in the everyday life both for the practical, psychological and emotional aspects. They felt also protected to express their COVID-19-related concerns and thoughts to the family. Moreover, the majority of the participants (62%) reported not having feelings of solitude than usual during the quarantine. Indeed, during the pandemic, participants have managed to maintain good relationships with others, such as family and friends. Furthermore, the majority of the participants (73,6%) did not receive and search support from other patients with the same disease.

Three aspects of coping strategies were analyzed: the faith, the using of meditation techniques and the ability to take pleasure from their activities and hobbies. Only the 29,5% of patients received support by the faith (19,3% of PLWH *vs* 40,5% of PLWOD, $p < 0.001$). Most of the participants (80,5%) did not use meditation techniques while the 71,3% continued to take pleasure from their activities and hobbies. Patients reported that found new activities, such as cooking, when their previous hobbies were suspended.

Stress And Experiences

Emotional experiences of the participants were analyzed to understand the stress of the patients during the quarantine (Table 4). A greater number of PLWOD were significant calmer than PLWH ($p = 0.039$) and, on the other hand, a greater number of PLWH were more restless than PLWOD ($p = 0.014$). Overall, 254 patients (78,4%) referred to have lived peacefully (72,5% of PLWH *vs* 84,7% PLWOD, $p = 0.026$).

Regarding daily habits, a major part of the participants (65,3%) reported not to have changed the quality of sleep (59,6% of PLWH *vs* 71,4% PLWOD, $p = 0.086$). Changes in nutritional supply were also analyzed: 63 patients (19,4%) referred an increase in food intake (25,2% of PLWH *vs* 13,4% of PLWOD, $p = 0.003$). Notwithstanding, only the 22,2% of patients reported weight increase (28,7% of PLWH *vs* 15,3% of PLWOD, $p = 0.013$). Moreover, PLWH smoked significant more than PLWOD (10,8% of PLWH *vs* 1,3% of PLWOD, $p < 0.001$).

The suicidal thoughts and the self-harming ideas were also analyzed among participants. Most of the patients (90,7%) did not experience intrusive thoughts. The introspection ability among participants was analyzed and most of patients (53,9%) reported that the quarantine was a period of personal development.

Lastly, it was asked to the PLWH group if the COVID-19 may be comparable with the HIV pandemic. Most of the patients (31,8%) referred that the two situations were not comparable.

Table 4
Stress and experiences

Characteristics	Observed		Total (N=324)	p- value
	PLWH (N=167)	PLWOD (N=157)		
Feelings during the quarantine: n (%)				
Calm				.039*
Yes	82 (49.1%)	95 (60.5%)	147 (45.4%)	
No	85 (50.9%)	62 (39.5%)	177 (54.6%)	
Indifferent				.530
Yes	5 (3.0%)	3 (1.9%)	8 (2.5%)	
No	162 (97.0%)	154 (98.1%)	316 (97.5%)	
Sad				.207
Yes	21 (12.6%)	13 (8.3%)	34 (10.5%)	
No	146 (87.4%)	144 (91.7%)	290 (89.5%)	
Sense of guilty				.579
Yes	0 (0.0%)	0 (0.0%)	0 (0.0%)	
No	167 (100%)	157 (100%)	324 (100%)	
Powerless				.116
Yes	23 (13.8%)	13 (8.28%)	36 (11.1%)	
No	144 (86.2%)	144 (91.7%)	288 (88.9%)	
Angry				.887
Yes	10 (6.0%)	10 (6.4%)	20 (6.2%)	
No	157 (94.0%)	147 (93.6%)	304 (93.8%)	
Nerves				.435
Yes	12 (7.2%)	8 (5.1%)	20 (6.2%)	
No	155 (92.8%)	149 (94.9%)	304 (93.8%)	
Troubled				.014*
Yes	20 (12.0%)	7 (4.46%)	27 (8.33%)	
No	147 (88.0%)	150 (95.5%)	297 (91.7%)	
Worried				.196

Characteristics	Observed		Total (N=324)	p- value
Yes	80 (47.9%)	64 (40.8%)	144 (44.4%)	
No	87 (52.1%)	93 (59.3%)	180 (55.6%)	
Anxious				.209
Yes	26 (15.6%)	17 (10.8%)	43 (13.3%)	
No	141 (84.4%)	140 (89.2%)	281 (86.7%)	
Frightened				.659
Yes	38 (22.8%)	39 (24.8%)	77 (23.8%)	
No	129 (77.3%)	118 (75.2%)	247 (76.2%)	
Panic				.421
Yes	3 (1.8%)	5 (3.19%)	8 (2.5%)	
No	164 (98.2%)	152 (96.8%)	316 (97.5%)	
Isolated				.957
Yes	22 (13.2%)	21 (13.4%)	43 (13.3%)	
No	145 (86.8%)	136 (86.6%)	281 (86.7%)	
Living with serenity the situation: n (%)				.026*
Not at all	10 (5.9%)	6 (3.8%)	16 (4.9%)	
A little	36 (21.6%)	18 (11.5%)	54 (16.7%)	
A lot	121 (72.5%)	133 (84.7%)	254 (78.4%)	
Feeling confident: n (%)				.290
Not at all	26 (15.6%)	18 (11.5%)	44 (13.6%)	
A little	57 (34.1%)	47 (29.9%)	104 (32.1%)	
A lot	84 (50.3%)	92 (58.6%)	176 (54.3%)	
Feeling pessimist: n (%)				.853
Not at all	98 (58.7%)	94 (59.9%)	192 (59.3%)	
A little	51 (30.5%)	44 (28.0%)	95 (29.3%)	
A lot	18 (10.8%)	19 (12.1%)	37 (11.4%)	
Change the quality of sleep: n (%)				.086
N-Miss	1	0	1	

Characteristics	Observed		Total (N=324)	p- value
Not at all	99 (59.6%)	112 (71.4%)	211 (65.3%)	
A little	37 (22.3%)	24 (15.3%)	61 (18.9%)	
A lot	30 (18.1%)	21 (13.4%)	51 (15.8%)	
Eating more: n (%)				.003*
Not at all	88 (52.7%)	111 (70.7%)	199 (61.4%)	
A little	37 (22.2%)	25 (15.9%)	62 (19.1%)	
A lot	42 (25.2%)	21 (13.4%)	63 (19.4%)	
Weight changes: n (%)				.013*
No	106 (63.5%)	121 (77.1%)	227 (70.1%)	
Loss weight	13 (7.8%)	12 (7.6%)	25 (7.7%)	
Gain weight	48 (28.7%)	24 (15.3%)	72 (22.2%)	
Characteristics	Observed		Total (N=324)	p- value
Smoking: n (%)				< .001*
Not smoke	93 (55.7%)	133 (84.7%)	226 (69.8%)	
Not at all	44 (26.3%)	18 (11.5%)	62 (19.1%)	
A little	12 (7.2%)	4 (2.5%)	16 (4.9%)	
A lot	18 (10.8%)	2 (1.3%)	20 (6.2%)	
Alcohol: n (%)				.026*
Not drink	109 (65.3%)	116 (73.9%)	225 (69.4%)	
Not at all	44 (26.3%)	39 (24.8%)	83 (25.6%)	
A little	8 (4.8%)	1 (0.6%)	9 (2.8%)	
A lot	6 (3.6%)	1 (0.6%)	7 (2.2%)	
Drugs: n (%)				.011*
Not use drugs	155 (92.8%)	156 (99.4%)	311 (96.0%)	
Not at all	11 (6.6%)	1 (0.6%)	12 (3.7%)	
A little	1 (0.6%)	0 (0.0%)	1 (0.3%)	
A lot	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Mood medications: n (%)				.627

Characteristics	Observed		Total (N=324)	p- value
Not at all	152 (91.0%)	141 (89.8%)	293 (90.4%)	
A little	9 (5.4%)	12 (7.6%)	21 (6.5%)	
A lot	6 (3.6%)	4 (2.5%)	10 (3.1%)	
Suicidal thoughts: n (%)				.198
Not at all	156 (93.4%)	138 (87.9%)	294 (90.7%)	
A little	7 (4.2%)	14 (8.9%)	21 (6.5%)	
A lot	4 (2.4%)	5 (3.2%)	9 (2.8%)	
Psychological support: n (%)				.586
Yes	23 (13.8%)	25 (15.9%)	48 (14.8%)	
No	144 (86.2%)	132 (84.1%)	276 (85.2%)	
Others were more scarred than the patients: n (%)				.840
N-Miss	2	2	4	
Not at all	35 (21.2%)	37 (23.9%)	72 (22.5%)	
A little	42 (25.5%)	37 (23.9%)	79 (24.7%)	
A lot	88 (53.3%)	81 (52.3%)	169 (52.8%)	
Period of reflection: n (%)				.104
N-Miss	0	1	1	
Not at all	40 (24.0%)	52 (33.3%)	92 (28.5%)	
A little	28 (16.8%)	29 (18.6%)	57 (17.6%)	
A lot	99 (59.3%)	75 (48.1%)	174 (53.9%)	
Covid-19 pandemic vs HIV pandemic: n (%)				
Not at all	103 (31.8%)	-		
A little	30 (9.3%)	-		
A lot	32 (9.9%)	-		
Note: * Significant values.				

Experience in hospital ward

PLWH have been using the hospital services for significant longer time than PLWOD (15,8 vs 9,65 years, $p < 0.001$). Furthermore, most of participants reported that the hygiene measures against COVID-19 adopted

by the Hospital were adequate (95.9%) and reported to feel protected from the infection in the clinic of origin both during the previous visits (96%) and at the moment of the interview (97,2%).

Regarding relationship with healthcare providers, 247 (77,7%) patients were not afraid about the possibility that the healthcare providers may be infected by COVID-19 and could not be reference points anymore (79,8% of PLWH *vs* 75,5%, NS). Moreover, 277 (86%) patients were not worried about the possibility that the medical staff could infect patients themselves. The majority of patients reported they would not have wanted to be informed about the sanitary situation of the ward before the visit (86,7%) and they felt supported by the healthcare providers during the COVID-19 pandemic (78,3%). Most of patients (90,3%) did not feel neglected by the sanitary system and they reported to be assisted as usual (Table S4 in the supplement materials).

Resilience

The resilience of the participants was investigated to study the differences between the groups in the capability to manage the critical psychological issue during the pandemic (Table S5 in the supplement materials). Participants had in general high level of resilience measured by RS (RS mean $\pm$ standard deviation: 59,7 $\pm$ 6,92). Regarding the total RS score, no significant differences were found between the groups (RS score: 59,7 in PLWH *vs* 59,6 in PLWOD) except in the item "I have enough energy to do what I should do" (RS10, $p = 0.002$) (Figure 1, item 10).

Linear Regression analysis

From the original 44 variables collected through the questionnaire, factor analysis allowed to identify latent factors defined based on the loading (Figure S1) as feeling of anger, need of attention, concern of the infection, psychological problems, anxiety/concern, coping and lockdown conditions. All the factors were included in the linear regression to test their effect on RS.

No significant differences were found between the groups in the Resilience Score ($p = 0.410$). However, regardless of chronic disease displayed a lower level of resilience who was unemployed (-3.52; 95% CI [-6.68 -0.35]; $p = 0.029$); who showed more anger feelings (-1.76; 95% CI [-2.67 -0.85]; $p < 0.001$), who showed more anxiety/concern (-1.56; 95% CI [-2.30 -0.82]; $p < 0.001$) and who had a history of psychological problems during the lockdown (-2.18; 95% CI [-3.17 -1.18]; $p < 0.001$). (Table S6 in the supplement materials).

Discussion

This study evaluates the emotional experiences, the psychological effects, and the resilience among PLWCD, comparing PLWH to PLWOD, during the pandemic of SARS-CoV-2 in a centre of North Italy.

Here, unlike previous studies [16, 45], in general we found that pandemic has not modified the psychological well-being of the participants. Patients in general underestimated the risk of belonging to a

frail category and the majority of responders were optimistic about the evolution of the emergency, also with the hope of vaccination. However, PLWOD were significantly more calm, less troubled, and more serene than PLWH, despite oncological patients were more worried than PLWH about the clinical outcomes of the COVID-19 infection. In fact, PLWH sustained that the ART was a protective factor for the infection, as initially published [44]. The sense of loneliness and isolation were mitigated by the possibility of using the new technologies [46], such as smartphone and online platforms of meeting. Noteworthy, a greater number of PLWH than PLWOD lived alone during the quarantine, but this aspect did not change the outcomes. In general, all the participants referred to be supported by the healthcare providers, put their trust in them and felt safe to go to the Hospital for their visits: hygienic measures and organization were adequate. These aspects probably allowed to maintain the adherence to the therapies and to the continuum of care, contrary to what was found in a previous study in HIV patients [45]. Our result may be explained by the fact that the care continuum in the Hospital during the lockdown was also guaranteed through telemedicine, which was a useful tool to preserve the care involvement [47]. We also explored the parallelisms between the COVID-19 and HIV pandemics and patients reported that the stigma due to the HIV positivity was not comparable at all with the COVID-19 positivity.

The quarantine did not upset drastically the habits of the participants, unlike a previous study [16]. Our results suggest that PLWH [16] and PLWOD were able to adapt themselves and their behaviours to the situation, indeed patients continued to take pleasure from their daily activities as usual and active coping strategies, such as passing time with new activities. These aspects were protective factors for the management of the chronic diseases and played an important role in the outcome of the patient's burden [20]. Some participants (PLWH > PLWOD) referred to eat more and therefore to gain weight, this is maybe linked to the reduction of physical activity and to the increase of cooking activities. Few differences in the habits were detected between the groups: PLWH drank alcohol and smoked cigarettes more than PLWOD, slightly in contrast to other studies [16]. It is important to consider that for PLWOD could be very difficult to differentiate the effects in the daily habits due to the therapies and due to COVID-19. Lastly, patients kept informed (PLWOD > PLWH) about the COVID-19 disease and understood the importance of the individual protection, such as facial masks and physical distancing, and paid attention to the COVID-19 symptoms. The period of the quarantine was an opportunity of reflection. Also, patients reported that the information obtained from the health care providers were preferred than the politicians' ones, in line with others [16].

Patients showed high level of resilience, with a ceiling effect in the RS score, and no significant differences were found between groups. Unlike a previous study [45], the participants' age did not influence the resilience score. In our study emerged that those who lived in a disadvantaged situation, those who had a history of psychological disorders and patients that felt negative emotions, such as angry, anxiety and concern, during the lockdown had lower values in the RS independently by the group.

Our study has some strength. To our knowledge, first: little is known about the emotional experience and the psychological effects of frailty patients that lived with an oncological disease during the pandemic and, second: we compared the emotional and psychological outcomes due to the pandemic in two different frailty chronic population. The study describes the importance of positive psychology in the

management of the critical situations and sheds a new light on resilience and coping strategies during acute crisis in patients that live with the chronicity. In particular, the presence of adaptive coping and high level of resilience in PLWCD could reduce drastically the burden due to the pandemic.

Some limitations need to be acknowledged. First, our work does not guarantee the generalisability of the results because the Italian situation cannot be compared to the others, in particular the work was conducted in a single centre included in service-rich city and with a free national health and welfare system. Second, we did not include a control group composed of healthy people. Additionally, the questionnaires were conducted in presence of a healthcare provider which raises potential response bias in the interviewed. Moreover, we did not collect data on disease stage in cancer patients. Since the majority of PLWOD was following a day-hospital care regime for prevention of recurrence, the underestimation of belonging to a frail category in PLWOD could be linked with the evolution of the disease. Last concern is the difference in the demographic characteristics and cultural factors [4] between the groups, such as the different number of foreigner people and age differences.

Methods

This observational study was conducted from December 16, 2020 to February 19, 2021 at the HIV and oncological outpatient clinics of ASST Spedali Civili Hospital, Brescia, Italy. The study involved the administration of an *ad hoc* descriptive anonymous questionnaire and a resilience scale (RS) [40]. The questionnaire was dealt with as a structured interview to increase the quality of data and the accessibility. The compilation requires about 20 minutes. The questionnaire and the RS were proposed to the patients during one of the routine medical contacts at the Spedali Civili Hospital.

This study was approved by Ethical Board of Brescia Province (procedure: NP 4364). It was conducted in accordance with the guidelines and standards of the Declaration of Helsinki (2013) and with the principles of Good Clinical Practice. The participation was voluntary and no written informed consent was needed. All the participants gave a verbal informed consent, participants that denied the consent were excluded.

The Questionnaire

We have constructed a questionnaire (Supplemental Material) aimed to explore: the emotional experience and the stress due to the sanitary emergency of COVID-19 in PLWH and PLWOD; the patient's perception about the adequacy of the clinical and psychological undertaking from the hospital during the COVID-19 pandemic; the clinical and demographic factors linked with a higher level of COVID-19 related stress (i.e. age, gender, disease).

The questionnaire was subdivided in clusters of questions: socio-demographic information and history, patients' family network and perceived support, feelings and experiences during the quarantine, patients' experience in their respective hospital clinic, management of the disease during the lockdown, patient's opinions about Coronavirus and sources of information used. The compilation of the questionnaire

requires about 15 minutes, and it includes both dichotomous questions which only have two possible responses, such as “Yes” or “No”, closed-ended questions and 3-points Likert-type scale (1 = not at all; 2 = a little; 3 = a lot).

The Resilience Scale

Resilience has been defined as the personal dynamic characteristic that moderates the negative effects of stress, that promotes the positive adaptation and effective coping strategies [41, 42]. The Italian version of the RS [43] (Supplemental Material) was administered to identify the capability and effective resources of patients in order to manage the critical psychological issue during the pandemic. The compilation of the RS requires about 5 minutes.

Statistical analysis

Continuous variables were reported as mean with standard deviation (mean $\pm$ SD) and were compared across the groups using the nonparametric Kruskal-Wallis test. Categorical variables were summarized through frequencies and percentages and were compared across the groups using the Chi-squared test. The Exploratory Factor Analysis (EFA) was computed to investigate the latent variables underlying the questions of the questionnaire regarding the emotional experiences and the life quality during the lockdown. The factor loading threshold cut-off value was set at $|0.32|$. The factors that emerged from the EFA and demographic data were included in a linear regression model with the score obtained from the RS. Statistical significance level was set at $\alpha = 0.05$; all statistical tests were two-tailed and 95% CI were computed for linear regression parameters. Statistical analysis was performed with R (version 4.0.2).

The study was approved by the Ethics Committee of University of Brescia and was conducted in accordance with the Declaration of Helsinki (2013) and with the principles of Good Clinical Practice. All participants gave a verbal consent, participants that gave not the consent were excluded.

Conclusion

Our results showed no significant psychological effects due to the pandemic, however PLWOD were more calm, less troubled, and more serene than PLWH. Few changes in the daily habits in the PLWH were found compared to PLWOD and PLWH showed higher distress than PLWOD according to comparable resilience score. High levels of social support and resilience were recorded in both groups. Last, patients with psychological problems, feelings of anger and concern deserve more attention as they show lower levels of resilience. Further investigations are needed to confirm this finding and to compare PLWCD with the general population, also considering age differences.

Availability of data and material Dataset generated and analysed during the current study are available in the Google Drive repository.

Declarations

Acknowledgements We express our gratitude to all the participants who took time to respond to our questionnaire.

Author Contributions Conceptualization: E.Q., S.C.; Methodology: S.C.; Data Collection: C.F., CM.V.; Formal analysis: S.R., S.C.; Supervision: E.F., S.C., F.C., A.B., S.C., E.Q.; Writing (original draft preparation): C.F.; Writing (review and editing): E.F., S.A., E.Q., A.B., S.R., F.C., C.F. All authors read and approved the final manuscript.

Additional Information

Funding This study was supported by “Gilead Fellowship program 2019” founded by Gilead Sciences Italy. Gilead Sciences had no role in the study design, statistical analyses, and data interpretation.

<https://www.fellowshipgileaditalia.it/fellowship-program>

Conflicts of Interest EF received speakers’ honoraria, research grants and advisory board fees from Viiv Healthcare, Janssen-Cilag, Gilead Sciences and Merck Sharp & Dohme. The remaining authors declare that they have no competing interests.

Availability of data and material Dataset generated and analysed during the current study are available in the Google Drive repository.

https://docs.google.com/spreadsheets/d/1vPAeR2CRaQXFmcK_oB23MsRY1o9AkzTB6HLeVyTbX8c/edit?usp=sharing

References

1. Anand KB, Karade S, Sen S, Gupta RM. SARS-CoV-2: Camazotz's Curse. *Med J Armed Forces India*. **76**(2):136-141 (2020).
2. Siewe Fodjo JN, et al. Impact of the COVID-19 Pandemic on the Medical Follow-up and Psychosocial Well-Being of People Living With HIV: A Cross-Sectional Survey. *J Acquir Immune Defic Syndr*. **85**(3), 257-262 (2020).
3. Wang C, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun*. **87**, 40-48 (2020).
4. Jones DL, et al. Mental Health, Coping, and Social Support Among People Living with HIV in the Americas: A Comparative Study Between Argentina and the USA During the SARS-CoV-2 Pandemic. *AIDS Behav*. **25**, 1–9 (2021).

5. Rizzi M, Castelli F, Latronico N, Focá E. SARS-CoV-2 invades the West. How to face a COVID-19 epidemic in Lombardy, Northern Italy? *Infez Med. Ahead of print Jun 1.* **28** (2), 133-134 (2020).
6. Xiong J, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord.* **277**, 55-64 (2020).
7. González-Sanguino C, et al. Mental health consequences during the initial stage of the 2020 Coronavirus pandemic (COVID-19) in Spain. *Brain Behav Immun.* **87**, 172-176 (2020).
8. Marbaniang I, et al. The burden of anxiety among people living with HIV during the COVID-19 pandemic in Pune, India. *BMC Public Health.* **20**(1), 1598 (2020).
9. Mazza C, et al. A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int J Environ Res Public Health.* **17**(9), 3165 (2020).
10. Li J, et al. Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. *World Psychiatry.* **19**(2), 249-250 (2020).
11. Mi T, et al. Mental Health Problems of HIV Healthcare Providers During the COVID-19 Pandemic: The Interactive Effects of Stressors and Coping. *AIDS Behav.* **25**(1), 18-27 (2021).
12. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med.* **383**(6), 510-512 (2020).
13. Budak JZ, Scott JD, Dhanireddy S, Wood BR. The Impact of COVID-19 on HIV Care Provided via Telemedicine-Past, Present, and Future. *Curr HIV/AIDS Rep.* **18**(2), 98-104 (2021).
14. Jiang H, Zhou Y, Tang W. Maintaining HIV care during the COVID-19 pandemic. *Lancet HIV.* **7**(5), e308-e309 (2020).
15. Pinto RM, Park S. COVID-19 Pandemic Disrupts HIV Continuum of Care and Prevention: Implications for Research and Practice Concerning Community-Based Organizations and Frontline Providers. *AIDS Behav.* **24**(9), 2486-2489 (2020).
16. Gwadz M, et al. Black and Latino Persons Living with HIV Evidence Risk and Resilience in the Context of COVID-19: A Mixed-Methods Study of the Early Phase of the Pandemic. *AIDS Behav.* **25**(5), 1340-1360 (2021).
17. Sun S, et al. Challenges to HIV Care and Psychological Health During the COVID-19 Pandemic Among People Living with HIV in China. *AIDS Behav.* **24**(10), 2764-2765 (2020).
18. Al-Quteimat OM, Amer AM. The Impact of the COVID-19 Pandemic on Cancer Patients. *Am J Clin Oncol.* **43**(6), 452-455 (2020).
19. Shiau S, Krause KD, Valera P, Swaminathan S, Halkitis PN. The Burden of COVID-19 in People Living with HIV: A Syndemic Perspective. *AIDS Behav.* **24**(8), 2244-2249 (2020).
20. Schreurs KM, de Ridder DT. Integration of coping and social support perspectives: implications for the study of adaptation to chronic diseases. *Clin Psychol Rev.* **17**(1), 89-112 (1997).
21. Marhefka S, Lockhart E, Turner D. Achieve Research Continuity During Social Distancing by Rapidly Implementing Individual and Group Videoconferencing with Participants: Key Considerations, Best Practices, and Protocols. *AIDS Behav.* **24**(7), 1983-1989 (2020).

22. Rockwell KL, Gilroy AS. Incorporating telemedicine as part of COVID-19 outbreak response systems. *Am J Manag Care*. **26**(4), 147-148 (2020).
23. Saag MS, et al. Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2018 Recommendations of the International Antiviral Society-USA Panel. *JAMA*. **320**(4), 379-396 (2018).
24. Mugavero MJ, Amico KR, Horn T, Thompson MA. The state of engagement in HIV care in the United States: from cascade to continuum to control. *Clin Infect Dis*. **57**(8), 1164-1171 (2013).
25. Krumme AA, et al. Depression, adherence and attrition from care in HIV-infected adults receiving antiretroviral therapy. *J Epidemiol Community Health*. **69**(3), 284-289 (2015).
26. Marconi VC, et al. Early warning indicators for first-line virologic failure independent of adherence measures in a South African urban clinic. *AIDS Patient Care STDS*. **27**(12), 657-668 (2013).
27. Alemu H, Haile Mariam D, Tsui A, Ahmed S, Shewamare A. Effect of depressive symptoms and social support on weight and CD4 count increase at HIV clinic in Ethiopia. *AIDS Care*. **24**(7), 866-876 (2012).
28. Wagner GJ, Ghosh-Dastidar B, Holloway IW, Kityo C, Mugenyi P. Depression in the pathway of HIV antiretroviral effects on sexual risk behavior among patients in Uganda. *AIDS Behav*. **16**(7), 1862-1869 (2012).
29. Rooks-Peck CR, et al. Mental health and retention in HIV care: A systematic review and meta-analysis. *Health Psychol*. **37**(6), 574-585 (2018).
30. Gonzalez A, Solomon SE, Zvolensky MJ, Miller CT. The interaction of mindful-based attention and awareness and disengagement coping with HIV/AIDS-related stigma in regard to concurrent anxiety and depressive symptoms among adults with HIV/AIDS. *J Health Psychol*. **14**(3), 403-413 (2009).
31. Chakraborty A, Hershow RC, Qato DM, Stayner L, Dworkin MS. Adherence to Antiretroviral Therapy Among HIV Patients in India: A Systematic Review and Meta-analysis. *AIDS Behav*. **24**(7), 2130-2148 (2020).
32. Wykowski J, Kemp CG, Velloza J, Rao D, Drain PK. Associations Between Anxiety and Adherence to Antiretroviral Medications in Low- and Middle-Income Countries: A Systematic Review and Meta-analysis. *AIDS Behav*. **23**(8), 2059-2071 (2019).
33. McClurg L. 'They know what a pandemic is': HIV survivors see similarities to AIDS epidemic. <https://www.npr.org/2020/08/03/898581035/they-know-what-a-pandemic-is-hiv-survivors-see-similarities-to-aids-epidemic>. (Accessed 1 April 2021).
34. Hargreaves J, Davey C. Group for lessons from pandemic HIV prevention for the COVID-19 response. Three lessons for the COVID-19 response from pandemic HIV. *Lancet HIV*. **7**(5), e309-e311 (2020).
35. Singer M. Pathogen-pathogen interaction: a syndemic model of complex biosocial processes in disease. *Virulence*. **1**(1), 10-18 (2010).
36. Halkitis, P. N., et al. Evidence for a syndemic in aging HIV-positive gay, bisexual, and other MSM: implications for a holistic approach to prevention and health care. *Annals of Anthropological Practice*. **36**(2), 365-386 (2012).
37. Halkitis PN, et al. Measurement model exploring a syndemic in emerging adult gay and bisexual men. *AIDS Behav*. **17**(2), 662-673 (2013).

38. Halkitis PN, Krause KD, Vieira DL. Mental Health, Psychosocial Challenges and Resilience in Older Adults Living with HIV. *Interdiscip Top Gerontol Geriatr.* **42**, 187-203 (2017).
39. Shippy RA, Karpiak SE. The aging HIV/AIDS population: fragile social networks. *Aging Ment Health.* **9**(3), 246-254 (2005).
40. Wagnild, G. M., & Young, H. M. Development and psychometric. *Journal of nursing measurement.* **1**(2), 165-17847 (1993).
41. Allen, R. S., Haley, P. P., Harris, G. M., Fowler, S. N., & Pruthi, R. Resilience: Definitions, ambiguities, and applications. *Resilience in aging*, 1-13 (2011).
42. Luthar SS, Cicchetti D, Becker B. The construct of resilience: a critical evaluation and guidelines for future work. *Child Dev.* **71**(3), 543-562 (2000).
43. Peveri, L. (2010). Resilienza e Regolazione delle Emozioni. Un Approccio Multimodale. Available online at: http://boa.unimib.it/bitstream/10281/7893/3/phd_unimib_707899.pdf
44. Del Amo J, et al. Incidence and Severity of COVID-19 in HIV-Positive Persons Receiving Antiretroviral Therapy: A Cohort Study. *Ann Intern Med.* **173**(7), 536-541 (2020).
45. Nguyen AL, et al. Living With HIV During the COVID-19 Pandemic: Impacts for Older Adults in Palm Springs, California. *AIDS Educ Prev.* **33**(4), 265-275 (2021).
46. Brown B, Taylor J, Fisher CB. Mitigating Isolation of People Aging With HIV During the COVID-19 Pandemic. *Public Health Rep.* **136**(4), 394-396 (2021).
47. Quiros-Roldan E, Magro P, Carriero C et al. Consequences of the COVID-19 pandemic on the continuum of care in a cohort of people living with HIV followed in a single center of Northern Italy. *AIDS Res Ther.* **17**(1), 59 (2020).

Figures

Figure 1

The radar plot of the means of each item in the Resilience Scale (RS). Each vertex represents each item of the RS.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FocFornarietalsupplementaryinformationSI.docx](#)