

# WITHDRAWN: lncRNA HAND2-AS1 is Down Regulated in Osteoarthritis and Regulates IL-6 Expression in Chondrocyte

**Zhenxing Si**

Second Affiliated Hospital of Harbin Medical University

**Shifeng Zhou**

First Affiliated Hospital of Harbin Medical University

**Zilong Shen**

The Second Affiliated Hospital of Haebin Medical University

**Feiyu Luan**

First Affiliated Hospital of Harbin Medical College

**Jinglong Yan**

[yanjinglong01@126.com](mailto:yanjinglong01@126.com)

Second Affiliated Hospital of Harbin Medical University <https://orcid.org/0000-0003-1583-5441>

---

## Research article

**Keywords:** Osteoarthritis, synovial fluid, lncRNA HAND2-AS1, Interleukin 6, chondrocytes

**Posted Date:** October 21st, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-93180/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Journal of Orthopaedic Surgery and Research on January 19th, 2021. See the published version at <https://doi.org/10.1186/s13018-021-02216-9>.

EDITORIAL NOTE:

The full text of this preprint has been withdrawn by the authors while they make corrections to the work. Therefore, the authors do not wish this work to be cited as a reference. Questions should be directed to the corresponding author.

# Abstract

**Background:** Osteoarthritis (OA) is a leading cause of disability. The incidence of OA is progressively rising due to the diminishing levels of physical activity and ever-expanding ageing population. However, the mainstay for OA treatment only can improve symptoms without delay the progression of this severe disease. This study aimed to explore the biological role and clinical function of lncRNA HAND2-AS1 in OA.

**Methods:** Blood samples and synovial fluid were collected from OA patients and normal subjects. HAND2-AS1 expression was detected by qRT-PCR and IL-6 expression was detected by ELISA; The plasma levels of HAND2-AS1 were also detected in different ages, stages and gender of OA patients and controls. Furthermore, the ROC curve was used to analyze whether HAND2-AS1 can distinguish OA patients from normal subjects. Also, Pearson correlation coefficient analysis was used to analyze the correlation between lncRNA HAND2-AS1 and IL-6. In addition, western blot was used to detect the IL-6 level upon HAND2-AS1 overexpression in chondrocytes and qRT-PCR was used to detect the HAND2-AS1 level after endogenous IL-6 treatment.

**Results:** HAND2-AS1 and IL-6 were dysregulated in plasma and synovial fluid of OA patients. The expression of HAND2-AS1 in plasma of OA patients was decreased with aging and progression. Furthermore, HAND2-AS1 downregulation effectively distinguished OA patients from the healthy controls. Overexpression of HAND2-AS1 inhibited IL-6 expression in chondrocytes, while treatment with exogenous IL-6 did not affect HAND2-AS1 expression.

**Conclusion:** HAND2-AS1 effectively distinguished OA patients from the healthy controls and regulates IL-6 expression in human chondrocytes.

Abbreviations: Long non-coding RNA, lncRNA; Osteoarthritis, OA; Interleukin 6, IL-6; Negative control plasmid, NC; Real-time quantitative polymerase chain reaction, qRT-PCR; Enzyme-linked immunosorbent assay, ELISA; Receiver operating characteristic curve, ROC;

**Trial registration:** ChiCTR, ChiCTR2000038635. Registered 11 February 2019, [http://www.chictr.org.cn/ ChiCTR2000038635](http://www.chictr.org.cn/ChiCTR2000038635)

## Introduction

Osteoarthritis (OA) is the most prevalent joint disorder and affects most commonly the knees, fingers, spine and hips. The hallmark of OA is progressive degradation of articular cartilage, synovitis, joint pain and alterations in periarticular tissues and subchondral bone.<sup>1,2</sup> The most common symptom that OA patients first present by far is pain, which is also the predominant symptom that leads to disability. Pain and loss of functional capacity are accompanied by an increased risk of additional comorbid conditions such as cardiovascular disease, diabetes or cancer.<sup>3,4</sup> Many risk factors such as an increase in life expectancy, obesity and population ageing have contributed to the increase in the incidence of OA.<sup>5</sup> The

mainstay for OA treatment includes pharmacological and nonpharmacological to improve function and reduce pain. However, these treatments for OA are directed at symptomatic response and do not affect structural alterations.<sup>6,7,8,9</sup> Although much effort has been devoted to developing new treatments for OA and promising results have been obtained from OA animal models, rare agents have completed clinical trials. Therefore, it's an urgent goal to elucidate the underlying mechanisms.<sup>10,11,12</sup>

The mechanism of OA is associated with acute and chronic inflammatory response and inflammation is commonly existed in OA patients' body. As an inflammatory disease, the OA progression and development are always accompanied by altered levels of inflammatory factors.<sup>5,13,14</sup> Interleukin 6 (IL-6) is a proinflammatory cytokine, which is upregulated frequently in OA patients and osteoarthritis animal models.<sup>15,16,17,18</sup> HAND2-AS1 is a newly defined lncRNA and has been proved play an important role in different types of tumors, such as prostate cancer,<sup>19</sup> ovarian carcinoma,<sup>20</sup> cervical cancer,<sup>21</sup> intraductal papillary mucinous neoplasm,<sup>22</sup> colorectal cancer,<sup>23</sup> liver cancer<sup>24</sup> and liver cancer stem cell.<sup>25</sup> However, the biological role and clinical function of HAND2-AS1 in OA are unclear. Here, we found that HAND2-AS1 is downregulated in OA, which is correlated to age and the progression stages of OA. Further, HAND2-AS1 may act as a biomarker that can effectively distinguish OA patients from normal subjects. What's more, the role of HAND2-AS1 in OA is likely achieved through the inhibition of IL-6. Our surprising findings provide a novel insight into how HAND2-AS1 regulates OA progression. And thus, HAND2-AS1 could play a critical role in the pathogenesis of OA.

## Materials And Methods

### Plasma specimens, synovial fluid, and cell line

Blood samples were collected from 67 clinically confirmed OA patients and 34 normal subjects who were admitted by the Ningxia Medical University from February 2019 to February 2020. Inclusion criteria for patients: 1) diagnosis of knee osteoarthritis; 2) All patients signed the informed consent forms. Exclusion criteria for patients: 1) Patients diagnosed with other clinical diseases; 2) Any treatment received within 3 months before admission; 3) Excluding the influence of other inflammatory diseases, such as pharyngitis, colds, etc; 4) Patients with other types of arthritis should be excluded.

To separate plasma, all blood samples were centrifuged at 1200g for 15 minutes in EDTA tubes at room temperature. Synovial fluid samples were collected from 21 OA patients and 11 normal subjects. The 67 patients with OA included 31 females and 36 males, ranging in age from 34 to 69 years (average  $46.5 \pm 4.2$  years). The 34 healthy individuals included 14 females and 20 males, ranging in age from 32 to 68 years old (average  $45.9 \pm 4.7$  years old). This study has passed the review of the Ethics Committee of the First Affiliated Hospital of Harbin Medical University. Human chondrocyte cell line was purchased from ATCC (Manassas) to perform in vitro cell experiments. Cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplied with 10% fetal bovine serum and 0.1 mg/mL G-418 and the culture conditions were 37°C and 5% CO<sub>2</sub>.

## Cell transfection

lncRNA HAND2-AS1 over-expression plasmid (HAND2-AS1) and negative control plasmid (NC) were designed and constructed by Sangon (China). Lipofectamine 2000 reagent (CA) was used to transiently transfect the human chondrocyte cell line.

## Western-blot analysis

Cells total proteins were extracted with radioimmunoprecipitation assay solution (Thermo Fisher Scientific). Gel electrophoresis was performed using 10% sodium dodecyl sulfate polyacrylamide and then gel transfer was performed with polyvinylidene difluoride membranes. The membranes were blocked for 1 hour in 5% nonfat milk at room temperature and then membranes were incubated with primary antibodies for 1 hour including IL-6 (1: 1600; Abcam, UK),  $\beta$ -actin (1: 4000; Abcam). Finally, the membranes were incubated with secondary antibody (1:1000, Beyotime, China) for 1 hour at room temperature. Signals were collected using ECL (Sigma-Aldrich) and Image J software was used for all data normalizations.

## Real-time quantitative polymerase chain reaction (qRT-PCR)

Total RNA was extracted using Trizol (Thermo Fisher Scientific) according to the instruction. Then reverse transcription was performed with superScript III Reverse Transcriptase (Thermo Fisher Scientific) and PCR reaction systems were prepared with SYBR Green Quantitative RT-qPCR Kit (Roche) according to the instructions. All primers used in this study were purchased from Sangon and as follows:

HAND2-AS1 Forward: Forward: 5'- GGAGTCACAGGCAGTCGTAGA -3',

HAND2-AS1 Reverse: 5'- GAAGGCACAGATCATTTCATGG -3';

$\beta$ -actin Forward: 5'-TTCCAGCCTTCCTTCCTGGG-3',

$\beta$ -actin Reverse: 5'-TTGCGCTCAGGAGGAGCAAT-3'.

Enzyme-linked immunosorbent assay (ELISA)

IL-6 Human ELISA Kit (Thermo Fisher Scientific) was used to detect the IL-6 level in plasma according to the instructions.

## Statistical analysis

Data were obtained from three biological replicates and presented as the mean  $\pm$  standard deviation. Receiver operating characteristic (ROC) curve was used to explore the diagnostic values of plasma lncRNA HAND2-AS1 for osteoarthritis, in which true positive cases were osteoarthritis patients and true negative cases were the healthy individuals. Pearson's correlation coefficient was used to analyze the correlations between HAND2-AS1 and IL-6. Differences in comparisons were analyzed using Student's t

test. Three or more group means by one- or two-way ANOVA and the significance was determined at  $*P < 0.05$ .

## Results

### **lncRNA HAND2-AS1 and IL-6 were dysregulated in plasma and synovial fluid of OA patients.**

Blood samples were collected from 67 clinically confirmed OA patients and 34 normal subjects, and the plasma was separated by centrifugation. The expression of lncRNA HAND2-AS1 in plasma was detected by qRT-PCR. As shown in Fig.1A, expression levels of plasma lncRNA HAND2-AS1 were significantly lower in OA patients comparing to the control group. The IL-6 level in plasma was detected by ELISA and the results showed that plasma levels of IL-6 were significantly higher in OA patients comparing to the control group (Fig.1B). Synovial fluid was collected from 21 clinically confirmed OA patients and 11 normal subjects and the expression of lncRNA HAND2-AS1 in synovial fluid was detected by qRT-PCR; The expression of IL-6 in synovial fluid was detected by ELISA. lncRNA HAND2-AS1 levels in synovial fluid were also significantly lower in OA patients comparing to healthy controls (Fig.1C) and IL-6 levels in synovial fluid were significantly higher in OA patients comparing to the control group (Fig.1D).

### **The expression of HAND2-AS1 in plasma of OA patients was decreased with aging.**

To investigate whether the expression of HAND2-AS1 altered with aging, OA patients and healthy controls were divided into three groups according to ages: 40-50-year-old group, 50-60-year-old group and 60-70-year-old group. The expression of HAND2-AS1 in plasma of these groups was detected by qRT-PCR. The results showed that the expression of HAND2-AS1 in the 50-60-year-old group was significantly lower than that in the 40-50-year-old group, while the expression of HAND2-AS1 in the 60-70-year-old group was significantly lower than that in the 50-60-year-old group in OA patients (Tab.1). However, in healthy control groups, no significant changes were observed in 40-50-year-old group, 50-60-year-old group and 60-70-year-old group (Tab.1).

### **The expression of HAND2-AS1 in plasma of OA patients was decreased with OA progression.**

To investigate whether the expression of HAND2-AS1 altered with the progression of OA, OA patients and healthy controls were divided into advanced stages and late stages according to the clinical diagnosis. The expression of HAND2-AS1 in plasma of these groups was detected by qRT-PCR. The results showed that the expression of HAND2-AS1 in the late stage was significantly lower than that in the advanced stage in OA patients. However, in healthy control groups, no significant changes were observed between advanced stage and late stage (Tab.2).

### **The expression of HAND2-AS1 in plasma of OA patients and healthy controls did not alter in different gender.**

since the incidence of OA in female is higher than that in male, we also determined whether the expression of HAND2-AS1 is related to gender in OA patients. OA patients were divided into male and

female two groups and the results showed that the level of HAND2-AS1 in OA patients was not significantly changed between the two groups. The same results were also observed in the healthy controls and these results suggested that different gender cannot affect the expression of HAND2-AS1 (Tab. 3).

### **HAND2-AS1 and IL-6 were inversely correlated in OA patients.**

To investigate whether the expressions of lncRNA HAND2-AS1 and IL-6 are related in OA patients and normal subjects, Pearson correlation coefficient analysis was used to analyze the correlation between HAND2-AS1 and IL-6 in plasma of OA patients and normal subjects (Fig.2A). HAND2-AS1 and IL-6 were significantly and inversely correlated in OA patients, while the expressions of HAND2-AS1 and IL-6 in normal subject plasma were analyzed by Pear After analysis of the poor correlation coefficient (Fig.2B).

### **HAND2-AS1 inhibited IL-6 in chondrocytes.**

To investigate whether HAND2-AS1 can inhibit IL-6 expression in chondrocytes, human chondrocyte cell line was transfected with HAND2-AS1 over-expression plasmid (HAND2-AS1) and negative control plasmid (NC). The over-expression of HAND2-AS1 was confirmed by qRT-PCR after transfection (Fig.3A) and the IL-6 level was detected by Western blot. The results showed that after HAND2-AS1 and NC transfection, the expression level of IL-6 protein in human chondrocyte cells was significantly lower than that in the NC control group (Fig.3B-C). Furthermore, 5, 10 and 20 ng of exogenous IL-6 were applied to human chondrocyte cells and qRT-PCR was used to detect HAND2-AS1 expression in these cells and the results showed that the expression of HAND2-AS1 in cells treated with exogenous IL-6 protein was not change significantly (Fig.3D), indicating that HAND2-AS1 regulated IL-6 expression in human chondrocytes.

## **Discussion**

OA has posed a significant burden on individuals and society since this severe musculoskeletal disease decreases mobility, productivity and quality of life and increases social expenditure and the use of healthcare services.<sup>26,27,28</sup> It has been estimated that up to 240 million people around the world are suffering from OA and due to the increasing levels of sedentary behavior, diminishing levels of physical activity and ever-expanding ageing population, the prevalence of OA is expected to become the most common form of musculoskeletal disease by 2040.<sup>29,30</sup> However, the disease-modifying osteoarthritis drug development is still elusive. HAND2-AS1 is an lncRNA with known functionality in the regulation of liver cancer, stomach cancer, endometrial cancer and colon cancer.<sup>31,32</sup> However, the biological role and clinical function of HAND2-AS1 in OA are unclear. We found that the downregulation of HAND2-AS1 is also involved in OA and the role of HAND2-AS1 in OA is likely achieved through the inhibition of IL-6, which is a key mediator of inflammatory responses in OA. Also, we found that the HAND2-AS1 expression is decreased along with age and the progression of OA. Most importantly, HAND2-AS1 may act as a biomarker that can effectively distinguish OA patients from normal subjects.

IL-6 is a proinflammatory cytokine and has been proved participate in many inflammatory diseases.<sup>33, 34, 35</sup> Many studies showed that inhibition of IL-6 is a promising way to the treatment of inflammatory diseases.<sup>36, 37, 38</sup> We observed that HAND2-AS1 can regulate IL-6 in OA patients, which may be an upstream inhibitor of IL-6. However, this regulation could not be observed in controls groups, suggest that the inhibition of IL-6 by HAND2-AS1 is likely indirect and there may exist pathological mediators between HAND2-AS1 and IL-6. We will further focus on the mechanism of the inhibition of IL-6 by HAND2-AS1.

Numerous lncRNAs have been proved play critical roles in OA and mediate the apoptosis of chondrocytes.<sup>39, 40</sup> Here we first reported the downregulation of lncRNA HAND2-AS1 in OA, which can effectively distinguish OA patients from the healthy controls, suggest that downregulation of HAND2-AS1 may be used to assist the diagnosis of osteoarthritis. Moreover, the expression of HAND2-AS1 is decreased along with the progression of OA and age, indicating that the level of HAND2-AS1 may reflect different stages of OA.

## **Declarations**

### **Ethics approval and consent to participate**

This study has passed the review of the Ethics Committee of the First Affiliated Hospital of Harbin Medical University. All patients signed the informed consent forms.

### **Consent for publication**

All authors are in agree with the content of the manuscript.

### **Availability of data and materials**

Not applicable

### **Competing interests**

The Author(s) declare that there is no conflict of interests regarding the publication of this paper

### **Funding**

Not applicable

### **Authors' contributions**

All authors have contributed significantly

### **Acknowledgements**

Not applicable

## References

1. Grandi FC, Bhutani N. Epigenetic Therapies for Osteoarthritis. *Trends Pharmacol Sci* 2020, **41**: 557-569.
2. Rice SJ, Beier F, Young DA, Loughlin J. Interplay between genetics and epigenetics in osteoarthritis. *Nat Rev Rheumatol* 2020, **16**: 268-281.
3. Emery CA, Whittaker JL, Mahmoudian A, Lohmander LS, Roos EM, Bennell KL, *et al*. Establishing outcome measures in early knee osteoarthritis. *Nat Rev Rheumatol* 2019, **15**: 438-448.
4. Hunter DJ, Bierma-Zeinstra S. Osteoarthritis. *Lancet* 2019, **393**: 1745-1759.
5. Conaghan PG, Cook AD, Hamilton JA, Tak PP. Therapeutic options for targeting inflammatory osteoarthritis pain. *Nat Rev Rheumatol* 2019, **15**: 355-363.
6. Jamshidi A, Pelletier JP, Martel-Pelletier J. Machine-learning-based patient-specific prediction models for knee osteoarthritis. *Nat Rev Rheumatol* 2019, **15**: 49-60.
7. Jones IA, Togashi R, Wilson ML, Heckmann N, Vangsness CT, Jr. Intra-articular treatment options for knee osteoarthritis. *Nat Rev Rheumatol* 2019, **15**: 77-90.
8. Meeson RL, Todhunter RJ, Blunn G, Nuki G, Pitsillides AA. Spontaneous dog osteoarthritis - a One Medicine vision. *Nat Rev Rheumatol* 2019, **15**: 273-287.
9. Si Z, Wang X, Kang Y, Wang X, Sun C, Li Y, *et al*. Heme Oxygenase 1 Inhibits Adult Neural Stem Cells Proliferation and Survival via Modulation of Wnt/beta-Catenin Signaling. *J Alzheimers Dis* 2020, **76**: 623-641.
10. Berenbaum F, Wallace IJ, Lieberman DE, Felson DT. Modern-day environmental factors in the pathogenesis of osteoarthritis. *Nat Rev Rheumatol* 2018, **14**: 674-681.
11. Leech RD, Eyles J, Batt ME, Hunter DJ. Lower extremity osteoarthritis: optimising musculoskeletal health is a growing global concern: a narrative review. *Br J Sports Med* 2019, **53**: 806-811.
12. Marshall M, Watt FE, Vincent TL, Dziedzic K. Hand osteoarthritis: clinical phenotypes, molecular mechanisms and disease management. *Nat Rev Rheumatol* 2018, **14**: 641-656.
13. Runhaar J, Bierma-Zeinstra SMA. Should exercise therapy for chronic musculoskeletal conditions focus on the anti-inflammatory effects of exercise? *Br J Sports Med* 2017, **51**: 762-763.
14. Storch H, Zimmermann B, Resch B, Tykocinski LO, Moradi B, Horn P, *et al*. Activated human B cells induce inflammatory fibroblasts with cartilage-destructive properties and become functionally suppressed in return. *Ann Rheum Dis* 2016, **75**: 924-932.
15. Klein K, Kabala PA, Grabiec AM, Gay RE, Kolling C, Lin LL, *et al*. The bromodomain protein inhibitor I-BET151 suppresses expression of inflammatory genes and matrix degrading enzymes in rheumatoid arthritis synovial fibroblasts. *Ann Rheum Dis* 2016, **75**: 422-429.
16. Latourte A, Cherifi C, Maillet J, Ea HK, Bouaziz W, Funck-Brentano T, *et al*. Systemic inhibition of IL-6/Stat3 signalling protects against experimental osteoarthritis. *Ann Rheum Dis* 2017, **76**: 748-755.

17. Nasi S, So A, Combes C, Daudon M, Busso N. Interleukin-6 and chondrocyte mineralisation act in tandem to promote experimental osteoarthritis. *Ann Rheum Dis* 2016, **75**: 1372-1379.
18. Si Z, Wang X, Zhang Z, Wang J, Li J, Li J, *et al.* Heme Oxygenase 1 Induces Tau Oligomer Formation and Synapse Aberrations in Hippocampal Neurons. *J Alzheimers Dis* 2018, **65**: 409-419.
19. Wei P, Yang J, Zhang D, Cui M, Li L. lncRNA HAND2-AS1 Regulates Prostate Cancer Cell Growth Through Targeting the miR-106a-5p/RBM24 Axis. *Onco Targets Ther* 2020, **13**: 4523-4531.
20. Gokulnath P, de Cristofaro T, Manipur I, Di Palma T, Soriano AA, Guarracino MR, *et al.* Long Non-Coding RNA HAND2-AS1 Acts as a Tumor Suppressor in High-Grade Serous Ovarian Carcinoma. *Int J Mol Sci* 2020, **21**.
21. Gong J, Fan H, Deng J, Zhang Q. LncRNA HAND2-AS1 represses cervical cancer progression by interaction with transcription factor E2F4 at the promoter of C16orf74. *J Cell Mol Med* 2020, **24**: 6015-6027.
22. Ding J, Li Y, Zhang Y, Fan B, Li Q, Zhang J, *et al.* Identification of key lncRNAs in the tumorigenesis of intraductal pancreatic mucinous neoplasm by coexpression network analysis. *Cancer Med* 2020, **9**: 3840-3851.
23. Jiang Z, Li L, Hou Z, Liu W, Wang H, Zhou T, *et al.* LncRNA HAND2-AS1 inhibits 5-fluorouracil resistance by modulating miR-20a/PDCD4 axis in colorectal cancer. *Cell Signal* 2020, **66**: 109483.
24. He J, Zhao H, Deng D, Wang Y, Zhang X, Zhao H, *et al.* Screening of significant biomarkers related with prognosis of liver cancer by lncRNA-associated ceRNAs analysis. *J Cell Physiol* 2020, **235**: 2464-2477.
25. Wang Y, Zhu P, Luo J, Wang J, Liu Z, Wu W, *et al.* LncRNA HAND2-AS1 promotes liver cancer stem cell self-renewal via BMP signaling. *EMBO J* 2019, **38**: e101110.
26. Jeon OH, David N, Campisi J, Elisseeff JH. Senescent cells and osteoarthritis: a painful connection. *J Clin Invest* 2018, **128**: 1229-1237.
27. Rausch Osthoff AK, Niedermann K, Braun J, Adams J, Brodin N, Dagfinrud H, *et al.* 2018 EULAR recommendations for physical activity in people with inflammatory arthritis and osteoarthritis. *Ann Rheum Dis* 2018, **77**: 1251-1260.
28. Roemer FW, Kwok CK, Hayashi D, Felson DT, Guermazi A. The role of radiography and MRI for eligibility assessment in DMOAD trials of knee OA. *Nat Rev Rheumatol* 2018, **14**: 372-380.
29. McAlindon TE, Bannuru RR. Osteoarthritis in 2017: Latest advances in the management of knee OA. *Nat Rev Rheumatol* 2018, **14**: 73-74.
30. Mobasher A, Rayman MP, Gualillo O, Sellam J, van der Kraan P, Fearon U. The role of metabolism in the pathogenesis of osteoarthritis. *Nat Rev Rheumatol* 2017, **13**: 302-311.
31. Huo W, Qi F, Wang K. Long non-coding RNA FER1L4 inhibits prostate cancer progression via sponging miR-92a-3p and upregulation of FBXW7. *Cancer Cell Int* 2020, **20**: 64.
32. Liu S, Zou B, Tian T, Luo X, Mao B, Zhang X, *et al.* Overexpression of the lncRNA FER1L4 inhibits paclitaxel tolerance of ovarian cancer cells via the regulation of the MAPK signaling pathway. *J Cell*


*Biochem* 2018.

33. Luis-Rodriguez D, Donate-Correa J, Martin-Nunez E, Ferri C, Tagua VG, Perez Castro A, *et al.* Serum urate is related to subclinical inflammation in asymptomatic hyperuricaemia. *Rheumatology (Oxford)* 2020.
34. Maurel M, Castagne R, Berger E, Bochud M, Chadeau-Hyam M, Fraga S, *et al.* Patterning of educational attainment across inflammatory markers: Findings from a multi-cohort study. *Brain Behav Immun* 2020.
35. Zeng Z, Yu H, Chen H, Qi W, Chen L, Chen G, *et al.* Longitudinal changes of inflammatory parameters and their correlation with disease severity and outcomes in patients with COVID-19 from Wuhan, China. *Crit Care* 2020, **24**: 525.
36. Garza KM, Zhang L, Borron B, Wood LB, Singer AC. Gamma Visual Stimulation Induces a Neuroimmune Signaling Profile Distinct from Acute Neuroinflammation. *J Neurosci* 2020, **40**: 1211-1225.
37. Lee WR, Lin YK, Alalaiwe A, Wang PW, Liu PY, Fang JY. Fractional Laser-Mediated siRNA Delivery for Mitigating Psoriasis-like Lesions via IL-6 Silencing. *Mol Ther Nucleic Acids* 2020, **19**: 240-251.
38. Shi W, Zhong J, Zhang Q, Yan C. Structural characterization and antineuroinflammatory activity of a novel heteropolysaccharide obtained from the fruits of *Alpinia oxyphylla*. *Carbohydr Polym* 2020, **229**: 115405.
39. Bai J, Zhang Y, Zheng X, Huang M, Cheng W, Shan H, *et al.* LncRNA MM2P-induced, exosome-mediated transfer of Sox9 from monocyte-derived cells modulates primary chondrocytes. *Cell Death Dis* 2020, **11**: 763.
40. Huynh NP, Gloss CC, Lorentz J, Tang R, Brunger JM, McAlinden A, *et al.* Long non-coding RNA GRASLND enhances chondrogenesis via suppression of the interferon type II signaling pathway. *Elife* 2020, **9**.

## Tables

Due to technical limitations, table 1,2,3 is only available as a download in the Supplemental Files section.

## Figures



**Figure 1**

lncRNA HAND2-AS1 and IL-6 were dysregulated in plasma and synovial fluid of OA patients. (A) qRT-PCR results showed the level of HAND2-AS1 in plasma of OA patients and controls. (B) ELISA results showed the level of IL-6 in plasma of OA patients and healthy controls. (C) qRT-PCR results showed the level of HAND2-AS1 in synovial fluid of OA patients and healthy controls. (D) ELISA results showed the level of IL-6 in synovial fluid of OA patients and healthy controls.




Figure 2

HAND2-AS1 and IL-6 were inversely correlated in OA patients. (A) Pearson's correlation coefficient was used to analysis the correlations between HAND2-AS1 and IL-6 in OA patients. (B) Pearson's correlation coefficient was used to analysis the correlations between HAND2-AS1 and IL-6 in healthy controls.




Figure 3

HAND2-AS1 inhibited IL-6 in human chondrocyte cells. (A) qRT-PCR results showed the level of relative HAND2-AS1 in human chondrocyte cells after HAND2-AS1 and NC transfection. (B-C) Western blot results showed the level of IL-6 in human chondrocyte cells after HAND2-AS1 and NC transfection. (D) qRT-PCR results showed the level of HAND2-AS1 in human chondrocyte cells after treatment with IL-6. Data are means  $\pm$  SEM of 3 independent experiments. \*P 0.05.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.JPG](#)
- [Table2.JPG](#)
- [Table3.JPG](#)