Assessment of risk factors for early childhood caries at different ages in Shandong, China and reflections on oral health education: a cross-sectional study

Zhifeng Wang (✉ kqwzf@sdu.edu.cn)

Shandong University

Meng Zhang

Shandong University

Xinyue Zhang

Shandong University

Yuan Zhang

Shandong University

Yanan Li

Shandong University

Chunchun Shao

Shandong University

Shijiang Xiong

Shandong University

Jing Lan

Shandong University

Research article

Keywords: Early childhood caries, Oral health education, Risk factor, Oral health care

Posted Date: February 21st, 2020

DOI: <https://doi.org/10.21203/rs.2.18718/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Oral Health on May 12th, 2020. See the published version at <https://doi.org/10.1186/s12903-020-01104-8>.

Abstract

Background: The high prevalence of early childhood caries (ECC) is widespread around the world, and oral health education (OHE) plays a vital role in preventing ECC. Numerous studies on ECC risk factor assessment have assisted us in enriching the content of OHE. The objective of this study was to further assess independent risk factors for ECC at different ages to provide evidence and insights for OHE.

Methods: Children aged 3-5 years old (N=1301) in Shandong Province were enrolled in this cross-sectional study. Data about oral health status and guardians' oral health knowledge, attitude, and practice (KAP) were extracted from the 4th National Oral Health Survey of China. The associations between ECC prevalence and various KAP variables were tested with chi-square tests, bivariate analysis and multivariable logistic regression analyses.

Results: The ECC prevalence in Shandong Province was 64.6%, and the dmft mean was 3.15. The independent variables with an increased risk for ECC were age, feeding method within 6 months of birth, bedtime sugar frequency, experience of toothache over the past year and dental visits. Feeding methods primarily contributed to the high risk of the 3-year-old group, while sugar habits mainly contributed to that of the 5-year-old group. The association between brushing and ECC was not confirmed in this study, and some positive knowledge and attitude variables were unexpectedly positively correlated with a high risk of ECC.

Conclusion: These data provide evidence to suggest that the ECC-related risk factors at different ages are inconsistent, which provides some insights for OHE. Specifically, in addition to promoting preventive dental visit, we should highlight the effects of feeding methods in the early stages of deciduous dentition and sugar habits in the late stages of deciduous dentition on ECC, as well as encourage supplemental training for food oral health practices.

Background

The high prevalence of early childhood caries (ECC) is a common phenomenon both in China and worldwide[1, 2], and this prevalence coupled with the related adverse effects on oral health and even general health[3, 4]. Hence, strengthening the prevention of ECC has been the most urgent and significant task for both dentists and society as a whole[5].

The current prevention of caries is mainly achieved through the following two important aspects: i) professional preventive measures implemented in dental clinics, including fluoride coating, pit and fissure sealing, etc.[6-8]; and research involving novel anti-caries materials is one of the hotspots of current basic research; and ii) self-oral health care (S-OHC) behaviour, which, in addition to brushing, gargling, and using flossing to remove dental plaque, refers to the control of high-risk factors related to caries, such as sugar consumption and feeding habits[7, 9, 10]. Access to this information is primarily achieved through oral health education (OHE), including dental clinics, school, community and network platforms[11].

OHE has always contributed to dental caries prevention[12, 13], as it brings intuitive guidance and assistance to human beings. OHE details are mainly based on studies involving ECC risk factor assessment, which could provide strong evidence and support for OHE. At present, many high-risk factors associated with dental caries have been demonstrated, such as poverty, race, eating habits, family economic status, education level, etc.[14-16]. However, few studies have separately analysed and assessed the risk factors for each age group; instead, they have considered all preschool children as a whole, which has hindered us from preventing dental caries more effectively by optimizing and enriching OHE.

Shandong Province is located in the lower reaches of the Yellow River and is bordered by the Bohai Sea in the north and the Yellow Sea in the east. The local residents primarily consume wheat and its products. The province's GDP ranks third in China, and it is one of the representative provinces in northern China. However, few studies have reported the oral health status of children in Shandong Province.

Therefore, this cross-sectional study is the first to report and analyse the oral health status and independent risk factors for children aged 3-5 years old in Shandong Province, China, and to pay special attention to the risk factors associated with ECC in different age groups. Those variables involving sociodemographic factors, guardians' knowledge, attitude, practice variables and children's oral health status were collected for statistical analyses. We hope that these data will provide guidance and evidence for improving and refining the contents of oral health education.

Methods

This cross-sectional study was conducted in 2015-2016 in Shandong Province and was part of the 4th National Oral Health Survey in China; it has been approved by the ethics committee of the Chinese Stomatological Association (NO. 2014-003).

Sample selection

Three- to five-year-old children living in Shandong Province were our target population. The sample size was determined by the equal allocation of the national plan, the calculating formula

[Please see supplementary files to access the equation.]

is based on estimated caries prevalence rate ($P=66.0\%$, data from the 3rd National Oral Health Survey, 2005), the design effect ($deff=4.5$), the significance level ($\alpha=5\%$), the margin of error ($\delta=10\%$) and the non-response rate (20%), a total of 13,392 children should be recruited for each age group in 31 provinces across the country[17]. According to this at least 1,296 sample data should ultimately be enrolled in this study. We incorporated the sample population according to the following multistage stratified sampling method:

1. The probability proportion to size (PPS) sampling method determined the two districts and two counties to be used for the survey.
2. Three kindergartens were randomly selected from each region.
3. Using cluster sampling and quota sampling, 36 children were selected to represent each age group from each kindergarten.

Inspector selection and training

Three inspectors from the Hospital of Stomatology were selected and trained. To ensure the consistency of the clinical practice inter- and intra-inspectors, each inspector randomly enrolled three participants for examination. Before calculating the Kappa value, every examination result had to be calibrated with a qualified inspector. The mean Kappa value of the intra-inspectors was determined through a re-examination of the samples, and the mean Kappa value of the inter-inspectors was calculated through the mutual examination of the samples. Clinical practice training ended when the Kappa value was greater than 0.8.

Informed consent and questionnaire

After the guardians had signed the informed consent forms, the children were scheduled for a standard oral examination. In addition, a questionnaire was completed by the guardians, and then the staff rechecked it for missing items, abnormal items, etc. The questionnaire contents mainly included the following:

1. Personal and family information.
2. OHC knowledge, mainly involved in the effect of brushing, bacteria, sugar, pit and fissure sealing, fluoride and other factors on the teeth and gums.
3. OHC attitude, mainly to investigate whether the guardian believes that oral health is extremely important.
4. Guardian-guided OHC practices for children, which mainly obtains information on the important risk factors that have been confirmed by other studies, including feeding methods, sugar consumption, brushing, toothaches, fluoride exposure, dental visits and other detailed information.

Caries assessment

A ball-end community periodontal index probe was used to assess dental status, according to the WHO guidelines[18]. Individuals who were recorded as having decayed teeth, filled teeth that still had caries, or missing teeth due to caries were included in the ECC group, and the individuals without decayed teeth were included in the caries-free group (CF). In addition, the written version of the examination results and treatment recommendations were sent to the children's guardians to increase the significance of the survey.

Statistical analysis

IBM SPSS Statistics version 17.0 was used to perform the statistical analyses. The prevalence of preschool children with ECC and the mean dmft count were calculated to assess the association of caries with selected KAP variables. Analyses of variance, [chi-square tests](#), Fisher's test and z tests for post hoc comparisons were performed to assess the statistical significance between different variables, all of which were analyzed using the caries-free group as a control. Multivariate logistic regression models were carried out based on the outcomes of ECC (any versus none) to evaluate the independent risk factors while also controlling for latent confounders found in the bivariate analyses ($P \leq 0.10$). All KAP variables included in the models were categorical variables. Some count variables were recoded into categorical variables if necessary, such as the income variable. Differences were considered significant at $P < 0.05$.

Results

A total of 2 districts and 2 counties were selected in Shandong Province. In total, 1,330 preschool children were invited to participate in the study, but due to the loss of 29 individuals' information, a total of 1,301 children's data were included in the final statistical analyses. Up to 64.6% of children aged 3-5 had ECC, and the mean dmft (standard deviation) was 3.15 ± 3.845 . The rates of the 3-, 4-, and 5-year-old age groups were 51.1%, 67.8%, and 73.9%, respectively, and the mean dmft values were 2.16 ± 3.096 , 3.21 ± 3.966 , and 4.01 ± 4.134 , respectively (see Table 1). While 96.4% of the decayed teeth were waiting to be treated.

Statistical analyses of ECC-related variables

The demographic characteristics are shown in Table 1. The ECC rates increased with age ($P = 0.000$), whereas gender, region, education background and income had no association with the rates of ECC in this study.

The distribution characteristics of the KAP variables are shown in Appendix Table S1. A total of 1,052 questionnaires (80.9%) were completed by the parents, and the rest were completed by other people ($P > 0.05$, data not shown). From the perspective of OHC knowledge (Q22a-h) and attitude (Q21a-f), 66.0%-98.8% of the individuals answered most of the questions correctly. While only 30.7% reported knowing that gum bleeding is not a normal phenomenon when brushing teeth (Q22a), approximately 16.0% and 27.7% reported knowing that pit and fissure sealing or fluoride, respectively, can protect teeth (Q22g, h), and 35.8% believed that the mother's oral health affects the child's oral health (21f). Those who reported knowing that bacteria (Q22d) and sugar (Q22e) can induce caries and that pit and fissure sealing can protect teeth (Q22g) and those who believe that teeth health is related to their own protection (Q21c) are unexpectedly more likely to experience ECC ($P < 0.05$). From the perspective of OHC practices (Q3-13), children with completely artificial feeding within 6 months of birth had the lowest rates of ECC compared with other feeding methods (Q3, 45.0% vs 65.3%-66.5%, respectively; $P = 0.005$). The frequency of eating sugary foods before going to bed at night was associated with increased rates of ECC (Q5, 58.6% vs 66.8% vs 73.8%; $P = 0.002$). In addition, 53.1% of all children ($n = 691$) had tooth-brushing habits,

with 98.8% using toothpaste; however, only 8.8% (n=60) reported knowing they were using fluoride toothpaste, and these children unexpectedly had the highest rates of ECC compared with other children who did not know the ingredients of their toothpaste (Q11, n=683, 80.0% vs 62.2%; $P=0.016$); in addition, the earlier the children started brushing their teeth, the lower rate of dental caries was (Q7, n=691, $P=0.033$). Children who had frequent toothaches or discomfort in the past 12 months (Q12) or children who had ever visited a dentist (Q13) generally had a higher caries rate than did children who had never experienced either event (Q12: 88.1%-100.0% vs 57.1% and Q13: 81.0% vs 60.7%, respectively; $P=0.000$). The number of brushing times per day were not associated with an increased number of caries in the current study.

Statistical analyses of ECC-related variables at different ages

Considering that age was a significant difference variable, the differences in ECC rates across the variables in each age group were analysed to eliminate the potential effects of age. The results of the bivariate analyses for the different ages are shown in Table 2. In the 3-year-old group, the children raised by parents who believe that sugar can induce dental decay, those who were primarily breastfed within 6 months of birth, those who had experienced toothache recently, or those who had visited a dentist were more likely to suffer from dental caries ($P<0.05$). In the 4-year-old group, girls generally had higher rates of caries (72.9% vs 62.5%; $P=0.025$), and children who had ever had a toothache or a dental visit still had the highest rates of ECC ($P<0.05$). In the 5-year-old group, the frequency of eating sweets before going to bed, toothache experience, dental visit experience and the cognitive status about bacteria being able to cause dental caries were associated with increased rates of ECC ($P<0.05$). Overall, having experienced toothache and having had dental visits (Q12 and 13) were both correlated with a high risk of ECC for each age group.

Logistic regression analyses

The results of the multivariable logistic regression analyses for ECC are shown in Table 3. Overall, the association between increasing age and the accumulation of ECC was still significant (OR: 2.44, 95% confidence interval (CI): 1.79-3.31 for the 5-year-old group; OR: 2.07, 95% CI: 1.55-2.78 for the 4-year-old group). Factors independently associated with an augmented risk for ECC include being breastfed completely within 6 months of birth, a high frequency of consuming sweets before bedtime (OR: 1.81, 95% CI: 1.09-3.02), having experienced a toothache over the past year (OR: 4.7, 95% CI: 3.11-7.11) and having had dental visits (OR: 1.75, 95% CI: 1.19-2.55). Unexpectedly, those families who reported knowing that sugar can induce caries were more likely to experience ECC (OR: 1.49, 95% CI: 1.07-2.06). However, toothbrushing was not associated with increased rates of ECC.

Table 4 shows the results of the multivariable logistic regression analyses used to identify risk factors independently associated with ECC in different age groups. For the 3-year-old group, those with a history of having been completely artificial fed within six months of birth were less likely to experience ECC compared with those who had been completely breastfed (OR: 0.28, 95% CI: 0.12-0.69) ; however, those with a history of having experienced toothache over the last year (OR: 5.83, 95% CI: 2.89-11.77) and positive cognition regarding which bacteria can cause dental caries (OR: 1.77, 95% CI: 1.02-3.07) were more likely to suffer from dental caries. The independent risk factors for the 4-year-old group only included gender and having had a toothache experience over the past year. For the 5-year-old group, the factors independently associated with an increased risk for ECC included having an increased frequency of bedtime sweet consumption, having had a toothache experience over the past year and having had dental visits. Brushing was still not an independent risk factor for ECC.

Discussion

As a populous province, Shandong Province has a population of nearly 100 million people. The current study collected eligible samples to represent the oral health status of this province and its cover areas with different dietary habits. Ultimately, out of a total of 1,330 children aged 3-5 years old, 1301 families provided complete and reliable data information, and the response rate was as high as 97.8%, which is very satisfying.

The caries prevalence rates for 3- to 5-year-old children in Shandong Province are 51.1%, 67.8% to 73.9%, respectively, which are all higher than the national levels (50.8%, 63.6%, 71.9%, respectively), while the dmft mean and the untreated rate of dental caries are lower than the national data (3.15 vs 3.35; 96.4% vs 96.9%)[19] . Other developing countries in Asia, such as Thailand (age 5: 78.5%)[20], Indonesia (age 0-5: 70.0%)[21], Myanmar (age 5-6: 81.3%) and Cambodia (age 3: 84.9%)[22] all have higher rates of ECC compared to those of Shandong Province. In developed countries such as Japan and Singapore, the ECC rates are down to 44.4% and 49%, respectively[23, 24], which are significantly lower than the worldwide result of 63% obtained by statistics[25]; these countries have also already achieved the WHO's goal of having less than a 50% prevalence rate of ECC for the year 2020[26]. These data alert us that there is still a long and hard way to go to reduce the prevalence of ECC in Shandong Province, China.

Age has been evidenced to be an independent risk factor for ECC by many studies[14, 22, 27], and this finding was corroborated in the current study. Most studies, however, evaluate 3- to 5-year-old children as whole, with few studies assessing the risk factors for each age. Therefore, we speculated that although the age span used in the current study is small, different ages may have different risk factors. Both the chi-square tests and the logistic regression model in the present study confirm this speculation (Table 2-4). Although different age groups share caries-related risk factors—the experience of toothache over the past year—the increased risk for ECC in children aged 3 is independently associated with the feeding method within 6 months after birth, while that of the 5-year-old children is primarily associated with the frequency of bedtime sweet consumption. This heterogeneity indicates that we should have different priorities in regard to OHE for preschool children of different ages.

The association between feeding method and ECC is intricate and involves a variety of factors, including the frequency, duration, amount, time and formula type of the feeding. Previous studies have reported that breastfeeding and its duration are not associated with an increased risk of ECC[14], yet studies have also found that both prolonged breastfeeding and high-frequency feeding in late infancy augment the risk of having dental caries[28, 29]. Interestingly, Van Palenstein et al.[30] reported that breastfeeding during daylight hours beyond the age of 12 months was not associated with ECC for 25- to 30-month-old infants; rather, the relevant factor is that of breastfeeding at night >2 times and exposing the child to >15 minutes per nocturnal feeding. Although the current study also provides evidence that breastfeeding has a greater adverse impact on ECC than that of artificial feeding, which is similar to the results of national studies completed 10 years ago[31], no more detailed information about the duration, frequency, time, and formula type were collected; thus, it is difficult to analyse and attain those specific factors that may have caused such outcomes. After all, the high cariogenicity of bottle feeding has also been confirmed by many studies[32]. One recent study has contributed evidence for the nonlinear association between breastfeeding duration and ECC and the auxiliary role of fluoridated water in reducing the negative effects of feeding on ECC[33]. This finding reminds us that more relevant variables need to be considered in future studies to elucidate the correlation between feeding methods and ECC.

Frequent exposure to dietary sugar and refined carbohydrates is well recognized to be associated with ECC[34-37], and bedtime sweet consumption is particularly serious[35, 38]. Excessive sugar consumption provides abundant nutrients for supragingival microorganisms and induces acid production, enamel-dentine demineralization, and caries formation[12, 39]. The limitation of sugar consumption has therefore been listed in a declaration that has gained worldwide support[40]. Interestingly, the data from the present study found that the increase in the frequency of bedtime sweet consumption was not an independent risk factor for ECC in children aged 3 or 4, it was only so for children aged 5 years. It is not possible from these data to conclude that there is no need to pay attention to the sugar consumption habits of those under 4 years old because sweet preference has been evidenced to be a high risk factor for ECC in children under 23 months of age in Japan[38]. Leroy et al.[41] and Schluter et al.[42] also confirmed that bedtime sweet consumption is significantly associated with ECC for children aged 3 and 4. Accordingly, it is possible that the outcomes of previous undesirable sugar consumption habits manifested in those 5 years old in the current study and are thus seen to be an independent risk factor for the current study sample. This is in full compliance with the actions of time factors in the pathogenesis of dental caries[12]. Therefore, even though sugar consumption habits in this study were only risk factors for the 5-year-old age group, knowledge transmission and behavioural interventions should be carried out early in life.

Previous studies have found that infrequent dental visits were correlated with an augmented risk for untreated caries[43, 44]. Some studies, however, have reported a positive association between the number of dental visits and ECC prevalence[14, 45]. This finding is consistent with the results of the current study. Unfortunately, this result goes against the goal of oral prevention. The design of the dental visit is expected to play a positive and energetic role in caries prevention[46]; however, 51.8% of families in the present study sought help from a dentist for treatment compared with only 7.8% who sought help for

prevention. An Australian study has reported that dental visits actually contribute 30.3% to the explanation of inequalities in ECC[16]. This result indicates that our awareness of prevention is still insubstantial and that the purpose of dental visits is incorrectly implemented as more for treatment and rarely for prevention. This definition is one of the priorities that we need to improve.

Toothbrushing has always been considered to be the primary and most effective way to clean plaque microbes and prevent dental caries. Many studies have confirmed the association between toothbrushing and ECC[47, 48]. Many studies have also verified that there is no correlation between these two variables[49, 50], which is consistent with the results of the present study. Abdelaziz et al.[51] reported that brushing was not an independent risk factor for ECC and S-ECC in preschool children. Edward Lo even found that supplemental training in regard to parental toothbrushing would not benefit the reduction of ECC for those children under 3 years of age who lived in a water-fluoridated area and had been provided with sufficient OHE[52]. However, the sure thing is the relationship between the earlier the brushing time and the lower the ECC risk[15, 31], the present study also confirmed this trend through a bivariate analyses. Honestly, the interpretation of these results as being relevant to such issues is further limited due to the deficiency of information about brushing details; for example, brushing duration, brushing compliance, floss assistance, and especially brushing efficiency assessment are not available in our data. It is therefore difficult to objectively assess the association between toothbrushing and ECC in the scenario of many factors being masked or even neglected. However, the special dietary habits of Shandong Province, in which wheat is the main food ingredient, may be one of the reasons why toothbrushing was not demonstrated to be an independent risk factor for ECC. Further studies should include these variables so that we can attain a clearer picture of the role of toothbrushing in ECC prevention.

The results of the present study actually reflect the current status and issues of oral health education in Shandong Province. First, there is a sharp contrast between the high awareness of oral health routine knowledge (66%-98.8%) and the low access to oral prevention knowledge (16%-35.8%), a negative association between some correct knowledge and attitude variables (Q22d/e: sugar or bacteria can induce ECC; Q21c: dental health is related to self-protection.) and ECC risk (Appendix Table S1, Table 2-4), and a positive correlation between frequency of toothache or dental visits and ECC. All of these results reflect that the knowledge and information provided by the current OHE in Shandong Province appears to be inefficient and that it underperforms in promoting the implementation of ECC prevention practices. However, many studies have confirmed that when a more systematic, proactive oral health intervention involving brushing training, referrals and dental care is implemented for a family with children, even if implemented by nondental professionals[53], it generally has a greater preventive effect against ECC[46, 54-57]. Therefore, a systematic and sound oral health care programme that involves clinical care, brushing training, referrals and so on is imperative[58], rather than just oral health education. Only in this way can we ensure the consistency between theoretical knowledge and actual action as much as possible, thereby reducing the incidence rates of ECC. In addition, the risk factors for ECC at different ages (3-5 years old) are not completely consistent. In the current study, the early stage of deciduous dentition (3 years old) is mainly related to feeding methods, while the later stage of deciduous dentition

(5 years old) is mainly associated with sugar-eating habits. These outcomes indicate that a more strategic and more targeted OHE method and content should be designed for children of different ages, which would also be more cost-effective. Finally, although a positive association between toothbrushing, the use of fluoride and the low risk of dental decay was not obtained in this study, the role of such an association in caries prevention is indeed indelible[15, 59, 60]. Therefore, assisting children to master this knowledge and these skills early in life is the responsibility of both dentists and parents.

In summary, based on the results of this study, when we conduct oral health education programmes, we should have different focuses for different age groups. In the early stages of deciduous dentition (less than 3 years old), in addition to routinely introducing oral health knowledge and cultivating oral health awareness, we should emphasize the impact of feeding habits in infancy and early childhood on dental health. It seems inconclusive which feeding method (breastfeeding vs artificial milk feeding) is more beneficial to dental health, but issues such as feeding methods and frequency of night milk consumption deserve to be emphasized. As children get older, the training and mastery of certain oral health behaviours should become a priority, including sugar consumption habits and toothbrushing habits.

The main advantage of this study is its large representative sample of children and its high response rate; also, this study is part of the 4th National Oral Health Survey of China and thus has highly reliable data sources. Furthermore, this study is the first to report the oral health status of children in Shandong Province over the past decade. Limitations of the study include the issues related to the OHC knowledge and attitude items designed for the questionnaire being relatively basic and traditional. Several major risk factors have been incorporated into the current study, unlike the monographic studies that have excavated and explored sufficient detailed information about one certain targeting factor associated with ECC[29, 33, 52]. We also have to acknowledge that some data on detailed information is limited, such as the frequency of bedtime feedings, brushing efficiency, floss assistance status, and cariogenic microbial flora, each of which is reported to be associated with ECC. Therefore, more information is waiting to be collected to further analyse the risk factors associated with ECC for each age group.

Conclusion

The data in the present study provide evidence that different risk factors are independently associated with an augmented risk for ECC at different ages. Feeding methods primarily contribute to the risk in the early stage of deciduous dentition, while sugar habits mainly contribute to the risk in the late stage of deciduous dentition. However, the contribution of toothbrushing and fluoride to caries prevention has not been confirmed. Of course, there is a need for more research to identify the detailed characteristics of the above-mentioned variables that cause ECC and to provide guidelines for children of different ages. In addition, the apparent gap between theoretical knowledge and actual behaviour in this study suggests that we need a more systematic oral health care programme that involves behavioural guidance and effect evaluations, rather than just OHE, to reduce the ECC prevalence rates.

List Of Abbreviations

ECC: early childhood caries; **CF: caries-free;** **KAP:** knowledge, attitude and practice; **OHC:** oral health care; **OHE:** oral health education; **WHO:** world health organization; **dmft:** decayed, missing and filled teeth; **RCT:** randomized controlled trial; **OR:** odd rates; **CI:** confidence interval.

Declarations

Ethics approval and consent to participate: This study had been approved by the Ethics Committee of the Chinese Stomatological Association (NO.2014-003) and the informed consent was obtained from a legal guardian for participants under 16 years old.

Consent for publication: Not applicable.

Availability of data and materials: The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: All phases of this study were supported by the Chinese Stomatological Association and School & Hospital of Stomatology, Shandong University. The design of this study and collection, analysis, and interpretation of data and the manuscript writing are funded by the Key Technology Research and Development Program of Shandong (CN), grant number is 2019GSF108278, and 2019GSF108184.

Authors' contributions: ZF W, J Land SJ X conceptualized and designed the study, reviewed and revised the manuscript. M Z collected and organized data, carried out the statistical analyses, drafted and revised the manuscript. Y Z instructed the methods and strategies of statistical analyses and reviewed and revised the manuscript. XY Z and YN L collected, organized and input data. CC S conducted the statistical analysis of the data. *All authors read and approved the final manuscript.*

Acknowledgements

Thank you very much to the Chinese Stomatological Association for the raw data of the survey they provided. Thank you also for the cooperation and help of various kindergartens.

References

1. **Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016.** *Lancet (London, England)* 2017, **390**(10100):1211-1259.
2. Meyer F, Enax J: **Early Childhood Caries: Epidemiology, Aetiology, and Prevention.** *Int J Dent* 2018, **2018**:1415873.
3. Casamassimo PS, Thikkurissy S, Edelstein BL, Maiorini E: **Beyond the dmft: the human and economic cost of early childhood caries.** *Journal of the American Dental Association (1939)* 2009,

140(6):650-657.

4. Kawashita Y, Kitamura M, Saito T: **Early childhood caries**. *Int J Dent* 2011, **2011**:725320-725320.
5. MA P, LMD M, RJ W, B D, R V, MR M *et al*: **Oral diseases: a global public health challenge**. 2019, **394**(10194):249-260.
6. White JR, Nagarajan N, Pop M: **Statistical methods for detecting differentially abundant features in clinical metagenomic samples**. *PLoS computational biology* 2009, **5**(4):e1000352.
7. Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, Ramos-Gomez F *et al*: **Dental caries**. *Nature Reviews Disease Primers* 2017, **3**(1):17030.
8. Stephen KW: **Use of fluoride**. *Lancet (London, England)* 1999, **354**(9172):74-75.
9. Tanaka S, Shinzawa M, Tokumasu H, Seto K, Tanaka S, Kawakami K: **Secondhand smoke and incidence of dental caries in deciduous teeth among children in Japan: population based retrospective cohort study**. *BMJ (Clinical research ed)* 2015, **351**:h5397.
10. Poulton R, Caspi A, Milne BJ, Thomson WM, Taylor A, Sears MR *et al*: **Association between children's experience of socioeconomic disadvantage and adult health: a life-course study**. *The Lancet* 2002, **360**(9346):1640-1645.
11. Khokhar MA, Khokhar WA, Clifton AV, Tosh GE: **Oral health education (advice and training) for people with serious mental illness**. *Cochrane Database of Systematic Reviews* 2016, 10.1002/14651858.CD008802.pub3(9).
12. Selwitz RH, Ismail AI, Pitts NB: **Dental caries**. *Lancet (London, England)* 2007, **369**(9555):51-59.
13. Arora A, Khattri S, Ismail NM, Kumbargere Nagraj S, Prashanti E: **School dental screening programmes for oral health**. *Cochrane Database of Systematic Reviews* 2017, 10.1002/14651858.CD012595.pub2(12).
14. Iida H, Auinger P, Billings RJ, Weitzman M: **Association between infant breastfeeding and early childhood caries in the United States**. *Pediatrics* 2007, **120**(4):e944-952.
15. Darmawikarta D, Chen Y, Carsley S, Birken CS, Parkin PC, Schroth RJ *et al*: **Factors associated with dental care utilization in early childhood**. *Pediatrics* 2014, **133**(6):e1594-1600.
16. Peres MA, Ju X, Mittinty M, Spencer AJ, Do LG: **Modifiable Factors Explain Socioeconomic Inequalities in Children's Dental Caries**. *Journal of dental research* 2019, **98**(11):1211-1218.
17. Du MQ, Li Z, Jiang H, Wang X, Feng XP, Hu Y *et al*: **Dental Caries Status and its Associated Factors among 3- to 5-year-old Children in China: A National Survey**. *The Chinese journal of dental research : the official journal of the Scientific Section of the Chinese Stomatological Association (CSA)* 2018, **21**(3):167-179.
18. Petersen PE, Baez RJ, World Health O: **Oral health surveys: basic methods 5th ed**, <https://apps.who.int/iris/handle/10665/97035>. Geneva: World Health Organization; 2013.
19. MQ D, Z L, H J, X W, XP F, Y H *et al*: **Dental Caries Status and its Associated Factors among 3- to 5-year-old Children in China: A National Survey**. 2018, **21**(3):167-179.

20. Srisilapanan P, Nirunsittirat A, Roseman J: **Trends over Time in Dental Caries status in Urban and Rural Thai Children.** *Journal of clinical and experimental dentistry* 2017, **9**(10):e1201-e1206.
21. Maharani D, Rahardjo A: **Mothers' Dental Health Behaviors and Mother-Child's Dental Caries Experiences: Study of a Suburb Area in Indonesia.** *MAKARA of Health Series* 2013, 10.7454/msk.v16i2.1632.
22. Turton B, Chher T, Sabbah W, Durward C, Hak S, Lailou A: **Epidemiological survey of early childhood caries in Cambodia.** *BMC oral health* 2019, **19**(1):107.
23. Morita A, Matsuyama Y, Isumi A, Doi S, Ochi M, Fujiwara T: **Association between grandparent co-residence, socioeconomic status and dental caries among early school-aged children in Japan: A population-based prospective study.** *Scientific reports* 2019, **9**(1):11345.
24. XL G, CY H, T L, D K, HB H, health XYJCd: **Dental caries prevalence and distribution among preschoolers in Singapore.** 2009, **26**(1):12-17.
25. Tinanoff N, Baez RJ, Diaz Guillory C, Donly KJ, Feldens CA, McGrath C *et al*: **Early childhood caries epidemiology, aetiology, risk assessment, societal burden, management, education, and policy: Global perspective.** *International journal of paediatric dentistry* 2019, **29**(3):238-248.
26. Hobdell M, Petersen PE, Clarkson J, Johnson N: **Global goals for oral health 2020.** *International dental journal* 2003, **53**(5):285-288.
27. Cabral M, Mota ELA, Cangussu MCT, Vianna MIP, Floriano FR: **Risk factors for caries-free time: longitudinal study in early childhood.** *Revista de saude publica* 2017, **51**:118.
28. Feldens CA, Rodrigues PH, de Anastacio G, Vitolo MR, Chaffee BW: **Feeding frequency in infancy and dental caries in childhood: a prospective cohort study.** *International dental journal* 2018, **68**(2):113-121.
29. Peres KG, Nascimento GG, Peres MA, Mittinty MN, Demarco FF, Santos IS *et al*: **Impact of Prolonged Breastfeeding on Dental Caries: A Population-Based Birth Cohort Study.** *Pediatrics* 2017, 10.1542/peds.2016-2943.
30. van Palenstein Helderma WH, Soe W, van 't Hof MA: **Risk factors of early childhood caries in a Southeast Asian population.** *Journal of dental research* 2006, **85**(1):85-88.
31. Sun X, Bernabe E, Liu X, Gallagher JE, Zheng S: **Early life factors and dental caries in 5-year-old children in China.** *Journal of dentistry* 2017, **64**:73-79.
32. Avila WM, Pordeus IA, Paiva SM, Martins CC: **Breast and Bottle Feeding as Risk Factors for Dental Caries: A Systematic Review and Meta-Analysis.** *PloS one* 2015, **10**(11):e0142922.
33. Ha DH, Spencer AJ, Peres KG, Rugg-Gunn AJ, Scott JA, Do LG: **Fluoridated Water Modifies the Effect of Breastfeeding on Dental Caries.** *Journal of dental research* 2019, **98**(7):755-762.
34. Ribeiro CCC, Silva M, Nunes AMM, Thomaz E, Carmo CDS, Ribeiro MRC *et al*: **Overweight, obese, underweight, and frequency of sugar consumption as risk indicators for early childhood caries in Brazilian preschool children.** *International journal of paediatric dentistry* 2017, **27**(6):532-539.

35. Fontana M, Jackson R, Eckert G, Swigonski N, Chin J, Zandona AF *et al*: **Identification of caries risk factors in toddlers.** *Journal of dental research* 2011, **90**(2):209-214.
36. Chen KJ, Gao SS, Duangthip D, Li SKY, Lo ECM, Chu CH: **Dental caries status and its associated factors among 5-year-old Hong Kong children: a cross-sectional study.** *BMC oral health* 2017, **17**(1):121.
37. Lin YC, Chang CS, Ho PS, Lee CH, Chen JH, Huang HL: **Immigrant(-)Native Differences in Sugar-Sweetened Beverage and Snack Consumption and Preventive Behaviors Associated with Severe Early Childhood Caries: A Large-Scale Survey in Taiwan.** *International journal of environmental research and public health* 2019, **16**(6):10.3390/ijerph16061047.
38. Nakayama Y, Mori M: **Association between nocturnal breastfeeding and snacking habits and the risk of early childhood caries in 18- to 23-month-old Japanese children.** *Journal of epidemiology* 2015, **25**(2):142-147.
39. Rudney JD, Jagtap PD, Reilly CS, Chen R, Markowski TW, Higgins L *et al*: **Protein relative abundance patterns associated with sucrose-induced dysbiosis are conserved across taxonomically diverse oral microcosm biofilm models of dental caries.** *Microbiome* 2015, **3**:69.
40. **Early Childhood Caries: IAPD Bangkok Declaration.** %J *International journal of paediatric dentistry.* 2019, **29**(3):384-386.
41. Leroy R, Hoppenbrouwers K, Jara A, Declerck D: **Parental smoking behavior and caries experience in preschool children.** *Community dentistry and oral epidemiology* 2008, **36**(3):249-257.
42. Schluter PJ, Durward C, Cartwright S, Paterson J: **Maternal self-report of oral health in 4-year-old Pacific children from South Auckland, New Zealand: findings from the Pacific Islands Families Study.** *Journal of public health dentistry* 2007, **67**(2):69-77.
43. Aligne CA, Moss ME, Auinger P, Weitzman M: **Association of pediatric dental caries with passive smoking.** *Jama* 2003, **289**(10):1258-1264.
44. Dye BA, Shenkin JD, Ogden CL, Marshall TA, Levy SM, Kanellis MJ: **The relationship between healthful eating practices and dental caries in children aged 2-5 years in the United States, 1988-1994.** *Journal of the American Dental Association (1939)* 2004, **135**(1):55-66.
45. Graves RC, Abernathy JR, Disney JA, Stamm JW, Bohannon HM: **University of North Carolina caries risk assessment study. III. Multiple factors in caries prevalence.** *Journal of public health dentistry* 1991, **51**(3):134-143.
46. Faustino-Silva DD, Colvara BC, Meyer E, Hugo FN, Celeste RK, Hilgert JB: **Motivational interviewing effects on caries prevention in children differ by income: A randomized cluster trial.** *Community dentistry and oral epidemiology* 2019, **47**(6):477-484.
47. Aida J, Ando Y, Oosaka M, Niimi K, Morita M: **Contributions of social context to inequality in dental caries: a multilevel analysis of Japanese 3-year-old children.** *Community dentistry and oral epidemiology* 2008, **36**(2):149-156.
48. Gao XL, Hsu CY, Xu YC, Loh T, Koh D, Hwarng HB: **Behavioral pathways explaining oral health disparity in children.** *Journal of dental research* 2010, **89**(9):985-990.

49. Begzati A, Berisha M, Meqa K: **Early childhood caries in preschool children of Kosovo - a serious public health problem.** *BMC public health* 2010, **10**:788.
50. Zhou Y, Lin HC, Lo EC, Wong MC: **Risk indicators for early childhood caries in 2-year-old children in southern China.** *Australian dental journal* 2011, **56**(1):33-39.
51. Abdelaziz W, Dowidar K, El Tantawi M: **Association of Healthy Eating, Juice Consumption, and Bacterial Counts with Early Childhood Caries.** *Pediatric dentistry* 2015, **37**:462-467.
52. Jiang EM, Lo EC, Chu CH, Wong MC: **Prevention of early childhood caries (ECC) through parental toothbrushing training and fluoride varnish application: a 24-month randomized controlled trial.** *Journal of dentistry* 2014, **42**(12):1543-1550.
53. George A, Sousa MS, Kong AC, Blinkhorn A, Patterson Norrie T, Foster J *et al*: **Effectiveness of preventive dental programs offered to mothers by non-dental professionals to control early childhood dental caries: a review.** *BMC oral health* 2019, **19**(1):172.
54. Dooley D, Moultrie NM, Heckman B, Gansky SA, Potter MB, Walsh MM: **Oral Health Prevention and Toddler Well-Child Care: Routine Integration in a Safety Net System.** *Pediatrics* 2016, **137**(1).
55. Okah A, Williams K, Talib N, Mann K: **Promoting Oral Health in Childhood: A Quality Improvement Project.** *Pediatrics* 2018, **141**(6).
56. Bartosova M, Svetlak M, Kukletova M, Borilova Linhartova P, Dusek L, Izakovicova Holla L: **Emotional stimuli candidates for behavioural intervention in the prevention of early childhood caries: a pilot study.** *BMC oral health* 2019, **19**(1):33.
57. Ismail A, Razak IA, Ab-Murat N: **The impact of anticipatory guidance on early childhood caries: a quasi-experimental study.** *BMC oral health* 2018, **18**(1):126.
58. Mouradian WE, Wehr E, Crall JJ: **Disparities in children's oral health and access to dental care.** *Jama* 2000, **284**(20):2625-2631.
59. Whelton HP, Spencer AJ, Do LG, Rugg-Gunn AJ: **Fluoride Revolution and Dental Caries: Evolution of Policies for Global Use.** *Journal of dental research* 2019, **98**(8):837-846.
60. Walsh T, Worthington HV, Glenny AM, Marinho VC, Jeroncic A: **Fluoride toothpastes of different concentrations for preventing dental caries.** *The Cochrane database of systematic reviews* 2019, **3**:Cd007868.

Tables

Table 1. Distribution and statistical significance by demographic variables for ECC.
(n=1301)

Variables	CF (n, (%))	ECC (n, (%))	<i>P</i>
gender			
male	235 (36.1)	416 (63.9)	0.616
female	226 (34.8)	424 (65.2)	
Age			
3	203 (48.9) ^a	212 (51.1) ^a	0.000
4	142 (32.2) ^b	299 (67.8) ^b	
5	116 (26.1) ^b	329 (73.9) ^c	
Region			
urban	227 (34.9)	424 (65.1)	0.343
weihai	111 (34.3)	213 (65.7)	
hedong, linyi	116 (35.5)	211 (64.5)	
rural	234 (36.0)	416 (64.0)	
weifang	106 (32.7)	218 (67.3)	
pingyi, linyi	128 (39.3)	198 (60.7)	
Education			
Elementary school and below	88 (34.9)	164 (19.5)	0.699
junior high school	192 (36.8)	330 (39.3)	
Senior high school and above	181 (34.3)	346 (41.2)	
Income			
less than 30,000	111 (34.0)	215 (27.4)	0.357
40,000-60,000	166 (38.0)	271 (34.5)	
more than 70,000	153 (33.8)	300 (38.2)	

CF: caries-free group; ECC: Early Childhood Caries group

^{a, b, c}: Differences of caries prevalence between row variables, the same mark represents no difference between the two variables.

Table 2. Differential KAP variables in different age groups. (n=1301)

Groups and variables	CF (n, (%))	ECC (n, (%))	P
For 3-year-old group (n=415)			
Knowledge			
Q22e Does eat sugar cause dental caries?			
correct answer	160 (46.2) ^a	186 (53.8) ^a	0.017
wrong answer or do not know	43 (62.3) ^b	26 (37.7) ^b	
Practice			
Q3_group Feeding method within 6 months of birth			
complete breastfeeding	122(46.6) ^a	140(53.4) ^a	0.028
mainly breastfeeding	34(47.9) ^{a,b}	37(52.1) ^{a,b}	
complete artificial feeding	23(74.2) ^b	8(25.8) ^b	
mainly artificial feeding	9(64.3) ^{a,b}	5(35.7) ^{a,b}	
half breastfeeding and half artificial feeding	15(40.5) ^{a,b}	22(59.5) ^{a,b}	
Q12 Has your child had a toothache or discomfort in the past 12 months?			
Never	179(54.4) ^a	150(45.6) ^a	0.000
Sometimes	11(18.0) ^b	50(82.0) ^b	
Often	0(0) ^{a,b}	4(100.0) ^{a,b}	
Not clear	13(61.9) ^a	8(38.1) ^a	
Q13 Did your child go to the hospital to see the teeth?			
Yes	24(35.8) ^a	43(64.2) ^a	0.023
Never	179(51.4) ^b	169(48.6) ^b	
For 4-year-old group (n=441)			
Gender			
male	81(37.5) ^a	135(62.5) ^a	0.025
female	61(27.1) ^b	164(72.9) ^b	
Practice			
Q12 Has your child had a toothache or discomfort in the past 12 months?			
Never	130(38.2) ^a	210(61.8) ^a	0.000
Sometimes	10(13.5) ^b	64(86.5) ^b	
Often	0(0) ^b	13(100.0) ^b	
Not clear	2(14.3) ^{a, b}	12(85.7) ^{a,b}	
Q13 Did your child go to the hospital to see the teeth?			
Yes	14(18.4) ^a	62(81.6) ^a	0.004
Never	128(35.1) ^b	237(64.9) ^b	
For 5-year-old group (n=445)			
Knowledge			
Q22d Can bacteria cause dental caries?			
correct answer	79(25.0) ^a	237(75.0) ^a	0.010
wrong answer	9(60.0) ^b	6(40.0) ^b	
do not know	28(24.6) ^a	86(75.4) ^a	
Practice			
Q5 Frequency of eating sweet before going to bed at night			
often	4(12.1) ^a	29(87.9) ^a	0.005
occasionally	56(22.5) ^a	193(77.5) ^a	
never	56(34.4) ^b	107(65.6) ^b	

Q12 Has your child had a toothache or discomfort in the past 12 months?			
Never	102(35.3) ^a	187(64.7) ^a	0.000
Sometimes	10(7.9) ^b	116(92.1) ^b	
Often	0(0) ^b	15(100) ^b	
Not clear	4(26.7) ^{a, b}	11(73.3) ^{a,b}	
Q13 Did your child go to the hospital to see the teeth?			
Yes	9(8.7) ^a	95(91.3) ^a	0.000
Never	107(31.4) ^b	234(68.6) ^b	

CF: caries-free group; ECC: Early Childhood Caries group

^{a, b}: Differences of caries prevalence between row variables, the same mark represents no difference between the two variables.

Table 3. Multivariable logistic regression model for ECC: children aged 3-5 years old, (n=1301).

Variables	OR	95% CI	P
age			
5 years old	2.44	1.79-3.31	0.000
4 years old	2.07	1.55-2.78	0.001
3 years old	1.00	NA	NA
22e_group Does eat sugar cause dental caries?			
Yes	1.49	1.07-2.06	0.017
No or do not know	1.00	NA	NA
Q3 Feeding method within 6 months of birth			
mainly breastfeeding	0.86	0.62-1.18	0.339
complete artificial feeding	0.42	0.25-0.69	0.001
mainly artificial feeding	0.81	0.45-1.47	0.488
half breastfeeding and half artificial feeding	0.84	0.56-1.25	0.378
complete breastfeeding	1.00	NA	NA
Q5 Frequency of eating sweet before going to bed at night			
often	1.81	1.09-3.02	0.023
occasionally	1.32	1.03-1.74	0.027
never	1.00	NA	NA
Q12 Does your child have toothache or discomfort within the past 12 months?			
Sometimes or often	4.70	3.11-7.11	0.000
Not clear	1.24	0.67-2.23	0.487
Never	1.00	NA	NA
Q13 Has your child ever seen a dentist?			
Yes	1.75	1.19-2.55	0.024
Never	1.00	NA	NA

OR: odd rates; CI: confidence interval; NA: not applicable.

Table 4. Multivariable logistic regression model for ECC at different age groups, (n=1301).

model 1 (For 3-year-old group; n=415)			
Variables	OR	95% CI	<i>P</i>
22e_group Does eat sugar cause dental caries?			
Yes	1.77	1.02-3.07	0.043
No or do not know	1.00	NA	NA
Q3 Feeding method within 6 months of birth			
mainly breastfeeding	0.88	0.50-1.53	0.646
complete artificial feeding	0.28	0.12-0.69	0.005
mainly artificial feeding	0.59	0.19-1.83	0.360
half breastfeeding and half artificial feeding	1.27	0.60-2.66	0.53
complete breastfeeding	1.00	NA	NA
Q12 Does your child have toothache or discomfort within the past 12 months?			
Sometimes or often	5.83	2.89-11.77	0.000
Not clear	0.66	0.26-1.68	0.386
Never	1.00	NA	NA
model 2 (For 4-year-old group; n=441)			
Variables	OR	95% CI	<i>P</i>
gender			
male	0.62	0.41-0.94	0.024
female	1.00	NA	NA
Q12 Does your child have toothache or discomfort within the past 12 months?			
Sometimes or often	4.73	2.36-9.49	0.000
Not clear	3.90	0.85-17.80	0.079
Never	1.00	NA	NA
model 3 (For 5-year-old group; n=445)			
Variables	OR	95% CI	<i>P</i>
Q5 Frequency of eating sweet before going to bed at night			
often	3.22	1.03-10.06	0.045
occasionally	1.64	1.03-2.61	0.036
never	NA	1.00	NA
Q12 Does your child have toothache or discomfort within the past 12 months?			
Sometimes or often	4.93	2.39-10.14	0.000
Not clear	1.25	0.38-4.14	0.716
Never	NA	1.00	NA
Q13 Did your child go to the hospital to see the teeth?			
Yes	2.54	1.17-5.53	0.018
Never	NA	1.00	NA

OR: odd rates; CI: confidence interval; NA: not applicable.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AppendixTableS1.docx](#)
- [Equation.docx](#)