

Incidence, Clinical Profile, and Risk Factors for Serious Bacterial Infections in Children Hospitalized With Fever in Ujjain, India

Ashish Pathak (✉ ashish.pathak@ki.se)

Karolinska Institutet <https://orcid.org/0000-0002-7576-895X>

Radika Upadhayay

Department of Pediatrics, Ruxmaniben Deepchand Gardi Medical College

Aditya Mathur

Department of Pediatrics, Ruxmaniben Deepchand Gardi Medical College

Sunil Rathi

Department of Pediatrics, Ruxmaniben Deepchand Gardi Medical College

Cecilia Stålsby Lundborg

Department of Global Public Health, Karolinska Institutet

Research article

Keywords: fever, serious bacterial infection, children, risk factors

Posted Date: December 15th, 2019

DOI: <https://doi.org/10.21203/rs.2.18748/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 21st, 2020. See the published version at <https://doi.org/10.1186/s12879-020-4890-6>.

Abstract

Background Fever is a cause for concern for both parents and the treating pediatrician and a common reason for antibiotic overuse. However, the proportion of children hospitalized for fever with serious bacterial infection (SBI) is uncertain. We aimed to evaluate the epidemiological, clinical, hematological, and biochemical risks for SBI among the children admitted for fever.

Method This prospective study was conducted in a rural teaching hospital in India on consecutive children, aged 3 months–12 years, presenting with fever 100°F (37.7°C) or higher. The presence of SBI was confirmed with one of the following criteria: (a) a positive blood culture; (b) roentgenographically confirmed pneumonia with high titres of C-reactive protein; (c) a culture-confirmed urinary tract infection; (d) enteric fever diagnosed clinically in addition to either a positive blood culture or high Widal titers; and (e) meningitis diagnosed clinically in addition to either a positive blood culture or cerebrospinal fluid culture. A predefined questionnaire was filled.

Results The study included 302 patients of whom 47% presented with SBI. The factors associated with confirmed SBI in bivariate analysis were history of previous hospitalization, history of chronic illness, history of medication in the previous 1 week, a partially immunized child, history of common cold, moderate-grade fever, toxic look, significant lymphadenopathy, absence of BCG scar, delayed development, irritability, breathlessness, respiratory distress, poor feeding, significant weight loss, suspected urinary tract infection, hyponatremia, hypokalemia, and abnormal leucocyte count. The final generalized logistic regression model revealed partially immunized child (RR 4.26), breathlessness (RR 1.80), weight loss (RR 2.28), and suspected urinary tract infection (RR 1.95) as risk factors for the increased risk of SBI.

Conclusion The study identified multiple risk factors for SBI. Pediatricians can be made aware of these risk factors. Further studies are warranted to identify age-specific risk factors for SBI because most clinicians depend on clinical signs and symptoms to identify SBI.

Background

Fever is a common clinical symptom in children and is one of the leading causes for medical consultation and hospital admissions [1]. Globally, 5.4 million children die before the age of 5 years and approximately 50% of the mortality is caused by infectious diseases, many of which present with fever [2].

The underlying infectious disorder causing fever in children can range from mild and self-limiting illness such as upper respiratory tract infection to more serious viral and bacterial illnesses [3]. Distinguishing between the benign self-limiting illness that is manageable at home and those that require hospitalization is the primary challenge for pediatricians and primary care physicians. The diagnosis remains challenging because a wide range of bacterial infections can cause fever in a child. The diagnostic challenge is exacerbated in resource-poor settings in terms of laboratory infrastructure and human

capacity [4]. The overlapping manifestations of serious bacterial infections (SBIs) further complicates diagnosis. The overreliance on clinical diagnosis is one of the reasons for irrational use of antibiotics [5].

In the state of Madhya Pradesh in India, where the present study was conducted, National Family Health Survey-4, undertaken in the year 2015–2016, reported that approximately 71% of the children less than 5 years of age were taken to a health facility for complaints of fever in 2 weeks preceding the survey [6]. Studies from India have reported febrile illness due to SBIs, such as pneumonia [7], typhoid fever [8], urinary tract infections (UTIs) [9], blood-stream infections [10], and meningitis [11]. However, as per our knowledge, there is lack of published studies from India on epidemiological or clinical predictors of SBIs among children hospitalized with fever. Clinical prediction rules complement practice guidelines and help physicians improve clinical decision-making [12, 13]. This study aimed to determine the incidence of SBIs in febrile children admitted in a rural hospital in India and to determine the clinical profile and risk factors associated with SBIs.

Methods

This was a prospective cohort study was done during the twelve months period from July 2015 to June 2016.

Study setting

This study was done in the Pediatrics ward of C.R. Gardi Hospital (CRGH), which is a teaching hospital attached to R.D. Gardi Hospital (RDGMC), Ujjain, in the state of Madhya Pradesh, in central India. Department of Pediatrics, RDGMC has 90 beds, distributed in two wards, out of a total of 630 beds in CRGH. A six-bedded pediatric intensive care unit is also attached to the department.

Provider characteristics

During the study period, the Department of Pediatrics had three units, each having 30 beds, distributed in two wards. Each unit had at least three residents and teaching faculty consisting of a professor, an associate professor, and an assistant professor. The experience of each professor was a minimum of 8 years and that of associate professor and assistant professor at least 5 and 3 years respectively.

Selection of the participants

Consecutive patients aged between 3 months to 5 years, admitted for fever and having an axillary temperature of 100°F (37.7 °C) or more taken with mercury thermometer at the time of admission were included in the study. Children admitted with fever but having chronic co-morbidities: malignancy, renal failure, hepatic failure, congestive cardiac failure, and bone marrow aplasia, children on immunosuppressive drugs such as steroids, and HIV positive children were not included in the study. Children transferred-in with fever from other hospitals with a diagnosed bacterial infection or a laboratory result suggestive of bacterial infection were not included, as it would not have allowed an independent

clinical assessment of cases. Children with osteomyelitis, cellulitis and patients with surgery and trauma were not included as these children are routinely admitted in wards other than pediatrics.

Data collection

All the admitted patients were screened for eligibility by the pediatric resident on-call as soon as possible after admission. The details of the study were discussed with the mother or caregiver accompanying the child fulfilling the inclusion criteria and a written consent obtained. After obtaining consent, a standardized pre-defined questionnaire containing clinical history and physical examination was filled in. The questionnaire also contained epidemiological, clinical, hematological and biochemical parameters of the cases. Patients were provided immediate routine or intensive care as per the department protocol. A senior consultant examined all patients within 24 hours of admission. Investigations were done at the discretion of the senior consultant.

Definitions

The socioeconomic classification of the families of the children was done using modified Kuppuswamy classification [14]. A child was said to have delayed developmental if after assessment the child was delayed for age and sex in any of gross motor, fine motor, language, or social domain. Significant lymphadenopathy was considered if size of inguinal nodes > 1.5 cm in diameter, or/and cervical /axillary > 1 cm in diameter on multiple sites, and/or presence of matted nodes, if lymphadenopathy was associated with focus of infection, and/or with systemic signs and symptoms [3]. Severe acute malnutrition (SAM) was assessed in children between the ages of 6 month to 5 years according to Indian Academy of Pediatrics consensus statement for diagnosis and management of SAM [15]. Suspected urinary tract infection was labeled among those children who were having complaint of burning micturition, pain in abdomen, frequency, urgency and urine routine or children below 2 years of age presenting with fever without identifiable source of infection and microscopy revealed > 15 leucocytes/mm³ [3]. Poor feeding was defined as when an infant has lack of interest in feeding or a problem receiving proper amount of nutrition [3]. Weight loss was defined as loss of > 10% of the actual weight of the individual during a period of 1 month and was labeled only if the weight was documented. Simple febrile seizure was a primary generalized, usually tonic-clonic attack associated with fever with seizures lasting for maximum of 15 minutes and did not recur within 24-hour period [3]. For electrolyte levels age appropriate values were considered [3]. Hyponatremia and hypernatremia were defined as serum sodium concentration below 130 mEq/L and above 150 mEq/L, respectively [3]. Hypokalemia and hyperkalemia were defined as potassium level below 3.5 mEq/L and above 5.5 mEq/L, respectively [3]. Abnormal leucocytes count was defined as leucocyte count in a child up-to 2 year of age < 6000 cells/mm³ and/or > 14000 cells/mm³. In children > 2 year of age, < 4000 cell/mm³ and/or > 12000 cells/mm³ [3].

Outcome measure

The unit of analysis was the child and not the number of febrile episodes. The primary outcome was proportion of children having SBI among the children presenting with fever. The presence of SBI was

confirmed by presence of at least one of the following criteria: 1) blood culture positive; 2) a child was considered to have bacterial pneumonia: if the child presented with breathlessness and had blood culture positive or if along with breathlessness, chest X-ray showed consolidation and C-reactive protein (CRP) value was more than 1000 µg/dl (X-ray chest were reported by a radiologist not part of the study); 3) a child was considered to have UTI if in a toilet trained child with suspected UTI midstream clean catch urine sample was culture positive and in non-toilet trained children with suspected UTI urine obtained by transurethral bladder catheterization was culture positive; all children were screened for UTI based on urine routine done on a spun urine sample 4) a child was diagnosed to have enteric fever when presentation was fever with malaise, headache, abdominal discomfort, coated tongue and Widal test positive with somatic antigen (O) and flagellar antigen (H) titers greater than 320 or blood culture was positive for S Typhi; 5) a child was considered to have bacterial meningitis in presence of clinical features suggestive of meningitis, with either blood culture and/or cerebrospinal fluid culture was positive. A fever was labeled as “viral or no confirmed bacterial infection” when the above diagnosis was ruled out, thus viral fever was a diagnosis of exclusion. The secondary outcome was to determine the epidemiological, clinical, hematological, and biochemical risk factors associated with SBI.

Laboratory methods

Two milliliter (ml) of blood was obtained for hematological and biochemical investigations and 3–4 ml for blood culture. The following investigation were done for all children included in the study: a) complete blood count done using five-part automated coulter counter using fluorescence flow cytometry (XS-800i, Sysmex India Pvt. Ltd, India). Peripheral smear examination was done for leucocyte morphology, immature to mature cell ratio and toxic granules, malarial parasite and red cell morphology; b) a quantitative C- reactive protein (CRP) (Vitros CRP slides, Vitros 250 Chemistry Analyzer, Ortho Clinical Diagnostics, Johnson & Johnson, USA); c) serum electrolytes: serum sodium, potassium, and calcium (Vitros 250 Chemistry Analyzer, Ortho Clinical Diagnostics, Johnson & Johnson, USA); d) Widal test was done using slide and tube agglutination test (Febrile Antigen Set, Span Diagnostic Ltd., India); e) blood and cerebrospinal fluid (CSF) culture was collected under sterile conditions in Bactec Peds Plus/F vial® and pathogens were isolated using the automated BacT/ALERT system (bio-Mérieux, Inc., Marcy l'Étoile, France); e) For doing urine cultures-urine sample was inoculated into the Blood Agar plate and Mac-Conkey agar plate using the semi quantitative method and incubated aerobically at 37 °C for 24 hours. A 0.5 McFarland suspension was prepared from pure culture of uro-pathogens in a nutrient broth and inoculated on Muller-Hinton agar. Antimicrobial susceptibility testing for blood and urine cultures was done using the Kirby Bauer disk diffusion method and results interpreted according to 2018 Clinical and Laboratory Standards Institute guidelines [16]. Extended-spectrum β-lactamase (ESBL) production was detected using a double-disc synergy test [17]. Multidrug-resistant isolates were defined as isolates having co-resistance to at least three antibiotic groups [18].

Sample size calculation

In the reported studies on SBI the cases varied, according to geographical area and the age of children included in the study, between 24 to 40% [19, 20]. We choose a proportion of 30% from above studies to

calculate the sample size. Sample size calculation was done to detect at least 15% difference around proportion of 0.30, with a power of 90 and two-sided alpha of 0.05. The estimated minimum sample size needed was 230 children. Assuming that 20% children will not be able to complete investigations for fever we increased the sample size by a similar proportion to 276 (230×0.2).

Data analysis

Data was entered in EpiData Entry (Version 3.1, Epi Data Software Association, Odense Denmark) and statistical analyses was performed using Stata (Version 13.0 Statacorp. Texas, USA). For continuous variables range, mean and standard deviation (SD) are presented. Categorical independent variables were investigated using Pearson chi-square with the dependent variable being SBI (yes/no). In case of cell count less than five or at least one cell-count equal to zero, the variable was excluded for further analyses. Pearson Chi-square test was used to test for each risk factor's association with SBI.

A generalized linear regression model (GLM) was used to examine the association of independent risk factors responsible for SBI. The independent risk factors were compared between neonates with and without SBI (binary outcome variable). The adjusted relative risk (RR) of SBI was calculated using multivariate predicted marginal proportions for logistic regression models and included the following independent variables as covariates: sex—male versus female; age—in months as continuous; partially immunized child—yes versus no; breathlessness—yes versus no; weight loss—yes versus no; and suspected UTI—yes versus no. The means along with the associated 95% confidence intervals (CI) and p values were reported from GLMs. A p value of < 0.05 was considered significant in the final model. For the final model, model discrimination was done using C-statistics.

Ethical considerations

The Institutional Ethical Committee approved the study (Approval number IEC Ref No 461/2015). The procedures followed were in accordance with the ethical standards set by the institutional ethics committee and with the Declaration of Helsinki.

Results

The total number of children admitted in the Pediatric ward during the study period, patients presenting with fever, number of patients excluded and reasons of exclusion, are shown in Fig. 1.

Causes of SBI

The distribution of diagnosis of 302 children presenting with fever according to presence or absence of SBI is shown in Table 1. Out of 302 febrile children, 142 children had SBI (47%). The SBI based on all positive cultures (blood, urine and CSF) was 30% ($n = 92$) and based on only positive blood culture was 26% ($n = 78$); but varied with clinical diagnosis. The most common diagnosis was bronchopneumonia (20%) with 93% SBI rate. Meningitis and enteric fever and had high SBI rates of 58% and 55%, respectively. The SBI in bronchopneumonia was diagnosed in 57 children. The diagnosis was based on

positive blood cultures in 33% (n = 19/57) and the remaining 67% (n = 38/57) were diagnosed as having SBI based on positive chest X-ray and CRP > 1000 µg/dl. Overall X-ray chest showed consolidation in 93% (n = 53/57) of cases of bronchopneumonia. The remaining cases were diagnosed based on breathlessness and positive blood culture (Table 1). In enteric fever SBI was diagnosed in 22 children. The diagnosis was based on clinical symptoms and a positive blood culture in 45% (n = 10/22) and in the remaining 55% (n = 12/22) based on clinical symptoms and positive Widal test (Table 1). Nearly one-fourth of suspected UTI were diagnosed SBI based on urine culture (Table 1). In our study 29/31 (94%) of the children presenting with severe acute malnutrition (SAM) and fever had confirmed SBI. Majority (86%) had evidence of bronchopneumonia, followed by sepsis (7%), UTI (5%) and meningitis (2%) (Table 1).

Table 1

Distribution of diagnosis of 302 children presenting with fever according to presence or absence of serious bacterial infection

Diagnosis	Total (n)	Outcome variable		Culture positive (n)	Commonest isolated 2 organism
		Serious bacterial infection	Serious bacterial infection		
	302 (%)	No (%)#	Yes (%)#		
Bronchopneumonia*	61(20)	4(7)	57*(93)	19(33)	P. aeruginosa n = 4, K. pneumoniae n = 3,
Viral fever	52(17)	52 (100)	0	0	-
Urinary tract infection	38(13)	29 (76)	9 (24)	9(100)	E. coli n = 7, P. aeruginosa n = 2
Severe acute malnutrition	31(10)	2(6)	29(94)	29(100)	E. coli n = 13, S. aureus n = 9
Febrile convulsion	25(8)	24(96)	1(4)	1(100)	S. aureus n = 1
Enteric fever	40(13)	18(45)	22(55)	10(45)	S. Typhi n = 10
Bronchiolitis	14(5)	14(100)	0(0)	0	-
Meningitis	12(4)	5(42)	7(58)	7(100)	S. pneumonia n = 5, S. aureus n = 2
Failure to thrive	10(3)	1(10)	9(90)	9(100)	P. aeruginosa n = 3, K. pneumoniae n = 2
Upper respiratory tract infection	8(2.5)	8(100)	0(0)	0	-
Infective endocarditis in cardiac disease	5(1.5)	0(0)	5(100)	5(100)	S. pneumonia n = 3, S. epidermidis n = 2,
Miliary tuberculosis	4(1.3)	1(25)	3(75)	3(100)	P. aeruginosa n = 3
Dysentery	2(0.7)	2(100)	0(0)	0	-
% -column percentage, %# -row percentage, * 38 cases were diagnosed X-ray chest positive and CRP > 1000 µg/dl					

Results of blood culture, CSF and urine cultures and antibiotic susceptibility testing

The overall culture positivity rate was 30% (92/302), with Gram-negative preponderance (65%). The most common Gram-negative bacteria isolated were Escherichia coli (n = 25), Pseudomonas aeruginosa (n =

17), followed by *Salmonella typhi* (n = 10) and *Klebsiella pneumoniae* (n = 5). The most common Gram-positive bacteria was *Staphylococcus aureus* (n = 17), followed by *Streptococcus pneumoniae* (n = 7). The in vitro antibiotic susceptibility pattern of the Gram-negative and Gram-positive pathogens are presented in Supplementary Table 1 and Table 2, respectively. High resistance for beta-lactam antibiotics (range 66–100%), ciprofloxacin (40–100%) and all orally used antibiotics was observed in three common Gram negative bacteria viz. *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. The ESBL rates were 75% for *E.coli* and 68% for *Klebsiella*. The MDR rates were 28% for *E.coli* and 33% for *Klebsiella*.

Table 2

Association of serious bacterial infection with co-factors related to demographic, socioeconomic factors, and past history in 302 children admitted with fever in Ujjain, India

Independent Variable	Outcome variable					
	Serious bacterial infection					
	Total(%)# n = 302	No (%) n = 160	Yes (%) n = 142	RR	95%CI	p value
Sex	180(60)	89(49)	91(51)	0	0.64–	0.143
Male	122(40)	71(58)	51(42)	.826	1.06	
Female						
Age in months	106(35)	53(50)	53(50)	R	R	-
Infant (3mo-12mo)	123(41)	68(55)	55(45)	0.808	0.48–	0.425
Toddler (> 12mo-36mo)	73(24)	39(53)	34(47)	0.871	1.36	0.652
Pre-school (> 36 mo-60 mo)					0.47–	
					1.58	
Socioeconomic status	70(23)	37(53)	33(47)	R	R	-
Upper middle	223(74)	119(53)	104(47)	0.97	0.57–	0.941
Lower middle	9(3)	4(44)	5(56)	1.40	1.67	0.636
Lower					0.34–	
					5.66	
H/o hospitalization (past 1 mo)	267(88)	157(59)	110(41)	2.21	1.86–	<
No	35(12)	3(9)	32(91)		2.64	0.001
Yes						
H/o chronic illness	295(98)	159(54)	136(47)	1.85	1.34–	<
No	7(2)	1(14)	6(86)		2.57	0.001
Yes						
H/o any medication (last one week)	266(88)	155(58)	111(42)	2.06	1.70–	<
No	34(11)	4(12)	30(88)		2.50	0.001
Yes						
Immunization status	107(35)	97(91)	10(9)	7.24	3.98–	<
Fully immunized	195(65)	63(32)	132(68)		13.17	0.001
Partially immunized						
H/o malaise	268(89)	146(54)	122(46)	1.29	0.94–	<
No	34(11)	14(41)	20(59)		1.76	0.001
Yes						
H/o fatigue	255(84)	139(56)	116(45)	1.21	0.90–	0.186
No	47(16)	21(45)	26(55)		1.62	
Yes						
H/o lethargy	284(94)	152(54)	132(46)	1.19	0.77–	0.418
No	18(6)	8(44)	10(56)		1.84	
Yes						

Independent Variable	Outcome variable Serious bacterial infection					
H/o restlessness	298(99)	158(53)	140(47)	1.06	0.39–	0.902
No	4(1)	2(50)	2(50)		2.85	
Yes						
H/o common cold	193(64)	128(66)	65(34)	2.09	1.66–	<
No	109(36)	32(29)	77(71)		2.64	0.001
Yes						

Factors associated with SBI- bivariate analysis and multivariate logistic regression analysis

The most common age group of children presenting with fever were those between 13–36 months (n = 123, 41%) with 91(51%) boys and 51(42%) girls. Most (74%) children belonged to lower-middle socio-economic class (Table 2). The duration of hospital-stay of the febrile children having SBI was longer. It was also calculated that with each day increase in stay beyond 7 days, the odds of presence of SBI increased by 1.57 times (95%CI 1.19-2.07; p value < 0.001). The factors on history and physical examination that were significantly associated with SBI on bi-variate analysis are shown in Table 3. The laboratory findings associated with SBI were hyponatremia, hypokalemia and abnormal leucocyte count and are presented in Table 4.

Table 3

Association of serious bacterial infection with signs and symptoms at presentation of 302 children admitted with fever in 302 children admitted with fever in Ujjain, India

Independent Variable	Outcome variable					
	Serious bacterial infection					
	Total(%)# n = 302	No (%) n = 160	Yes (%) n = 142	RR	95%CI	p value
Fever at admission	134(44)	86(64)	48(36)	R	R	-
Low grade fever (99.9–100°F)	154(51)	67(44)	87(56)	2.32	1.44–3.74	<
(37.72–37.77 °C)	14(5)	7(50)	7(50)	1.79	0.593–5.41	0.001
Moderate fever (100.1–103°F)						0.301
(37.83–39.44 °C)						
High grade fever (103.1–105°F)						
(39.50–40.55 °C)						
Toxic look	118(39)	79(67)	39(33)	1.69	1.27–2.25	<
No	184(61)	81(44)	103(56)			0.001
Yes						
Significant lymphadenopathy	245(81)	145(59)	100(41)	1.80	1.45–2.24	<
No	57(19)	15(26)	42(74)			0.001
Yes						
BCG scar	249(82)	151(61)	98(39)	2.10	1.73–2.56	<
Present	53(18)	9(17)	44(83)			0.001
Not present						
Generalized swelling	297(98)	159(54)	138(46)	1.72	1.09–2.71	0.019
No	5(2)	1(20)	4(80)			
Yes						
Independent Variable	Outcome variable					
	Serious bacterial infection					
	Total(%)# n = 302	No (%) n = 160	Yes (%) n = 142	RR	95%CI	p value
Delayed development	284(94)	158(56)	126(44)	2.00	1.62–2.46	<
No	18(6)	2(11)	16(89)			0.001
Yes						

%#-column percentage, %- row percentage, BCG- Bacillus Calmette–Guérin, UTI-Urinary Tract Infection

Independent Variable	Outcome variable					
	Serious bacterial infection					
Irritability No Yes	241(80) 61(20)	143(60) 17(28)	98(41) 44(72)	1.77	1.42–2.20	< 0.001
Seizures No Yes	262(87) 40(13)	134(51) 26(65)	128(49) 14(35)	0.71	0.46–1.11	0.137
Breathlessness No Yes	214(71) 88(29)	142(66) 18(20)	72(34) 70(80)	2.36	1.90–2.93	< 0.001
Respiratory distress No Yes	220(73) 82(72)	140(64) 20(24)	80(36) 62(76)	2.07	1.67–2.57	< 0.001
Poor feeding No Yes	250(83) 52(17)	147(59) 13(25)	103(41) 39(75)	1.82	1.46–2.25	< 0.001
Weight loss No Yes	263(87) 39(13)	157(60) 3(8)	106(40) 36(92)	2.29	1.92–2.72	< 0.001
Vomiting No Yes	251(83) 51(17)	131(52) 29(57)	120(48) 22(43)	0.90	0.64–1.26	0.554
Gastrointestinal complains No Yes	258(85) 44(15)	132(51) 28(64)	126(49) 16(36)	0.74	0.49–1.12	0.159
Suspected UTI No Yes	173(57) 129(43)	133(77) 27(21)	40(23) 102(79)	3.41	2.56–4.55	< 0.001
%#-column percentage, %- row percentage, BCG- Bacillus Calmette–Guérin, UTI-Urinary Tract Infection						

Table 4

Association of serious bacterial infection with laboratory findings at presentation of 302 children admitted with fever in Ujjain, India

Independent Variable	Outcome variable Serious bacterial infection					
	Total(%)# n = 302	No (%) n = 160	Yes (%) n = 142	RR	95%CI	p value
Hyponatremia						
No	255 (84)	144 (56)	111 (44)	1.51	1.18–1.94	0.001
Yes	147 (16)	16 (34)	31 (66)			
Hypernatremia						
No	185 (61)	96(52)	89 (48)	0.94	0.73–1.20	0.636
Yes	117 (39)	64 (55)	53 (45)			
Hypokalemia						
No	267 (88)	149 (56)	118 (44)	1.55	1.19–2.01	0.001
Yes	35 (12)	11 (31)	24 (69)			
Hyperkalemia						
No	272 (90)	143 (53)	129 (47)	0.91	0.56–1.40	0.679
Yes	30(10)	17 (57)	13 (43)			
Hypocalcemia						
No	284 (94)	154 (54)	130 (46)	1.45	1.02–2.06	0.035
Yes	18 (6)	6 (33)	12 (67)			
Abnormal leucocyte count						
No	22 (7)	18 (82)	4 (18)	2.71	1.10–6.62	0.029
Yes	280 (93)	142 (51)	138 (49)			
%#-column percentage, %- row percentage						

The details of the final model are presented in Table 5. The ROC of the final fitted model was 0.964, which reflects excellent model fit. In the final logistic regression model the following variables were found to be statistically significant associated with SBI were: unimmunized versus partly immunized status (RR 4.26), breathlessness (RR 1.80), presence of weight loss (RR 2.28), and suspected UTI (RR 1.95).

Table 5
Multivariate logistic regression model for serious bacterial infection
for 302 children admitted with fever in Ujjain, India

Independent Variable	Outcome variable Serious bacterial infection		
	adjusted RR	95%CI	p value
Sex	0.87	0.62–1.24	0.464
Age in month (continuous)	1.00	0.99–1.01	0.262
Partly immunized child	4.26	2.17–8.37	< 0.001
Breathlessness	1.80	1.25–2.61	0.001
Weight loss	2.28	1.13–3.40	< 0.001
Suspected UTI	1.95	1.32–2.88	0.001

Discussion

This is the first study from central Indian province of Madhya Pradesh that reports the SBI rates among children. The study addresses a significant concern that general practitioner and pediatricians face while treating children presenting with fever. The study identified factors associated with risk for SBI like unimmunized versus partly immunized status (RR 4.26), breathlessness (RR 1.80), presence of weight loss (RR 2.28), and suspected UTI (RR 1.95), which can be looked for in future studies on fever in hospitalized children.

Equal distribution of sex of the children presenting with fever is naturally expected as was found in our study (Table 3). Almost equal distribution of sex was reported in other studies [21, 22]. Most children with SBIs were in the 3–12 month age group (Table 3). Similar findings were reported in an Indian study, which reported 77% of the children with fever were infants [21].

None of the patients in our study diagnosed with bronchiolitis presented SBIs. Similar results were reported in children less than 3 months of age [22]. However, most (93%) patients with radiologically confirmed pneumonia had a confirmed SBI. Similar findings were reported in a study in England, wherein 92% of the patients presented microbiologically and radiologically proven pneumonia [23]. However, another study reported low percentages of SBIs in children with pneumonia [22]. In our study, out of the 38 patients presenting with UTI, only 9 (24%) had confirmed SBI. A low proportion of SBI of up to 18% [20]

was reported among children with UTI. However, other studies have reported high SBI rates of 30% and more [23]. In our study, 29/31 (94%) of the children presenting with severe acute malnutrition (SAM) and fever had confirmed SBI. Majority (86%) of the patients presented evidence of bronchopneumonia, followed by sepsis (7%), UTI, and meningitis. A high proportion of bronchopneumonia responsible for SBI in patients with SAM has been reported by other studies from India, Zambia, and East Africa [24–26]. An East African study reported a higher rate of sepsis in patients with SAM than that in the present study [24]. The synergism between malnutrition and infectious disease is well known. Protein and vitamin deficiency inhibit the formation of specific antibodies and also cause impairment of the pulmonary defense mechanism [27]. The increased incidence and severity of infections in malnourished children is due to limited production and diminished functional capacity of B-cell and T-cell components of the immune system [28]. In our case series, of the 25 patients with simple febrile convulsion, only 1 (4%) had confirmed SBI. A similar low risk for SBI in first-time seizures was reported [22].

One of the problems we faced in the diagnosis of enteric fever was a low rate of culture positivity (19%). Low rates of culture positivity in suspected enteric fever has been reported from other tertiary care hospitals in India. The main cause for such low rates is antibiotic use prior to culture [8].

Compared with patients without SBIs, those with confirmed bacterial infections require a longer hospital stay; this has been reported in our study [29] and other settings [22]. A history of previous hospitalization increased the odds for SBI 15-fold (Table 2). Another study reported a 3-fold increase in the risk of SBI in patients with a history of hospitalization [30]. Furthermore, SBI risk in chronic diseases, such as tuberculosis, HIV infection, and diabetes, was reported [21]. As observed in this study, the increased risk of SBI in partially immunized or unimmunized children compared with fully immunized children have been reported earlier [21, 22]. The chances of infection were the least among adolescents who were fully immunized (9.1%) compared with unimmunized children (33.7%) [31]. Our study found that children with developmental delay had increased risk of SBI. A study showed that 70% of the children with developmental delay had SBI [32]. Possible reasons could be a lack of verbal ability in developmentally delayed children and nonspecific nature of presenting complaints of SBI.

Children with bacterial infections presented common symptoms, such as breathlessness in 80%, poor feeding in 75%, irritability in 75%, weight loss in 92%, suspected UTI in 79%, and common cold in 71% patients (Table 4). A high prevalence (50%-64%) of respiratory symptoms in children with SBI has been reported [32, 33]. Up to 78% of febrile children reported SBI with poor feeding and 76% children reported SBI with irritability [30]. Suspected UTI was reported as the main presenting complaint in approximately 70% of the patients diagnosed with SBI [30]. A sick-looking child is a common general examination finding in children with confirmed bacterial infections [19, 32, 34]. In our study, on general examination, we observed that 83% of the children who had bacterial infection had no BCG scar mark. Another Indian study [21] observed that 60% of the children with bacterial infection were without BCG scar mark.

Hyponatremia, hypokalemia, and hypocalcemia were reported in sick children [35]. However, studies reporting SBI and electrolyte abnormalities are lacking [36]. Abnormal leucocyte counts have been

reported in other studies as an indicator for SBI [22, 37].

Common pathogens identified in the present study are similar to those reported by other studies [5, 8, 38–42]. However, the prevalence of extended-spectrum beta-lactamase and multidrug-resistant pathogens was high but should be interpreted with caution in view of the small sample size. The problem of antimicrobial resistance has been documented in many settings including India, where many reports of high prevalence of antibiotic-resistant bacteria have emphasized commensal bacteria [43] as well as pathogenic bacteria [30].

Conclusions

The study identified multiple risk factors for SBI. Pediatricians can be made aware of these risk factors. The results of this study should help develop clinical prediction rules that can supplement the current clinical practice guidelines. Further studies are warranted to identify age-specific risk factors for SBI as most clinicians depend on clinical signs and symptoms to identify SBIs.

Abbreviations

SBI serious bacterial infection

UTI urinary tract infection

SAM severe acute malnutrition

CRP C-reactive protein

OR odds ratios

CI confidence intervals

ROC receiver-operating-characteristics curves

Declarations

Ethics approval and consent to participate The institutional ethical committee of RD Gardi Medical College, Ujjain, Madhya Pradesh, India approved the study (Approval number IEC Ref No 461). An informed written consent was obtained from parents or guardians on behalf of child participants in the local language, Hindi.

The procedures followed were in accordance with the ethical standards set by the institutional ethics committee and that of the Declaration of Helsinki.

Consent to publish Not applicable.

Availability of data and materials The datasets used and/or analysed during the current study is available from the corresponding author on reasonable request.

Competing interests The authors declare that they have no competing interests.

Funding The study was conducted by post-doctoral LR Akerham's Foundation for Clinical Research grant (2012) to Dr Ashish Pathak from Department of Women and Children's Health, International Maternal and Child Health Unit, Uppsala University, Uppsala, Sweden and internal funds available to Department of Pediatrics, RD Gardi Medical College, Ujjain, India.

Authors' contributions AP, RU, AM, SR, and CSL participated in the conception and design of the study. AP and RU collected the data. AP, RU, and AM performed the statistical analysis and drafted the manuscript. AP, RU, AM, SR, and CSL revised the paper critically for substantial intellectual content. All authors read and approved the final manuscript.

Acknowledgements We are thankful to all the study participants for taking part in the study. We would also like to thank the Dean, Dr MK Rathore, and Medical Director, Dr VK Mahadik, RD Gardi Medical College, Ujjain for the support that was provided us in undertaking the study.

Authors' Information

AP is professor and head of unit in Department of Pediatrics, RD Gardi Medical College. RU is resident doing her MD Pediatrics. AM is a public health expert and research associate in Department of Pediatrics. SR is senior professor in the Department of Pediatrics, RD Gardi Medical College, Ujjain. MD is professor and head of the Department of Pediatrics. CSL is professor and research group leader at Karolinska Institutet, Stockholm, Sweden.

References

1. NICE: Feverish Illness in Children: Assessment and Initial Management in Children Younger than 5 Years. National Institute for Health and Care Excellence Clinical guideline No. 160. 2013 NICE, London. [Available from: <https://www.nice.org.uk/CG047>]
2. Child GBD, Adolescent Health C, Reiner RC, Jr., Olsen HE, Ikeda CT, Echko MM, Ballestreros KE, Manguerra H, Martopullo I, Milllear A et al: Diseases, Injuries, and Risk Factors in Child and Adolescent Health, 1990 to 2017: Findings From the Global Burden of Diseases, Injuries, and Risk Factors 2017 Study. *JAMA Pediatr* 2019, 173(6):e190337.
3. Kliegman R GJJS, Blum N, Shah S, Wilson K, Beherman R (ed.): *Nelson Textbook of Pediatrics*, 21st Edition edn: Elsevier 2019.
4. Kanchanachitra C, Lindelow M, Johnston T, Hanvoravongchai P, Lorenzo FM, Huong NL, Wilopo SA, dela Rosa JF: Human resources for health in southeast Asia: shortages, distributional challenges,

- and international trade in health services. *Lancet* 2011, 377(9767):769-781.
5. Pathak A: Lessons from diagnosis-prescribing and antibiotic resistance surveillance in Ujjain, India : the lull before the storm. Sweden Karoloska Institutet, Stockholm, Sweden; 2012. [Available from: <https://openarchive.ki.se/xmlui/handle/10616/40948>]
 6. NHFS-4: National Family Health Survey-4-Ministry of Health and Family Welfare, Government of India, International Institute for Population Sciences, Mumbai. In.; 2015-16. [Available from:<http://rchiips.org/nfhs/NFHS-4Reports/India.pdf/>].
 7. Singh AJ, Kaur A: Minor injuries in ninth class school children of Chandigarh and rural Haryana. *Indian Pediatr* 1996, 33(1):25-30.
 8. Zaki SA, Karande S: Multidrug-resistant typhoid fever: a review. *J Infect Dev Ctries* 2011, 5(5):324-337.
 9. Goel S, Mukherjee SB: Urinary Tract Infection ,A Tale of 50 Years. *Indian Pediatr* 2016, 53(1):57-58.
 10. Dharmapalan D, Shet A, Yewale V, Sharland M: High Reported Rates of Antimicrobial Resistance in Indian Neonatal and Pediatric Blood Stream Infections. *J Pediatric Infect Dis Soc* 2017, 6(3):e62-e68.
 11. Batra P, Gupta S, Gomber S, Saha A: Predictors of meningitis in children presenting with first febrile seizures. *Pediatr Neurol* 2011, 44(1):35-39.
 12. Leroy S, Bressan S, Lacroix L, Andreola B, Zamora S, Bailey B, Da Dalt L, Manzano S, Gervaix A, Galetto-Lacour A: Refined Lab-score, a Risk Score Predicting Serious Bacterial Infection in Febrile Children Less Than 3 Years of Age. *Pediatr Infect Dis J* 2018, 37(5):387-393.
 13. Nijman RG, Vergouwe Y, Moll HA, Smit FJ, Weerkamp F, Steyerberg EW, van der Lei J, de Rijke YB, Oostenbrink R: Validation of the Feverkidstool and procalcitonin for detecting serious bacterial infections in febrile children. *Pediatr Res* 2018, 83(2):466-476.
 14. Sharma R, Saini NK: A Critical Appraisal of Kuppuswamy's Socioeconomic Status Scale in the Present Scenario. *J Family Med Prim Care* 2014, 3(1):3-4.
 15. Indian Academy of P, Dalwai S, Choudhury P, Bavdekar SB, Dalal R, Kapil U, Dubey AP, Ugra D, Agnani M, Sachdev HP et al: Consensus Statement of the Indian Academy of Pediatrics on integrated management of severe acute malnutrition. *Indian Pediatr* 2013, 50(4):399-404.
 16. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; 24th informational supplement. CLSI M100-S24. Clinical and Laboratory Standards. Wayne, PA: CLSI. In.; 2015.
 17. Jarlier V, Nicolas MH, Fournier G, Philippon A: Extended broad-spectrum beta-lactamases conferring transferable resistance to newer beta-lactam agents in Enterobacteriaceae: hospital prevalence and susceptibility patterns. *Rev Infect Dis* 1988, 10(4):867-878.
 18. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B et al: Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012, 18(3):268-281.

19. De S, Williams GJ, Hayen A, Macaskill P, McCaskill M, Isaacs D, Craig JC: Accuracy of the "traffic light" clinical decision rule for serious bacterial infections in young children with fever: a retrospective cohort study. *BMJ* 2013, 346:f866.
20. Chipwaza B, Mhamphi GG, Ngatunga SD, Selemani M, Amuri M, Mugasa JP, Gwakisa PS: Prevalence of bacterial febrile illnesses in children in Kilosa district, Tanzania. *PLoS Negl Trop Dis* 2015, 9(5):e0003750.
21. Bharti B, Bharti S, Verma V: Severe pneumonia in a remote hilly area: integrated management of childhood illness. *Indian J Pediatr* 2006, 73(1):33-37.
22. Benigno Miguel ME, Herrera Labarca Patricio: Children less than 3 months hospitalised due to acute febrile syndrome: 5 years clinical experience. *Rev chil pediatr [online]* 2015, 86(4):270-278.
23. Nademi Z, Clark J, Richards CG, Walshaw D, Cant AJ: The causes of fever in children attending hospital in the north of England. *J Infect* 2001, 43(4):221-225.
24. Sunguya BF, Koola JI, Atkinson S: Infections associated with severe malnutrition among hospitalised children in East Africa. *Tanzan Health Res Bull* 2006, 8(3):189-192.
25. Kumar R, Singh J, Joshi K, Singh HP, Bijesh S: Co-morbidities in hospitalized children with severe acute malnutrition. *Indian Pediatr* 2014, 51(2):125-127.
26. Irena AH, Mwambazi M, Mulenga V: Diarrhea is a major killer of children with severe acute malnutrition admitted to inpatient set-up in Lusaka, Zambia. *Nutr J* 2011, 10:110.
27. Fonseca W, Kirkwood BR, Victora CG, Fuchs SR, Flores JA, Misago C: Risk factors for childhood pneumonia among the urban poor in Fortaleza, Brazil: a case-control study. *Bull World Health Organ* 1996, 74(2):199-208.
28. Schaible UE, Kaufmann SH: Malnutrition and infection: complex mechanisms and global impacts. *PLoS Med* 2007, 4(5):e115.
29. Lindsjo C, Sharma M, Mahadik VK, Sharma S, Stalsby Lundborg C, Pathak A: Surgical site infections, occurrence, and risk factors, before and after an alcohol-based handrub intervention in a general surgical department in a rural hospital in Ujjain, India. *Am J Infect Control* 2015, 43(11):1184-1189.
30. Gielen AC, McDonald EM, Shields W: Unintentional home injuries across the life span: problems and solutions. *Annu Rev Public Health* 2015, 36:231-253.
31. Abrahamsen SK, Haugen CN, Rupali P, Mathai D, Langeland N, Eide GE, Morch K: Fever in the tropics: aetiology and case-fatality - a prospective observational study in a tertiary care hospital in South India. *BMC Infect Dis* 2013, 13:355.
32. Brent AJ, Lakhanpaul M, Thompson M, Collier J, Ray S, Ninis N, Levin M, MacFaul R: Risk score to stratify children with suspected serious bacterial infection: observational cohort study. *Arch Dis Child* 2011, 96(4):361-367.
33. Hildenwall H, Amos B, Mtove G, Muro F, Cederlund K, Reyburn H: Causes of non-malarial febrile illness in outpatients in Tanzania. *Trop Med Int Health* 2016, 21(1):149-156.

34. Craig JC, Williams GJ, Jones M, Codarini M, Macaskill P, Hayen A, Irwig L, Fitzgerald DA, Isaacs D, McCaskill M: The accuracy of clinical symptoms and signs for the diagnosis of serious bacterial infection in young febrile children: prospective cohort study of 15 781 febrile illnesses. *BMJ* 2010, 340:c1594.
35. Sedlacek M, Schoolwerth AC, Remillard BD: Electrolyte disturbances in the intensive care unit. *Semin Dial* 2006, 19(6):496-501.
36. Gonzalez CF, Finberg L, Bluestein DD: Electrolyte Concentration during Acute Infections. Observations in Infants and Children. *Am J Dis Child* 1964, 107:476-482.
37. De S, Williams GJ, Hayen A, Macaskill P, McCaskill M, Isaacs D, Craig JC: Value of white cell count in predicting serious bacterial infection in febrile children under 5 years of age. *Arch Dis Child* 2014, 99(6):493-499.
38. Larru B, Gong W, Vendetti N, Sullivan KV, Localio R, Zaoutis TE, Gerber JS: Bloodstream Infections in Hospitalized Children: Epidemiology and Antimicrobial Susceptibilities. *Pediatr Infect Dis J* 2016, 35(5):507-510.
39. Musicha P, Cornick JE, Bar-Zeev N, French N, Masesa C, Denis B, Kennedy N, Mallewa J, Gordon MA, Msefula CL et al: Trends in antimicrobial resistance in bloodstream infection isolates at a large urban hospital in Malawi (1998-2016): a surveillance study. *Lancet Infect Dis* 2017, 17(10):1042-1052.
40. Pathak A, Marothi Y, Kekre V, Mahadik K, Macaden R, Lundborg CS: High prevalence of extended-spectrum beta-lactamase-producing pathogens: results of a surveillance study in two hospitals in Ujjain, India. *Infect Drug Resist* 2012, 5:65-73.
41. Rudinsky SL, Carstairs KL, Reardon JM, Simon LV, Riffenburgh RH, Tanen DA: Serious bacterial infections in febrile infants in the post-pneumococcal conjugate vaccine era. *Acad Emerg Med* 2009, 16(7):585-590.
42. Thacker N, Pereira N, Banavali SD, Narula G, Vora T, Chinnaswamy G, Prasad M, Kelkar R, Biswas S, Arora B: Epidemiology of blood stream infections in pediatric patients at a Tertiary Care Cancer Centre. *Indian J Cancer* 2014, 51(4):438-441.
43. Pathak A, Chandran SP, Mahadik K, Macaden R, Lundborg CS: Frequency and factors associated with carriage of multi-drug resistant commensal *Escherichia coli* among women attending antenatal clinics in central India. *BMC Infect Dis* 2013, 13:199.

Figures




Figure 1

Study flow chart showing recruitment process of patients presenting with fever.