

# Determinant of Pneumonia Among Children Aged 2-59 Months in Public Health Facilities in Worabe Town, Southern Ethiopia: Facility-Based Case Control Study

**Roza kumdin Seramo**

Silite Zone Health Department, Southern Ethiopia

**Shikur Mohammed Awol**

Saint Paul's Hospital Millennium Medical College

**Yasin Awol Wabe** (✉ [awolyasin30@gmail.com](mailto:awolyasin30@gmail.com))

Worabe Comprehensive Specialized Hospital

**Musa Mohammed Ali**

Hawassa University, College of Medical and Health Science, School of medical laboratory science

---

## Research Article

**Keywords:** Pneumonia, Determinants, Children 2-59 months, Worabe, Ethiopia

**Posted Date:** October 5th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-938293/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

## Introduction:

Childhood pneumonia remains a major health problem in developing countries, including Ethiopia with significant morbidity and mortality. The determinants of pneumonia in children under –five years of age children are numerous and vary widely vary across the regions of the world. Taking the significance of the problem and variability of risk factors, a study is needed to identify the potential determinants of pneumonia in children under-five children years of age.

## Methods

A facility-based unmatched case-control study was conducted among 435 children (145 cases and 290 controls) aged 2-59 months at public health facilities in Worabe town from December 28, 2016 to January 30, 2017. Data were collected with a pretested interviewer-administered questionnaire, and entered into Epi info and transferred to SPSS version 22 for analysis. Univariate and multivariable logistic regression was employed with a 95% CI, and a *p-value* of < 0.05 was used as a determinant of pneumonia.

## Results

Among the factors assessed in this study, stunting [AOR=3.6, 95% CI: 1.9-6.9], carrying the child on the back during cooking [AOR= 2.0, 95% CI: 1.2-3.2], absence of chimney in the cooking room [AOR= 2.2, 95% CI: 1.3-3.7], having a history of asthma [AOR= 5.0, 95% CI: 2-12], and a previous upper respiratory tract infection [AOR= 3.7, 95% CI:2.3-6.1] were found to be determinants of pneumonia.

## Conclusions

Children with stunting, a previous history of asthma, acute upper respiratory tract infection and carrying the child on back during cooking were at higher risk of pneumonia. Therefore, all health institutions should promote early treatment and provision of health education about the health risk of child exposure to biomass fuel smoke and early health-seeking for childhood illnesses.

## Introduction

Pneumonia is a respiratory tract infection that affects children. *Streptococcus pneumoniae*, *Haemophilus influenzae* type b (Hib), and respiratory syncytial virus are among the common pathogens involved [1]. *S. pneumoniae* is the most common cause of bacterial pneumonia in children under five years of age in the developing world. Microbes that are commonly found in a child's upper respiratory tract can cause lower respiratory tract infection. They may also spread via air-borne droplets from cough or sneeze or may spread through blood, especially during and shortly after birth [2].

Pneumonia is the major killer of children under the age of five years than any other diseases known to affect children. Childhood pneumonia causes substantial mortality and morbidity among children under-five, with developing nations carrying the highest pneumonia burden [3, 4].

Every year, nearly 1 million children die of pneumonia worldwide and approximately about 15% of all deaths occur in children under age five. This means a loss of over 2,500 children lives every day or 100 children every hour. It is most prevalent in sub-Saharan Africa and South Asia, Nigeria, Pakistan, the Democratic Republic of Congo and Ethiopia, accounting for 50% of total deaths [5–7]. In Ethiopia, pneumonia is the leading cause of death, accounting for 18% of all under-five mortality burdens [8]. As a result, Ethiopia ranks the fifth (62 deaths in 1000) among 15 countries in the world and has the highest death rate from

clinical pneumonia in children under five [9]. It also causes millions of morbidity burdens and costs the health services program, as health services are passed on to cure high pneumonia morbidity cases, with an estimated 3,370,000 children encountering pneumonia annually, which contributes to 20% of all causes of deaths killing over 40,000 under-five children every year [10–11].

Ethiopia follows the same strategy and has adopted the World Health Organization (WHO) Integrated Management of Childhood Illness (IMNCI) guidelines since 2001. Moreover, with the objective of increasing access to life-saving interventions, Ethiopia made a policy breakthrough of introducing community-based treatment of pneumonia through health extension workers in 2010 [12]. Since then, 38,000 health extension workers from nearly 15,000 health posts have been equipped with the skills and supplies to treat pneumonia at the community level using the integrated community case management (ICCM) approach [13].

Previous studies indicated that some of most determinants for pneumonia include the educational status of parents, age of the mother, family child caring practice, family income, age and sex of the child, parental smoking or indoor air pollution from biomass fuel smoke, absence window in the kitchen, overcrowding, parental asthma, household history of pneumonia, malnutrition, lack exclusively breastfeeding, lack of zinc supplementation, comorbidity condition such as diarrhea, measles, acute upper respiratory infection (AURTI), previous history of asthma, and wide variation across the regions of the world [1, 10, 14–37].

Contributing factors of pneumonia need to be studied to better inform health care providers and policy-makers to consider pneumonia prevention strategies. Although there are studies conducted in Ethiopia, the risk factors for pneumonia differ in different societies and study populations, and the findings vary [37]. Taking the significance of the problem and variability of risk factors, research is needed to identify the potential determinants of pneumonia in children under five children years of age. Above all, there was no previous study in this study area that could show the determinants of pneumonia. Therefore, the aim of this study was to identify determinants of pneumonia among children 2-59 aged months in public health facilities in Worabe town Sothern Ethiopia.

## **Methods And Materials**

### **Study setting**

The study was conducted in public health facilities in Worabe town, Silte zone, Southern Nation Nationalities and People's Region. Worabe town, the capital city of Silte zone is located 175km and 198km far from Addis Ababa and the regional capital, Hawassa, respectively. The town has a total population of 29,600, of which 4618 are children under five years old, which is projected from the 2007 census report. There are two public health facilities in Worabe town and 6 private clinics. All of them were given IMNCI service according to the town health office report. The town implemented urban health extension programs [38].

### **Study design and period**

An unmatched case-control study was conducted from December 28, 2016, to January 30, 2017.

### **Source population**

All children were aged two months to five years in public health facilities of Worabe town.

### **Study population**

Children who were 2-59 months old visited the selected health facilities during the study period. Cases were children aged 2-59 months who visited pediatric units, registered and classified for pneumonia as defined by the Federal Democratic Republic

of Ethiopian Ministry of Health IMNCI guideline adopted from WHO [39]. A control group was defined as children aged 2-59 months who attended the public health system in Worabe town with a diagnosis of nonpneumonia.

### **Sample size determination**

A sample size of 435 (145 cases and 290 controls) was determined using Epi-Info version 3.5.4 statistical software with assuming two side confidence level (CI) 95%, power=80%, ratio control to case 2:1 and taking a history of AURTI as a predictors of pneumonia with percent of control exposed 22.4%, 1.80 Odds ratio from a case control study [21] and an estimated nonresponse rate of 10%.

### **Sampling technique**

All public health facilities in the town were purposively included based on patient load and the presence of accessible trained staff on IMNCI. The sample size was distributed to each health facility based on the average number of daily case loads.

According to the zonal health bureau Health Management Information System report preceding five months prior to the study, the average number of daily pneumonia patients was seven under-five pneumonia cases per day in Worabe comprehensive specialized hospital and two per day in Worabe Health Center [40]. Based on this, the sample size allocated for the Worabe health center was one case and two controls on a daily basis with a total of 21 cases and 42 controls, and the sample size allocated for Worabe Comprehensive Specialized Hospital (WCSH) was six cases and twelve controls on a daily basis with a total of 124 cases and 248 controls.

**Selection of cases:** All cases (diagnosed and recorded as pneumonia/severe pneumonia) were considered in the study until the required sample size was reached/fulfilled.

**Selection of controls:** As the control-to-case ratio was 2:1, two children who did not have pneumonia and visited selected health facilities for different services at the time of data collection were randomly selected by systematic random sampling after the cases were identified.

### **Eligibility criteria**

#### **Inclusion criteria**

The study included all children who were between 2 and 59 months of age, those who were residents of Worabe town for a minimum of six months, and visited the pediatric unit of WCSH and Worabe Health Center during the study period.

#### **Exclusion criteria**

Children with the following conditions were excluded from the study: cardiac disease, cough that lasted >15 days (suspected of pulmonary tuberculosis), cough because of the recent history of aspiration of a liquid or a foreign body and caregiver who did not have any information about the child at the time of data collection.

#### **Study variables**

The variables included in this questionnaire were both independent and dependent variables for which an association was going to be assessed.

#### **Independent variables**

Sociodemographic factors, Parental factors such as educational and occupational status, parental smoking, age of the mother, family size, and family- caring practice (parental/home maid, place of child during cooking and family income).

Child factors included age and sex, immunization status, a pre-existing illness such as a history of diarrhea, AURTI, and acute lower respiratory tract infection/pneumonia in the last 2 weeks and asthma.

**Environmental factors:** type of fuel used for cooking, crowding status, place of cooking, parental asthma, and household history of tuberculosis and pneumonia

**Nutritional factors in the child:** undernutrition, breastfeeding status of the child for the first 6 months and duration of breastfeeding, age of complementary feeding started and zinc supplementation.

### **Dependent variables**

Pneumonia among children 2-59 months of age.

### **Operational definitions**

**Pneumonia:** A child aged 2-59 months with cough and/or difficult breathing less than two weeks of duration plus fast breathing and/or chest in drawing.

**Fast breathing** is defined as:

- For children in the age interval of 2 months to 11 months, 50 breaths per minute or more
- For children in the age interval of 12 months to 5 years, 40 breaths per minute or more

**History of acute upper respiratory tract infection (AURTI):** a child who had a history of ear infection, common cold, tonsillitis, or pharyngitis in the last fifteen days prior to data collection.

### **Data collection tools and procedures**

A structured questionnaire was developed based on a review of previously published studies and adapted to the local situation with certain modifications [17-28]. The questionnaires included information on the possible risk factors for pneumonia, including sociodemographic factors, environmental/home-based factors, nutritional factors, immunization status, pre-existing illness and child care practices. Data were collected by IMNCI-trained nurses working in under-five clinics who received two days of training regarding the research. After the study participants were identified as cases and controls, mothers/primary care takers were interviewed based on the interviewer-administered pretested structured questionnaire, but the data collectors were blinded to the status of the respondent.

**Anthropometric measurements:** Weight and height of the child were taken at the beginning of the interview by data collectors. A suspended scale of 25 kg capacity graduated at 0.1 kg was used for weighing infants and children. After taking the weight of every child, the scale was checked for its accuracy. The scale reading was taken to the nearest 0.1 kg. Length measurements in the lying position were taken for children less than two years of age, and height measurements were taken for children 2-5 years of age. The anthropometric data were analysed in terms of weight for age, length for age, and weight for length using WHO Anthrosoftware to prepare for SPSS. The WHO (2006) growth standard was used to report anthropometric measurements result by Z- score, and the global acute malnutrition standard was used to classify the child's nutritional status as normal, stunted, wasted or underweight.

### **Data quality management**

Before conducting the study, data collectors were trained for two days. The questionnaire was pretested on a 5% sample size at Kbit Primary Hospital to ensure the validity and reliability of the survey tools. After collecting the pretest data, it was checked for potential problems related to the tool, such as any difficult question that was understandable or unclear to reply and corrective measures were taken.

### **Data processing and analysis**

The collected data were checked for completeness, coded and entered into Epi Info version 7 and exported to the statistical package for social sciences (SPSS) version 22 for analysis. The entered data were cleaned and checked for consistency and extent of outliers. Different statistical assumptions and appropriate corrections were made prior to analysis. Descriptive analyses were carried out for each of the independent variables. Binary logistic regression analysis was used to test the association between the independent and dependent variables. Bivariate analysis was performed for each of the independent variables with the outcome variable. Variables that had a  $p$ -value  $< 0.2$  on bivariate analysis were taken as candidates for multivariable logistic regression model analysis to identify their independent effects. Finally, variables with  $p$ -values less than 0.05 on multivariable logistic regression analysis were considered statically significant factors for the outcome variable. The strength of the association between the dependent variable and independent variables was expressed by odds ratios (ORs) with 95% confidence intervals.

## Results

### Socio-demographic characteristics

A total of 435 (145 cases and 290 controls) children aged 2 to 59 months participated in the study, making the respondent rate 100% for both groups. The highest proportion of cases (57.2%) and controls (62.4%) were in the age group of 2-11 months, with mean ages of 14.6 (SD± 12.8) and 12.6 months (SD± 12.7) for cases and controls, respectively. Approximately 52% of cases and 51% of controls were male. In the bivariable logistic regression analysis, none of the sociodemographic variables except family size illustrated an association with the occurrence of pneumonia in children ( $p=0.1$ ) at a  $p$ -value  $<0.2$  significance level (Table 1).

Table 1

Sociodemographic characteristics of children aged 2 to 59 months in public health facilities in Worabe town, Silte zone, 2017(N =145 cases and 290 controls).

Variables	Case		Control		Total		P-value	COR(95% C.I)
	Frequency	percent	Frequency	percent	frequency	percent		
<b>Age of the child</b>								
2-11 months	83	57.2	181	62.4	264	60.7%	0.200	0.7[0.4,1.2]
12-23months	31	21.4	61	21	92	21.1%	0.500	0.8[0.4,1.5]
24-59months	31	21.4	48	16.6	79	18.2%		1
<b>Sex of the child</b>								
Male	76	52.4	148	51	224	51	0.800	1.2(0.7,1.6)
Female	69	47.6	142	49	211	49		1
<b>Age of the mother</b>								
18-24	38	26.2	95	32.8	133	31	0.5	0.8(0.5,1.5)
25-31	79	54.5	138	47.6	217	50	0.6	1.2(0.7,1.9)
≥32	28	19.8	57	19.7	85	19		1
<b>Educational status of mother</b>								
Not attended formal education	51	35.2	80	27.6	131	30.1	0.7	1(0.6,2.2)
Primary	51	35.2	130	44.8	181	41.6	0.29	0.7(0.4,1.3)
Secondary	24	16.6	46	15.9	70	16	0.86	0.9(0.4,1.9)
Higher education	19	13.1	34	11.7	53	12.2		1
<b>Educational status of father</b>								
Not attended formal education	44	30.3	71	24.5	115	26.4	0.4	1.3(0.7,2.4)
Primary	53	36.6	121	41.7	174	40	0.8	0.9(0.5,1.6)
Secondary	21	14.5	41	14.1	62	14.3	0.8	1(0.5,2.2)
Higher	27	18.6	57	19.6	84	19.3		1

<b>Mother's occupation</b>												
House wife	88	60.7	204	70.3	292	67.2	0.2	0.5(0.2,1.4)				
Civil servant	17	11.7	33	11.4	50	11.5	0.4	0.6(0.2,1.9)				
Merchant	33	22.8	45	15.5	78	17.9	0.7	0.8(0.3,2.5)				
Others	7	4.8	8	2.8	15	3.4		1				
<b>Father's occupation</b>												
Merchant	51	35.2	124	42.8	175	40.3	0.9	0.9(0.5,1.9)				
Daily labourers	44	30.3	71	24.4	115	26.4	0.3	1.5(0.7,2.9)				
Civil servant	31	21.4	50	17.2	81	18.6	0.3	1.5(0.7,2.9)				
Others	19	13.1	45	15.5	64	14.7		1				
<b>Family monthly income</b>												
<1000	42	29	47	16	23	15.9	64	22.1	87	20	0.4	0.8(0.4,1.4)
1000-1500	27	18.6	39	13.4	66	15.2	0.2	1.4(0.8,2.7)				
1600-2500	53	36.6	99	34.1	152	34.9	0.7	1.1(0.7,1.8)				
>2500	42	29	88	30.3	130	29.9		1				
<b>Family size</b>												
2-3	26	17.9	69	23.8	95	21.8	0.1*	0.8(0.5,1.3)				
4-5	67	46.2	135	46.6	202	46.5	0.4	0.6(0.4,1.1)				
6 and above	52	35.9	86	29.7	138	31.7		1				
<b>Family child care practice</b>												
Parental	116	80	228	78.6	344	79	0.7	1(0.7,1.8)				
Home maid	29	20	62	21.4	91	21		1				
<b>Key:</b> *Candidate variable for multivariate analysis at P<0.2, COR: crude odds ratio, CI: confidence interval, 1: reference group.												

## Nutritional, pre-existing illness and vaccination status of the children

The majority (79.3%) of cases and 85.5% of controls were exclusively breastfed for the 1st six months of life. Regarding the nutritional status of the children, 11.9% cases and 9% controls, 26.9% cases and 18.6% controls were underweight and stunted respectively. More than 90% of cases and 95% controls received pentavalent and pneumococcal conjugate vaccines.

The study indicated that, variables like Zinc supplementation ( $p=0.01$ ), stunting ( $p=0.001$ ), wasting ( $p=0.06$ ), exclusive breastfeeding for the first 6 months ( $p=0.1$ ), child with previous history of AURTI ( $p<0.001$ ) and diarrhoea ( $p=0.06$ ) in the preceding two weeks and history of asthma in the child ( $p<0.001$ ) were associated with pneumonia in the bivariate logistic regression at  $p<0.2$  (Table 2).

Table 2

Nutritional status, vaccination status and pre-existing illness among children aged 2-59 months in public health facilities in Worabe town, Silte zone, 2017 (N = 145 cases and 290 controls).

Variables	Case		Control		Total		P- value	COR(95% C.I)
	frequency	percent	Frequency	percent	frequency	percent		
<b>Breast feeding status of the child during the first 6 months</b>								
Exclusive BF	115	79.3	247	85.2	362	83.2	0.1*	1.2(0.9,1.6)
Mixed feeding	30	20.7	43	14.8	73	16.8		1
<b>Zinc Supplementation</b>								
Supplemented	56	38.6	77	26.6	133	30.6	0.01*	1.7[1.1,2.7]
Not supplemented	89	61.4	213	73.4	302	69.4		1
<b>Weight for age</b>								
Underweight	17	11.7	26	9	43	9.9	0.4	1.3(0.7,2.6)
Normal	128	88.3	264	91	392	90.1		1
<b>Height for age</b>								
Stunted	36	24.8	23	7.9	59	13.6	0.00*	3.8(2.2,6.8)
Not stunted	109	75.2	267	92.1	376	86.4		1
<b>Weight for height</b>								
Wasted	29	20	38	13.1	67	15.4	0.06*	1.7(0.9,2.8)
Not wasted	116	80	252	86.9	368	84.6		1
<b>Dose the child received Pentavalents vaccine</b>								
Yes	134	92.4	277	95.5	411	94.5	0.2	0.6(0.3,1.3)
No	11	7.6	13	4.5	24	5.5		1
<b>Get measles vaccine</b>								
Yes	76	52.4	128	44.1	203	46.7	0.70	1.5(0.9,2.3)
No	13	9	28	9.7	50	11.5	0.50	1.3(0.7,2.5)
N/A(age <9 months)	56	38.6	134	46.2	182	41.8		1
<b>Diarrhoea in last 2 weeks</b>								
Yes	44	30.3	54	18.6	98	22.5	0.06*	1.9[1.2,3]
No	101	69.7	236	81.4	337	77.5		1
<b>AURTI in the last 2 weeks</b>								
Yes	68	46.9	51	17.6	119	27.4	0.00*	4[2.7,6.5]
No	77	53.1	239	82.4	316	72.6		1

**Key:** \*Candidate variable for multivariate analysis at P<0.2, COR: crude odds ratio, CI: confidence interval, 1: reference group.

Variables	Case		Control		Total		P- value	COR(95% C.I)
	frequency	percent	Frequency	percent	frequency	percent		
<b>Previous history of asthma in the child</b>								
Yes	23	15.9	10	3.4	33	7.6	0.00*	5.3[2.4,11.4]
No	122	84.1	280	96.6	402	92.4	1	
<b>Pneumonia among family in last 2 weeks</b>								
Yes	34	23.4	61	21	95	21.8	0.6	1.2(0.7,1.9)
No	111	76.6	229	79	340	78.2	1	
<b>Parental asthma</b>								
Yes	7	4.8	13	4.5	20	4.6	0.8	1.1(0.4,2.8)
No	138	95.2	277	95.5	415	95.4	1	
<b>Key:</b> *Candidate variable for multivariate analysis at $P < 0.2$ , COR: crude odds ratio, CI: confidence interval, 1: reference group.								

## Environmental factors associated with pneumonia

More than half of the cases (57.2%) and controls (54.8%) lived in households that used wood as a source of cooking fuel. The highest proportion of cases (63.4%) and controls (63.8%) had mothers cooking their food in the kitchen. Approximately 61% of cases and 44% controls stayed near or on their mothers'/care takers back while she was cooking. The bivariate logistic regression analysis revealed that crowding ( $p=0.04$ ), type of cooking fuel ( $p=0.02$ ), absence of chimneys in the cooking room ( $p < 0.01$ ) and place of child during cooking ( $p < 0.01$ ) had significant associations with pneumonia among children at the  $p < 0.2$  significance level (Table 3).

Table 3

Environmental characteristics of children aged 2-59 months in public health facilities in Worabe town, Silte zone, 2017 (N = 145 cases and 290 controls).

Variables	Case		Control		Total		P-value	COR(95% C.I)
	frequency	percent	Frequency	percent	frequency	percent		
<b>Type of cooking fuel mostly used the family</b>								
Wood	83	57.2	159	54.8	242	55.6	0.40	1.4(0.8,2.3)
Charcoal	21	14.5	41	14.1	62	14.2	0.20	1.4(0.7,2.6)
Kerosene	10	6.9	8	2.8	18	4.1	0.02*	3.3(1.2,9.1)
Electricity	31	21.4	82	28.3	113	26		1
<b>Place of cooking the family mainly used</b>								
In living room	53	36.6	105	36.2	158	36.3	0.70	1.1(0.7,1.8)
In kitchen attached to the living room	37	25.5	64	22.1	101	23.2	0.40	1.3(0.8,2.1)
In separate kitchen	55	37.9	121	41.7	176	40.5		1
<b>Chimney in the cooking room</b>								
No	101	69.7	140	48.3	241	55.4	<0.01*	2.5(1.6,3.7)
Yes	44	30.3	150	51.7	194	44.6		1
<b>Wind in the living room</b>								
Yes	124	85.5	261	90	385	88.5	0.20	1.5(0.8,2.8)
No	21	14.5	29	10	50	11.5		1
<b>Place of the child during cooking(usual location)</b>								
Near or carried on back	93	64.1	127	43.8	220	50.6	<0.01*	2.3[1.5,3.5]
Outside of cooking room	52	35.9	163	56.2	215	49.4		1
<b>Smoker among household residents</b>								
Yes	11	7.6	19	6.6	30	6.8	0.70	1.2(0.5,2.5)
No	134	92.4	271	93.4	405	93.2		
<b>Crowding status</b>								
Crowded	60	41.4	91	31.4	151	35	0.04*	1.5[1.1,2.3]
Not crowded	85	58.6	199	68.6	284	65		1
<b>Key:</b> *Candidate variable for multivariate analysis at P<0.2, COR: crude odds ratio, CI: confidence interval, 1: reference group.								

## Factors associated with pneumonia among children aged 2-59 months

Logistic regression analysis showed that stunting, previous history of asthma diagnosis in the child, place of the child during cooking, absence of chimney in the cooking room and history of AURTI in the last 2 weeks prior to data collection were determinants of pneumonia. The findings revealed that the odds of stunting were 3.6 times higher among children with

pneumonia than controls (AOR=3.6, 95% CI: 1.8, 7; P<0.001). On the other hand, the odds of carrying the child on the back while cooking/besides the cooking mother were twice as high as those of compared to children outside of cooking the room among cases than controls (AOR= 2, 95% CI: 1.2, 3.3; P=0.006).

Similarly, the odds of children living in households with no chimney in the cooking room were 2.2 times higher among cases than controls (AOR= 2, 95% CI: 1.3, 3.7; P=0.003).

The odds of having a previous history of asthma were fivefold higher among cases than controls (AOR= 5, 95% CI: 2, 12; P<0.001). Likewise, the odds of having a history of AURIT in the last 2 weeks prior to data collection were 3.7 times higher among children with pneumonia than among children without pneumonia (AOR= 3.7, 95% CI: 2.3, 6.1; P<0.001) (Table 4).

Table 4

Determinants of pneumonia among children aged 2-59 months in public health facilities in Worabe town, Southern Ethiopia, 2017 (N = 145 cases and 290 controls).

Variables	Cases No (%)	Controls No (%)	COR (95% C.I.)	AOR (95% C.I.)	P-value
<b>Breast feeding status of the child during the first 6 months</b>					
Exclusive BF	115(79.3%)	247(85.2%)	1.5(0.9,1.6)	0.8(0.4,1.5)	0.4
Mixed feeding	30(20.7%)	43(14.8%)	1.0	1	
<b>Zinc supplementation</b>					
Supplemented	56(38.6%)	77(26.6%)	1.7[1.2,2.7]	1.4(0.8,2.3)	0.2
Not supplemented	89(61.4%)	213(73.4%)	1.0	1	
<b>Height for age</b>					
Stunted	36(24.8%)	23(7.9%)	3.8[2.2,6.8]	3.6[1.8,7]*	<0.001
Not stunted	109(75.2%)	267(92.1%)	1.0	1	
<b>Crowding status</b>					
Crowded	60(41.4%)	91(31.4%)	1.5[1.1,2.3]	1.2(0.7,2)	0.6
Not crowded	85(58.6%)	199(68.6%)	1.0		
<b>Place of child during cooking</b>					
Near or carried on back	93(64.1%)	127(43.8%)	2.3[1.5,3.5]	2.0[1.2,3.3]*	0.006
Outside of cooking room	52(35.9%)	163(56.2%)	1.0	1	
<b>Type of cooking fuel mostly used the family</b>					
Wood	83(57.2%)	159(54.8%)	1.4(0.8,2.3)	1.3(0.7,2.4)	0.500
Charcoal	21(14.5%)	41(14.1%)	1.4(0.7,2.6)	0.9(0.4,1.9)	0.800
Kerosene	10(6.9%)	8(2.8%)	3.3(1.2,9.1)	2.3(0.7,7.3)	0.200
Electricity	31(21.4%)	82(28.3%)	1.0	1	
<b>Weight for height</b>					
Wasted	29(20%)	38(13.1%)	1.7(0.9,2.8)	1.4(0.7,2.7)	0.300
Not wasted	116(80%)	252(86.9%)	1.0	1	
<b>Family size</b>					
2-3	26(17.9%)	69(23.8%)	0.6(0.4,1.1)	0.7(0.3,1.6)	0.400
4-5	67(46.2%)	135(46.6%)	0.8(0.5,1.3)	0.9(0.5,1.6)	0.700
6 and above	52(35.9%)	86(29.7%)	1.0	1	
<b>Chimney in the cooking room</b>					
No	101(69.7%)	140(48.3%)	2.5(1.6,3.7)	2.2[1.3,3.7]	0.003*
Yes	44(30.3%)	150(51.7%)	1.0	1	

**Key:** \*\*Variable significant at P<0.05, COR: crude odds ratio, AOR: adjusted odds ratio, CI: confidence interval, 1: reference group.

Variables	Cases No (%)	Controls No (%)	COR (95% C.I.)	AOR (95% C.I.)	P-value
<b>History of diarrhoea in the last 2 weeks</b>					
Yes	44(33.3%)	54(18.6%)	1.9[1.2,3]	1.5(0.9,2.7)	0.1
No	101(69.7%)	236(81.4%)	1.0	1	
<b>History of AURTI in the last 2 weeks</b>					
Yes	68(46.9%)	51(17.6%)	4[2.7,6.5]	3.7[2.3, 6.1]	<0.001*
No	77(53.1%)	239(82.4%)	1.0	1	
<b>History of asthma in the child</b>					
Yes	23(15.9%)	10(3.4%)	5.3[2.4,11.4]	5.0[2, 12]	<0.001*
No	122(84.1%)	280(96.6%)	1.0	1	
<b>Key:</b> **Variable significant at P<0.05, COR: crude odds ratio, AOR: adjusted odds ratio, CI: confidence interval, 1: reference group.					

## Discussion

In this study, preexisting respiratory illnesses, such as a history of asthma in the child and a previous history of AURTI, undernutrition (stunting), absence of chimney in the cooking room and carrying the child on back during cooking, were found to be significant risk factors associated with pneumonia among children aged 2-59 months.

The odds of carrying the child on the back while cooking/besides the cooking mother was 2 times greater than that of children outside of the cooking room among cases than controls. This might be due to indoor air pollution from biomass fuel smoke, as the majority of the households use wood and charcoal for cooking with no chimneys in the cooking area. This result is consistent with a cross-sectional study conducted in north west Ethiopia that found that a child who carries back during cooking was five times more likely to develop pneumonia than their counterparts [17]. Similar findings were obtained in different countries that reported indoor air pollution as risk factor for pneumonia [15, 25, 28, 29, 30]. However, two studies conducted in Ethiopia contradicted the current study results (those reported an absence of a significant association between the place of the child during cooking and the occurrence of pneumonia in children) [18, 23]. The variation of the results could be due to differences in study setting and methodology.

Stunting was also another factor identified to be significantly associated with pneumonia, indicating that the odds of stunting were 3.6 times higher among children with pneumonia than controls, which this is in line with a study conducted in Northwest Ethiopia and Bangladesh [17, 19]. The possible reason might be that stunting shows long-term malnutrition, which weakens the child's immunity and makes the child vulnerable to pneumonia. From different prospects of different studies, malnutrition (undernutrition) weakens the respiratory muscles needed to clear secretions in the respiratory tract, which intern predisposes to pneumonia [1, 24, 41].

In this study, there was the strongest association between pneumonia and a history of asthma, indicating that children who had a previous history of asthma had approximately fivefold increased odds of developing pneumonia compared to their counterparts. According to the WHO 2008 bulletin report, concomitant diseases such as asthma were likely risk factors for the occurrence of childhood pneumonia [15]. Other studies conducted in different parts of the world also reported that a history of asthma is associated with the occurrence of pneumonia episodes [21, 36]. In a study conducted in the Philippines, 55.4% of children who developed consolidated pneumonia had asthma as an underlying illness [27]. However, in the present study, it was unable to conclude whether uncontrolled asthma or an underlying asthmatic condition was a risk factor for pneumonia or whether pneumonia was a trigger of asthma attack. Further study is need to define a causal relationship between asthma and pneumonia.

Finally, a history of AURTI in a child in the last two weeks preceding the current pneumonia was identified to put a child 3.7 times at risk of pneumonia compared to their counterpart. This is in line with the study conducted in southwest Ethiopia, which indicates that the risk of pneumonia is more than 5 times for children who had previous AURTIs [23], and a similar finding in the Netherlands showed a strong relation between the occurrence of community-acquired pneumonia and an increasing number of previous AURTIs (the risk of pneumonia was 2.46 times & 1.8 times more likely if the child had three and two episodes of AURTIs respectively, in the past) [21]. However, a study conducted in Ethiopia reported no significant association between occurrences of pneumonia and preceding infection of AURTI in children [22]. A possible explanation might be that AURTI increases susceptibility to infections that intern predispose patients to pneumonia or that there might be descending infection from the upper to lower respiratory tract.

### **Limitations of the study**

The diagnosis of pneumonia was based on the clinical WHO IMNCI classification guidelines, which introduced misclassification recall bias; the study did not consider private health facilities and did not consider the complete population.

## **Conclusions**

This study suggests that children stunting, a previous history of asthma and upper respiratory tract infection, the absence of chimney in the cooking room and carrying the child on back during cooking; are strongly associated with an increased risk of childhood pneumonia. Therefore, the zonal health bureau, in collaboration with the agricultural office needs to improve the nutritional status of children, and the town administrative health office, in collaboration with the health facilities should provide health education to the community about the health risk of child exposure to biomass fuel smoke and to have chimneys in the cooking room.

## **Abbreviations**

AURTI  
Acute Upper Respiratory Tract Infection  
ICCM  
Integrated Community Case Management  
IMNCI  
Integrated Management of Neonatal and Childhood Illness  
WCSH  
Worabe Comprehensive Specialized Hospital  
WHO  
World Health Organization.

## **Declarations**

### **Competing interests**

The authors declare that they have no competing interests

### **Funding**

No funding was received for this study.

### **Availability of data and materials**

All necessary data are available within the paper.

## Ethical approval and consent to participant

The study was ethically approved by the Institutional Review Board of Saint Paul's Millennium Medical College Department of Public Health. An official permission letter was obtained from the study site. The objectives, expected outcomes, benefits, and risks of the study were explained to mothers/guardians/caregivers of the study participants. Data were collected after written informed consent was obtained. The study was conducted in accordance with the Declaration of Helsinki.

## Consent for publication

Not applicable

## Authors' contributions

<sup>RKS</sup> Designed the study, was involved in data collection, data analysis and interpretation, wrote the paper and prepared the manuscript for publication. <sup>YAW&MMA</sup> Data analysis; and interpretation and prepared the initial draft of the manuscript for publication. <sup>SMA</sup> Had oversight of all the stages of the research, critically reviewed the final draft of the paper and the prepared manuscript for publication. The final manuscript was read and approved by all authors.

## Acknowledgement

We would like to thank the Zone Health Bureau, Worabe Health Centre, Worabe Comprehensive Specialized Hospital, Saint Paul's Hospital Millennium Medical College Department of Public Health for facilitating the study. We also acknowledge all study participants for their willingness to take part in the study.

## References

1. WOH and UNICEF. Pneumonia and Diarrhea: Tackling the deadliest diseases for the world' poorest children. New York, June 2012.
2. United Nation Children's Fund: World Pneumonia Day (November 12), Facts about Childhood Pneumonia. New York, 2015.
3. Li Liu, Hope, L. *et al.* Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000., **379**, 2151–2161 (2012).
4. Wang, H. *et al.* Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990-2013: a systematic analysis for the Global Burden of Disease Study. 2013. *Lancet* 2014; 384: 957–79
5. World Pneumonia day: Universal access for pneumonia prevention and care, 2014.
6. WHO media centre: pneumonia fact sheet, Sep 2016
7. UNICEF: EVERY BREATH COUNTS, Every 35 seconds, a child dies from pneumonia. Addis Ababa, Ethiopia – January 31, 2016
8. World Health Organization and UNICEF. *Fulfilling the Health agenda for the Women and Children countdown to 2015: Maternal, Newborn and child survival. The 2014 report* (Geneva, WHO,, 2014).
9. United Nations Children's Fund. *Pneumonia and Diarrhea Progress Report* (IVAC, Bangladesh, 2013).
10. Christa, L. F. *et al.* *Childhood Pneumonia and Diarrhoea Global burden of childhood pneumonia and diarrhoea Lancet*, **381**, 1405–1416 (2013).
11. World Health Organization and UNICEF & Handbook *IMCI, Integrated management of childhood illness* (WHO, Geneva, 2005).
12. Federal Ministry of Health (FMOH): National plans on Integrated Community Case management of common childhood illness. Ethiopia, 2010.
13. United Nation Children's Fund, Ethiopia: Central Data Base, October 2014.

14. Levine, O., Dinleyici, E. C. & Pneumonia The forgotten killer of children. *Annotated*. 2010);4(10);27–28.
15. Rudan, I., Boschi-Pinto, C., Biloglav, Z., Mulholland, K. & Campbell, H. Epidemiology and Etiology of childhood pneumonia. *Bull World Health Organ*, **86** (5), 408–416 (2008).
16. Rudan, I. *et al.* Epidemiology and aetiology of childhood pneumonia in 2010: estimates of incidence, severe morbidity, mortality, underlying risk factors and causative pathogen in 192 countries. *Journal of Global Health*. 2013;3(1);1–14, . Doi: 10.7189/jogh.03.010401.
17. Abeje, G., Wubshet, M. & Asres, G. Prevalence of pneumonia among under-five children in Este town and the surrounding rural kebele, Northwest Ethiopia; A community based cross-sectional study. *Science Journal of Public Health*, **2** (3), 150–155 (2014).
18. Teshome, A. Prevalence of pneumonia and factors associated among children 2-59 months old in Wondo Genet district, Sidama zone, SNNPR, Ethiopia. *Curr Pediatr Res*, **21** (1), 19–25 (2017).
19. Azad k. Risk factors for acute respiratory infections (ARI) among children under five years in Bangladesh. *Journal of Scientific Research*, **1** (1), 72–81 (2009).
20. Fatmi & White, F. A comparison of 'cough and cold' and pneumonia: risk factors for pneumonia in children under 5 years revisited. *International Journal of Infectious Diseases*, **6** (4), 295–301 (2002).
21. Teepe, J., Grigoryan, L. & Verheij, T. J. Determinants of community-acquired pneumonia in children and young adults in primary care. *Eur Respir Journal*. 2010;35(5);1113–1117
22. Dadi, A. F., Kebede, Y. & Birhanu, Z. Determinants of pneumonia in children aged two months to five years in urban areas of Oromia Zone, Amhara Region, Ethiopia. *Open Access Library Journal*. 2014;1–10: e1044. <http://dx.doi.org/10.4236/oalib.1101044>.
23. Geleta, D., Tessema, F. & Ewnetu, H. Determinants of community acquired pneumonia among children in Kersa District, Southwest Ethiopia: Facility based case control study. *J Pediatr Neonatal Care*. 2014;5(2);00179. DOI: 10.15406/jpnc.2016.05.00179
24. Pore, P. D., ghattargi, C. H. & Rayate, M. V. Study for risk factors of acute respiratory infection (ARI) in under five in solapur. *National Journal of Community Medicine*. 2010; 1(2).
25. Taksande, A. M. & Yeole, M. Risk factors for acute respiratory infection (ARI) in under-fives in a rural hospital of Central India. *J Pediatr Neonat Individual Med*, **5** (1), e050105 (2016).
26. Gedefaw, M., Berhe, R., Gedefaw, M. & Berhe, R. Determinates of Childhood Pneumonia and Diarrhea with Special Emphasis to Exclusive Breastfeeding in North Achefer District, Northwest Ethiopia: A case control study. *Open Journal of Epidemiology*, **5** (2), 107–112 (2015).
27. Kosai, H. *et al.* Incidence and risk factors for childhood pneumonia-Like Episodes in Biliran Island, Philippines A community-based study. *PLoS One*, **10** (5), e0125009 <https://doi.org/10.1371/journal.pone.0125009> (2015).
28. Dherani, M. *et al.* Indoor air pollution from unprocessed solid fuel use and pneumonia risk in children aged under-five years: a systematic review and meta-analysis. *Bulletin World Health Organ*, **86**, 390–398 (2008).
29. Bautista, L., Correa, A., Baumgartner, J., Breyse, P. & Matanoski, G. Indoor charcoal Smoke and acute respiratory infections in Young Children in the Dominican Republic. *American Journal of Epidemiology*, **169**, 572–580 (2009).
30. Bruce, N. *et al.* Pneumonia case-finding in the RESPIRE Guatemala indoor air pollution trial: standardizing methods for resource-poor settings. *Bull World Health Organ*, **85**, 535–544 (2007).
31. Roux, D. M., Myer, L., Nicol, M. P. & Zar, H. J. Incidence and severity of childhood pneumonia in the first year of life in a South African birth cohort: the Drakenstein Child Health Study. *Lancet Glob Health*, **3**, e95–103 (2015).
32. Maliha, N., Zafar, F. & Arif, A. Effect of Behavioral Risk Factors on Development of Pneumonia among Children Under-5 Years of Age: A Case-Control Study. *Malaysian Journal of Paediatrics and Child Health Online Early*, **1** (07), 1–13 (2011).
33. Zheng, X. H. *et al.* Home risk factors for childhood pneumonia in Nanjing, China. *Chin Sci Bull*, **58**, 4230–4236 <https://doi.org/10.1007/s11434-013-5686-5> (2013).

34. Schmidt, W. P., Cairncross, S., Barreto, M. L., Clasen, T. & Genser, B. Recent diarrhoeal illness and risk of lower respiratory infections in children under the age of 5 years. *Int J Epidemiol*, **38**, 766–772 (2009).
35. Rahman, S. S., Khatun, A., Azhar, B. S., Rahman, H. & Hossain, S. A Study on the Relationship between Nutritional Status and Prevalence of Pneumonia and Diarrhoea among Preschool Children in Kushtia: *Pediatrics Research International Journal*. 2014:39.
36. Hoving, M. F. & Brand, P. L. causes of recurrent pneumonia in children in a general hospital. *J Paediatr Child Health*. 2013; 49, E208–212. Doi: 10.1111/jpc.12114 PMID: 23438187.
37. Bernardo, L., Horta, Cesar, G. & Victora Short-term effects of breastfeeding: A systematic review on the benefits of breastfeeding on diarrhoea and pneumonia mortality. 2013
38. Worabe Town Administrative Health Unit: Annual plan performance report, 2008 EFY
39. Integrated management of. *newborn and childhood illness (IMNCI): chart booklet for health workers* (Ministry of Health, 2015).
40. Silte Zone Health Department: Health Management Information System report from Hamle (July) to Sane (June) 2008 EFY.
41. Mathew, J. L. *et al.* Acute respiratory infection and pneumonia in India: A systematic review of literature for advocacy and action: UNICEF-PHFI series on newborn and child health, India. *Indian Pediatr*, **48**, 191–218 [PubMed]. (2011).