

Soil Quality Evaluation Under Different Land Use Types in Kersa Sub-watershed, Eastern Ethiopia

Yared Mulat¹Kibebew Kibret²Bobe Bedadi²Muktar Mohammed³

¹Hawassa University, Wondo Genet College of Forestry and Natural Resources, department of natural resource management, P.O.Box 128, Shashemene, Ethiopia.

Email address of the corresponding author: yaredappo@gmail.com

²School of Natural Resources Management and Environmental Sciences, Haramaya University, P.O.Box 138, Dire Dawa, Ethiopia.

³Oda-Bultum University, Department of Forest Resources Management, P. O. Box 226, Chiro, Ethiopia.

Abstract

Background:soil quality, which can be inferred using indicators that interact synergistically, is affected by land use types and agricultural management practices. This study assessed the status of soil quality under three adjacent land uses (cultivated, grazing, and fallow) in Kersa sub watershed (622 ha). Soil samples were collected from the surface soil (0-20 cm depth) of the identified land uses with three replications and the soil quality parameters were analysed. A minimum data set of soil quality indicators were selected from physical, chemical, and biological parameters using the literature review and expert opinion method. Linear scoring functions were used to give unit less score for the selected data sets, which were then integrated into a soil quality index (SQI).

Results:The results revealed that bulk density, aggregate stability, pH, cation exchange capacity (CEC), available P, and soil organic carbon(SOC) had a significant difference in SQI among the different land uses. The soil quality indices were 0.69 for grazing land, 0.62 for cultivated land, and 0.59 for the fallow land. The SQI of all the land uses fall in the intermediate soil quality ($0.55 < \text{SQI} < 0.70$) class.

Conclusion:In almost all the quality indicators assessed, the grazing land was superior to the cultivated and fallow lands. Therefore, implementing management practices that enhance soil quality is imperative for sustainable agricultural production in the study area.

Keywords: Soil quality ,Minimum data set, soil quality indicator, Land use

1. BACKGROUND

Soils in agricultural production are vital part of ecological system which produces food and fiber for human utilization. Currently, sustainable and productive agriculture is highly related with soil quality (Karlen *et al.*, 1997). Soil quality refers to the capacity of soil to function within ecosystem boundaries to sustain biological productivity, maintain environmental quality and, promote plant and animal health (Doran and Parkin, 1994). Soil quality (SQ) addresses the issues of productivity and sustainability simultaneously that's why currently it becomes so indispensable for developing countries.

Land use type and agricultural management can be considered as the major factors that affect soil quality as a result of the change it brings on the soil physical, chemical, and biological properties (Caravaca *et al.*, 2002). These changed properties, in turn, affect land productivity. That is the reason why Hartemink (2003) stated that soil degradation is the principal component of land degradation, and almost all land degradation caused by soil degradation. Land degradation is a serious treat in Sub-Saharan Africa, where nearly 67 percent of the total land is degraded from light to severe extent (Million *et al.*, 2019). It is more severe in highlands of Ethiopia where 85 percent of the population lives. Ethiopia is among the sub-Saharan African (SSA) countries with very high soil degradation rates most of it in the form of nutrient mining (Ermias *et al.*, 2016 ; Misra *et al.*, 2003).

In Ethiopia, total cultivated land has reached approximately 12 million hectares in mid-2013, but most of the soils are highly degraded (ATA, 2013). Out of the total land, the highlands of Ethiopia cover about 50% of the land area with 95% of the cropping land accounting for over 90% of the country's economy (IFPRI, 2010). This shows that the overwhelming majority areas in the highlands are being put under intensive cropping. Moreover, new and often marginal and sloppy lands are being brought into crop production. Such land use changes have contributed to soil degradation and soil loss by deteriorating the soil physical and chemical properties (Karlton *et al.*, 2013). Land use practices affect the distribution and supply of soil nutrients by directly altering soil properties and by influencing biological transformations in the rooting zone. Furthermore, the high dependence on agriculture which is characterized by uncertain rainfall, poor management, and steep terrains, aggravates SQ degradation.

Soil quality degradation in Kersa sub-watershed, which is part of East Hararghe highlands, is uttered by the pressing problems of accelerated soil erosion, low productivity of agricultural lands, and deforestation (Bobe,2004; Kibebew, 2014). In the study area, land is used principally by smallholders for subsistence agriculture which is characterized by low yields and productivity. Since there is no as such sustainable soil quality management practices in the study area, farmers always seek for more land, whereby even the steep slopes and marginal lands are brought under cultivation to meet the increasing demand for food by the growing population. Hence, knowledge on SQ related to different land uses is imperative for developing management scenarios for sustainable use of the soil resource. This helps the famers' endeavor towards sustainable and productive use of soils as well as improving the SQ.

Therefore, assessing and monitoring SQ in soils that are currently under different land use types can provide effective tools for determining the properties of degraded soils and revealing sustainable land practices for land managers, and decision makers to make informed decisions against SQ degradation (Tesfahune, 2014). However, as indicated by Diack and Stott (2001) soil quality cannot be measured directly, but can be inferred by measuring soil physical, chemical, and biological properties that used as quality indicators .Thus, universally accepted soil quality index (SQI), which considers the soils' physical, chemical and biological properties, and variability, is of paramount importance for sustainable utilization and site-specific management of soil resources. Despite such importance of soil quality index (SQI) assessment, few studies have been made to quantify the proper indicators for evaluating and monitoring soil quality in Ethiopia and most of them are conducted in the northern part of Ethiopia (Tekluet *et al.*, 2007; Gebreyesus, 2014;Aweke *et al.*, 2015;Yoseph *et al.*,2017). No attempt has been made to generate minimum data set and evaluate soil quality in the eastern part of Ethiopia.

Hence, assessing the soil quality status is imperative to design better soil management practices that enhance productivity and environmental sustainability. In addition, it helps planners and decision makers to evaluate which land use system is the most sustainable and to take appropriate remedial measures considering potentials and constraints of the different land uses in the Kersa sub-watershed. Against this backdrop, the present study hypothesized that the different land use types in Kersa sub-watershed are assumed to possess different soil quality due to differences in their physical, chemical, and biological soil attributes. Therefore, the objectives of this study were to establish minimum data set

(MDS) of soil quality indicators by selecting appropriate scoring functions for each indicator and evaluate soil quality of different land use types using SQI.

1.2. MATERIALS AND METHODS

1.2.1. Description of the Study Area

The study area, Kersa sub-watershed, is located in Kersa district, East Hararghe zone of Oromia National Regional State between 9°26' 28" N to 9°27' 50"N, and 41°52' 0 "E to 41°53'50"E (Figure 1.1) with altitude that ranges from 1968 to 2127 meters above sea level. The total area of the watershed is 622 hactar.




Figure 1.1. Map of Kersa sub-watershed

Based on twenty one years (1995-2015) of data obtained from Ethiopian National Meteorology Agency (NMA), the study area receives a mean annual rainfall of 665 mm (Appendix Table1). The rainfall pattern in the area is bi-modal with high amount of rainfall occurring during the main rainy season between July to September and the short rainy season stretching from March to June (Figure 1.2). The highest mean rainfall is received in August. Similarly, based on 18 (1997-2014) years climate data obtained from NMA, the mean minimum and maximum annual air temperatures of the area are 12 and 24 °C, respectively, with mean annual air temperature of 18 °C (Appendix Tables 2 and 3).




Figure 1.2. Mean monthly rainfall and mean monthly minimum and maximum temperatures of the study area.

According to the geological map of Ethiopia, first published in 1973 at a scale of 1:2000000, the geology of Kersadistrictis covered by Adigrat formation constituted by sandstones and shell.Hamanlei series formation that contains Oxfordian limestone covers the lower part of the landscape, while the upper part of the landscape is covered by the lower complex undifferentiated precamberian rock. Moreover, Mohr (1964) indicated that Hararghe highlands lie over the crystalline bedrock composed mainly of granitic rock and gneiss material. Although the study area encompasses different land use types, the dominant land uses observed during the study were grazing, cultivated, and fallow land use types (Table 1).

Table 1.1. Description of land use types observed within Kersa sub-watershed in eastern Ethiopia

No	Land use	Description
1	Grazing	Land used as communal grazing land for cattle and it is managed through controlled system whereby livestock is confined in a stall and fed with cut-and-carry system.
2	Cultivated	This includes land used for cultivation of crops under rain fed conditions. The main cropping system is mixed cropping where khat (<i>Catha edulis</i>) is intercropped with sorghum. Only small amount of organic matter is returned to the soil, while almost no crop residue is returned to the soil due to its use for other purposes such as animal feed, fuel wood, source of cash, and construction material (e.g., sorghum).
3	Fallowing	This includes land that has been once under intensive cultivation but is now relieved, due probably for rehabilitation, since 2012.

The farming system of the area is predominantly subsistence farming based on mixed crop-livestock production. Livestock are integral part to the farming system, supplying draught power for cultivation, food and income to households. The major rain fed field crops grown are sorghum and maize often intercropped with common bean and khat. Besides, around homesteads, the vegetation is dominated by *Eucalyptus globules* and *Eucalyptus camaldulensis* trees.

M2	weight of the dish, stable aggregates and sand (g)
M3	weight of the dish and sand (g)
(M2 – M3)	weight of stable aggregates
(M3 – M1)	weight of sand
W	total weight of the sample.

Infiltration was determined in the field with double ring infiltrometer. Bulk density was determined from undisturbed (core) soil samples collected using core sampler (Black and Hartge, 1986). Particle density was determined by the graduated cylinder method (Issamand Antoine, 2007). Finally, total porosity was estimated from bulk density (BD, g/cm³) and particle density (PD, g/cm³) values using the following formula:

$$\text{Total porosity(\%)} = \left(1 - \frac{\text{BD}}{\text{PD}}\right) \times 100$$

The major soil chemical and biological properties analyzed were soil pH, organic carbon (OC), soil respiration, available P, exchangeable bases (Ca, Mg, Na, and K), and cation exchange capacity (CEC). Soil pH and electrical conductivity were measured using a 1:2.5 soil to water ratio as described in van (Reeuwijk, 1993). Organic carbon was determined using the wet oxidation method (Walkley and Black, 1934). Soil respiration was determined by Solvita soil respiration test kit as described by Woods End Laboratories (2015) which intended to show respiration under natural field moisture condition. Available phosphorus was determined by using the Olsen method (Olsen *et al.*, 1954).

Cation exchange capacity and exchangeable basic cations were extracted by 1M ammonium acetate (pH 7). Soil CEC was measured after leaching the ammonium acetate extracted (ammonium ion saturated) soil samples with 10% sodium chloride solution. The Na⁺ ions replace the NH₄⁺ ions in the sample and the amount of ammonium ion in the percolate was determined by the Micro-Kjeldahl procedures and reported as CEC method van Reeuwijk (1993). In the extract, exchangeable Ca and Mg were determined by atomic absorption spectrophotometer (AAS), while exchangeable K and Na were read by flame photometer. Extractable micronutrients (Fe, Mn, Cu, and Zn) were extracted by diethylenetriaminepenta-acetic acid (DTPA) method and were read using atomic absorption spectrophotometer (AAS) (Lindsay and Norvell, 1978).

1.2.4. Calculation of Soil Quality Index

Soil quality indexing follows three basic steps. These steps are indicator selection, interpretation/scoring, and integration into soil quality index value.

1.2.4.1. Indicator selection

The general criteria used for SQ indicator selection included influence on soil production and sensitivity to environmental changes, ease and cost of sampling and analysis method of the indicators to be selected. As a result selection of minimum data sets (MDS) of indicators was based on method of expert opinion and reviews of previous studies. This is because Andrews *et al.* (2002) stated that expert opinion based and statistical methods using principal component analysis (PCA) provided similar representative indicators as related to the measurement of environmental and production goals.

1.2.4.2. Indicator interpretation

The selected indicators were transformed into unit less scores ranging from 0 to 1 in linear technique according to Diack and Stott (2001). Scores ranging from 0 to 1 were assigned to the indicators included in the MDS by applying “more is better”, “less is better” or “optimum is better” function. The scores were assigned based on soil function after deciding the linear relationship of the anticipated response (i.e., “less is better”, “more is better” or “optimum better”). The score for each indicator was calculated after establishing the baseline, the lower, and the upper threshold values. The baseline value of the soil property corresponds to the score equal to 0.5. In this study, threshold or limit values were assigned to each soil quality indicator from natural ecosystems, best managed systems, and values adopted from literatures (Table 3.2).

The equations of the score curves are as follows:

$$f(x) = \begin{cases} 0.1 & X \leq L \\ 0.9 * \frac{X-L}{U-L} + 0.1 & L \leq X \leq U \\ 1 & X \geq U \end{cases} \quad \text{Eq. 1}$$

$$f(x) = \begin{cases} 1 & X \leq L \\ 1 - 0.9 * \frac{X-L}{U-L} & L \leq X \leq U \\ 0.1 & X \geq U \end{cases} \quad \text{Eq. 2}$$

Where $f(x)$ is the linear score; x is the soil property value; and L and U are the lower and upper threshold values, respectively. Eq. 1 was used for the “more is better” scoring function, whereas Eq. 2 was used for the “less is better” function. For the “optimum is

better” function, were scored as “more is better” for the increasing part and then scored as “less is better” for the decreasing part.

1.2.4.3. Soil quality indexing

Soil quality indicators are integrated into an index (SQI) by summing the scores from individual indicators obtained on each land use types and dividing by the total number of indicators as described by Mastoet *al.* (2008):

$$SQI = \left(\frac{\sum_{i=1}^n S_i}{n} \right) \quad \text{Eq.3}$$

Where SQI is the soil quality index, S_i is the linear scored value of individual indicators, and n is the number of indicators included in the dataset. A SQI value close to 1 refers to the best functioning soil, while zero (lower threshold) value refers to severely degraded soil. In relation to this, Marzaioliet *al.* (2010) divided SQI into three grades. $SQI < 0.55$ was regarded as low soil quality; $0.55 < SQI < 0.70$ was regarded as intermediate soil quality, and $SQI > 0.70$ as high soil quality.

1.2.5. Statistical Analysis

Measured data were subjected to analysis of variance (ANOVA) using the General Linear Model (GLM) procedure in which land use types were considered as independent variables (factors) and the selected soil properties as dependent variables. LSD was performed at $P < 0.05$ level using statistical package SAS, Version 9.2.

1.3 RESULTS AND DISCUSSION

1.3.1. Effects of Land Use Types on selected soil Quality Parameters

1.3.1.1. Soil physical quality indicators

Soil texture attributes (sand, clay) and basic infiltration rate (cm hr^{-1}) exhibited no statistically significant difference among the different land use types in Kersa sub-watershed ($P > 0.05$) (Table 1.2). Soil texture is an inherent soil property that is controlled primarily by soil formation processes (Brejda *et al.*, 2000b) and it is not influenced considerably by land use types and soil management in the short run. Rather, it is the major intrinsic factor that affects other soil properties such as infiltration albeit infiltration affected by land use types. As a result, it was rational that texture and infiltration rate exhibited statistically no significant difference among land use types in Kersa sub-watershed.

Bulk density, porosity, and aggregate stability were found to be statistically different in the different land use types. Table 1.2 indicates that bulk density was significantly higher in fallow land as compared to the grazing and cultivated lands. The highest bulk density in the fallow land could be a reflection of the low clay and amount of organic matter content which could also result in poor soil aggregation. The poor soil structure affects water intake and retention, which eventually results in soil quality degradation. On the other hand, the lower bulk density in grazing land due to the effect of high organic matter, which can give better structure or aggregation. Similar finding was reported by Zhang *et al.* (2008).

Baseline and threshold reference value for bulk density determined by Harris *et al.* (1996) for natural ecosystems takes into account 1.5 g/cm^3 as baseline value above which there is an increasing probability of soil compaction (Table 1.3). In reference to this value, soil bulk density of the soils under the three land uses was within the optimal range. The total porosity (%) of soils under the different land uses was significantly variable and ranged between 57-52 percent. The highest total porosity values were recorded under the grazing land use due to the low bulk density and high organic matter content.

Water stable aggregates were significantly affected by land use types ($P < 0.05$). Higher percentage of aggregate stability was recorded in soils of grazing land (66.3%) as compared to the other land use types (Table 1.2). This could be attributed to the relatively

higher soil organic matter content. The finding is in conformity with Peng *et al.*(2015) who reported the contribution of soil organic matter in improving the formation of soil structure . Similarly Yoseph *et al.*(2017) also reported that soil aggregate stability was significantly lower in crop land soils than in the soils under other land uses types.

Table 1.2. Mean values of soil quality indicators among different land use types in Kersa sub-watershed

LUT	Sand (%)	Clay (%)	TP (%)	BIR (cm hr ⁻¹)	WSA (%)	BD (g/cm ³)	Exchangeable properties (Cmol(+)/kg soil)				Av.P (mgkg ⁻¹)	OC (%)	TN (%)	pH (H ₂ O)	SBR (CO ₂ /min/day)	Fe(mgkg ⁻¹)	Zn(mg kg ⁻¹)
							Ca	K	Mg	CEC							
Grazing	54.0 ^a	29.0 ^a	57.56 ^a	2.20 ^a	66.30 ^a	1.18 ^b	23.33 ^a	2.64 ^a	7.20 ^a	39.00 ^a	10.12 ^a	1.87 ^a	0.18 ^a	7.29 ^a	7.6 ^a	6.60 ^a	1.68 ^a
Cultivated	54.0 ^a	29.3 ^a	56.40 ^a	2.13 ^a	59.00 ^b	1.30 ^{ab}	22.50 ^a	0.94 ^b	6.10 ^b	31.5 ^b	6.43 ^b	1.05 ^b	0.14 ^b	7.25 ^a	6.0 ^b	4.60 ^b	0.92 ^b
Fallow	51.0 ^a	28.0 ^a	52.2 ^b	2.16 ^a	53.00 ^c	1.33 ^a	19.6 ^b	0.80 ^b	5.70 ^b	29.40 ^c	5.63 ^b	0.57 ^c	0.08 ^c	6.72 ^b	3.7 ^c	4.30 ^c	0.86 ^b
Mean	53	23.3	55.30	2.20	59.44	1.27	21.83	1.46	6.35	33.31	7.34	1.16	0.136	7.09	5.8	5.16	1.16
CV	11.8	28.8	1.11	12.30	2.02	5.16	2.09	12.41	3.99	2.06	20.90	1.35	5.45	3.41	10.4	1.93	5.43
LSD	12.4	13.4	1.23	0.52	2.40	0.13	0.91	0.36	0.51	1.376	3.10	0.03	0.01	0.48	1.1987	0.19	0.13

LUT: Land Use Type, OC: organic carbon; AV. P: available Phosphorus; BD: Bulk Density, SBR: Soil Basal Respiration; BIR: Basic infiltration rate; WSA: Water Stable Aggregate; TP: Total Porosity; CEC: Cation Exchange Capacity.

Means followed by different letters within a column are significantly different at 0.05 levels.

1.3.1.2. Soil chemical and biological quality indicators

Soil pH was significantly ($P < 0.05$) affected by land use type in which the lowest pH was recorded under the fallow land (Table 1.2). According to soil pH ratings of Tekalign (1991), the overall pH range of the studied soils falls under neutral soil reaction ranges, which are a in favorable of most macro and micro nutrients and activities of microorganisms.

The significantly high pH in the grazing land could be attributed to limited removal of basic cations through erosion and leaching. However, removal of basic cations by crop through intensive cultivation, and continuous use of acid forming inorganic fertilizers could be the reason for the lower soil pH in cultivated and fallow land use types. The results are in agreement with the reports of Yihenew *et al.* (2015) who reported lower pH values for soils of cultivated lands and this ascribed to intensive cultivation and continuous use of acid forming inorganic fertilizers.

The effect of land uses on available P was significant ($P < 0.05$). Accordingly, significantly higher available P was recorded in soils under the grazing land as compared to those under cultivated and fallow land. The high organic matter content in the grazing land soils, which releases phosphorus during its mineralization, might contribute positively. These results are also in consonance with Solomon *et al.* (2001) and Wakene and Heluf (2003) who reported that soil P could be affected by vegetation cover in the land use. Nevertheless, the results of the present study were in contrast to that of Alemayehu and Sheleme (2013) who obtained relatively higher available P under cultivated land than that of grass land soils. The authors attributed the higher concentration of available P found under cultivated land to long term manure and house refuse applications .

Available P is below the base line limit (Table 3. 2). This low available P status might be attributed to low P fertilizer application and losses associated with erosion and removals during harvesting and continuous cropping and limited or no biomass return to the soil. The findings of this study are in agreement with the findings of Girma and Endalkachew(2013) who indicated that low available phosphorus might be related to continuous cropping, surface erosion, and absence of biomass addition to the soils.

Analysis of variance revealed significant differences in CEC of soils under different land use types ($P < 0.05$) (Appendix table 12). The highest and lowest CEC values were

recorded under grazing and fallow land uses, respectively. The high CEC in grazing land could be related to the combined effect and contribution of organic matter and types and amount of clay content in the soil. The result is in accordance with Bekele *et al.* (2006) who revealed that CEC of soils changes due to change in land uses. Furthermore, Yihenew *et al.* (2015) and Alemayehu and Sheleme (2013) pointed out that high cation exchange capacity (CEC) under grasslands was the result of high organic matter and clay content.

Soil organic carbon (SOC) showed significant variation with land use type ($P < 0.05$). Accordingly, the highest organic carbon content was recorded under grazing land (1.87%), while the lowest was observed in soils under the fallow land (Table 3.1). The lower soil organic carbon content under cultivated and fallow land could be related to low amount of organic material being returned to the soil system and high rate of oxidation of soil organic matter as a result of continuous cultivation for long period of time. Correspondingly, the research under Ambo Ber, Northern Ethiopian by Yoseph *et al.* (2017) indicated that the grassland had significantly higher soil organic carbon than cropland. Nevertheless, the current finding is contrary to earlier report by Boke (2004) who obtained relatively higher content of organic carbon in cultivated land than that of grassland.

The base line limit of OC is 3.5% (Table 1.3). In reference to this limit, the OC content was far below the base line limit in all land use types which indicates high biological soil quality degradation observed in the study soils. In line with this, Gebayese *et al.* (2014) emphasized that the lower soil OC could result in poor aggregate stability and, thus, aggravates soil degradation.

1.3.2. Soil Quality Index (SQI) of Different Land Use Types

The selected minimum data set indicators for this study to calculate the SQI consisted of clay, sand, bulk density, aggregate stability, soil pH, cation exchange capacity, SOC, and AvP. These indicators have also been consistently reported (Girmay and Singh, 2012; Aweke *et al.*, 2013; Cardoso *et al.*, 2013). The scoring function of “more is better” used parameters like water stable aggregates, organic carbon, clay content, CEC, and available P. On the other hand, bulk density was used as “less is better” scoring function; while sand content and soil pH were used for “optimum is better” scoring function.

The results ANNOVA of the integrated soil quality index for the land uses were 0.69 for grazing land, 0.62 for cultivated land, and 0.59 for fallow land, all of which are classified

into intermediate soil quality ($0.55 < \text{SQI} < 0.70$) class and statistically significant ($P < 0.05$) (Table 1.4). The soil quality index was found to be significantly higher in the grazing land use type as compared to the cultivated and fallow land uses. This reveals that the soils under the grazing land use type in kersa watershed are better off in terms of soil functioning and soil health (Andrews *et al.*, 2003). Higher soil quality index for grazing land is due to the controlled grazing system. In addition, grass roots contribute organic matter through their extended roots below surface which are not easily decomposed (Guo and Gifford 2002). Thus, highly interrelated variables of high OC, good aggregate stability, low bulk density, higher CEC, and available P yielded high soil quality index in the grazing land. On the other hand, the cultivated and fallow land use types scored intermediate but with relatively lower SQI values, which indicate the need for a judicious control of soil quality in the respective land use types (Nakajima *et al.*, 2015).

In line with the findings of the current study, Girmay and Singh (2012) reported intermediate soil quality index values for grazing lands at Maileba and Gum Selassa catchments in northern Ethiopia. On the other hand, Gebreyesus (2014) reported soil quality value index that is in the range of 'degraded' for uncultivated marginal land systems in Mai-Negus catchment of northern Ethiopia. The findings also agreed with the results of similar previous studies conducted in the world (Ngo-Mbogbaet *et al.*, 2015) where grasslands improved soil quality compared to continuous cultivation of crops.

Table 1.3. Soil quality indicators, scoring function values and references used for evaluating the soil quality indices in Kersa sub-watershed

Source of limits	Scoring curve	Threshold		baseline	Optimum	Slope at baseline	Source of Threshold
		Lower	Upper				
Clay	More is better	0	30	15	----	0.266	GebreyesusBrhane (2014)
Sand	Optimum	0	60	Lo-30Up-50	36	0.44	GebreyesusBrhane (2014)
Porosity (%)	Optimum	20	80	Lo -40 Up- 60	50	0.1280	Karlenet <i>al.</i> , (1994a, b)
Agregate stability %	More is better	10	60	30		0.296	Harris <i>et al.</i> (1996)
Bulk densiy(g/cm ³)	Less is better	1.0	2.2	1.5		-2.62	Harris et al. (1996)
Av.P(mg kg ⁻¹)	More is better	5	29	15		0.433	Mausbach and Seybold (1998)
CEC (cmol+ kg ⁻¹)	More is better	6	46	20		0.245	GebreyesusBrhane (2014)
OC (%)	More is better	1	6.5	3.5		1.046	Kay and Angers (1999)
pH	Optimum	4.5	9.5	6.5	5.3	1.3012	Karlenet <i>al.</i> ,(1994a, b)

The ANOVA result revealed that soil properties that explained the greatest proportion of the total variance in the present study included organic carbon, water stable aggregates, pH, CEC, available Phosphorous and bulk density (Table 1.4). These scored parameters made a significant difference among the different land use types ($P < 0.05$). As a result, there was no need of analysing all soil quality indicators to monitor the change in the study sub-watershed these can lead to rapid and economical of assessment. However, clay and sand content was the least important parameter in explaining the variability of soil quality since it did not statistically differ among land uses. This result is in agreement with the findings of Awekeet *al.* (2015) who identified soil organic carbon, water stable aggregates, and cation exchange capacity and Girmay and Singh (2012) also identified bulk density and available phosphorous as soil properties influencing the integrated soil quality index in northern Ethiopia. Other similar findings were also reported by Andrews *et al.* (2002) and Gebreyesus (2015).

Table 1.4. Means of soil quality parameters' scores in the different land use types in Kersa sub-watershed

LU	Clay	Sand	WSA	OC	CEC	pH	BD	AV.P	SQI
GR	0.93a	0.91a	1.00a	0.24a	0.84a	0.51b	0.86a	0.29a	0.69a
CL	0.90a	0.91a	0.98b	0.11b	0.67b	0.50b	0.78ab	0.15b	0.62b
FA	0.92a	0.87a	0.87b	0.03c	0.63c	0.59a	0.75b	0.12b	0.59b
Mean	0.91	0.89	0.95	0.12	0.71	0.53	0.53	0.18	0.63
LSD	0.27	0.18	0.03	0.02	0.04	0.08	0.09	0.11	0.034
CV	14.9	10.4	1.72	2.03	2.16	8.17	6.16	29.2	2.24

LU: Land Use, GR: Grazing Land; CL: Cultivated land, FA:Fallow Land, TP: Total Porosity, WSA: water stable aggregate, CEC: Cation Exchange Capacity, BD: Bulk density,SQI: soil quality index, AvP: available Phosphorus

Means followed by different letters within a column are significantly different from each other at 0.05 levels.

1.4. CONCLUSION

Soil quality parameters are affected by several physical, chemical, and biological parameters of the different land use types. Accordingly, statistically significant differences in soil quality of the different land uses were observed in the study area. The soil quality index of all land uses falls under intermediate soil quality ($0.55 < \text{SQI} < 0.70$) class. The major parameters that affected the soil quality index among the different land use types were bulk density, aggregate stability, pH, CEC, available P, and soil organic carbon. In most of the quality indicators assessed, the land under controlled grazing resulted in better quality as compared to the cultivated and fallow lands. Thus, the result of this study highlighted the potential of the land use types in improving or maintaining soil quality. Hence, locally appropriate soil fertility, productivity, and integrated soil conservation practices and enhancing of soil organic matter and other attributes should be identified and implemented. These will contribute significantly to the maintenance of soil quality for sustainable agricultural production in the study area.

1.5. Declarations

Authors' contributions

YA: collected, analyzed, interpreted the data and made the final write up which was part of his Doctoral thesis in Soil Science at Haramaya University, Ethiopia. KK, BB and MM, as co-authors edited the final manuscript. All authors read and approved the final manuscript.

Author details

¹Wondo Genet College of Forester and Natural Resource, Hawassa University, Hawassa, Ethiopia.

²School of Natural Resources Management and Environmental Sciences, Haramaya University, Dire Dawa, Ethiopia.

³Oda-Bultum University, Department of Forest Resources Management, P. O. Box 226, Chiro, Ethiopia

Acknowledgements

The project was funded Hawassa University and Haremaya University. Thus we would like to thank these institutions. We would also like to thank the anonymous reviewers who contributed significantly to the improvement of the article.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Not applicable.

Consent for publication

Not applicable.

Ethics approval and consent to participate

“Not applicable”.

Funding

Hawassa and Haremaya University provided fund for carrying out this study.

1.6. References

Agricultural Transformation Agency (2013) Annual report transforming agriculture in Ethiopia. Addis Ababa, Ethiopia.

Alemayehu K, and Sheleme B (2013) Effects of different land use systems on selected soil properties in South Ethiopia. *Journal of soil science and environ. Management*, 4:100-107.

Andrews S, Flora, C. B., Mitchell, J. P. and Karlen, D. L. (2003) Growers' perceptions and acceptance of soil quality indices. *Geoderma*, 114, 187-213.

Andrews S, Karlen D., Mitchell J (2002) A comparison of soil quality indexing methods for vegetable production systems in Northern California. *Agric. Ecosyst. Environ.*, 90:25-45.

Awdenege M, Melku D, Fantaw Y, (2013) Land use effects on soil quality indicators: A case study of abo-wonso southern Ethiopia. *Applied and Environmental Soil Sci.*, 32:234-242.

Aweke G, Singh B, Lal R (2013) Organic carbon and nitrogen associated with soil aggregates and particle sizes under different land uses in Tigray, Northern Ethiopia, *Land Degrad Develop.* doi:10.1002/ldr.2261, 2013.

Aweke G, Singh B, Lal R (2015) Soil quality indices for evaluating smallholder agricultural land uses in Northern Ethiopia. *Sustainability*, 7: 2322-2337

Bekele L, Kleja, D, Nilsson I, Olsson M (2006) Soil carbon sequestration under different exotic tree species in the southwestern highlands of Ethiopia. *Geoderma*, 136, 886-898.

Black G, Hartge K.H (1986) Bulk density in methods of soil analysis. In: A. Klute, (editor), Part 1. Physical and mineralogical methods, American Society of Agronomy-Soil Science Society of America, Madison, Wisconsin, USA. pp186.

Black G, Hartge K (1986) Particle density. In: Klute, A. (ed.). *Methods of soil analysis*. Madison. American Society of Agronomy, 1: 377-382.

Bobebedadi (2004) Evaluation of soil erosion in the Hararghe region of Ethiopia using soil loss models, rainfall simulation and field trials (Ph.D dissertation).University of Pretoria, Pretoria, South Africa.

Boke S. (2004) Soil phosphorus fractions as influenced by different cropping systems in Andosols and Nitosols in Kembata-Tembaro and Wolayta Zones SNNPRS (MSc Thesis). Alemaya University, (AU) Ethiopia.

Bouyoucus G (1962) Hydrometer method improvement for making particle size analysis of soils. Agron.Journal. 54: 179-186.

Brejda J , Moorman T, Karlen B, Dao T (2000b) Identification of regional soil quality factors and indicators: I. Central and southern high plains. Soil Sci. Soc. Am. J., 64:2115-2124.

Caravaca F, Masciandaro G ,Ceccanti B (2002) Land use in relation to chemical and biochemical properties in semiarid Mediterranean environment. Soil Till. Res., 68:23-30.

Cardoso, Vasconcellos R, Bini D, Miyauchi, M, Dos Santos C, Diack M , Stott D (2001) Development of soil quality index for the Chalmers silty clay loam from the Midwest USA, pp 550.

Diack M , Stott D (2001) Development of soil quality index for the Chalmers silty clay loam from the Midwest USA, In D. E. Stott, R. H. Mohtar, and G. C. Steinhardt (eds.), Sustaining the global farm, pp. 550_555.

Doran W ,Parkin T (1994) Defining and assessing soil quality. Sustainable Environment, Journal. 39:287-296.

Ermias T, Woldeamlak B , Belay S (2016) Effects of land use and land cover on selected soil quality indicators in the headwater area of the Blue Nile basin of Ethiopia. Environ Monit Assess., 188: 83-95.

Gebreyesus B (2014) Soil quality assessment strategies for evaluating soil degradation in Northern Ethiopia,” Applied and Environmental Soil Science, 14:64-78.

Girma A ,Endalkachew W (2013) Soil Properties, and Soil Organic Carbon Stocks of Tropical Andosol under Different Land Uses .Open Journal of soil science

GirmayG , Singh B (2012) Changes in soil organic carbon stocks and soil quality: land-use system effects in northern Ethiopia. Act aAgriculturaeScandinavica and Section B-Soil and Plant Science, 62: 519-530.

Guo L, and Gifford M (2002)Soil carbon stocks and land use change: a meta analysis. Global Change Biology, 8:345-360.

Harris R, KarlenL ,Mulla D (1996) A conceptual framework for assessment and management of soil quality andhealth, inMethods for Assessing Soil Quality, J.W. DoranandA.J. Jones, Eds., vol. 49, pp. 61–82, Soil Science Society of America,Madison,Wis, USA.

HarteminkA (2003) Soil fertility decline in tropics with case studies on plantations walling ford: CAB international /ISRIC.

IFPRI (International Food Policy Research Institute) (2010) Fertilizer and soil fertility potential in Ethiopia Constraint and Opportunities for Enhancing the System.Working Paper. Addis Ababa, Ethiopia.

Issam I, Antoine H (2007) Methods of analysis for soils of arid and semi-arid region. FAO,

Kandeler E. (1996) Aggregate stability. In: Schiner et al(eds): Methods in Soil Biology. Berlin, Springer-Verlag: 390–395.

Karlen D, Mausbach M, Doran W, Cline H, Harris R, Schuman G (1997) Soil quality: a concept, definition, and framework for evaluation. Soil Sci. Soc.of American Journal, 61: 4-10.

Karlen D, Peterson G , Westfall D (2014a) Soil and water conservation: Our history and future challenges. Soil Sci. Soc. Am. J., 78:1493-1499.

Karlen D, Wollenhaupt, N.C., Erbach, D.C., Berry, E.C., Swan, J.B., Eash, N.S. and Jordahl, J.L. 1994a. Crop residue effects on soil quality following 10-years of no-till corn. Soil Tillage Res., 31, 149-167.

Karlen L , Wollenhaupt , Erbach D, Berry C, Swan J, Eash N ,Jordahl, J (1994b) Long-term tillage effects on soil quality. *Soil Tillage Res.*, 32: 313-327.

Karlton E , Tekalign M, Taye B, Samuel G , Selamyihun K (2013) Towards improved fertilizer recommendations in Ethiopia Nutrient indices for categorization of fertilizer blends from woreda soil inventory data: A discussion paper. Ethiopian Agricultural Transformation Agency (ATA)/ Ethiopian Soil Information System (EthioSIS), Addis Ababa, Ethiopia.

Kay B,Angers D (1999) Soil structure, in *Handbook of Soil Science*, M. E. Summer, ed., pp. 229-269, CRCPress, NewYork, NY, USA,

Kibebew K (2014) Report on Characterization of agricultural soils in Cascape intervention woredas in Eastern Region of Ethiopia ,Haramaya University.PP1-234

MarzaioliR ,D'Ascoli R , De Pascale R , Rutigliano F (2010) Soil quality in a Mediterranean area of Southern Italy as related to different land use types. *Journal Appl.Soil Ecol.*, 44: 205-212.

MastoR ,Chhonkar P, Singh D, Patra A (2008) Soil quality response to long- term nutrientand crop management on a semi-arid Inceptisol. *Agric. Ecosyst. Environ.*, 118:130-142.

Mausbach M , Seybold C (1998) Assessment of soil quality,” in *Soil Quality and Agricultural Sustainability*, R. Lal, Ed., pp. 33-43, Sleeping Bear Press, Chelsea, Mich, USA.

Misra R, Lesschen P, Smaling , Roy, R (2003) Assessment of soil nutrient balance: approaches and methodologies (Vol. 14, *FAO Fertilizer and Plant Nutrition Bulletin*), Rome, Italy.

Ngo-Mbogba M, Yemefack M , Nyeck B (2015) Assessing soil quality under different land cover types within shifting agriculture in South Cameroon. *Journal Soil Tillage Res.*, 150: 124-131.

Olsen S , Cole C, Watanabe S, Dean (1954) Estimation of available phosphorous in soils by extraction with sodium bicarbonate. *USDA Circular*, 939: 1-19.

Peng X, Yan X , Zhou H , Zhang Y, Sun H (2015) Assessing the contributions of sesquioxides and soil organic matter to aggregation in an ultisol under long-term fertilization. *Soil Tillage Res.*, 146, 89-98.

Reeuwijk Van (1993) Procedures for soil analysis.4th ed. International Soil Reference and Information Center, the Netherlands.

Million Sileshi, Ruben K ,Khamaldin M,Stafen S.(2019) Determinants for adoption of physical soil and water conservation measures by smallholder farmers in Ethiopia, *International soil and water conservation research*,<https://doi.org/10.1016/j.iswer.2019.08.002>.

Solomon D, Lehmann M, Tekaligh M, Fritzsche F., Zech, W (2001) Sulfer fractions in particle size separation of the humid Ethiopia highland as influenced by land use changes. *Geoderma*, 102: 45-59.

Tekalign T (1991) Soil, plant, water, fertilizer, animal manure and compost analysis. International Livestock Research Center for Africa, Addis Ababa, Ethiopia.Working Document No.13.

Teklu E, ItannaF ,Stahr K (2007)Indexing soil quality: A new paradigm in soil science research. *Aust. J. Soil Res.* 45:129-137.

Teshome Y, Shelem B, Kibebew K (2016) Characterization and classification of soils of abobo area, western Ethiopia. *Applied and Environmental Soil Science*. Article ID 4708235.

Wakene N and Heluf G (2003) Forms of phosphorus and status of available micronutrients under different land-use systems of alfisols in Bako area of Ethiopia. *Ethiopian J. Nat. Res.*, 5: 17-37.

Woods End Laboratories Inc (2016) Woods End Laboratories Inc. Mt Vernon. USA.

Yihenew G, Fantanesh A ,Solomon A (2015) The effect of land use types, management practices and slop classes on selected soil physico-chemical properties in Zikre watershed, North-western Ethiopia. *Open Journal of environment system Research*,4:1-7

Yoseph I, Delelegn T, WitoonP ,Amila B, BirruY,Tesfaye W ,Hans G, Douglas L (2017) Changes in land use alter soil quality and aggregate stability in the highlands of northern Ethiopia. Scientific Reports.7: 13602

Zhang Y, Zhao Y, Shi X , Lu X,Yu D ,Wang H , Sun D, Darilek J (2008) Variation of soil organic carbon estimates in mountain region: a case study from south western china. Geoderman, 146:449-456.

