Effect of Below-Knee Compression Garments on Delayed-Onset Muscle Soreness (DOMS): A Randomized Controlled Laboratory Study With Multi-Parametric Analysis

Thilo Hotfiel

Erlangen University Hospital Department of Surgery: Universitätsklinikum Erlangen Chirurgische Klinik

Svenja Höger

University Hospitals Erlangen Department of Radiology: Universitätsklinikum Erlangen Radiologisches Institut

Armin M. Nagel

University Hospitals Erlangen Department of Radiology: Universitätsklinikum Erlangen Radiologisches Institut

Michael Uder

University Hospitals Erlangen Department of Radiology: Universitätsklinikum Erlangen Radiologisches Institut

Wolfgang Kemmler

Institute of Physics Friedrich Alexander University Erlangen

Raimund Forst

Friedrich Alexander University Erlangen Nuremberg Orthopedic-Rheumatology Department: Friedrich-Alexander-Universität Erlangen-Nürnberg
Orthopädisch-Rheumatologische Abteilung

Martin Engelhardt

Center for Musculoskeletal Surgery Osnabrück

Casper Grim

Center for Musculoskeletal Surgery Osnabrück

Rafael Heiss (✉ rafael.heiss@uk-erlangen.de)

University Hospitals Erlangen Department of Radiology: Universitätsklinikum Erlangen Radiologisches Institut <https://orcid.org/0000-0002-2897-5411>

Original Research Article

Keywords: Sports injuries, muscle injuries, muscle fatigue, regeneration, recovery

Posted Date: October 26th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-94383/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: The impact of compression on overexertion-related muscle injuries such as delayed-onset muscle soreness (DOMS) is a subject of controversy. Identifying the mechanisms, optimal conditions, and application settings under which compression therapy may alter muscle metabolism and improve recovery is crucial. The purpose of this study was to investigate the effects of wearing below-knee compression garments during exercise and a post-exercise period of 6 hours on clinical, functional, and morphological outcomes in DOMS in the lower limb.

Methods: Eighteen healthy volunteers (age: 24.1 ± 3.6 years, BMI 22.7 ± 2.7 kg/m²) were enrolled in this controlled laboratory study. Repeated measures were acquired at baseline (T_0), 6 h (T_1), and 48 h (T_2) after a standardized eccentric and plyometric exercise protocol, with the participants wearing a compression garment (21–22 mmHg) on a randomly selected calf during and for the first 6 h after exercise. Magnet resonance imaging (MRI) was performed using a 3T scanner for quantification of intramuscular edema (T2 signal intensity (SI), T2 time, and manual volume segmentation); jump height, calf circumference, ankle dorsiflexion (DF), creatine kinase (CK) levels, and muscle soreness were assessed for clinical and functional evaluation.

Results: DOMS was confirmed in all participants after 48 h, with a significant increase in muscle soreness ($p < 0.001$) and CK levels ($p = 0.001$), decrease in jump height ($p < 0.01$), and the presence of intramuscular edema ($p < 0.01$) in both the compressed and non-compressed limbs. No differences between the compressed and non-compressed limbs were observed for muscle soreness and jump height at any point in time. Similarly, MRI T2 SI, T2 time, soreness, and manual segmentation revealed no significant effect of the compression treatment. The assessment of calf circumference and DF showed no statistically significant changes at any point of time in either the compression or non-compression limb ($p = 1.0$).

Conclusions: The continuous application of below-knee compression garments during combined eccentric and plyometric exercise and until 6 h post-exercise has no significant effect on clinical signs of DOMS, jump performance, or the development of intramuscular edema observed in MRI.

Background

High-performance sports and the importance of achieving peak physical capacity and performance have led athletes and coaches to continually seek treatments or techniques that may improve regeneration and recovery (1–3). The rate and quality of recovery has been considered to be essential for both elite and recreational athletes during repetitive training or competition (1–4).

A wide variety of interventions aiming to either prevent or relieve the development of muscle soreness, thereby accelerating recovery from performance-limiting conditions, have been reported (2). Compression garments are traditionally applied for vascular purposes (e.g., deep venous and lymphatic disease) or surgery, and their clinical benefits are well documented (5–7). Recently, compression garments have increasingly been used by those participating in sporting activities (6, 8, 9). Existing systematic reviews have concluded that compression garments are effective in enhancing recovery from overexertion-related muscle injuries (primarily within 48 h of injury), whereas the majority of previous studies failed to support performance-enhancing effects (1, 8–12). Various mechanisms have been hypothesized to explain the potential beneficial effects of compression garments, including altered muscle biomechanics, leading to reduced power expenditure and microtrauma; improvements in proprioception; elevated tissue pressure; and even placebo effects (8, 13). A limitation of most existing studies is that they have focused on a variety of exercise types, outcome parameters, or types of compression, including different wearing time. This leads to a broad heterogeneity of results and a paucity of data on fundamental physiological mechanisms occurring upon the application of compression therapy for muscular issues (12). However, identifying the mechanisms, metabolic conditions, and application modes under which compression may alter muscle metabolism and improve recovery is crucial. Hence, there is a need for a highly standardized multiparametric approach to assess clinical, functional, and morphological changes. The results might be of high clinical relevance for the understanding of compression principles, the optimization of its application, and the development of further therapeutic regimens. We hypothesized that compression garments (21–22 mmHg) are able to reduce intramuscular edema and microstructural damage and thereby improve functional performance and recovery in cases of delayed-onset muscle soreness (DOMS). The purpose of our study was to investigate the effects of wearing below-knee compression garments during and for 6 h after a standardized plyometric and eccentric exhausting exercise using clinical, functional, and morphological outcome measurements.

Methods

Study design:

At baseline (T_0), 6 h (T_1), and 48 h (T_2) after a standardized exercise protocol, MRI, creatine kinase (CK) levels, jump height, calf circumference, ankle dorsiflexion (DF), and muscle soreness were assessed. After the baseline measurements, an exercise intervention (described below) was performed with participants wearing a compression garment (21–22 mmHg, 75% polyamide, 25% elastane) on a randomly selected calf. Compression garments were worn continuously during the exercise intervention and for the following 6 h. The garments were removed immediately before the beginning of the second measurements (T_1).

Participant recruitment and study population:

Healthy individuals between 20 and 30 years of age were recruited from medical and sports facilities. Inclusion criteria were lack of signs, symptoms, or history of chronic disease; no current acute or overuse injuries of the lower limb; and no history of muscle injury. All volunteers were asked to forego alcohol and sports for 72 h before and 48 h after the first MRI acquisition. Exclusion criteria were any symptoms of lower-limb muscle soreness in the 3 months prior to the study and regular training habits that included eccentric or plyometric exercises.

Ethical approval:

The study was approved by the local institutional Ethics Committee and written informed consent was obtained from all volunteers (Ref. No. 33_16 B; University of Erlangen-Nürnberg).

Exercise Intervention:

All volunteers performed an established standardized training that included plyometric and eccentric exercises of both lower legs while wearing a compression garment on a randomly selected calf (left or right side). All exercises were monitored by an experienced strength and conditioning coach. Each participant warmed up using two sets of heel raises (15 repetitions per set), with a 20 s break between sets. Immediately after the warm-up, the plyometric exercise was performed. Volunteers completed 5 sets of 10 repetitions each of drop jumps from a 0.25 m box (5 s between jumps, 1 min between sets). Volunteers were asked to hop off from the box with both feet, land with both feet together, and perform a maximal vertical jump with the shortest possible contact time to the ground. After a 2 min break, a previously described eccentric exercise was performed on a specifically manufactured slant plate (-35°) (14, 15). To increase the load, each volunteer wore a weighted vest bearing approximately 40% of his/her body weight during the entire eccentric exercise (Figure 1). All participants performed 5 sets of 50 repetitions each and rested 60 s between sets, with the last set being performed until muscle fatigue, so that no further repetition of eccentric exercise was possible.

Imaging:

MRI was performed on a 3T scanner (Magnetom Prisma; Siemens Healthcare; Erlangen, Germany) with an 18-channel body array coil. An axial T2-weighted turbo inversion recovery magnitude (TIRM) sequence (total acquisition time (AT), 4:53 min; inversion time (IT), 260 ms; echo time (TE), 69 ms; repetition time (TR), 5120 ms; resolution, 0.8 x 0.8 x 4.0 mm) and a coronal T2-weighted TIRM sequence (AT, 4:14 min; IT, 260 ms; TE, 68 ms; TR, 5870 ms; resolution, 0.9 x 0.9 x 4.0 mm) of both lower legs were acquired. An additional axial T2 mapping (AT, 9:35 min; TE, 10.6–190.8 ms; TR, 3170 ms; resolution, 0.7 x 0.7 x 8.0 mm) was applied to quantify the tissue water content. In addition, we performed a T1-weighted turbo spin-echo sequence to depict the anatomy and morphology of the lower leg.

To analyze the extent of edema, the T2 signal intensity and T2 time value (ms) were assessed according to previously published investigations (16). The compartments of the lower leg were differentiated by defining specific regions of interest in the T1-weighted images corresponding to the anatomic margin of the medial (MGM) and lateral (LGM) head of the gastrocnemius muscle, the soleus muscle (SM), and the tibialis anterior (TA) muscle and copying these over the T2-weighted TIRM and T2 mapping sequence images. In addition, the volume of total intramuscular edema in the MGM was quantified using a threshold-based manual segmentation (16).

For grading of the exercise-induced muscle damage, a modification of the Peetrans classification (17) was used: grade 0 indicated a negative MRI without any visible pathology, grade I indicated edema but no architectural disorganization, grade II indicated architectural disruption indicating partial tear, and grade III indicated total muscle rupture. All subjects with grade I lesions in axial TIRM images were considered to have induced DOMS. The MRI images were analysed blindly by an experienced radiologist using a professional image-processing program (syngo.via VB10; Siemens AG).

Jump height:

The power of both lower legs was assessed separately by jumps with full knee extension after familiarization with the movement pattern. Starting in an upright position with both arms fixed to the upper body and 90° flexion in the hip and knee of the contralateral leg, avoiding additional impulse movements, volunteers were asked to jump as quickly and explosively as possible in order to perform the highest possible jump. Tests were performed on a force platform (KMP Newton GmbH; Stein, Germany). The software provided by the manufacturer automatically calculates jumping height based on ground reaction forces. Mean jump height was determined using three repeated measurements.

Calf circumference:

The maximum calf circumference was measured using a tape measure while the participant was sitting on the edge of a table with free-hanging lower legs (14). The location of measurement was labeled with a permanent marker at baseline to ensure the same position for follow-up measurements.

Ankle dorsiflexion:

Passive DF of both ankle joints was assessed manually with a goniometer while the participant was sitting on the edge of a table with free-hanging lower legs (18).

Muscle soreness:

Self-reported muscle soreness was evaluated using a 10 cm visual analogue scale for pain (0, no pain; 10, worst pain). Pain scores were reported for each leg individually at rest and during activity (going down stairs) (14).

Creatine kinase (CK) levels:

Blood CK levels were measured at baseline and at 6 h and 48 h after the eccentric exercise. Approximately 5 mL blood was collected by vein puncture from an antecubital vein into serum tubes. CK measurement was conducted using the UV test according to the IFCC method (37°C; Cobas 6000; Roche Diagnostics; Mannheim, Germany).

Blinding:

Test assistants and outcome assessors were kept unaware of which of the participants' legs were supplied with compression garments and were not allowed to ask about this.

Statistical analysis:

All assessed parameters were checked for normality with the Shapiro–Wilk test. In cases in which the data were normal, a repeated-measures analysis of variance, with time and treatment (compression versus non-compression) as the repeated independent variables, was performed for each of the dependent variables of interest: CK, T2 signal intensity, T2 time, calf circumference, jump height, and ankle DF. When the data were not normally distributed, a Friedmann test with a post hoc Dunn–Bonferroni test was performed. Muscle soreness scores were analyzed using the Friedmann test. All statistical tests were performed using SPSS Version 23 (IBM Corporation; Armonk, NY, USA), and P values of less than .05 were considered to indicate statistical significance.

Results

A total of 18 healthy volunteers (8 women, 10 men; mean age: 24.1 ± 3.6 ; range, 18–29 years, BMI 22.7 ± 2.7 kg/m²) were included in this study. All recruited volunteers successfully performed the plyometric and eccentric exercises described and completed the measures strictly according to the study protocol.

The mean CK value was 174.5 ± 131.7 U/L at baseline and 4423 ± 7489.3 U/L 48 h after the exercise ($p = 0.001$). Moreover, 48 h after the exercise, all subjects presented grade I lesions in the MGM according to the modified Peetrans classification (17, 19), indicating the induction of DOMS (20). Therefore, all 18 volunteers were included in further evaluation.

Muscle soreness significantly increased from baseline (T_0) to 48 h follow-up (T_2) in both the compressed (0 ± 0 versus 4.7 ± 2.3 , $p < 0.001$) and non-compressed (0 ± 0 versus 4.8 ± 2.0 , $p < 0.001$) limbs when climbing stairs, whereas no differences were observed at 6 h (T_1). No difference in muscle soreness was observed between the compressed and non-compressed limbs at either follow-up time (T_1 and T_2).

Similarly, for jump height, the compression treatment had no statistically significant effect at any time point ($p = 1.00$). A significant reduction in jump height appeared for both the compressed (8.5 ± 2.2 cm versus 6.6 ± 1.7 cm, $p = 0.008$) and non-compressed (8.1 ± 2.0 cm versus 6.8 ± 2.2 cm, $p = 0.003$) leg at T_2 . The assessment of calf circumference and range of motion showed no statistically significant changes at any time point, nor did wearing the compression garment ($p = 1.0$). In contrast, T2 SI and T2 time, representing intramuscular edema, demonstrated a significant increase in the MGM in both the compressed (T2 SI: 28.5 ± 13.1 versus 83.1 ± 70.2 , $p = 0.005$; T2 time: 36.3 ± 3.9 ms versus 57.7 ± 27.9 ms, $p < 0.001$) and non-compressed (T2 SI: 30.5 ± 10.2 versus 76.2 ± 49.3 , $p = 0.011$; T2 time: 36.3 ± 3.3 versus 54.9 ± 23.8 , $p = 0.003$) lower leg after 48 h (T_2). No effect of wearing a compression garment appeared for T2 SI ($p = 1.0$) or T2 time ($p = 1.0$) of the MGM (Figure 2). Comparable results were achieved using manual segmentation of the volume of total intramuscular edema in the MGM, which showed a significant increase for both the compressed ($p = 0.025$) and non-compressed ($p = 0.003$) legs at T_2 , with wearing a compression garment having no impact ($p = 1.0$; Figure 3). Additional results are presented in Table 1.

Discussion

To the best of our knowledge, this study is the first to investigate the effects of wearing below-knee compression garments (continuously during exercise and for the 6 h post-exercise) on both morphological and functional changes in skeletal muscles by means of a multiparametric analysis.

The underlying approach of using an exercise-induced DOMS model indicating ultrastructural muscle injuries has been successfully established in previous studies (14, 16, 21, 22) and was performed here with eccentric and plyometric exercises. The protocol used in this study can be considered effective, as we were able to observe significant increases in CK levels, muscle soreness, and intramuscular edema in all participants, indicating the development of clinical signs of DOMS after 48 h. DOMS, an ultrastructural muscle injury, is classified as an overexertion muscle disorder (23). It is

one of the most common reasons for impaired muscle performance in sports and is associated with muscle soreness, reduced strength, and range of motion, and is frequently observed in both professional and recreational athletes (20, 24-26).

The major finding of our study was that the continuous application of compression during combined plyometric and eccentric exercises and throughout a post-exercise period of 6 h showed no influence on muscle edema, soreness, or jump height. These results contradict hypothesis, which predicted altered morphological and functional changes in muscles subjected to compression treatment. The understanding of how and under which conditions muscle tissue responds to compression is essential knowledge for improved recovery and prevention. Several exercise-, athlete-, and compression-related aspects must be contemplated.

DOMS was previously considered to be the result of exhausting or prolonged muscle activity, particularly after eccentric exercise and/or unfamiliar and not-well-coordinated sporting activities (20). DOMS is associated with ultrastructural damage to the skeletal muscle tissue, involving protein degradation (27); loss of myofibrillar integrity, with Z-band streaming and disruption; inflammatory processes (28); and changes in microcirculation and vascular permeability (29).

A recent meta-analysis demonstrated a significant effect of exercise modality on the effects of compression at all time points (30). The same authors found compression to be most effective for improving recovery from exercise that elicits a large degree of muscle damage (30); just as demonstrated in our study. Recovery from resistance or plyometric exercise has been demonstrated to derive the greatest benefits from compression (30, 31). Indeed, the wide range in exercise designs, including differences in timing, contraction forms, and intensity, has contributed to apparently inconsistent findings; thus, specific recommendations for the ideal exercise regime supporting the use of compression have not yet been elucidated (11). The protocol applied in this study consists of an exhausting exercise protocol leading to DOMS. However, no significant effects could be observed for compression on either the functional or morphological level. In this study, plyometric forms of contractions have been included in order to target high-speed plyometric contractions. Notably, in sports, there are no isolated heavy loaded eccentric contractions that induce a “pure eccentric overload” as applied traditionally in numerous DOMS models, indicating that these models do not simulate valid and realistic sports conditions (14, 20). Instead, during athletic activities such as running, change of directions, or jumps, eccentric contractions are shorter and part of the entire stretch-shortening cycle (9, 32). Remarkably, particularly for high-speed plyometric conditions, there is evidence to suggest that compression garments may function through a biomechanical support of the muscle-tendon unit, resulting in greater mechanical efficiency and promoting a lower energy expenditure (33, 34) and reduced microtrauma and muscular damage (13). From a biomechanical perspective, it has been further postulated that compression is able to reduce muscle vibrations and oscillations during repetitive landing loads, making plyometric contractions susceptible to compression (35). However, only few studies have investigated real-time effects of compression on mechanical variables.

Heterogeneous strategies are discussed for the ideal form of compression. The manifestation of overexertion muscle injuries and DOMS is due to a complex sequence of local and systemic physiological responses (36, 37). The earliest clinical manifestations of DOMS commonly begin 6–12 h post-exercise and increase progressively in the context of an inflammatory response until reaching a peak pain level 48–72 h after exercise. The inflammatory process is associated with electrolyte imbalances, leukocyte accumulation, and an upregulation of circulating pro-inflammatory cytokines, accompanied by intramuscular edema, compartment swelling, nociceptor activation, and/or pain sensation. It is thought that applying compression generates an external pressure gradient that attenuates changes in osmotic regulation and reduces the spread of edema and swelling (11).

Based on pathophysiological foundations, studies on compression focus on different time-dependent application modes, including “during exercise” (i.e., biomechanical support) and “post-exercise” treatment aiming to limit the inflammatory processes mentioned above and thereby preventing the development and clinical signs of DOMS (2). A meta-analysis found that several studies described recovery-enhancing effects for wearing compression garments both during exercise and afterwards, whereas other studies were not able to confirm these findings (30). Most of the scientific studies reporting benefits of post-exercise compression consider compression periods between 24 h and 72 h after exercise, which may not be routinely applicable due to discomfort caused by the compression garments. Our study aimed to address sports-specific conditions in both the exercise protocol and the wearing regime of compression garments, which was supposed to represent an ordinary day of intense training or competition: The continuous application during exercise and during the post-exercise period of exemplary 6 h (i.e., during the return trip after competition or in the evening after training) is practiced by many sports teams and athletes. The results presented elucidate a tendency toward an exposure time-dependent effect of compression for the post-exercise period after exhausting eccentric and plyometric exercises.

Study limitations and future directions:

While our study increased the existing knowledge on compression in sports, some limitations should be mentioned. We acknowledge that our measurements may not be transferable to real sport-specific conditions. The preceding exercise intervention involving eccentric and plyometric exercises allows us to relate data to standardized exhausting muscle activity, an integral part of many sports. However, we were not able to investigate effects of compression on sporting activities in which different metabolic demands interact and may influence physiological responses continuously. Further, we used each subject as their own control. We chose an intra-individual control group because inter-individual differences in DOMS expression have occurred in our preliminary tests and have been described in several studies (38, 39). However, one calf may influence the outcome of the other calf due to a systemic inflammatory healing response. Nevertheless, the findings may be significant for the practical

implementation of compression garments. This study contributes to the ongoing controversial debate about when and how long compression should be applied. Hence, the remarkable findings obtained with this multiparametric approach can be valuable for further investigations. Longitudinal data assessments including multiple and frequent multiparametric measures during an observational period up to 72 h must be implemented in order to determine exact application times supporting recovery-enhancing effects in DOMS.

Conclusions

This is the first study investigating the effects of wearing below-knee compression garments during exercise and a post-exercise period of 6 h on clinical, functional, and morphological outcomes in DOMS. The continuous application of below-knee compression garments during combined eccentric and plyometric exercises and for the following 6 h has no statistically significant effects on clinical signs of DOMS, jump performance, or the development of intramuscular edema observed in MRI.

Abbreviations

AT – Acquisition time

BMI – Body mass index

CK – Creatine kinase

DF – Dorsiflexion

DOMS – Delayed-onset muscle soreness

IFCC - International Federation for Clinical Chemistry reference

IT - Inversion time

LGM – Lateral head of the gastrocnemius muscle

MGM – Medial head of the gastrocnemius muscle

MRI - Magnet resonance imaging

SI – Signal intensity

SM – Soleus muscle

TA – Tibialis anterior muscle

TE – Echo time

TIRM – Turbo inversion recovery magnitude

TR – Repetition time

Declarations

Ethics approval and consent to participate: The local Ethics Committee approved the methods of the study with no requirements (Ref. No. 33_16 B; University of Erlangen-Nuremberg, Erlangen, Germany). All patients were informed about the purpose, benefits, and risks of the investigation prior to signing an institutionally approved informed consent document to participate in the study.

Consent for publication: Not applicable

Availability of data and materials: The authors fully support data sharing and collaboration. Data are available upon request for researchers who provide a methodologically sound proposal beginning 3 months and ending 5 years after article publication. Data include the study protocol, informed consent form, ethical approval, DICOM files, and statistical analysis for all of the deidentified participant data collected during the trial.

Competing interests: The study was funded by Bauerfeind AG (Zeulenroda-Triebes, Germany). The funders had no role in study design, data collection, or analysis; decision to publish; or preparation of the manuscript. Martin Engelhardt and Casper Grim are consultants of Bauerfeind AG (Zeulenroda-Triebes, Germany) outside the submitted work. Thilo Hotfiel, Svenja Höger, Armin M. Nagel, Michael Uder, Wolfgang Kemmler, Raimund Forst and Rafael Heiss declare that they have no competing interests

Funding: The study was funded by by Bauerfeind AG (Zeulenroda-Triebes, Germany).

Authors' contributions: TH, RH, ME, and WK designed the study. WK and SH monitored the data collection and prepared the data for further analysis. AN and RH were responsible for the conceptualization of MRI software and hardware parameters. MRI images were assessed by RH and MU. CG, ME, RF, MU, RH, and TH interpreted the data. All authors have made major contributions to drafting and writing the manuscript and approved the manuscript for publication.

Acknowledgements: *Not applicable*

References

1. Hotfiel T, Mayer I, Huettel M, Hoppe MW, Engelhardt M, Lutter C, et al. Accelerating Recovery from Exercise-Induced Muscle Injuries in Triathletes: Considerations for Olympic Distance Races. *Sports (Basel)*. 2019;7(6).
2. Heiss R, Lutter C, Freiwald J, Hoppe MW, Grim C, Poettgen K, et al. Advances in Delayed-Onset Muscle Soreness (DOMS) - Part II: Treatment and Prevention. *Sportverletz Sportschaden*. 2019;33(1):21-9.
3. Kellmann M, Bertollo M, Bosquet L, Brink M, Coutts AJ, Duffield R, et al. Recovery and Performance in Sport: Consensus Statement. *International journal of sports physiology and performance*. 2018;13(2):240-5.
4. Kellmann M. Preventing overtraining in athletes in high-intensity sports and stress/recovery monitoring. *Scandinavian journal of medicine & science in sports*. 2010;20 Suppl 2:95-102.
5. Lim CS, Davies AH. Graduated compression stockings. *CMAJ*. 2014;186(10):E391-8.
6. Reich-Schupke S, Surhoff S, Stücker M. Pressure profiles of sport compression stockings. *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology : JDDG*. 2016;14(5):495-506.
7. Rabe E, Partsch H, Hafner J, Lattimer C, Mosti G, Neumann M, et al. Indications for medical compression stockings in venous and lymphatic disorders: An evidence-based consensus statement. *Phlebology*. 2018;33(3):163-84.
8. Vercruyssen F, Gruet M, Colson SS, Ehrstrom S, Brisswalter J. Compression Garments, Muscle Contractile Function, and Economy in Trail Runners. *International journal of sports physiology and performance*. 2017;12(1):62-8.
9. Mota GR, Simim MAM, Dos Santos IA, Sasaki JE, Marocolo M. Effects of Wearing Compression Stockings on Exercise Performance and Associated Indicators: A Systematic Review. *Open Access J Sports Med*. 2020;11:29-42.
10. Born DP, Sperlich B, Holmberg HC. Bringing light into the dark: effects of compression clothing on performance and recovery. *International journal of sports physiology and performance*. 2013;8(1):4-18.
11. Hill J, Howatson G, van Someren K, Leeder J, Pedlar C. Compression garments and recovery from exercise-induced muscle damage: a meta-analysis. *Br J Sports Med*. 2014;48(18):1340-6.
12. Marques-Jimenez D, Calleja-Gonzalez J, Arratibel I, Delextrat A, Terrados N. Are compression garments effective for the recovery of exercise-induced muscle damage? A systematic review with meta-analysis. *Physiol Behav*. 2016;153:133-48.
13. Trenell MI, Rooney KB, Sue CM, Thompson CH. Compression garments and recovery from eccentric exercise: A P-31-MRS study. *J Sport Sci Med*. 2006;5(1):106-14.
14. Heiss R, Kellermann M, Swoboda B, Grim C, Lutter C, May MS, et al. Effect of Compression Garments on the Development of Delayed-Onset Muscle Soreness: A Multimodal Approach Using Contrast-Enhanced Ultrasound and Acoustic Radiation Force Impulse Elastography. *The Journal of orthopaedic and sports physical therapy*. 2018;48(11):887-94.
15. Hotfiel T, Kellermann M, Swoboda B, Wildner D, Golditz T, Grim C, et al. Application of Acoustic Radiation Force Impulse (ARFI) Elastography in Imaging of Delayed Onset Muscle Soreness (DOMS): A Comparative Analysis With 3T MRI. *J Sport Rehabil*. 2017;1:29.
16. Heiss R, Hotfiel T, Kellermann M, May MS, Wuest W, Janka R, et al. Effect of Compression Garments on the Development of Edema and Soreness in Delayed-Onset Muscle Soreness (DOMS). *Journal of sports science & medicine*. 2018;17(3):392-401.
17. Ekstrand J, Healy JC, Walden M, Lee JC, English B, Hagglund M. Hamstring muscle injuries in professional football: the correlation of MRI findings with return to play. *Br J Sports Med*. 2012;46(2):112-7.
18. Pasurka M, Lutter C, Hoppe MW, Heiss R, Gaulrapp H, Ernstberger A, et al. Ankle flossing alters periarticular stiffness and arterial blood flow in asymptomatic athletes. *J Sports Med Phys Fitness*. 2020.
19. Peetrons P. Ultrasound of muscles. *Eur Radiol*. 2002;12(1):35-43.
20. Hotfiel T, Freiwald J, Hoppe MW, Lutter C, Forst R, Grim C, et al. Advances in Delayed-Onset Muscle Soreness (DOMS): Part I: Pathogenesis and Diagnostics. *Sportverletz Sportschaden*. 2018;32(4):243-50.
21. Hotfiel T, Kellermann M, Swoboda B, Wildner D, Golditz T, Grim C, et al. Application of Acoustic Radiation Force Impulse Elastography in Imaging of Delayed Onset Muscle Soreness: A Comparative Analysis With 3T MRI. *J Sport Rehabil*. 2018;27(4):348-56.
22. Jakeman JR, Byrne C, Eston RG. Lower limb compression garment improves recovery from exercise-induced muscle damage in young, active females. *Eur J Appl Physiol*. 2010;109(6):1137-44.

23. Mueller-Wohlfahrt HW, Haensel L, Mithoefer K, Ekstrand J, English B, McNally S, et al. Terminology and classification of muscle injuries in sport: the Munich consensus statement. *British journal of sports medicine*. 2013;47(6):342-50.
24. Kim SK, Kim MC. The affect on delayed onset muscle soreness recovery for ultrasound with bee venom. *Journal of physical therapy science*. 2014;26(9):1419-21.
25. Mizuno S, Morii I, Tsuchiya Y, Goto K. Wearing Compression Garment after Endurance Exercise Promotes Recovery of Exercise Performance. *Int J Sports Med*. 2016;37(11):870-7.
26. Pearcey GE, Bradbury-Squires DJ, Kawamoto JE, Drinkwater EJ, Behm DG, Button DC. Foam rolling for delayed-onset muscle soreness and recovery of dynamic performance measures. *Journal of athletic training*. 2015;50(1):5-13.
27. Ulbricht A, Gehlert S, Leciejewski B, Schiffer T, Bloch W, Hohfeld J. Induction and adaptation of chaperone-assisted selective autophagy CASA in response to resistance exercise in human skeletal muscle. *Autophagy*. 2015;11(3):538-46.
28. Lewis PB, Ruby D, Bush-Joseph CA. Muscle soreness and delayed-onset muscle soreness. *Clin Sports Med*. 2012;31(2):255-62.
29. Yanagisawa O, Sakuma J, Kawakami Y, Suzuki K, Fukubayashi T. Effect of exercise-induced muscle damage on muscle hardness evaluated by ultrasound real-time tissue elastography. *Springerplus*. 2015;4:308.
30. Brown F, Gissane C, Howatson G, van Someren K, Pedlar C, Hill J. Compression Garments and Recovery from Exercise: A Meta-Analysis. *Sports medicine (Auckland, NZ)*. 2017;47(11):2245-67.
31. Hettchen M, Glöckler K, von Stengel S, Piechle A, Lötzerich H, Kohl M, et al. Effects of Compression Tights on Recovery Parameters after Exercise Induced Muscle Damage: A Randomized Controlled Crossover Study. *Evidence-based complementary and alternative medicine : eCAM*. 2019;2019:5698460.
32. Nicol C, Avela J, Komi PV. The stretch-shortening cycle : a model to study naturally occurring neuromuscular fatigue. *Sports medicine (Auckland, NZ)*. 2006;36(11):977-99.
33. Bringard A, Perrey S, Belluye N. Aerobic energy cost and sensation responses during submaximal running exercise positive effects of wearing compression tights. *International journal of sports medicine*. 2006;27(5):373-8.
34. Kemmler W, von Stengel S, Köckritz C, Mayhew J, Wassermann A, Zapf J. Effect of compression stockings on running performance in men runners. *Journal of strength and conditioning research*. 2009;23(1):101-5.
35. Stickford AS, Chapman RF, Johnston JD, Stager JM. Lower-leg compression, running mechanics, and economy in trained distance runners. *International journal of sports physiology and performance*. 2015;10(1):76-83.
36. Hoppeler H. Eccentric Exercise. *Physiology and Application in Sport and Rehabilitation*. London: Routledge. 2015;Vol. 8).
37. Böning D. Muskelkater. *Dtsch Z Sportmed*. 2000;51(2), 63-64.
38. Kraemer WJ, Bush JA, Wickham RB, Denegar CR, Gómez AL, Gotshalk LA, et al. Influence of compression therapy on symptoms following soft tissue injury from maximal eccentric exercise. *The Journal of orthopaedic and sports physical therapy*. 2001;31(6):282-90.
39. Yanagisawa H, Kurita R, Kamata T, Yoshioka K, Kato D, Iwasawa A, et al. Effect of the sp(2)/sp(3) Ratio in a Hybrid Nanocarbon Thin Film Electrode for Anodic Stripping Voltammetry Fabricated by Unbalanced Magnetron Sputtering Equipment. *Analytical sciences : the international journal of the Japan Society for Analytical Chemistry*. 2015;31(7):635-41.

Tables

Table 1

Range of motion, circumference of the lower leg, jump height, T2 signal intensity, and T2 relaxation time within the medial head of the gastrocnemius muscle (MGM), total volume of intramuscular edema within the MGM, subjective pain at rest and while climbing stairs of the non-compressed and compressed lower leg before (baseline, T0) and 6 h (T1) and 48 h (T2) after plyometric and eccentric training. P values < 0.05 are bolded.

	Non-compression					Compression					Non-compression vs. Compression		
	Baseline (t0)	6 hours (t1)	48 hours (t2)	p value (t0/t1)	p value (t0/t2)	Baseline	6 hours	48 hours	p value (t0/t1)	p value (t0/t2)	p value (t0/t0)	p value (t1/t1)	p value (t2/t2)
ROM [°]	38.0 ± 5.9	36.1 ± 7.9	33.2 ± 8.9	1.00	1.00	38.6 ± 6.8	37.4 ± 8.7	32.4 ± 8.5	1.00	1.00	1.00	1.00	1.00
Circumference [cm]	36.9 ± 2.4	36.7 ± 2.4	37.3 ± 2.6	1.00	1.00	36.7 ± 2.4	36.4 ± 2.4	37.4 ± 2.5	1.00	0.182	1.00	0.693	1.00
Jump Height [cm]	8.1 ± 2.0	8.2 ± 2.3	6.8 ± 2.2	1.00	0.027	8.5 ± 2.2	8.0 ± 2.5	6.6 ± 1.7	1.00	0.008	1.00	1.00	1.00
T2 signal intensity	30.5 ± 10.2	36.7 ± 14.5	76.2 ± 49.3	1.00	0.011	28.5 ± 13.1	32.4 ± 13.4	83.1 ± 70.2	1.00	0.005	1.00	1.00	1.00
T2 relaxation time [ms]	36.3 ± 3.3	39.1 ± 5.8	54.9 ± 23.8	1.00	0.007	36.3 ± 3.9	38.4 ± 5.5	57.7 ± 27.9	1.00	< 0.001	1.00	1.00	1.00
Volume of edema [ml]	12.1 ± 8.9	17.8 ± 11.0	106.4 ± 108.9	1.00	0.003	15.4 ± 13.0	18.3 ± 20.9	117.9 ± 135.0	1.00	0.025	1.00	1.00	1.00
Pain at rest	0.0 ± 0.0	0.4 ± 0.9	1.0 ± 1.5	1.00	0.223	0.0 ± 0.0	0.1 ± 0.3	0.7 ± 1.2	1.00	0.272	1.00	1.00	1.00
Pain at climbing stairs	0.0 ± 0.0	1.0 ± 1.1	4.8 ± 2.0	0.223	< 0.001	0.0 ± 0.0	0.8 ± 0.9	4.7 ± 2.3	1.00	< 0.001	1.00	1.00	1.00

Figures

Figure 1

Procedure of the standardized eccentric exercise of the calf muscles: (A) start position, with heel raised; (B) eccentric contraction (-35°); (C) composites of the entire setting.

Figure 2

Axial T2-weighted turbo inversion recovery magnitude (A, C, E) and axial T2 mapping (B, D, F) images of the lower leg at baseline (A, B) and at 6 h (C, D) and 48 h (E, F) after standardized exercise of the calf muscles. Muscle edema is apparent in the medial gastrocnemius muscle in the post-exercise images after 48 h. This participant wore the compression garment on the left calf for 6 h after exercise. The calibration tubes contain 40 mmol/L NaCl (*) and 40 mmol/L NaCl with 5% agarose (#).

Figure 3

Total volume of intramuscular edema was assessed in the coronal T2-weighted TIRM images. Representative images before (A, C, E) and after (B, D, F) color-coded manual segmentation of the edematous medial gastrocnemius muscle are shown for measurements at baseline (A, B) and at 6 h (C, D) and 48 h (E, F) after standardized exercise. Hyperintense vessels were segmented in the baseline images and subtracted from the segmented volume in the post-exercise images.