Prevalence of and Factors Associated With Lactational Mastitis in Eastern and Southern Africa: An Exploratory Analysis of Community-Based Household Surveys

Mariame O. Ouedraogo (✉ m.ouedraogo@mail.utoronto.ca)

University of Toronto Dalla Lana School of Public Health

Lenka Benova

Institute of Tropical Medicine Antwerp

Tom Smekens

Institute of Tropical Medicine Antwerp

Gezahegn G. Sinke

Children Believe

Abraha Hailu

Children Believe

Herbert B. Wanyonyi

Amref Health Africa

Madalitso Tolani

Amref Health Africa

Caristus Zumbe

Amref Health Africa

Ibukun-Oluwa O. Abejirinde

University of Toronto Dalla Lana School of Public Health

Research

Keywords: Lactational Mastitis, Sub-Saharan Africa, Prevalence, Associated Factors, Exploratory Analysis, Maternal Morbidity, Newborn Health

Posted Date: October 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-944609/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at International Breastfeeding Journal on March 28th, 2022. See the published version at <https://doi.org/10.1186/s13006-022-00464-x>.

Abstract

Background

Lactational mastitis is an extremely painful and distressing inflammation of the breast, which can seriously disrupt breastfeeding. Most of the evidence on the frequency of this condition and its risk factors is from high-income countries. Thus, there is a crucial need for more information on lactational mastitis and its associated factors in Sub-Saharan Africa (SSA).

Methods

We used data from representative, community-based cross-sectional household surveys conducted with 3,315 women from four countries (Ethiopia, Kenya, Malawi, and Tanzania) who reported ever-breastfeeding their last child born in the two years before the survey. We first estimated country-specific and pooled prevalence of self-reported lactational mastitis and examined mastitis-related breastfeeding discontinuation. Additionally, we examined factors associated with reporting mastitis in the pooled sample using bivariate and multivariable logistic regression accounting for clustering at the country level and post-stratification weights.

Results

The prevalence of self-reported lactational mastitis ranged from 3.1% in Ethiopia to 12.0% in Kenya. Close to 17.0% of women who experienced mastitis stopped breastfeeding because of mastitis. The adjusted odds of self-reported lactational mastitis were approximately two-fold higher among women who completed at least some primary school compared to women who had no formal education. Study participants who delivered by caesarean section had 1.46 times higher odds of reporting lactational mastitis than women with a vaginal birth. Despite wide confidence intervals, our models also indicate that young women (15 – 24 years) and women who practiced prelacteal feeding had higher odds of experiencing lactational mastitis than older women (25+ years) and women who did not give prelacteal feed to their newborn.

Conclusion

The prevalence of lactational mastitis in four countries of SSA might be somewhat lower than estimates reported from other settings. We found that lactational mastitis may result in breastfeeding discontinuation in approximately 1 in 5 woman-infant dyads affected by the condition. Factors such as younger age, higher education, caesarean section, and prelacteal feeding may be associated with experiencing mastitis. Further studies should explore the risk and protective factors for lactational mastitis in African contexts and address the negative consequences on breastfeeding.

Introduction

Mastitis is an inflammatory condition of the breast, commonly associated with lactation; hence the name lactational mastitis[1]. Definitions of lactational mastitis vary somewhat across the literature; however, generally, they combine breast symptoms (e.g., nipple pain, redness, and swelling) and flu-like symptoms (e.g., fever and chills) experienced during the breastfeeding period[2]. Lactational mastitis is an extremely painful and distressing condition, estimated to occur in up to 20% of breastfeeding women, and is most frequently reported to occur in the first month of the postpartum period[2].

To date, most of the evidence on the occurrence of lactational mastitis has emerged from high- and upper-middle-income countries; the three high-quality studies included in a recent meta-analysis of incidence were conducted China, Iran and Australia[2]. In this meta-analysis, it was estimated that around a quarter of women breastfeeding up to 6 months of age develop lactational mastitis[2]. In contrast, high quality data on mastitis from low-income countries, particularly Sub-Saharan Africa (SSA), remains limited. To date, studies conducted in SSA indicate a risk of lactational mastitis ranging between 0.5% and 2.6%[2, 3]. In contrast, the experience of other breast problems (e.g., blocked ducts and breast engorgement) was reported to be higher (i.e., 15%)[3].

In addition to the limited evidence on the occurrence of lactational mastitis in SSA countries, minimal information is available on the risk factors of lactational mastitis. In high-income countries, associations between mastitis and anatomical and breastfeeding factors (e.g., nipple damage, attachment difficulties, and poor positioning) and obstetric history (e.g., primiparity) were found[2]. Evidence surrounding the importance of socio-demographic factors in experiencing lactational mastitis has been heterogeneous[1]. Studies have also shown that women who exclusively breastfed were less likely to experience breast health problems, including mastitis, than women who did not exclusively breastfeed their newborn[3, 4]. Education and counseling of mothers about optimal breastfeeding techniques have also been suggested to address poor positioning and attachment and ultimately mastitis[5].

Recognizing the factors associated with lactational mastitis is important, given that they can hinder optimal breastfeeding practices and discourage continued breastfeeding. Mastitis can result in premature supplementary feeding and breastfeeding cessation and ultimately impact the growth and development of infants by increasing their exposure to diarrhea and common childhood illnesses, as well as the risk of health problems later in childhood and adolescence [6–9]. Breastfeeding is also associated with maternal health, including positive mental health benefits of achieving own breastfeeding goals and lower lifetime incidence of type II diabetes, breast, and ovarian cancer[8, 10–12].

Given the role that lactational mastitis could play in influencing breastfeeding practices, this study aims to investigate the prevalence and factors associated with lactational mastitis in four countries in SSA. In addition, the present study also documents the breastfeeding habits of women who experienced lactational mastitis.

Methods

Using cross-sectional data from population-based household surveys, this study aimed to investigate the prevalence of lactational mastitis across four Eastern and Southern Africa countries and explore factors associated with it.

Study settings

This study used household-level data collected as part of the endline evaluation of the Canada-Africa Initiative to Address Maternal, Newborn and Child Mortality (CAIA-MNCM) project. The CAIA-MNCM project, funded by Global Affairs Canada, aimed to reduce maternal and child mortality in targeted areas of Ethiopia, Kenya, Malawi, and Tanzania and was implemented by a consortium of non-governmental organizations (Amref Health Africa, Children Believe, WaterAid, and The Hospital for Sick Children's Centre for Global Child Health) between 2016 and 2020.

In Ethiopia, the CAIA-MNCM project was implemented in the Oromia Zone of the Amhara region. Kenya rolled out the project in four sub-counties of Siaya County (Alego-Usonga, Bondo, Ugenya, and Ugunja). In Malawi, the project targeted Chikwawa, Neno, and Mwanza districts in the Southern Region and Ntchisi in the Central Region. Finally, in Tanzania, the CAIA-MNCM project took place in Geita and Nyang'hwale districts located in the Geita Region.

Data sources

For this study, we used data from the CAIA-MNCM project endline household surveys conducted between February and July 2020 in the four countries of interest. These cross-sectional surveys aimed to provide data representative of the zones, districts, and counties where the project took place. The household surveys were designed to establish and understand changes in community-level RMNCH, nutrition, and WASH indicators. A total of 6,000 households (Ethiopia: N = 1,000; Kenya: N = 1,000; Malawi: N = 2,000; Tanzania: N = 2,000) were targeted for participation. Although the CAIA-MNCM project was implemented in four districts in Malawi, the quantitative data collection was only conducted in two districts - Chikwawa and Ntchisi. We assumed that the results of Chikwawa would apply to Mwanza and Neno, which are located in the same region.

For the household surveys in all four countries, we adopted a two-stage clustered design similar to that employed by the Demographic and Health Surveys (DHS)[13]. The first stage or primary sampling units were the enumeration areas (EA) from the most recent national population census. A sample from the list of all EAs covering the target population was selected using probability proportional to size selection. In the second stage of sampling, a fixed number of households was randomly selected per sampled EA. The selection of households varied among countries: in Ethiopia, we carried out a household listing exercise with the assistance of village leaders and community health workers. In Kenya and Tanzania, we used the household listings recorded by community health workers (CHWs) since CHWs are expected to update household listings at least once a year. In Malawi, we used a random walk approach; while the initial plan was to perform a systematic random selection of households, the random walk approach was adopted, given the time and financial constraints.

Study population

The household surveys targeted all women of reproductive age (15–49 years) as well as mothers or primary caretakers of children under five years. In addition, a 25% sample of households was selected for interviewing fathers of children under five years of age. In total, across the four countries, we interviewed 6,298 women of reproductive age, 5,914 mothers and caregivers of children under five, and 924 fathers of children under five.

Analysis for this study was restricted to the sample of women of reproductive age who had a live birth in the two years before the survey and reported ever breastfeeding their last child born within the two years preceding the survey. A total of 311 women in Ethiopia, 205 in Kenya, 1,339 in Malawi, and 1,460 in Tanzania were eligible.

Description of variables

The outcome variable of interest was self-reported lactational mastitis at any point during the breastfeeding period related to the last-born child under two years of age. Lactational mastitis was determined by asking the following sequence of questions: *“At any point since you started breastfeeding [NAME] or within a month of stopping breastfeeding, did you experience the following: breast tenderness or pain, redness of any part of the breast, breast lump?”* and, if women responded positively, *“While you had those symptoms, did you also experience high temperature, flu-like symptoms?”* These questions were developed based on a set of items commonly used in studies of self-reported mastitis[2]. Women who responded positively to both questions were categorized as having experienced lactational mastitis during their last pregnancy. In addition to collecting information on women's self-report of mastitis, we inquired about the frequency of symptoms of mastitis and whether the mastitis-related symptoms resulted in stopping breastfeeding.

The explanatory variables were selected after a thorough review of variables used in past studies, which have been documented in a recent systematic review and meta-analysis[2]. Variables included maternal age at the time of the survey (categorised into five-year intervals: 15–19, 20–24,

25–29, 30–34, 35–39, 40–44, and 45–49 years), time since birth (< 6 months, 6–24 months), parity at the time of index birth (nulliparous, multiparous), highest education level (none, less than primary, completed primary, some secondary, higher than secondary), household wealth index constructed following principal component analysis steps defined by the DHS Program [13] and categorised into reasonably equally-sized quartiles (poorest, poorer, richer, richest) within each country, index birth by caesarean section (yes, no), and counseling and observation of breastfeeding during postnatal care (PNC) of the newborn (immediate, delayed, absent/no).

We also derived two variables to measure breastfeeding within the first six months of life. The first variable, applied to the entire sample, was based on the proportion of participating women who reported their newborns not receiving any liquids to their newborns within the first three days of life (prelacteal feeding). The second variable measuring exclusive breastfeeding during the first six months of life was only applied to a sub-sample of participants whose last-born child was < 6 months of age at time of the survey. It consisted of measuring the proportion of women who indicated not giving liquids and foods other than breastmilk to their newborn in the 24 hours before their interview. Both variables were dichotomous (yes, no).

Statistical analysis

All data cleaning and preparation were done on SAS/STAT® software, version 9.4 of the SAS system (SAS Institute, Cary, NC, USA). We conducted the analyses on RStudio version 1.4.1106[14].

We conducted an initial descriptive analysis to highlight the characteristics of study participants by country. Descriptive statistics from Malawi and Tanzania were based on weighted samples to account for different probabilities of household selection in each district. We also examined the country-specific distribution of self-reported mastitis symptoms and the prevalence of self-reported mastitis.

To increase our statistical power in analysis of risk factors, we pooled data from the four countries. We calculated post-stratification weights in order to account for the countries' differing population sizes. For each country, we estimated the population fraction and the sample fraction. For the population fraction, we divided each country's population size by the sum of the four countries' populations. For the sample fraction, we divided each country's sample size by the total household survey sample size. The final country-specific post-stratification weights were obtained by dividing the population fraction by the sample fraction. An extra step was taken in Malawi and Tanzania, where we calculated district-specific population and sample fractions. These were subsequently multiplied by the country post-stratification weights.

We examined the variables associated with self-reported lactational mastitis through bivariate and multivariable logistic regression models. All results are presented as odds ratios (ORs) with corresponding 95% confidence intervals (95% CI), summarizing the direct effect of the variables on lactational mastitis, instead of their total effects [15]. All models accounted for post-stratification weights and clustering of observations at the country level. The final multivariable logistic regression models excluded parity and counselling on breastfeeding during newborn PNC visits due to collinearity with maternal age and observation of breastfeeding during PNC, respectively.

Given that lactational mastitis is more commonly experienced in the first few months following birth and to limit the risk of recall bias, we conducted additional sensitivity analysis restricting the logistic regression to women whose last-born child was < 6-months old at the time of the survey (see Additional file 1). There was a total of 762 women in this sensitivity analysis (Ethiopia: N = 104, Kenya = 47, Malawi: N = 232, Tanzania: N = 379).

Results

Characteristics of the Study Participants

In all four countries, most study participants were age 20–29 years (Table 1). Ethiopia had a large proportion of participants who did not attend formal schooling (67.5%), while in the other countries, most women attended or completed primary school (54.1% in Kenya, 33.4% in Malawi, and 51.4% in Tanzania). In all countries, the majority of participants had more than one child. The caesarean section (c-section) rate was highest in Kenya (11.2%) and the lowest in Ethiopia (2.9%). In Kenya, Malawi, and Tanzania, 72.2%, 58.1%, and 45.9% of women, respectively, indicated being counseled on breastfeeding during their newborn PNC visit, which occurred in the first 48 hours following birth. Similarly, in all countries except Ethiopia, many women reported that the PNC provider observed their breastfeeding within the first 48 hours following birth.

Table 1
Background characteristics of the participating women, by country

	Ethiopia N (%)	Kenya N (%)	Malawi ^a N (%)	Tanzania ^a N (%)
Total sample size	311	205	1,338	1460
Maternal age				
15–19	11 (3.5)	12 (5.8)	97 (8.1)	109 (6.6)
20–24	83 (26.8)	40 (19.5)	399 (29.2)	467 (31.2)
25–29	89 (28.6)	58 (28.3)	340 (24.3)	385 (27.1)
30–34	69 (22.2)	60 (29.3)	246 (18.5)	261 (18.3)
35–39	52 (16.7)	26 (12.7)	187 (14.4)	140 (10.3)
40–44	6 (1.9)	8 (3.9)	53 (4.2)	75 (5.1)
45–49	1 (0.3)	1 (0.5)	16 (1.3)	23 (1.4)
Time since last live birth				
< 6 months	104 (33.4)	47 (22.9)	238 (17.8)	389 (26.6)
6–24 months	207 (66.6)	158 (77.1)	1100 (82.2)	1071 (73.4)
Highest education level				
Never attended school	210 (67.5)	5 (2.4)	283 (23.4)	410 (29.0)
Some primary	78 (25.1)	39 (19)	360 (28.5)	129 (9.9)
Completed primary	20 (6.4)	111 (54.1)	474 (33.4)	773 (51.4)
Some secondary	3 (0.9)	40 (19.5)	217 (14.5)	144 (9.2)
Higher	0 (0)	10 (4.9)	3 (0.25)	4 (0.4)
Household wealth index quartile				
Poorest	68 (21.9)	41 (20.1)	397 (29.6)	343 (21.9)
Poorer	117 (37.6)	54 (26.5)	379 (29.5)	373 (22.8)
Richer	34 (10.9)	50 (24.5)	271 (20.4)	360 (25.9)
Richest	92 (29.6)	59 (28.9)	288 (20.5)	384 (29.3)
Parity^b				
Nulliparous	66 (23.0)	40 (20.8)	312 (25.4)	280 (19.4)
Multiparous	221 (77.0)	152 (79.2)	809 (74.6)	1063 (80.6)
Most recent birth by caesarean section^c				
Yes	9 (2.9)	23 (11.2)	107 (6.9)	53 (3.1)
No	302 (97.11)	182 (88.8)	1229 (93.1)	1407 (96.9)
Counselling on breastfeeding during PNC of newborn				
Yes, immediate (48 hours after birth)	74 (23.8)	148 (72.2)	777 (58.1)	670 (45.9)
Yes, delayed (> 48 hours after birth)	28 (9.0)	24 (11.7)	376 (28.1)	361 (24.7)
No ^d	209 (67.2)	33 (16.9)	185 (13.8)	429 (29.4)
Observation of breastfeeding during PNC of newborn				

^a Weighted frequencies and percentages; ^b Parity data missing for 24 women in Ethiopia, 13 in Kenya, 137 in Malawi, and 117 in Tanzania; ^c Caesarean section data missing for three women in Malawi who did not respond; ^d Includes participants who did not receive counselling on/observation of breastfeeding within their PNC and those who did not receive PNC at all.

	Ethiopia N (%)	Kenya N (%)	Malawi ^a N (%)	Tanzania ^a N (%)
Yes, immediate (48 hours after birth)	66 (21.2)	150 (73.2)	734 (54.9)	684 (46.8)
Yes, delayed (> 48 hours after birth)	28 (9.0)	24 (11.7)	335 (25.0)	351 (24.0)
No ^d	217 (69.8)	31 (15.1)	269 (20.1)	425 (29.1)
^a Weighted frequencies and percentages; ^b Parity data missing for 24 women in Ethiopia, 13 in Kenya, 137 in Malawi, and 117 in Tanzania; ^c Caesarean section data missing for three women in Malawi who did not respond; ^d Includes participants who did not receive counselling on/observation of breastfeeding within their PNC and those who did not receive PNC at all.				

Prevalence of self-reported lactational mastitis and characteristics of women reporting lactational mastitis

The prevalence of self-reported lactational mastitis since the birth of their most recent child ranged from 3.1% in Ethiopia to 12.0% in Kenya (Table 2). Among women with self-reported lactational mastitis, 61.0% indicated having experienced mastitis-related symptoms more than once. In addition, about 17.0% of women with lactational mastitis reported stopping breastfeeding due to mastitis; this ranged from 9.2% in Tanzania to 20.5% in Malawi (data not shown in table).

As shown for the pooled sample in Table 3, self-reported lactational mastitis was most commonly observed among participants aged between 15 and 24 years (8.0%). About 7.3% of women who were 6–24 months postpartum reported lactational mastitis, compared to 5.2% of women who were within the first six months of the postpartum period. Close to 4.0% of women who never attended formal education experienced lactational mastitis, while 9.3% of respondents part of the poorest wealth quantile indicated mastitis. We found no evidence of important differences in the proportion of self-reported mastitis by parity nor by counselling and observation of breastfeeding during PNC. We also observed that 11.3% of participating women who gave birth by c-section had self-reported lactational mastitis, compared to 6.5% of women who had a vaginal delivery. In addition, reporting of lactational mastitis was higher among women who reported their newborns receiving prelacteal feed (13.7%) compared to 6.4% among women who did not. Similar bivariate associations between explanatory factors and experience of lactational mastitis were found in the sub-group analysis of women < 6 months postpartum (see Additional file 1, Table A1).

Table 2
Distribution of self-reported lactational mastitis symptoms and estimated prevalence of self-reported lactational mastitis among the study participants, by country and pooled (N = 3,314)

	Ethiopia N (%)	Kenya N (%)	Malawi ^a N (%)	Tanzania ^a N (%)	Pooled N (%)
Mastitis symptoms					
Breast tenderness or pain	10 (3.1)	54 (25.9)	149 (11.0)	207 (14.0)	382 (11.5)
Redness of any part of the breast	10 (3.1)	24 (11.5)	50 (3.7)	80 (5.4)	186 (5.6)
Breast lump	13 (4.1)	32 (15.4)	50 (3.7)	89 (6.0)	237 (7.1)
In addition to at least one of the symptoms above					
Fever	9 (60.0)	21 (35.6)	56 (32.0)	94 (39.3)	183 (41.3)
Flu-like symptoms	7 (46.7)	24 (39.3)	59 (33.7)	106 (44.3)	189 (41.8)
Prevalence of mastitis					
	10 (3.1)	25 (12.0)	73 (5.4)	124 (8.4)	222 (6.7)
^a Weighted frequencies and percentages					

Table 3
Reporting of lactational mastitis by women according to selected explanatory factors, pooled sample (N = 222 women with self-reported lactational mastitis)

	Weighted Frequency of women reporting lactational mastitis	Weighted % of women reporting lactational mastitis	P-value
Maternal age ^a			
15–24 years	84	8.0	0.1418
25–35 years	117	6.4	
36–49 years	21	4.7	
Time since birth			
< 6 months	49	5.2	< 0.0001
6–24 months	173	7.3	
Highest education level			
Never attended school	52	3.9	0.0045
Some primary	56	8.3	
Completed primary	86	8.8	
Some secondary or higher	28	9.1	
Household wealth index quartile			
Poorest	68	9.3	0.4062
Poorer	62	6.0	
Richer	49	8.1	
Richest	43	4.6	
Parity			
Nulliparous	58	7.9	0.4007
Multiparous	162	6.8	
Most recent birth by caesarean section			
Yes	19	11.3	0.0845
No	203	6.5	
Counselling on breastfeeding during PNC of newborn			
Yes, immediate (48 hours after birth)	101	7.2	0.9306
Yes, delayed (> 48 hours after birth)	32	6.5	
No ^b	89	6.3	
Observation of breastfeeding during PNC of newborn			
Yes, immediate (48 hours after birth)	104	7.5	0.7010
Yes, delayed (> 48 hours after birth)	31	6.5	
No ^b	87	6.0	
Prelacteal feed			
Yes	18	13.7	0.1181
^a Maternal age was collapsed into broader categories due to sample size limitations; ^b Includes participants who did not receive counselling on/observation of breastfeeding within PNC and those who did not receive PNC at all.			

	Weighted Frequency of women reporting lactational mastitis	Weighted % of women reporting lactational mastitis	P-value
No	204	6.4	
^a Maternal age was collapsed into broader categories due to sample size limitations; ^b Includes participants who did not receive counselling on/observation of breastfeeding within PNC and those who did not receive PNC at all.			

Binary logistic regression analyses of factors associated with self-reported lactational mastitis

The bivariate logistic regression analyses (Table 4) indicated that being 25–35 years was associated with lower odds of reporting lactational mastitis compared to being aged between 15 and 24 years. On the other hand, having at least some primary education was associated with higher odds of self-reported lactational mastitis, with the stronger association being reported when comparing women who never attended school with their counterparts who attended at least some secondary school (OR = 2.74; 95% CI: 1.19–6.31). Although not statistically significant, women who were nulliparous had 1.17 times higher odds of self-reported lactational mastitis than multiparous women. Women who gave birth by c-section were 84% more likely to self-report experiencing mastitis than women who delivered vaginally. Prelacteal feeding was associated with increased odds of self-reported lactational mastitis. The confidence interval was however large, suggesting some variability and thus moderate evidence that women who practice prelacteal feeding have higher likelihood of experiencing lactational mastitis.

The multivariable logistic regression models (Table 4) show that the odds of lactational mastitis were 46% greater in women 6–24 months postpartum than in women who were in the first five months of the postpartum phase. Additionally, the odds of self-reported mastitis were approximately two-fold higher among women who completed at least some primary school. Birth by c-section was significantly associated with higher odds of self-reported lactational mastitis (aOR:1.46; 95% CI: 1.07–2.00). Despite not being statistically significant, the models also show that younger maternal age and giving prelacteal feed were associated with higher likelihood of lactational mastitis.

Even though wide confidence intervals were noted due to small sample size, the direction of the ORs between the considered explanatory variables and self-reported lactational mastitis were also confirmed in our sub-group analyses of women whose last-born child was < 6 months at the time of the survey. In these analyses, we observed that compared to women who were exclusively breastfeeding their newborns, women were not had higher odds of lactational mastitis (aOR = 2.55; 95%CI: 0.66–9.80, Additional file 1).

Table 4
Bivariate and multivariable logistic regression of selected explanatory factors and lactational mastitis, pooled sample (N = 3,314)

	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)
Age^a		
15–24 years	1.0	1.0
25–35 years	0.79 (0.64–0.96)	0.88 (0.68–1.13)
36–49 years	0.56 (0.24–1.31)	0.71 (0.33–1.54)
Time since birth		
< 6 months	1.0	1.0
6–24 months	1.42 (1.38–1.47)	1.46 (1.37–1.55)
Highest education level		
Never attended school	1.0	1.0
Some primary	2.21 (1.24–3.95)	2.19 (1.18–4.06)
Completed primary	2.38 (1.01–5.60)	2.26 (0.96–5.29)
Some secondary or higher	2.46 (1.17–5.19)	2.74 (1.19–6.31)
Household wealth index quartile		
Poorest	1.0	1.0
Poorer	0.63 (0.26–1.51)	0.65 (0.28–1.49)
Richer	0.86 (0.79–0.93)	0.75 (0.66–0.85)
Richest	0.47 (0.15–1.52)	0.42 (0.13–1.35)
Parity		
Multiparous	1.0	^b
Nulliparous	1.17 (0.81–1.69)	
Most recent birth by caesarean section		
No	1.0	1.0
Yes	1.84 (0.91–3.74)	1.46 (1.07–2.00)
Counselling on breastfeeding during PNC of newborn		
Yes, immediate (48 hours after birth)	1.0	^b
Yes, delayed (> 48 hours after birth)	1.15 (0.41–3.21)	
No ^c	1.04 (0.65–1.68)	
Observation of breastfeeding during PNC of newborn		
Yes, immediate (48 hours after birth)	1.0	1.0
Yes, delayed (> 48 hours after birth)	1.28 (0.57–2.88)	1.07 (0.38–3.03)
No ^c	1.10 (0.76–1.58)	0.99 (0.53–1.89)
Prelacteal feed		
No	1.0	1.0
Yes	2.31 (0.78–6.82)	2.15 (0.70–6.61)

^aMaternal age was collapsed into broader categories due to sample size limitations; ^bExcluded from the final model due to collinearity; ^cIncludes participants who did not receive counselling on/observation of breastfeeding within PNC and those who did not receive PNC at all.

Discussion

In this study, we estimated the prevalence of lactational mastitis and its associated factors in representative samples of breastfeeding women in Ethiopia, Kenya, Malawi, and Tanzania. It aimed to address the dearth of evidence on lactational mastitis in LMICs, particularly in Sub-Saharan Africa.

Analysis showed a pooled prevalence of mastitis among breastfeeding women of 6.7%, with the lowest prevalence noted in Ethiopia (3.1%) and the highest in Kenya (12.0%). More than half of women who experienced mastitis reported multiple episodes. Our estimates are somewhat lower than those reported in previous studies conducted in other settings. For instance, a small prospective cohort study of nulliparous women in Australia reported an incidence of lactational mastitis of 20% [16]. Another small cohort study conducted in Iran found that about 19% of women experienced mastitis [17]. In contrast, in China, a lower proportion of breastfeeding women reported experiencing at least one episode of mastitis (6.3%) [18]. Our pooled prevalence estimate of lactational mastitis is also higher than incidences noted in HIV-infected and uninfected women in South Africa (1% and 0.5%, respectively) [3], but closer to what has been observed in Nepal (8.0%) [19]. The prevalence we identified in Kenya is similar to that noted in a cross-sectional assessment of Ghanaian women at a breast care center (11.8%) [20].

Several reasons can explain the differences between our findings and those observed in other countries. First, there are considerable differences in the definition and measurement of mastitis across studies. For instance, some studies have considered diagnosis by health care providers [18] or the incidence of breast and systemic symptoms within a specific time frame [16], while our study was based on self-reported symptoms of mastitis during up to two years preceding participation in the household surveys. A recent systematic review and meta-analysis demonstrated substantial variability in the definitions of mastitis used in studies, but also in the population of focus (i.e., general population versus hospital-based population) [2]. Additionally, prospective cohort studies can provide more accurate measures of mastitis incidence. As such, our study, which estimated self-reported prevalence retrospectively, may be underestimating the true frequency of lactational mastitis in the four countries. Finally, differences between the countries considered in this study and other settings may be due to variabilities in maternal gut bacteria, thus leading to distinct milk microbiome, and differences in breastfeeding practices [2].

An important result of our analysis is that 17.0% of women who indicated experiencing mastitis stopped breastfeeding their infants because of mastitis-related symptoms. Continued breastfeeding despite mastitis symptoms has been observed in other settings, as it represents one way to alleviate symptoms and prevent the development of other breast complications [21]. Another explanation is that the lack of access to affordable and safe alternatives to breast milk in SSA may result in women preferring or being advised to continue breastfeeding [21]. Further quantitative and qualitative studies into the beliefs, attitudes, and practices of women experiencing mastitis and other clinical issues related to mastitis will be important to understand the enabling factors supporting continued breastfeeding.

Breastfeeding women who had some education and delivered by c-section had higher odds of reporting experiencing mastitis. These results mirror those found in other studies [19, 22]. Women who have some education may be working in formal settings preventing them from establishing and maintaining regular feeding practices to decrease the risk of milk stasis. There is also a possibility that women with higher levels of education were more likely to have understood the survey questions and therefore more likely to report experiencing mastitis symptoms.

The link between c-section and mastitis observed in this study is consistent with findings of a prospective cohort study conducted in Nepal which reported that women who gave birth by c-section were at higher risk of mastitis [19]. A potential explanation raised by the authors is that c-section is associated with delayed initiation of breastfeeding and therefore increased risk of milk stasis, a major risk factor for mastitis [19]. Other studies have also identified that, compared to women who had a vaginal birth, women who delivered by c-section have a higher proportion of breastfeeding difficulties [23].

The relationship of mastitis and household wealth quantile is interesting to note, as we expected wealth and education to be highly correlated and thus related to lactational mastitis in a similar way. Very few studies have looked at changes in the frequency of mastitis by a measure of socioeconomic status (SES), such as household wealth. However, Vogel et al. found that high SES was associated with higher odds of reported mastitis (OR = 1.13) [24], which is in accordance with the relationship we observed between education and mastitis. Another study conducted in Australia highlighted that women from wealthier households had lower odds of lactational mastitis [25]. There are possible pathways through which high SES could lead to a reduced likelihood of lactational mastitis. For instance, compared to women from poorer households, those from richer households may have greater access to health services and supportive counseling on breastfeeding and information to prevent mastitis. An alternative explanation is that women from wealthier households may have access to water for handwashing, thus limiting the transfer of bacteria to the breast during the breastfeeding process. Future studies are needed to elucidate the pathways through which SES could influence the risk of lactational mastitis.

While the confidence intervals for ORs for the relationships of maternal age and prelacteal feeding practices with mastitis were wide in our study, the direction of the ORs suggests that they may be important factors associated with mastitis. These potential determinants of mastitis should be further investigated in future studies with larger sample sizes. The available literature supports findings that lactational mastitis is higher in young women and women who did not practice prelacteal feedings than in their counterparts who are older and who did not engage in prelacteal feeding respectively [4, 19].

Strengths and Limitations

The main strength of this study is that we used large representative household surveys across four SSA countries to address the scarcity of information on the prevalence of lactational mastitis and related breastfeeding discontinuation in these settings. We also explored associations that have not been considered in the past in those countries. In addition, from the four countries, we had high participation rates (more than 95% in each country), thus limiting the risks of non-response bias.

Still, there are important limitations to be considered. First, the random walk approach used in Malawi may affect the representativeness of the data. Also, as our surveys took place in specific districts, counties, and zones within the four countries, there remains a possibility that our results would only apply to those specific areas. Another limitation is that the associations are based on cross-sectional data; we, therefore, cannot interpret any of the relationships presented here as causal. There are chances of reverse causality. For instance, mastitis could explain why women do not exclusively breastfeed during the first six months after birth. Still, for some explanatory variables such as c-section and prelacteal feeding, we can reasonably conclude that they occurred before mastitis. Another limitation is that all the variables included in this study were based on self-report, hence increasing the chances of recall bias and misclassification. Although we do not think that women are likely to underreport experiencing mastitis, to appraise potential bias in reporting, we compared mastitis among women < 6 months postpartum and those 6–24 months postpartum. We found that the odds of ever experiencing mastitis were higher among women 6–24 months, consistently with the longer time at risk of the condition. We also attempted to overcome the risk of recall bias by restricting our sensitivity analyses to women less than six-month postpartum. Some of the variables considered in this study may have also been underestimated or overestimated because of social desirability bias or challenges understanding the questions. We attempted to ease the comprehension of the questions by administering the surveys in local languages and training the data collectors on the technical interpretation of survey terms. Also, the majority of the questions (except from the mastitis questions) were framed based on standardized DHS and Multiple Indicator Cluster Survey (MICS) questionnaires.

Because of sample size limitations, we could not conduct country-specific logistic regression analyses. Similarly, while the directions of the ORs from the sensitivity analyses were aligned to those reported for the full sample, the sample size limited our ability to get precise ORs. Future studies will be needed to further document the risk factors of lactational mastitis early in the postpartum period. Finally, our surveys did not include information on other key factors such as breast characteristics and history of mastitis. More studies should explore additional variables capturing potential anatomical, socio-demographic, individual and health facility factors related to the occurrence and experience of lactational mastitis.

Conclusions

In this study, we have for the first time the prevalence of lactational mastitis in four countries of Eastern and Southern Africa. We found that between 3 and 12 in 100 women with a recent birth self-reported experiencing the condition, and more than half of them more than once. Across the four countries, we observed variability in the occurrence of mastitis among breastfeeding women. Our study suggests that younger maternal age, higher education, birth by c-section, and prelacteal feeding are associated with higher likelihood of lactational mastitis. However, further investigations into the occurrence of lactational mastitis and its predisposing factors in Africa, particularly among women up to six-months postpartum, are required to ensure optimal breastfeeding practices. Additionally, more studies on the experiences of women with lactational mastitis are needed, including the factors enabling continued breastfeeding after a mastitis episode and the support and care women prefer and desire, for breastfeeding in general, but also for mastitis in particular.

Abbreviations

AOR: Adjusted Odds Ratio

CAIA-MNCM: Canada-Africa Initiative to Address Maternal, Newborn, and Child Mortality

CHWs: Community Health Workers

CI: Confidence Interval

C-section: Caesarean section

DHS: Demographic and Health Survey

EA: Enumeration Area

MICS: Multiple Indicator Cluster Survey

LMICs: Low- and Middle-Income Countries

OR: Odds Ratio

PNC: Postnatal Care

SES: Socio-economic status

SSA: Sub-Saharan Africa

Declarations

Ethics approval and consent to participate

This study was nested within the CAIA-MNCM project, a multi-country project aiming at examining key reproductive, maternal, newborn, and child health indicators. Approvals for the CAIA-MNCM project were obtained from the Amhara Public Health Institute Research Ethics Review Committee in Ethiopia (R/T/T/D/3/634), the Amref Health Africa's Ethics and Scientific Review Committee (ESRC) in Kenya (P724/2019), the Malawi's National Health Science Research Committee (NHSRC) (19/08/2387), The United Republic of Tanzania's National Institute for Medical Research (R.8a/Vol. IX/3275), and The Hospital for Sick Children Research Ethics Board (REB) (Ethiopia: 66299, Kenya: 65962, Malawi: 65821, Tanzania: 66199). The authors confirm that all methods were performed in accordance with relevant guidelines and regulations. All participating women provided informed consent.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during this study cannot be made available, as the study participants did not agree to share their data with organizations or research groups outside of those part of the CAIA-MNCM consortium.

Competing interests

The authors declare that they have no competing interests.

Funding

This research study was nested within the Canada-Africa Initiative to Address Maternal, Newborn and Child Mortality, which was supported by Global Affairs Canada, under the Partnerships for Strengthening Maternal, Newborn, and Child Health initiative (project number: D-001988). LB is partly funded by the Research Foundation – Flanders (FWO) as part of her Senior Postdoctoral Fellowship.

Authors' contributions

LB, IOA, and MOO conceived the study and designed the analytical plan. GG, AH, HBW, MT, MOO, and IOA managed the data collection efforts in the four countries. MOO performed all data preparation and analyses, with TS providing guidance on the statistical approach. MOO, LB, and IOA wrote the manuscript with input from all authors. All authors interpreted the findings and contributed to the writing and completion of the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

The study team acknowledges the women who participated in the surveys for their time and participation, as well as all the Country project staff for their contribution to the survey data collection.

References

1. World Health Organization. Mastitis Causes and Management. 2000.
https://apps.who.int/iris/bitstream/handle/10665/66230/WHO_FCH_CAH_00.13_eng.pdf;jsessionid=3E4F318C50B6712C3E4E94F4FCD65319?sequence=1. Accessed 2 Jul 2021.
2. Wilson E, Woodd SL, Benova L. Incidence of and Risk Factors for Lactational Mastitis: A Systematic Review. *Journal of Human Lactation*. 2020;36:673–86.

3. Bland RM, Becquet R, Rollins NC, Coutoudis A, Coovadia HM, Newell ML. Breast health problems are rare in both HIV-infected and HIV-uninfected women who receive counseling and support for breast-feeding in South Africa. *Clin Infect Dis*. 2007;45:1502–10. doi:10.1086/523320.
4. Coovadia HM, Rollins NC, Bland RM, Little K, Coutoudis A, Bennish ML, et al. Mother-to-child transmission of HIV-1 infection during exclusive breastfeeding in the first 6 months of life: an intervention cohort study. *Lancet*. 2007;369:1107–16. doi:10.1016/S0140-6736(07)60283-9.
5. Tiruye G, Mesfin F, Geda B, Shiferaw K. Breastfeeding technique and associated factors among breastfeeding mothers in Harar city, Eastern Ethiopia. *International Breastfeeding Journal*. 2018;13:1–9. doi:10.1186/s13006-018-0147-z.
6. Lamberti LM, Fischer Walker CL, Noiman A, Victora C, Black RE. Breastfeeding and the risk for diarrhea morbidity and mortality. *BMC Public Health*. 2011;11 SUPPL. 3:1–12. doi:10.1186/1471-2458-11-S3-S15.
7. Lamberti LM, Zakarija-Grković I, Fischer Walker CL, Theodoratou E, Nair H, Campbell H, et al. Breastfeeding for reducing the risk of pneumonia morbidity and mortality in children under two: A systematic literature review and meta-analysis. *BMC Public Health*. 2013;13 SUPPL.3. doi:10.1186/1471-2458-13-S3-S18.
8. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: A meta-analysis. *BMC Public Health*. 2014;14. doi:10.1186/1471-2458-14-1267.
9. Sankar MJ, Sinha B, Chowdhury R, Bhandari N, Taneja S, Martinez J, et al. Optimal breastfeeding practices and infant and child mortality: A systematic review and meta-analysis. *Acta Paediatrica International Journal of Paediatrics*. 2015;104:3–13. doi:10.1111/apa.13147.
10. Chowdhury R, Sinha B, Sankar MJ, Taneja S, Bhandari N, Rollins N, et al. Breastfeeding and maternal health outcomes: a systematic review and meta-analysis. *Acta Paediatrica (Oslo, Norway: 1992)*. 2015;104 Suppl 467:96. doi:10.1111/APA.13102.
11. Krol KM, Grossmann T. Psychological effects of breastfeeding on children and mothers. *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz*. 2018;61:977. doi:10.1007/S00103-018-2769-0.
12. Kendall-Tackett K. A new paradigm for depression in new mothers: the central role of inflammation and how breastfeeding and anti-inflammatory treatments protect maternal mental health. *International Breastfeeding Journal* 2007. 2007;2:1. doi:10.1186/1746-4358-2-6. 2:1–14.
13. ICF. The DHS Program website. <https://www.dhsprogram.com/>. Accessed 18 Feb 2021.
14. RStudio. RStudio: Integrated development environment for R. 2021. <http://www.rstudio.com/>.
15. Westreich D, Greenland S. The table 2 fallacy: Presenting and interpreting confounder and modifier coefficients. *Am J Epidemiol*. 2013;177:292–8. doi:10.1093/aje/kws412.
16. Cullinane M, Amir LH, Donath SM, Garland SM, Tabrizi SN, Payne MS, et al. Determinants of mastitis in women in the CASTLE study: A cohort study. *BMC Family Practice*. 2015;16:1–8. doi:10.1186/s12875-015-0396-5.
17. Zarshenas M, Zhao Y, Poorarian S, Binns CW, Scott JA. Incidence and Risk Factors of Mastitis in Shiraz, Iran: Results of a Cohort Study. *Breastfeeding Medicine*. 2017;12:290–6. doi:10.1089/bfm.2016.0153.
18. Tang L, Lee AH, Qiu L, Binns CW. Mastitis in Chinese breastfeeding mothers: A prospective cohort study. *Breastfeeding Medicine*. 2014;9:35–8. doi:10.1089/bfm.2013.0032.
19. Khanal V, Scott JA, Lee AH, Binns CW. Incidence of Mastitis in the Neonatal Period in a Traditional Breastfeeding Society: Results of a Cohort Study. *Breastfeeding Medicine*. 2015;10:481–7. doi:10.1089/bfm.2015.0080.
20. Ohene-Yeboah M. Breast pain in Ghanaian women: clinical, ultrasonographic, mammographic and histological findings in 1612 consecutive patients - PubMed. *West Afr J Med*. 2008;20. <https://pubmed.ncbi.nlm.nih.gov/18689298/>. Accessed 5 Jul 2021.
21. de Allegri M, Sarker M, Hofmann J, Sanon M, Böhler T. A qualitative investigation into knowledge, beliefs, and practices surrounding mastitis in sub-Saharan Africa: What implications for vertical transmission of HIV? *BMC Public Health*. 2007;7:22. doi:10.1186/1471-2458-7-22.
22. Kinlay JR, O'Connell DL, Kinlay S. Risk factors for mastitis in breastfeeding women: Results of a prospective cohort study. *Aust N Z J Public Health*. 2001;25:115–20.
23. Hobbs AJ, Mannion CA, McDonald SW, Brockway M, Tough SC. The impact of caesarean section on breastfeeding initiation, duration and difficulties in the first four months postpartum. *BMC Pregnancy Childbirth*. 2016;16:1–9. doi:10.1186/s12884-016-0876-1.
24. Vogel A, Hutchison BL, Mitchell EA. Mastitis in the first year postpartum. *Birth*. 1999;26:218–25. doi:10.1046/j.1523-536x.1999.00218.x.
25. Amir LH, Garland SM, Lumley J. A case-control study of mastitis: nasal carriage of *Staphylococcus aureus*. *BMC Family Practice*. 2006;7:57. doi:10.1186/1471-2296-7-57.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile115Sept2021.docx](#)