Entanglement and Bell Inequalities : No Correlation

Eugene Scarzafava (✉ scarzafava@comcast.net)

EXCO Research (unaffiliated) <https://orcid.org/0000-0002-9627-0675>

Article

Keywords: Bell Inequalities, EPR-Bohm, EPR, entanglement, non-locality, action-at-a-distance, entangled particles

Posted Date: October 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-94476/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Entanglement and Bell Inequalities : No Correlation

E. A. Scarzafava

(Dated: October 26, 2020)

Abstract

Claims that entanglement, or non-local interaction (“spooky action-at-a-distance”), has been experimentally proven are false. The claims are based entirely on the fact that the results of experiments performed to test that issue (EPR-Bohm experiments) violate various theoretical inequalities (Bell Inequalities).

The various Bell Inequalities used to interpret the results of EPR-Bohm experiments are shown to be inappropriate for such experiments. The inequalities do not correctly provide for the correlation between the particles in each pair, which is the essence of such experiments; and therefore the inequalities are irrelevant. Claims that the results of such experiments, because they violate the inequalities, require the conclusion that the measurement of one particle in an “entangled” pair can instantly affect the state of its partner, even at great distances, are untrue and need to be modified.

I. BACKGROUND

In 1935 Einstein, Poldolsky and Rosen[1] (EPR) presented a paper in which they concluded that the quantum mechanical description of reality given by wave mechanics is not complete. They did this by first showing that either (1) the wave function description is not complete or (2) physical quantities corresponding to two non-commuting operators cannot have simultaneous reality. They described a thought experiment in which two systems start out with interactions between them, but subsequently can no longer interact. Then, starting with the assumption that the wave function does give a complete description, they showed that the two physical quantities of non-commuting operators can have simultaneous reality. The negation of alternative (1) leads to the negation of alternative (2), and thus their stated conclusion. They stressed the fact that by simultaneously real they are referring to quantities that either one or the other, but not not both simultaneously, of the two quantities can be predicted. They posed the locality condition that the process of measurement carried out on the first system does not disturb the second system in any way.

It is difficult to carry out such experiments, but in 1957 Bohm and Aharonov[2] described experiments that can be performed that test the concept of the EPR paradox. In one case, two electrons in a singlet spin state are separated by a method that does not alter the total spin and, when they are sufficiently far apart, measurements are made on the two systems. In another case, two photons are created with opposite momentum and with each photon in a polarization state that is orthogonal to the other. The experiments were still difficult to carry out with the technology available at that time, but preliminary results appeared to favor the view that measurements on one system affected measurements on the correlated system that was spatially distant from the first. They did allow for the possibility that quantum theory is not complete and may be an approximation arising from “a deeper subquantum-mechanical level”.

In 1964 Bell[3] derived a set of inequalities that he claimed must apply to any system, even with hidden variables, if that system were to meet the requirements of locality (i.e., measurements on one system can not affect, nor be affected by measurements on the other). Using the example of of an experiment based on the Bohm-Aharonov experiment[2] (EPR-Bohm experiments), and imposing his locality requirement, Bell developed inequalities that were incompatible with the statistical predictions of quantum mechanics.

Subsequent experiments, such as those using an optical analog of the Stern-Gerlach filters of Bell's paper, have shown dramatic violations of variations of Bell's inequalities, while being in good agreement with quantum mechanical predictions.[4] This has led to the conclusion that experiments show that a measurement on either member of an "entangled" pair has an instantaneous effect on the state of the other, regardless of the distance between them (spooky action-at-a-distance). That view is reflected in college textbooks and material intended to educate and inform the general public.[5] In the next section we show that view to be incorrect, because inappropriate localization requirements were used in the development of the inequalities.

II. CRITIQUE OF THE BELL INEQUALITIES

In this paper it is shown that violation of the Bell Inequalities does not prove non-local, action at a distance behavior for "entangled" particles. The methods used to impose locality in the development of the inequalities make them meaningless in the interpretation of the experimental results of EPR-Bohm experiments. There are several variations of inequalities used to interpret the EPR-Bohm experiments, all of which are referred to generically as Bell Inequalities. In this section it is shown that they all use localization methods that are incorrect for such experiments.

In 1978 Clauser and Shimony[6] provided an excellent review of the then current Bell Inequalities. This paper relies heavily on their review. In Section 3.3 of Clauser and Shimony they discuss the locality concept and provide a very reasonable definition:

Suppose a pair of correlated systems, which have a joint state λ , separate. They then continue to evolve perhaps in an inherently stochastic way, and given λ , a and b , one can define probabilities for any particular outcome at either apparatus. We allow that, given λ , these two probabilities may each depend on the associated (local) apparatus parameter, a or b respectively, and of course upon λ , but we assume that these probabilities are otherwise independent of each other.

This definition of locality seems very common-sensical. It says that the outcome (or the probability of outcomes) of a measurement performed on one part of a composite system is independent of what aspects of the other component the

77 experimenter chooses to measure. It by no means excludes the possibility of
 78 obtaining knowledge concerning system 2 from an examination of system 1. The
 79 state λ contains information common to both systems, and a measurement on
 80 one of these presumably reveals some of this. Nor does it prevent a measurement
 81 performed on one component of a composite system from locally disturbing that
 82 component. What it does prescribe, in essence, is that the measured value of a
 83 quantity on one system is not causally affected by what one chooses to measure
 84 on the other system, since the two systems are well separated (e.g. space-like
 85 separated) when the measurements are performed.[6]

86 It is difficult to design experiments to test the original Bell Inequalities; so experiments
 87 have been designed to be interpreted by variations of the inequalities, many of which are
 88 discussed in the following subsections.

89 **A. Clauser and Horne**

90 We start with Clauser and Horne[7]. This is because it most simply demonstrates the
 91 localization problem with Bell Inequalities and because Clauser-Horne(CH) type Bell In-
 92 equalities were commonly used for EPR-Bohm experiments. In this description of their de-
 93 velopment many of the details and explanations are glossed over, but much of it is straight
 94 from their paper.

95 Clauser and Horne, describe a typical EPR-Bohm experiment, using an experimental
 96 arrangement in which a source of coincident two-particle emissions is viewed by two analyzer-
 97 detector assemblies, 1 and 2. Each apparatus has an adjustable parameter, a at apparatus
 98 1 and b at apparatus 2, where a and b are angles specifying the orientation of the analyzers,
 99 e.g., the axes of linear polarizers for photons or the directions of the field gradients of Stern-
 100 Gerlach magnets for fermions.

101 During a period of time, while the adjustable parameters have the values a and b , the
 102 source may emit N of the two-particle systems of interest. Let $N_1(a)$ and $N_2(b)$ be the
 103 number of counts at detectors 1 and 2, respectively, and let $N_{12}(a, b)$ be the number of
 104 simultaneous counts from the two detectors (coincident counts). When N is sufficiently large
 105 the ensemble probabilities are given by Clauser and Horne's Equations (1):

$$p_1(a) = N_1(a)/N$$

$$p_2(b) = N_2(b)/N$$

$$p_{12}(a, b) = N_{12}(a, b)/N$$

Consider one of the two-component emissions from the source and consider the state specification of that system at a time intermediate between its emission and its impingement on either apparatus, and denote that state by λ . Let the probabilities of a count being triggered at apparatus 1 and 2 be $p_1(\lambda, a)$ and $p_2(\lambda, b)$, respectively, and let $p_{12}(\lambda, a, b)$ be the probability that both counts are triggered. Then, since every system in the ensemble may not have the same initial state, let $\rho(\lambda)$ be the normalized probability density characterizing the ensemble of emissions. Then the ensemble probabilities given in their Equations (1), above are Clauser and Horne's Equations (2)

$$p_1(a) = \int_{\Gamma} d\lambda \rho(\lambda) p_1(\lambda, a),$$

$$p_2(b) = \int_{\Gamma} d\lambda \rho(\lambda) p_2(\lambda, b),$$

$$p_{12}(a, b) = \int_{\Gamma} d\lambda \rho(\lambda) p_{12}(\lambda, a, b)$$

where Γ is the space of the states λ .

This is where the crucial error in their development is made. In the equation for the ensemble probability for the coincident counts they make the substitution, their Equation (2')

$$p_{12}(\lambda, a, b) = p_1(\lambda, a) p_2(\lambda, b) \tag{1}$$

and they justify this as a reasonable locality condition, which leads to

$$p_{12}(a, b) = \int_{\Gamma} d\lambda \rho(\lambda) p_1(\lambda, a) p_2(\lambda, b) \tag{2}$$

However, this factorization is definitely not a reasonable locality condition. It is an unacceptable one, because it erases the correlation between the two particles that were emitted. It defeats the whole purpose of the EPR-Bohm experiments. The substitution is,

of course, mathematically at fault in that the sum of the product of two quantities is not, in general, equal to the product of their sums. Thus also, $N_{12}(a, b)$, which is the sum of all coincident counts, would not, in general, be equal to the product of $N_1(a)$ with $N_2(b)$, which are the sums of all counts at each detector.

A great deal of time, effort and ingenuity is expended by the experimenters to construct the sophisticated apparatus used to create correlated particle pairs and to ensure that their spin or polarization states are not disturbed by external means until they encounter the detectors. The two particles in a given pair are correlated (complementary) to each other, but do not specifically share that correlation with any other particles. Thus one needs to be careful in how one counts, especially in accounting for the sums of coincident counts.

[Note: It could be said that correlation between the particles is included in Equation (2) through the use of λ , which can represent shared variables. Each particle shares a common set of values or instructions and thus two particles effectively are correlated when each separately interacts with its detector. This position fails on several levels. (I) The correlation (complementarity) is specific for each separate pair of particles. It does not apply to particles in the aggregate. (II) No use is made of any potential correlation in deriving the inequalities. (III) The correlation of interest is the fact that the spins or polarizations of the particles are complementary, because they were created in a singlet state. Hidden variables may explain why each particle exhibits a certain value when measured, but are not the source of the correlation of interest. (IV) Use of this factored probability is equivalent to deterministic models, such as those discussed in Section II.B, below, which make no use of hidden variables and yield precisely the same inequalities. Disjoint subspaces also do not provide for complementarity in the measurements for each pair.]

Clauser and Shimony[6] in their Section 3.6 and Equations (3.21) assume from symmetry considerations that

$p_1(a)$ is independent of b

$p_2(b)$ is independent of a

and $p_{12}(a, b) \equiv p_{12}(|a - b|)$

158 That is a much more reasonable expression for $p_{12}(a, b)$ than Equation (2) above, because
 159 it implies correlation between the particles. If the spins or polarizations of the particles
 160 in a pair are complementary, the probability of triggering both detectors for a given pair
 161 necessarily depends on the relative orientation of detector a when it interacts with particle
 162 1 and detector b when it interacts with particle 2. To arbitrarily ignore the dependence
 163 of coincidence on the relative orientation of the detectors by replacing it with the product
 164 of probabilities of each particle triggering its separate detector, regardless of the relative
 165 orientation of the detectors, essentially erases the complementary correlation.

166 However, Clauser and Horne use Equation (2), which denies the correlation between
 167 the particles at the time of the measurement, to obtain their inequality. They prove a
 168 mathematical lemma and, using that lemma, establish the inequality

$$\begin{aligned} -1 \leq p_1(\lambda, a)p_2(\lambda, b) - p_1(\lambda, a)p_2(\lambda, b') + p_1(\lambda, a')p_2(\lambda, b) \\ + p_1(\lambda, a')p_2(\lambda, b') - p_1(\lambda, a') - p_2(\lambda, b) \leq 0 \end{aligned} \quad (3)$$

169 where a and a' are two orientations of analyzer 1, and b and b' are two orientations of
 170 analyzer 2. There is nothing wrong with this, but then they use Equation (2) to obtain their
 171 famous inequality

$$\begin{aligned} -1 \leq p_{12}(a, b) - p_{12}(a, b') + p_{12}(a', b) + p_{12}(a', b') \\ - p_1(a') - p_2(b) \leq 0 \end{aligned} \quad (4)$$

172 The use of this inequality to interpret the results of EPR-Bohm experiments is, for the
 173 reasons stated above, meaningless.

174 B. Wigner, Belinfante and Holt

175 Clauser and Shimony[6], in Section 3.7 discuss Bell Inequalities developed independently
 176 by Wigner[8], Belinfante[9], and Holt[10]. Their method consists of subdividing the space
 177 Λ of states of a two-component system into subspaces corresponding to various possible
 178 values of the observables of interest, and then performing calculations on the measures of
 179 these subspaces. Clauser and Shimony show how their method can be used to derive the
 180 CH Inequality (Equation (4), above). That should be enough to disqualify them from being

181 used for EPR-Bohm experiments, but it is instructive to see why they are equivalent, using
 182 Clauser and Shimony's analysis.

183 They start with the assumption that the detection or non-detection of component 1
 184 is completely determined by the parameter a of the first analyzer and the state of the
 185 composite system, but is independent of the parameter b of the other analyzer, and likewise
 186 for component 2. They then assume that determinism applies and exhaustively subdivide
 187 the space Λ into 16 mutually disjoint subspaces $\Lambda(ij;kl)$, where each letter can take the
 188 value 0 or 1, with 1 denoting detection and 0 non-detection.; with i and j referring to the
 189 results if the parameter of the first analyzer is chosen respectively to be a or a' ; and with k
 190 and l referring to the results if the parameter of the second analyzer is chosen respectively to
 191 be b or b' . If a probability measure ρ is assumed to be given on Λ , then $\rho(ij;kl)$ is defined to
 192 be the probability that the composite state is in $\Lambda(ij;kl)$. All the $\rho(ij;kl)$ are non-negative,
 193 and because the 16 subspaces are disjoint and exhaustive, we have:

$$\sum_{ijkl} \rho(ij;kl) = 1$$

194 We now define $p_1(a)$ to be the probability that component 1 will be detected if the parameter
 195 is chosen to be a ; $p_2(b)$ to be the probability that component 2 will be detected if
 196 its parameter is chosen to be b ; and $p_{12}(a,b)$ to be the probability of joint detection of
 197 both components if the two parameters are chosen respectively to be a and b . Analogous
 198 definitions are given for the values of the other parameters. Then, for example,

$$p_{12}(a,b) = \rho(11;11) + \rho(11;10) + \rho(10;11) + \rho(10;10)$$

199

$$p_{12}(a,b') = \rho(11;11) + \rho(11;01) + \rho(10;11) + \rho(10;01)$$

200 etc., and, for example,

$$\begin{aligned} p_2(b) &= \rho(11;11) + \rho(11;10) + \rho(10;11) + \rho(10;10) \\ &\quad + \rho(01;11) + \rho(01;10) + \rho(00;11) + \rho(00;10) \end{aligned}$$

201 Using these, and equivalent equations for the other probabilities, Clauser and Shimony show
 202 that Equation (4), above, is recovered. This deterministic approach, because of its disjoint
 203 subspaces, also does not allow for correlation between each pair of particles; so it is not
 204 surprising that it is equivalent to the CH Inequality, a fact that is supported by Fine[11]
 205 (see Section II.F below), and, thus is inappropriate for EPR-Bohm experiments

C. Bell

The concept of Bell Inequalities was introduced by Bell[3] in 1964. Bell's paper dealt with an EPR-Bohm experiment using fermions in a singlet state moving in opposite directions. Measurements are made by Stern-Gerlach magnets on selected components of the spins $\vec{\sigma}_1$ and $\vec{\sigma}_2$. If measurement of the component $\vec{\sigma}_1 \cdot \vec{a}$ where $\vec{a}$ is some unit vector, yields the value -1 then, according to quantum mechanics, measurement of $\vec{\sigma}_2 \cdot \vec{a}$ must yield the value -1 and vice versa. Since this is true even when the measurements are made at places remote from one another, it implies the possibility of a more complete specification of the state, as proposed by EPR, which he denotes by λ , where λ is quite general, denoting whatever is needed for a complete specification.

The result A of measuring $\vec{\sigma}_1 \cdot \vec{a}$ is then determined by $\vec{a}$ and λ , and the result B of measuring $\vec{\sigma}_2 \cdot \vec{b}$ in the same instance is determined by $\vec{b}$ and λ ; so

$$A(\vec{a}, \lambda) = \pm 1 \text{ and } B(\vec{b}, \lambda) = \pm 1 \quad (5)$$

Then he states that if $\rho(\lambda)$ is the probability distribution of λ , the expectation value of the product of the two components $\vec{\sigma}_1 \cdot \vec{a}$ and $\vec{\sigma}_2 \cdot \vec{b}$ is

$$P(\vec{a}, \vec{b}) = \int d\lambda \rho(\lambda) A(\vec{a}, \lambda) B(\vec{b}, \lambda) \quad (6)$$

which he states is equivalent to the quantum mechanical expectation value

$$\langle \vec{\sigma}_1 \cdot \vec{a} \vec{\sigma}_2 \cdot \vec{b} \rangle = -\vec{a} \cdot \vec{b}$$

The factorization in Equation (6), as pointed out by Clauser and Shimony in their Section 3.1 Equation (3.4), is an expression of locality for a deterministic hidden-variables theory. As discussed in Section A, above, it is not an acceptable locality condition.

Thus, there is no need to discuss Bell's development further, because the inequalities are inappropriate, are never used anyway and the development contains some dubious steps. In 1971 Bell[12] offered an updated and improved proof, but it contains the same factorization as Equation (6) and thus also is inappropriate.

229 In a paper describing the “S-Matrix Interpretation of Quantum Theory” Stapp[13] dis-
 230 cusses the completeness of quantum theory with the use of an EPR-Bohm experiment. Two
 231 fermions in a singlet spin state, traveling in opposite directions, each pass through a Stern-
 232 Gerlach type device. The axes of the two devices (A_1 and A_2) are denoted by D_1 and D_2 .
 233 They are both normal to the c.m. line of flight, and $\theta(D_1, D_2)$ is the angle between them.
 234 Two different settings, D'_1 and D''_1 of D_1 are considered; and two different settings, D'_2 and
 235 D''_2 of D_2 are considered. Altogether four alternative combinations of settings are consid-
 236 ered. Let j label the individual experiment, i.e., a single pair of particles in a singlet state.
 237 Then let $n_{1j}(D_1, D_2)$ be defined to be +1 or -1, according to whether the theory predicts
 238 that the particle from the j th pair that passes through A_1 is deflected up or down, when the
 239 settings of the axes are D_1 and D_2 . The numbers $n_{2j}(D_1, D_2)$ are defined analogously for the
 240 other particle. Based on the condition that the theory gives predictions for various possible
 241 alternative settings, for each individual pair j , the numbers $n_{1j}(D_1, D_2)$ and $n_{2j}(D_1, D_2)$ are
 242 defined for all four of the arguments D_1 and D_2 .

243 Stapp then discusses the fact that, based on his predictability condition, the eight num-
 244 bers $n_{ij}(D_1, D_2)$ are defined for each experiment j . Of these eight numbers only two can
 245 be compared to experiment. The other six correspond to the three alternative experiments
 246 that could have been performed but were not. He then states that if the experimenters
 247 had actually adjusted the mechanical devices to give the alternative experimental setup,
 248 then the alternative experiments would have had certain definite results. Then the numbers
 249 $n_{1j}(D_1, D_2)$ and $n_{2j}(D_1, D_2)$ can be defined by the results that the experiments would have
 250 had if they had actually been performed. (This is what is called “counterfactual definite-
 251 ness” and Eberhard[14] refers to as “conjugate events”.) This is an interesting concept and
 252 potentially could be used to introduce correlation into the theory, because there is a corre-
 253 lation between $n_{1j}(D_1, D_2)$ and $n_{2j}(D_1, D_2)$ for each j . It’s not clear how to do that, and
 254 there is no indication that Stapp attempted to do so.

255 Stapp then states that the following relationship with quantum mechanics should hold
 256 with increasing accuracy as N increases:

$$\frac{1}{N} \sum_{j=1}^N n_{1j}(D_1, D_2) n_{2j}(D_1, D_2) = -\cos\theta(D_1, D_2) \quad (7)$$

It is clear that this suffers from the same factorization problem of Equation (1) in Section II.A, above. No provision is made for a correlated $n_{12j}(D_1, D_2)$. Because there is no correlation between the particles, the method is meaningless for the interpretation of EPR-Bohm experiments. Clauser and Shimony point out that in 1978 Stapp[15] responded to some critics of his earlier paper and also provided an equivalence theorem that shows his approach is, in fact, equivalent to Equation (2) in Section II.A, above, confirming that it is an inappropriate measure.

E. Eberhard

1. Eberhard 1977

In 1977 Eberhard[16] published a paper purporting to derive Bell Inequalities without using the concept of hidden variables. He considers an EPR-Bohm experiment with two measuring locations, A and B, at a distance from each other, with a knob, a, on apparatus A and a knob, b, on apparatus B. N events are recorded, with a measurement on both A and B. There are only two outcomes for each measurement, +1 or -1. The j th event corresponds to a response α_j in A and β_j in B. Each term α_j and β_j is either +1 or -1. He defines correlation by a statistical mean

$$C = \frac{1}{N} \sum_j \alpha_j \beta_j = \langle \alpha_j \beta_j \rangle$$

where the brackets around a quantity designates the statistical mean of that quantity over j. C is equal to the fraction of events when α_j and β_j have the same signs minus the fraction where they have opposite signs. There are two positions $a^{(1)}$ and $a^{(2)}$ of the knob a and two positions, $b^{(1)}$ and $b^{(2)}$ for the knob b. He defines:

$$C^{(1,1)} = \langle \alpha_j \beta_j \rangle, \quad a = a^{(1)}, b = b^{(1)}$$

$$C^{(2,1)} = \langle \alpha_j \beta_j \rangle, \quad a = a^{(2)}, b = b^{(1)}$$

$$C^{(1,2)} = \langle \alpha_j \beta_j \rangle, \quad a = a^{(1)}, b = b^{(2)}$$

$$C^{(2,2)} = \langle \alpha_j \beta_j \rangle, \quad a = a^{(2)}, b = b^{(2)}$$

and then defines for event j:

$$\gamma_j = \alpha_j^{(1)} \beta_j^{(1)} + \alpha_j^{(2)} \beta_j^{(1)} + \alpha_j^{(1)} \beta_j^{(2)} - \alpha_j^{(2)} \beta_j^{(2)} \quad (8)$$

Each product $\alpha_j \beta_j$ is equal to ± 1 ; so therefore

$$\begin{aligned} \frac{1}{N} \sum_j \gamma_j = \langle \gamma \rangle &= \langle \alpha_j^{(1)} \beta_j^{(1)} \rangle + \langle \alpha_j^{(2)} \beta_j^{(1)} \rangle \\ &+ \langle \alpha_j^{(1)} \beta_j^{(2)} \rangle - \langle \alpha_j^{(2)} \beta_j^{(2)} \rangle \leq 2 \end{aligned} \quad (9)$$

and finally

$$C^{(1,1)} + C^{(2,1)} + C^{(1,2)} - C^{(2,2)} \leq 2 \quad (10)$$

which is the famous CHSH[17] inequality.

[It should be noted that the CHSH Inequality is obtained using a “correlation function”, $P(a, b) = \sum_{\Gamma} A(a, \lambda) B(b, \lambda) \rho(\lambda) d\lambda$ which is a factored form that actually denies correlation of the particles. They don’t even refer to a quantity like $AB(a, b, \lambda)$; so, clearly the CHSH Inequality suffers the same problem as the CH Inequality discussed in Section II.A, above, and is inappropriate for EPR-Bohm experiments.]

The statistical means used in the development by Eberhard also do not provide for correlation between the particles, which is why he reaches the same inappropriate inequality.

2. Eberhard 1992

In 1992 Eberhard[14] presented an alternative inequality based on an entirely different approach. The initial state is prepared as a superposition of two photons a and b, with correlated planes of polarization. One state is defined with photon a polarized vertically and photon b polarized horizontally while the other state has photon a polarized horizontally and photon b vertically. The angle of the detection device for photon a is denoted α and that of photon b as β .

The Bell Inequalities are concerned with expectation values of quantities that can be measured in four different experimental setups defined by specific values $\alpha_1, \alpha_2, \beta_1$ and β_2 of α and β . The setups are referred to by the symbols $(\alpha_1, \beta_1), (\alpha_1, \beta_2), (\alpha_2, \beta_1)$, and (α_2, β_2) , where the first index designates the value of α and the second index the value of β .

In each setup, the “fate” of photon a and the “fate” of photon b is referred to by an index (o) for photons detected in the ordinary beam, (e) for photons detected in the extraordinary beam, or (u) for photons undetected.. Therefore there are nine types of events: (o,o), (o,u),

306 (o,e), (u,o), (u,u) (u,e), (e,o), (e,u) and (e,e), where the first index designates the fate of
 307 photon a and the second index the fate of photon b. He defines conditions:

308 (i) The fate of photon a is independent of the value of β , i.e., is the same in an event of
 309 the sequence corresponding to setup (α_1, β_1) as in the event with the same event number
 310 k for (α_1, β_2) ; also same fate for a in (α_2, β_1) and (α_2, β_2) ; this is true for each k for these
 311 sequences.

312 (ii) The fate of photon b is independent of the value of α , i.e., is the same in event k for
 313 sequences (α_1, β_1) and (α_2, β_1) ; also same fate for b in sequences (α_1, β_2) and (α_2, β_2) .

314 These are perfectly reasonable locality conditions, but they only apply for each given k,
 315 not for the sums over all k. Note that the relative orientations of the detectors for a given
 316 k will not, in general, be the same as that for any other k's.

317 Following an argument first used by Stapp[13], when four sequences are found satisfying
 318 conditions (i) and (ii), the four events with the same event number k in the four sequences
 319 are called “conjugate events”. He then uses the notation $n_{f_1 f_2}(\alpha, \beta)$, where f_1 is the fate
 320 (o,e,or u) of photon a and f_2 is the fate of photon b, for the number of events with that
 321 outcome. For example, $n_{oo}(\alpha_1, \beta_1)$ is the number of events with the outcome o for photon a
 322 and with the outcome o for photon b when α is α_1 and β is β_1 .

323 [Note: This is the number of events with that outcome, not an event for a single
 324 value of k, and that the concept of “conjugate event” does not really apply.
 325 Nevertheless, Eberhard proceeds to develop his inequality using these totals,
 326 which for notational purposes he calls boxes.]

327 He starts by considering $n_{oo}(\alpha_1, \beta_1)$. He then subtracts from that total some of the boxes
 328 for which events, had they been for the same k, would have been conjugate to events that
 329 go into the box $n_{oo}(\alpha_1, \beta_1)$.

$$\begin{aligned} n_{oo}(\alpha_1, \beta_1) - n_{uo}(\alpha_2, \beta_1) - n_{eo}(\alpha_2, \beta_1) \\ - n_{ou}(\alpha_1, \beta_2) - n_{oe}(\alpha_1, \beta_2) \end{aligned} \quad (11)$$

330 Because the remainder shown in Expression(11), above must be less than or equal to the
 331 number in box $n_{oo}(\alpha_2, \beta_2)$, he obtains:

$$\begin{aligned} n_{oo}(\alpha_1, \beta_1) - n_{uo}(\alpha_2, \beta_1) - n_{eo}(\alpha_2, \beta_1) - n_{ou}(\alpha_1, \beta_2) \\ - n_{oe}(\alpha_1, \beta_2) \leq n_{oo}(\alpha_2, \beta_2) \end{aligned} \quad (12)$$

332 , or his inequality:

$$\begin{aligned}
& n_{oe}(\alpha_1, \beta_2) + n_{ou}(\alpha_1, \beta_2) + n_{eo}(\alpha_2, \beta_1) + n_{uo}(\alpha_2, \beta_1) \\
& + n_{oo}(\alpha_2, \beta_2) - n_{oo}(\alpha_1, \beta_1) \geq 0
\end{aligned} \tag{13}$$

333 He then claims that Equation (13) is equivalent to Equation (4) of Section II.A, above. He
334 also points out that if the o's and e's in Equation (13) are interchanged one obtains a similar
335 inequality, which, if averaged with Equation (13) yields an inequality essentially identical to
336 the CHSH Inequality mentioned in Section II.E.1, above. That is not surprising, considering
337 his use of boxes, rather than reserving the correlation for individual pairs. Clearly, this
338 makes his inequality inappropriate for EPR-Bohm experiments for the reasons discussed
339 above.

340 3. *Giustina et.al*

341 Another form of Eberhard's inequality was developed by Giustina, et.al.[18]. In an
342 experiment, one records measurements of singles counts, S (number of detection events
343 on one side), and C (number of detected pairs), for the four combination of settings
344 $(\alpha_1, \beta_1), (\alpha_1, \beta_2), (\alpha_2, \beta_1)$ and (α_2, β_2) .

345 The number of events for which one of the outcomes is undetected follows from the mea-
346 sured rates. For a given measurement length the measured coincident counts are $C_{kl}(\alpha_i, \beta_j)$
347 and the single counts are $S_k^\alpha(\alpha_i)$ and $S_l^\beta(\beta_j)$, where k and l are o or e. The terms in Eber-
348 hard's inequality, Equation (13) above, are then given by the following measured quantities:

$$\begin{aligned}
& n_{oo}(\alpha_1, \beta_1) = C_{oo}(\alpha_1, \beta_1) \\
& n_{oe}(\alpha_1, \beta_2) = C_{oe}(\alpha_1, \beta_2) \\
& n_{ou}(\alpha_1, \beta_2) = S_o^\alpha(\alpha_1) - C_{oo}(\alpha_1, \beta_2) - C_{oe}(\alpha_1, \beta_2) \\
& n_{eo}(\alpha_2, \beta_1) = C_{eo}(\alpha_2, \beta_1) \\
& n_{uo}(\alpha_2, \beta_1) = S_o^\beta(\beta_1) - C_{oo}(\alpha_2, \beta_1) - C_{eo}(\alpha_2, \beta_1) \\
& n_{oo}(\alpha_2, \beta_2) = C_{oo}(\alpha_2, \beta_2)
\end{aligned}$$

354 Inserting these expressions into Equation (13) above yields:

$$\begin{aligned}
& -C_{oo}(\alpha_1, \beta_1) + S_o^\alpha(\alpha_1) - C_{oo}(\alpha_1, \beta_2) \\
& + S_o^\beta(\beta_1) - C_{oo}(\alpha_2, \beta_1) + C_{oo}(\alpha_2, \beta_2) \geq 0
\end{aligned} \tag{14}$$

,where the coincidence counts $C_{oe}(\alpha_1, \beta_2)$ and $C_{eo}(\alpha_2, \beta_1)$ have dropped out. Quite cleverly, the resulting inequality now contains only directly available detection events related to the ordinary beams of α and β . This inequality was used by Giustina, et. al., in their experiment and was used in the recent NIST experiment[4]. Obviously it suffers from the same lack of correlation as the Eberhard[14] inequality above, upon which it is based.

F. Summary

The list of Bell Inequalities discussed above is not exhaustive and each is not dealt with rigorously, just enough to give a flavor of the development and to show that it does not provide for correlation between the correlated particles. It also shows that it does not matter whether a deterministic hidden-variables model or a factorized stochastic model is used. They all erase the correlation between the particles in a pair.

This is not surprising in light of a 1982 paper by Fine[11] which proves that every deterministic hidden-variables model is equivalent to a factorizable stochastic model. (See also Appendix A of Hall[19].) The factorized probability makes these models unsuited, even irrelevant, for use in interpreting the results of EPR-Bohm experiments. The probability for the factorized model

$$p_{12}(a, b) = \int_{\Lambda} d\lambda \rho(\lambda) p_1(\lambda, a) p_2(\lambda, b) \quad (15)$$

correctly accounts for the locality of the measuring devices and the fact that a measurement at one detector does not affect, and is not affected by, the measurement at the other detector, which is all that EPR requires. Unfortunately, however, this arbitrary localization condition is too severe, in that it also effectively erases the correlation that is specific to each particle pair. For the EPR-Bohm experiments, $p_{12}(\lambda, a, b)$ is definitely not equal to $p_1(\lambda, a) p_2(\lambda, b)$, except for an individual pair. Thus one must conclude, for the reasons expressed in the discussion of Equation(1) and Equation(2) of Section II.A above, that the Bell Inequalities are inappropriate for use in interpreting the results of EPR-Bohm experiments.

III. CONCLUSION

It is shown that none of the Bell Inequalities have any relevance for EPR-Bohm experiments. The extreme locality conditions they use do not account for the correlation between

the particles in a pair; but it is the correlation between the particles in a pair that is the characteristic feature of such experiments. The only locality condition required by EPR is that a measurement on one of the particles is not affected by and can not affect the measurement made on the second particle. There is therefore no reason why, in an EPR-Bohm experiment, the inequalities should not be violated; in fact one should expect that they would be violated, because they express conditions that are decidedly untrue if the experimenters did their jobs well.

In light of this it is incorrect to claim that violation of Bell Inequalities requires the conclusion that the measurement of one particle in an “entangled” pair can instantly affect the state of its partner, even at a great distance. This action at a distance may be true, but that is not proved by the results of any EPR-Bohm experiments yet performed. It’s also possible that Einstein, Podolsky and Rosen are correct in claiming that there is simultaneous reality for the quantities being measured and thus quantum mechanics provides an incomplete description.

Because the experiments have been performed at space-like separations, there are really two possible interpretations: either

(1) the spin/polarization state of each particle has simultaneous reality, and a measurement on one, though related to, does not affect a measurement on the other, or

(2) contrary to all of our hard-earned current understanding of space-time, the spin/polarization state of each particle only becomes real at the time of measurement and the measurement of one immediately determines the state of the other, even when they are space-like separated.

Although each interpretation is, perhaps, possible, it is erroneous to claim that violation of Bell Inequalities requires the second.

[1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).

[2] D. Bohm and Y. Aharonov, Phys. Rev. **108**, 1070 (1957).

[3] J. S. Bell, Physics **1**, 195 (1964).

[4] A. Aspect, P. Grangier, and G. Roger, Phys. Rev. Lett. **49**, 91 (1982); A. Aspect, J. Dalibard, and G. Roger, *ibid.* **49**, 1804 (1982); L. K. Shalm, E. Meyer-Scott, B. G. Christensen, P. Bierhorst, M. A. Wayne, M. J. Stevens, T. Gerrits, S. Glancy, D. R. Hamel, M. S. Allman,

411 *et al.*, *ibid.* **115**, 250402 (2015).

412 [5] D. J. Griffiths, *Introduction to Quantum Mechanics* (Benjamin Cummings, 2004); B. Greene,
413 *The Fabric of the Cosmos* (Vintage Books, 2005); D. I. Kaiser, *How the Hippies Saved Physics*
414 (W. W. Norton, 2011); L. Ost, *NIST Team Proves 'Spooky Action at a Distance' is Real*
415 (2015), url-<http://www.nist.gov/pml/div686/20151105loophole.cfm>; D. Powell, *Discover* **37**,
416 58 (2016).

417 [6] J. F. Clauser and A. Shimony, *Rep. Prog. Phys.* **41**, 1881 (1978).

418 [7] J. F. Clauser and M. A. Horne, *Phys. Rev. D* **10**, 526 (1974).

419 [8] E. P. Wigner, *Am. J. Phys.* **38**, 1005 (1970).

420 [9] F. J. Belinfante, *A Survey of Hidden-Variables Theories* (Oxford: Pergamon, 1973).

421 [10] R. A. Holt, *Atomic Cascade Experiments*, Ph.D. thesis, Harvard University (1973).

422 [11] A. Fine, *Phys. Rev. Lett.* **48**, 291 (1982).

423 [12] J. S. Bell, *Foundations of Quantum Mechanics* (New York: Academic, 1971) pp. 171–81.

424 [13] H. P. Stapp, *Phys. Rev. D* **3**, 1303 (1970).

425 [14] P. H. Eberhard, *Phys. Rev. A* **47**, R747 (1993).

426 [15] H. P. Stapp, *Found. of Phys.* **9**, 1 (1979).

427 [16] P. H. Eberhard, *Nuovo Cim.* **38B**, 75 (1977).

428 [17] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).

429 [18] M. Giustina, A. Mech, S. Ramelow, B. Wittmann, J. Kofler, J. Beyer, A. Lita, B. Calkins,
430 T. Gerrits, *et al.*, *Nature* **497**, 227 (2013).

431 [19] M. J. W. Hall, *Phys. Rev. A* **84**, 022102 (2011).