Information-Motivation-Behavioral Skills Model of Consistent Condom Use among Transgender Women in Shenyang, China

Huwen Wang

Shanghai Jiao Tong University School of Public Health

Ruijie Chang

Shanghai Jiao Tong University School of Public Health

Qiuming Shen

Shanghai Jiao Tong University School of Public Health

Lhakpa Tsamlag

Shanghai Jiao Tong University School of Public Health

Shuxian Zhang

Shanghai Jiao Tong University School of Public Health

Yue Shi

Shanghai Jiao Tong University School of Public Health

Tiecheng Ma

China Love Aid

Zixin Wang

Chinese University of Hong Kong

Rui She

Chinese University of Hong Kong

Ying Wang

Shanghai Jiao Tong University School of Public Health

Joseph TF Lau

Chinese University of Hong Kong

Yong Cai (✉ caiyong202028@hotmail.com)

Shanghai Jiao Tong University School of Public Health <https://orcid.org/0000-0002-5706-7906>

Research article

Keywords: Transgender women, Condom use, Information-Motivation-Behavioral Skills model, HIV/AIDS, China

Posted Date: December 12th, 2019

DOI: <https://doi.org/10.21203/rs.2.18841/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The Information-Motivation-Behavioral Skills (IMB) model has received consistent empirical support in the context of HIV prevention among various key populations, but not yet among transgender women (TGW). None effective interventions has been carried out among TGW so far to control their high prevalence of unprotected anal intercourse. The intent of the current study is to examine the application of the IMB model to clarifying the association between condom use correlates and condom use frequency among TGW in China.

Methods: Using snowball sampling, we recruited 198 self-identified TGW in Shenyang, China from April 2017 to July 2017. Participants were required to complete a questionnaire assessing their background characteristics and IMB model constructs. Structural equation modeling (SEM) was conducted to demonstrate the utility of the IMB model. **Results:** The consistent condom use (CCU) rate was 47.0%.

Results of SEM indicated that HIV-preventive motivation (comprising condom use attitude and subjective norms; $\beta = 0.823$, $P < 0.001$) and behavioral skills (including condom use skills and self-efficacy; $\beta = 0.979$, $P = 0.004$) were related to more frequent condom use, whereas HIV knowledge was unrelated to condom use ($\beta = 0.052$, $P = 0.540$).

Conclusions: The low CCU rate suggested that TGW in China were at high risk of HIV infection and transmission and a key intervention population. HIV-preventive interventions for this population should focus on enhancing motivation and strengthening behavioral skills to increase condom use frequency and reduce HIV infection.

Background

Transgender women (TGW) are those whose gender identity (woman) is different from their birth-assigned sex (male) [1]. Despite the scarcity of this population (0.5–0.9% of the population are identified as transgender worldwide [2–4]), their contribution to HIV prevalence is considerable. The overall HIV prevalence (self-reported and laboratory confirmed) for TGW worldwide is 18.8–28% [1, 5–8]; this group has a 49-fold higher odds of infection compared with other reproductive age adults [6], and twice the odds compared with men who have sex with men (MSM) [5]. In China, TGW are not covered by routine HIV surveillance [9], but several small-sample studies suggest TGW's high HIV prevalence rate. The number might even be underestimated because of low rates of HIV testing among this particular population in China [9]. For example, in a cross-sectional study among 220 TGW sex workers conducted in 2014, 16.8% self-reported as HIV positive. Among those who self-reported as HIV negative or unknown HIV serostatus, and were willing to take up HIV testing ($n = 90$), 9.1% were detected to be HIV positive [10].

Among all risky sexual behaviors, unprotected anal intercourse (UAI; anal intercourse without condoms) has the highest risk for HIV acquisition and transmission among TGW [11, 12]. Pinkerton et al. found that inconsistent or non-condom users were 10–20 times more likely to acquire HIV than consistent condom users [13]. Promoting consistent condom use among key population has been verified to be a promising

risk reduction strategy. Smith et al. estimated that condom effectiveness of reducing HIV incidence was 72.3% (95% confidence intervals [CI]: 60.7–80.5%) for receptive anal sex, 62.9% (95% CI: 46.3–74.3%) for insertive anal sex, and 70.5% (95% CI: 58.2–79.2%) for any receptive or insertive anal sex [12]. A meta-analysis by the World Health Organization of five cohort studies [14] assessed condom effectiveness among MSM and transgender people in developed countries and found that the relative effect on HIV transmission of condom use was a relative risk of 0.36 (95% CI: 0.20–0.67) and that consistent condom use (CCU) reduced HIV transmission by 64% (95% CI: 32–80%).

However, none effective interventions has been carried out among TGW so far. UAI prevalence remains extremely high among TGW, even higher than other sex/gender minorities. A cross-sectional study conducted by Shan et al. in two Chinese province found that 81.5% TGW reported not always using condoms with stable partners, and 70.9% not with casual partners (Shan et al., 2018). According to Poteat et al., TGW had more than twice the odds as MSM of reporting condomless receptive anal sex with male partners (Poteat et al., 2017). The high prevalence of UAI highlights the urgent need for targeted and effective prevention interventions targeting this population.

Theory-based intervention is widely believed to be more likely to effectively change behavior and maintain such behavior change than non-theory-based intervention. Among extant theory and models, the Information-Motivation-Behavioral Skills (IMB) model has received consistent empirical support in the context of HIV prevention among various key populations, including heterosexual people living with HIV [15], sexually transmitted infection (STI) clinic patients [16], female sex workers [17], senior high school students [18, 19], college students [20], male street laborers [21], unmarried female migrants [22], and MSM [23].

The model posits that HIV-preventive behavior is a function of individuals' knowledge (information) of HIV transmission and prevention, their motivation to carry out HIV prevention (attitudes toward and subjective norms regarding HIV-preventive acts), and their skills in conducting specific preventive behaviors (possession of the requisite skills and a sense of self-efficacy) [24, 25]. The model further assumes that information and motivation may have direct (when complicated or novel behavioral skills are unnecessary to behavior) or indirect effects (through behavioral skills) on behavior [24, 25].

Fisher et al. pointed out that the IMB model constructs were highly generalizable determinants of HIV-preventive behavior in diverse populations of interest, but these constructs should have content that is specific to particular target populations as well as to particular HIV-preventive behaviors [24]. To the best of our knowledge, there have been no tests of the utility of the IMB model of condom use among TGW. It is therefore important to conduct a test before using it to guide the development of interventions targeting this population.

Based on the above-mentioned evidence, we intended to examine the application of the IMB model to clarifying the association between condom use correlates and condom use frequency among TGW in China. Theoretical evidence was expected to be provided to guide the development of interventions targeting increasing condom use frequency in this population. We hypothesized that information and

motivation would affect condom use directly and indirectly through behavioral skills and would be independent from each other.

Methods

Study site

Subjects were recruited in Shenyang (provincial capital of Liaoning Province), China between April 2017 and July 2017. According to the sixth national population census of the People's Republic of China [26], Shenyang is the largest city in Northeast China by urban population, with 8.1 million resident population. The number of MSM population in Shenyang ranks the second in China with an estimated 140,000 MSM in 2006 [27]. MSM population accounted for 81.2% of the city's annual newly reported HIV cases [28], while the national average proportion is only 28.3% [29].

Participants

The inclusion criterion were: 1) older than 18 years; 2) male birth assigned sex; 3) self-identified as woman; 4) reporting having sexual intercourse experiences in the last 6 months; 5) agreeing to participate in the study and willing to provide written informed consent.

Recruitment

Snowball sampling [30, 31] was used for recruitment because TGW in China are hard to reach and hidden due to the universal discrimination. The participants were recruited through a non-governmental organization (NGO), China Love Aid, which dedicated to improving the physiological and psychological health of TGW. With the assistance of the NGO, five participants were recruited as the 'seeds'. The five seeds then each recruited a maximum of five participants, thus, aggregating the snowball until saturation reached. All eligible participants were asked to read and sign informed consent documents for voluntary participation after being informed of the following: all data collected were anonymous and strictly confidential and would be used only for research; refusal would not affect their right to use any service; they could quit at any time without being questioned.

Data collection

A total of 198 eligible TGW provided written informed consent and participated in this study. They were then asked to complete the questionnaires during a face-to-face interview in a private room by trained investigators, who were staff of the NGO, to ensure the quality and completeness of the response. The questionnaires were anonymous and took approximately 30 minutes to complete. All subjects were paid 200 CNY (about 30 USD) as remuneration after completion of the above procedure. The ethics committee

of the School of Public Health, Shanghai Jiao Tong University, approved the recruitment procedure and the study protocol.

Measures

Background Characteristics

The following background characteristics were collected, including age, educational level, marital status, monthly income, and length of residence in Shenyang.

Information

To assess knowledge deficits and misconceptions about HIV risks and self-protective behaviors, the 18-item true–false Brief HIV Knowledge Questionnaire [32] (Cronbach's $\alpha = 0.683$) was administered. The HIV knowledge score was obtained by summing the number of correct responses. Higher scores indicated greater information.

Motivation

Motivation to engage in condom use was measured in accord with the principal constructs of the Theory of Reasoned Action [33] and the Health Belief Model [34]. Two scales were developed by the authors to provide two separate indicators of motivation. A total score was obtained by summing scores for individual items. Higher scores on the two scales indicated stronger motivation.

The 6-item, Likert type Attitude Scale (Cronbach's $\alpha = 0.528$) assessed subjects' attitudes toward condom use. For favorable statements ("I think condoms can prevent the transmission of HIV"; "I think condoms should be used during anal intercourse"; "I will try to persuade my partner into using condoms"), the strongly agree response is given a weight of 5 and the strongly disagree response a weight of 1, with other responses weighted accordingly. For unfavorable statements ("I think using condoms during anal intercourse is uncomfortable for both partners"; "I think condoms are unreliable in preventing HIV"; "In my opinion, condoms are too much trouble"), the scoring is reversed.

The 4-item Subjective Norms Scale (Cronbach's $\alpha = 0.806$) assessed perceptions of what others (family members, friends, peers who are engaged in the same work, and significant others) think about the participants' consistent condom use during anal sexual intercourse, where 1 = strongly disapprove, and 5 = strongly approve.

Behavioral Skills

To assess subjects' HIV-preventive behavioral skills, the 6-item Condom Influence Strategy Questionnaire (CISQ-S) [35] (Cronbach's $\alpha = 0.873$) and the 6-item Self-Efficacy Scale (Cronbach's $\alpha = 0.874$) developed by the authors were administered. Each scale represented a separate indicator of behavioral skills. Higher scores on both scales indicated better behavioral skills.

The CISQ-S assessed subjects' condom use negotiation skills and unprotected sex refusal skills. Questions were answered in a Likert format that ranged from 1 = never to 5 = always. An example item from the scale is "I would like to discuss condom use with my partner".

The Self-Efficacy Scale assessed their ability to engage in condom use effectively by these questions: "I can carry a condom with me in case I need it"; "I can use a new condom every time I have anal sex"; "I can use condoms throughout my anal sex"; "I can avoid the slipping of the condom during anal sex"; "I can discuss condom use with my partner even after drinking alcohol"; "I can negotiate using condoms with my partner no matter who he is". Participants answered on a 5-point Likert scale that ranged from 1 = definitely can't do to 5 definitely can do.

Condom Use

The questionnaire contained three frequency rating items (all scaled from 1 = never to 4 = always) to assess subjects' condom use with their regular sexual partners (defined as those who were in a stable relationship such as boyfriends that did not involve transactional sex), casual sexual partners (defined as those who were not regular partners where transactional sex was not involved), and transactional sexual partners (defined as those with whom transactional sex was involved) during the last 6 months. Similar definition has been widely used in other studies (e.g. [36]). To compare background characteristics between subjects with and without CCU, a dichotomous variable were created, with 1 indicating always using condom with all types of sexual partners, and 0 indicating not always using condom with at least one type of partners. Besides, a total condom use frequency was obtained by calculating the average frequency of use with the three types of sexual partners. Higher scores indicated more frequent condom use.

Statistical analysis

Descriptive analyses were conducted using IBM SPSS Statistics 23.0 (IBM Corp., Armonk, NY, USA). Comparisons were conducted between subjects showing CCU and those not showing CCU to detect potential risk factors using Student's t-test and Pearson's chi-square test. Spearman's correlation analysis was conducted to examine correlations among IMB model constructs. Confirmatory factor analysis (CFA) and structural equation modeling (SEM) were conducted using IBM SPSS AMOS 21.0 (IBM Corp., Chicago, IL, USA). The CFA measurement model included two latent constructs (motivation, behavioral skills) that predicted its indicators (motivation: condom use attitude, subjective norms; behavioral skills: condom use skills, self-efficacy). Once an acceptable measurement model had been established, a

structural equation model was built to examine correlates of condom use among TGW based on the IMB model. Model fit was assessed using the ratio of chi-square values to degrees of freedom (χ^2/df), the comparative fit index (CFI), and the root mean square error of approximation (RMSEA) [37]. A χ^2/df ratio of 3 or less, a CFI greater than 0.90, and an RMSEA lower than 0.08 indicated acceptable model fit [37, 38].

Results

Background Characteristics of Subjects

A total of 198 TGW completed the questionnaire (Table 1). They had an average age of 33.5 years (standard deviation = 9.6; range: 18–62 years). Most participants had a junior high school education or above (92.4%), were single (77.3%), and had a monthly income of 3001–6000 CNY (48.5%). Only 35.4% were local residents, with the rest 64.6% migrating to Shenyang. Nearly half the subjects (47.0%) reported CCU; the CCU rates with regular sexual partners, casual sexual partners, and transactional sexual partners were 26.3%, 28.8%, and 59.1%, respectively. CCU subjects were significantly younger (31.9 vs. 34.9 years; $P = 0.030$) and tended to be better educated than non-CCU subjects (62.3% vs. 40.0% with senior high school level or above; $P = 0.005$).

Table 1. Background characteristics of all subjects and comparisons between CCU and non-CCU subjects

Characteristics	All subjects (N=198) (n[%])	Subjects with CCU (n=93) (n[%])	Subjects without CCU (n=105) (n[%])	P value
Age (years) (mean $\pm$ SD) (range)	33.5 $\pm$ 9.6 (18-62)	31.9 $\pm$ 9.4 (18-59)	34.9 $\pm$ 9.6 (18-62)	0.030
Educational level				0.005
Primary school and below	15 (7.6)	8 (8.6)	7 (6.7)	
Junior high school	83 (41.9)	27 (29.0)	56 (53.3)	
Senior high school	50 (25.3)	27 (29.0)	23 (21.9)	
College and above	50 (25.3)	31 (33.3)	19 (18.1)	
Marital status				0.164
Single	153 (77.3)	77 (82.8)	76 (72.4)	
Married	11 (5.6)	5 (5.4)	6 (5.7)	
Divorced	34 (17.2)	11 (11.8)	23 (21.9)	
Monthly income (CNY)				0.525
$\leq$ 3000	46 (23.2)	23 (24.7)	23 (21.9)	
3001-6000	96 (48.5)	40 (43.0)	56 (53.3)	
6001-12000	41 (20.7)	22 (23.7)	19 (18.1)	
> 12000	15 (7.6)	8 (8.6)	7 (6.7)	
Length of residence in Shenyang				0.237
Local residents	70 (35.4)	31 (33.3)	39 (37.1)	
Living locally $\leq$ 1 year	19 (9.6)	13 (14.0)	6 (5.7)	
Living locally 1-5 years	50 (25.3)	24 (25.8)	26 (24.8)	
Living locally > 5 years	59 (29.8)	25 (26.9)	34 (32.4)	
CCU	93 (47.0)			
With regular sexual partner (n = 115)	52 (26.3)	-	-	-
With casual sexual partner (n = 117)	57 (28.8)	-	-	-
With transactional sexual partner (n = 170)	117 (59.1)	-	-	-

CCU: consistent condom use; SD: standard deviation.

P value < 0.05 was considered significant (in bold).

Bivariate Associations between Variables

The bivariate correlations among the IMB model constructs are shown in Table 2. HIV knowledge did not correlate with any model variables (attitude toward condom use: $r = 0.124$, $P = 0.083$; subjective norms: $r = 0.040$, $P = 0.571$; condom use skills: $r = 0.038$, $P = 0.594$; self-efficacy: $r = 0.027$, $P = 0.708$; condom use frequency: $r = -0.030$, $P = 0.670$). Condom use frequency was associated with all variables except HIV knowledge and subjective norms ($r = 0.030$, $P = 0.673$). Apart from the aforementioned correlations, all the variables were significantly associated with the other ones.

Table 2. Spearman's correlation analysis of IMB model constructs

IMB variables						
HIV knowledge	1.000					
Attitude toward condom use	0.124	1.000				
Subjective norms	0.040	0.291***	1.000			
Condom use skills	0.038	0.330***	0.288***	1.000		
Self-efficacy	0.027	0.365***	0.312***	0.683***	1.000	
Condom use frequency	-0.030	0.228**	0.030	0.491***	0.442***	1.000

*** $P < 0.001$. IMB: Information-Motivation-Behavioral Skills.

Confirmatory Factor Analysis

The model fit indices indicated an acceptable model fit: $\chi^2 = 0.046$, $df = 1$, χ^2/df ratio = $0.046 < 3$, CFI = $1.000 > 0.90$, and RMSEA = $0.000 < 0.08$. Given the overall fit of the data to the model, no additional paths or co-variances were added.

Table 3 shows the ranges, medians, interquartile ranges, and factor loadings for the IMB model constructs. The median percentage of correct responses to HIV knowledge was 61.1% (11.0/18). The median scores for attitude toward condom use, subjective norms, condom use skills, self-efficacy, and condom use frequency was 23.0/30 (76.7%), 16.0/20 (80.0%), 26.0/30 (86.7%), 28.0/30 (93.3%), 3.5/4 (87.5%), respectively. All factor loadings in the model were significant ($P < 0.001$), indicating that the hypothesized latent construct predicted its proposed manifest indicators well.

Table 3. Ranges, medians, interquartile ranges, and factor loadings for the IMB model constructs

Items	Range	Median (25 th -75 th quartile)	FL
Information			
HIV knowledge	0-18	11.0 (9.0-13.0)	-
Motivation			
Attitude toward condom use	6-30	23.0 (22.0-26.0)	0.567
Subjective norms	4-20	16.0 (14.0-18.0)	0.542
Behavioral skills			
Condom use skills	6-30	26.0 (21.0-29.3)	0.872
Self-efficacy	6-30	28.0 (24.8-30.0)	0.883
Behavior			
Condom use frequency	1-4	3.5 (3.0-4.0)	-

IMB: Information-Motivation-Behavioral Skills; FL: factor loading, all significant at $P < 0.001$.

The IMB Model

The model was depicted in Figure 1. The model fit indices indicated an acceptable model fit: $\chi^2 = 11.258$, $df = 5$, χ^2/df ratio = $2.252 < 3$, CFI = $0.980 > 0.90$, and RMSEA = 0.080 . Condom use frequency was strongly associated with behavioral skills ($\beta = 0.979$, $P = 0.004$). Motivation had no direct effect on condom use frequency but had a reliable effect on behavioral skills ($\beta = 0.823$, $P < 0.001$), which in turn significantly affected condom use frequency. Contrary to the IMB model, knowledge was not associated with either behavioral skills ($\beta = -0.075$, $P = 0.372$) or condom use frequency ($\beta = 0.052$, $P = 0.540$) and neither was it a covariate of motivation ($r = 0.143$, $P = 0.172$). In combination, these variables accounted for 38.7% of the variation in condom use frequency. Path coefficients were showed in detail in Table 4.

Table 4. Path coefficients of the IMB model

Path	Estimate	Std. Estimate	S.E.	P value
Information↔Motivation	0.727	0.143	0.532	0.172
Information→Behavioral skills	-0.115	-0.075	0.129	0.372
Information→Condom use frequency	0.013	0.052	0.021	0.540
Motivation→Behavioral skills	2.477	0.823	0.577	< 0.001
Motivation→Condom use frequency	-0.266	-0.546	0.186	0.152
Behavioral skills→Condom use frequency	0.158	0.979	0.054	0.004

Std. Estimate: standardized estimate; S.E.: standard error.

P value < 0.05 was considered significant (in bold).

Discussion

The results of this study demonstrated the utility of the IMB model for understanding condom use among TGW in China. The IMB model fit the data well and accounted for 38.7% of the variance in condom use frequency. The findings supported some, but not all, of the hypothesized associations between information, motivation, behavioral skills and condom use.

Results showed that the CCU rate (47.0%) was much higher than that reported in previous Chinese research (18.5% with stable partners, and 29.1% with casual partners) [39]. We attributed it to the higher proportions of subjects reporting transactional sexual intercourse in the current sample compared to the previous study; 170/198 (85.9%) have had transactional sexual intercourse in the previous 6 months. Cambou et al. discovered that unprotected anal intercourse was more prevalent in partnerships with a greater degree of interpersonal commitment like stable partners [40]. So condom use with transactional sexual partners was normally more frequent than stable and casual partners, which buffered the overall rate. Variable CCU rates with different types of sexual partners detected in the current study proved this speculation; CCU rates with regular sexual partners, casual sexual partners, and transactional sexual partners were 26.3%, 28.8%, and 59.1%, respectively. Anyway, the overall CCU rate in China was much lower than that reported in previous systematic research among US transgender population (61.8%) [1], indicating the severity of high risk sexual behavior among Chinese TGW and appealing to urgent and effective intervention.

Furtherly, a SEM was conducted to demonstrate the utility of the IMB model in clarifying the association among information, motivation, behavioral skills and condom use. Inconsistent with our hypothesis, motivation was not directly associated with condom use. However, in accord with some previous studies [19, 41], the effect of motivation on condom use was completely mediated by behavioral skills. More specifically, TGW who are highly motivated (who have a positive attitude and strong subjective norms about condom use) are likely to acquire the requisite skills (condom use skills and self-efficacy) for condom use. The results highlighted the crucial role of behavioral skills in promoting condom use.

While condom use is the most reliable method to reduce HIV risk, it is also a complex sexual behavior. Previous studies found that using a condom conveys potentially undesirable implicit messages such as lacking trust, lacking commitment or being HIV/STI-infected [42]. As such, complicated skills, such as condom use negotiation, were needed to convert one's safer-sex intentions into practical safer-sex behaviors (e.g. condom use). The important role of behavioral skills on condom use has been proved among other populations in previous studies. A meta-analysis of psychosocial correlates associated with heterosexual condom use found that partner communication about condom use was the most highly correlated factor in association with condom use [43]. The intervention effect of behavioral skills has also been verified among other populations. Gause et al. concluded through their meta-analysis that participants who were exposed to HIV prevention interventions targeting improved partner communication used condoms more frequently [42]. It's reasonable to speculate based on our findings and present studies that intervention targeting improving TGW's condom use behavioral skills can significantly elevate frequency of condom use among this population.

Consistent with the IMB model, correlation between motivation and information was insignificant, identifying motivation and information as independent constructs [11]. The path from motivation to condom use by way of behavioral skills can exist independently. It indicated that well-informed was an unnecessary prerequisite for an individual to be well-motivated.

Contrary to the IMB model, information showed no association with behavioral skills and condom use. Previous tests of the IMB model have found inconsistencies in the relationship between information and behavioral skills [19, 21, 22] and between information and condom use [16–18, 20, 23, 24]. Some researchers ascribed it to the speculation that information might impact initial behavioral change rather than maintain such behavior over time [22, 44]. Others speculated that one's information might have limited impact on dyadic behavioral change such as condom use, which involves communication and negotiation with his partner [22, 45]. In the current study, the lack of association between information and behavioral skills may be further ascribed to these two constructs not being measured at the same level of specificity [16]; information comprised knowledge about HIV transmission and self-protective behaviors, while behavioral skills consisted of condom use negotiation skills and unprotected sex refusal skills. Nevertheless, results of information provided some hints for future intervention. Subjects showed obvious deficits and misconceptions about HIV risks and self-protective behaviors (11.0/18) in the current study, indicating that the extant health propaganda and education on HIV prevention in China remains insufficient among TGW population.

The present study has some limitations that should be acknowledged. Firstly, cross-sectional study limits our ability to make causal inference. Longitudinal studies are needed to further verify the causal mechanism. Secondly, as the subjects were all from Shenyang, Liaoning Province, it may not be possible to generalize the findings to TGW from other locations. Thirdly, some IMB model studies among other populations have used it to compare different groups (such as patients reporting prior STI treatment versus those not reporting such treatment (Scott-Sheldon et al., 2010)) to define a more specific target subgroups. The CCU rate showed substantial variation across different types of sexual partners in this study, indicating possible disparity of the modeling results. Limited to sample size, we failed to make further explorations. Future research is recommended to make clear the mechanisms underlying how information, motivation, and behavioral skills affect condom use with different types of sexual partners among TGW.

Conclusions

The current study is the first to examine the determinants of condom use frequency in TGW using the IMB model. Findings indicated that interventions for this population should focus simultaneously on enhancing motivation and strengthening behavioral skills regarding condom use to substantially increase HIV-preventive behavior. Besides, although information was not significant in the model, participants' scores on information indicated that health propaganda and education on HIV prevention among TGW population in China are far from sufficient and need further improvements.

List Of Abbreviations

IMB	Information-Motivation-Behavioral Skills
TGW	transgender women
SEM	structural equation modeling
CCU	consistent condom use
UAI	unprotected anal intercourse
CI	confidence intervals
MSM	men who have sex with men
STI	sexually transmitted infection
NGO	non-governmental organization
CISQ-S	Condom Influence Strategy Questionnaire
CFA	confirmatory factor analysis
χ^2/df	chi-square values to degrees of freedom
CFI	comparative fit index
RMSEA	root mean square error of approximation
FL	factor loading
Std. Estimate	standardized estimate
S.E.	standard error

Declarations

Ethics approval and consent to participate: All procedures performed in studies involving human participants were in accordance with the ethical standards of the ethics committee of the School of Public Health, Shanghai Jiao Tong University, and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

Consent for publication: Not applicable.

Availability of data and materials: The datasets used during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: The National Natural Science Funds of China (71603166, 71673187). The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' contributions: YC, HW, RC, ZW, RS, YW and JL contributed to the design of the study. HW, RC, QS, LT, SZ, YS and TM made great efforts to recruit eligible participants and to collect data. HW analyzed and interpreted data and drafted the manuscript. YC, ZW, RS, YW and JL substantively revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements: The authors are grateful to the study participants for their contribution. The authors would also like to express deep gratitude to China Love Aid for providing assistance with recruitment and investigation. Thanks to Diane Williams, PhD, from Liwen Bianji, Edanz Editing China (www.liwenbianji.cn/ac), for English language editing of a draft of this manuscript.

References

1. Becasen JS, Denard CL, Mullins MM, Higa DH, Sipe TA: **Estimating the Prevalence of HIV and Sexual Behaviors Among the US Transgender Population: A Systematic Review and Meta-Analysis, 2006-2017.** *Am J Public Health* 2018:e1-e8.
2. Conron KJ, Scott G, Stowell GS, Landers SJ: **Transgender health in Massachusetts: results from a household probability sample of adults.** *Am J Public Health* 2012, **102**(1):118-122.
3. Kuyper L, Wijzen C: **Gender identities and gender dysphoria in the Netherlands.** *Arch Sex Behav* 2014, **43**(2):377-385.
4. Van Caenegem E, Wierckx K, Elaut E, Buysse A, Dewaele A, Van Nieuwerburgh F, De Cuypere G, T'Sjoen G: **Prevalence of Gender Nonconformity in Flanders, Belgium.** *Arch Sex Behav* 2015, **44**(5):1281-1287.
5. Poteat T, Ackerman B, Diouf D, Ceesay N, Mothopeng T, Odette KZ, Kouanda S, Ouedraogo HG, Simplicie A, Kouame A *et al*: **HIV prevalence and behavioral and psychosocial factors among transgender women and cisgender men who have sex with men in 8 African countries: A cross-sectional analysis.** 2017, **14**(11):e1002422.
6. Baral SD, Poteat T, Stromdahl S, Wirtz AL, Guadamuz TE, Beyrer C: **Worldwide burden of HIV in transgender women: a systematic review and meta-analysis.** *The Lancet Infectious diseases* 2013, **13**(3):214-222.
7. Herbst JH, Jacobs ED, Finlayson TJ, McKleroy VS, Neumann MS, Crepaz N: **Estimating HIV prevalence and risk behaviors of transgender persons in the United States: a systematic review.** *AIDS Behav* 2008, **12**(1):1-17.
8. Meerwijk EL, Sevelius JM: **Transgender Population Size in the United States: a Meta-Regression of Population-Based Probability Samples.** *Am J Public Health* 2017, **107**(2):e1-e8.
9. Tang S, Tang W, Meyers K, Chan P, Chen Z, Tucker JD: **HIV epidemiology and responses among men who have sex with men and transgender individuals in China: a scoping review.** *BMC Infectious Diseases* 2016, **16**(1):588.
10. Cai Y, Wang Z, Lau JT, Li J, Ma T, Liu Y: **Prevalence and associated factors of condomless receptive anal intercourse with male clients among transgender women sex workers in Shenyang, China.**

11. Fisher J, Fisher W: **The Information-Motivation-Behavioral skills model of AIDS risk behavior change: Empirical support and application.** *Understanding and preventing HIV risk behavior: Safer sex and drug use* 1996.
12. Smith DK, Herbst JH, Zhang X, Rose CE: **Condom effectiveness for HIV prevention by consistency of use among men who have sex with men in the United States.** *J Acquir Immune Defic Syndr* 2015, **68**(3):337-344.
13. Pinkerton SD, Abramson PR: **Effectiveness of condoms in preventing HIV transmission.** *Soc Sci Med* 1997, **44**(9):1303-1312.
14. Caceres CF: **Prevention and treatment of HIV and other sexually transmitted infections among men who have sex with men and transgender people: recommendations for a public health approach.** *World Health Organization* 2010:69-83.
15. Christiana NS, Sibylle NS, Daniel G, Tom P, Victoria G, Christophe R, Rickenbach M, ., Dias SF, Daniela R: **Condom use with steady partners among heterosexual people living with HIV in Europe: testing the information-motivation-behavioral skills model.** *Aids Patient Care Stds* 2010, **24**(12):771-780.
16. Scott-Sheldon LA, Carey MP, Venable PA, Senn TE, Coury-Doniger P, Urban MA: **Predicting condom use among STD clinic patients using the Information - Motivation-Behavioral Skills (IMB) model.** *Journal of Health Psychology* 2010, **15**(7):1093.
17. Zhang H, Liao M, Nie X, Pan R, Wang C, Ruan S, Zhang C, Tao X, Kang D, Jiang B: **Predictors of consistent condom use based on the Information-Motivation-Behavioral Skills (IMB) model among female sex workers in Jinan, China.** *BMC Public Health* 2011, **11**(1):113.
18. Cai Y, Ye X, Shi R, Xu G, Shen L, Ren J, Huang H: **Predictors of consistent condom use based on the Information-Motivation-Behavior Skill (IMB) model among senior high school students in three coastal cities in China.** *BMC Infectious Diseases* 2013, **13**(1):262.
19. Ybarra ML, Korchmaros J, Kiwanuka J, Bangsberg DR, Bull S: **Examining the Applicability of the IMB Model in Predicting Condom Use Among Sexually Active Secondary School Students in Mbarara, Uganda.** *AIDS and Behavior* 2013, **17**(3):1116-1128.
20. Liu Z, Wei P, Huang M, Liu YB, Li L, Gong X, Chen J, Li X: **Determinants of Consistent Condom Use among College Students in China: Application of the Information-Motivation-Behavior Skills (IMB) Model.** *Plos One* 2014, **9**(9):e108976.
21. Van HN, M PD, Debattista J: **Predictors of condom use behaviour among male street labourers in urban Vietnam using a modified Information-Motivation-Behavioral Skills (IMB) model.** *Culture Health & Sexuality* 2016, **18**(3):321.
22. Shen Q, Wang Y, Wang Z, Wang Y, Luo M, Wang S, Jiang X, Yang Y, Cai Y, Shang M: **Understanding condom use among unmarried female migrants: a psychological and personality factors modified information-motivation-behavioral skills model.** *BMC Public Health* 2019, **19**(1):223.
23. Jiang H, Chen X, Li J, Tan Z, Cheng W, Yang Y: **Predictors of condom use behavior among men who have sex with men in China using a modified information-motivation-behavioral skills (IMB) model.**

BMC Public Health 2019, **19**(1):261.

24. Fisher JD, Fisher WA, Williams SS, Malloy TE: **Empirical tests of an information-motivation-behavioral skills model of AIDS-preventive behavior with gay men and heterosexual university students.** *Health Psychology Official Journal of the Division of Health Psychology American Psychological Association* 1994, **13**(3):238-250.
25. Fisher JD FW: **Changing AIDS-risk behavior.** *Psychological Bulletin* 1992, **111**:19.
26. Statistics NBo: **Sixth National Population Census of the People's Republic of China-Liaoning Province.** 2010.
27. Liang D, Li J: **The preliminary study of civil non-profit organizations in the programe of MSM health education.** *Chin J AIDS STD* 2006, **12**:271-272.
28. Zhang J, Xu J-j, Song W, Pan S, Chu Z-x, Hu Q-h, Yu H, Mao X, Jiang Y-j, Geng W-q *et al*: **HIV Incidence and Care Linkage among MSM First-Time-Testers in Shenyang, China 2012–2014.** *AIDS and Behavior* 2018, **22**(3):711-721.
29. Ncaids N: **Update on the AIDS/STD epidemic in China and main response in control and prevention in December, 2015.** *Chin J AIDS STD* 2016, **22**(2):69.
30. Magnani R, Sabin K, Saidel T, Heckathorn D: **Review of sampling hard-to-reach and hidden populations for HIV surveillance.** *Aids* 2005, **19 Suppl 2**:S67-72.
31. Valerio MA, Rodriguez N, Winkler P, Lopez J, Dennison M, Liang Y, Turner BJ: **Comparing two sampling methods to engage hard-to-reach communities in research priority setting.** *Bmc Medical Research Methodology* 2016, **16**(1):146.
32. Michaelp C, Schroder KE: **Development and psychometric evaluation of the brief HIV Knowledge Questionnaire.** *Aids Education & Prevention* 2002, **14**(2):172-182.
33. Ajzenl., FishbeinM: **Understanding Attitudes and Predicting Social Behavior: An Illustration of Applied Social Research.** 1980.
34. Janz NK, Becker MH: **The Health Belief Model: A Decade Later.** *Health Education Quarterly* 1984, **11**(1):1-47.
35. Noar SM, Morokoff PJ, Harlow LL: **Condom Negotiation in Heterosexually Active Men and Women: Development and Validation of a Condom Influence Strategy Questionnaire.** *Psychology & Health* 2002, **17**(6):711-735.
36. Wang Z, Wu X, Lau J, Mo P, Mak W, Wang X, Yang X, Gross D, Jiang H: **Prevalence of and factors associated with unprotected anal intercourse with regular and nonregular male sexual partners among newly diagnosed HIV-positive men who have sex with men in China.** *HIV Medicine* 2017, **18**(9):635-646.
37. Kline RB: **Principles and practice of structural equation modeling.** *Journal of the American Statistical Association* 2011, **101**(12).
38. Mcdonald RP, Moon-Ho Ringo H: **Principles and practice in reporting structural equation analyses.** *Psychological Methods* 2002, **7**(1):64.

39. Shan D, Yu MH, Yang J, Zhuang MH, Ning Z, Liu H, Liu L, Han MJ, Zhang DP: **Correlates of HIV infection among transgender women in two Chinese cities.** *Infect Dis Poverty* 2018, **7**(1):123.
40. Cambou MC, Perez-Brumer AG, Segura ER, Salvatierra HJ, Lama JR, Sanchez J, Clark JL: **The Risk of Stable Partnerships: Associations between Partnership Characteristics and Unprotected Anal Intercourse among Men Who Have Sex with Men and Transgender Women Recently Diagnosed with HIV and/or STI in Lima, Peru.** *PLOS ONE* 2014, **9**(7):e102894.
41. Bryan AD, Fisher JD, Benziger TJ: **HIV prevention information, motivation, behavioral skills and behaviour among truck drivers in Chennai, India.** *Aids* 2000, **14**(6):756-758.
42. Gause NK, Brown JL, Welge J, Northern N: **Meta-analyses of HIV prevention interventions targeting improved partner communication: effects on partner communication and condom use frequency outcomes.** *Journal of Behavioral Medicine* 2018, **41**(4):423-440.
43. Sheeran P, Abraham C, Orbell S: **Psychosocial correlates of heterosexual condom use: A meta-analysis.** *Psychological Bulletin* 1999, **125**(1):90-132.
44. Jill J, Montgomery S, Emmons C-A, Kessler R, Ostrow D, Wortman C, O'Brien K, Eller M, Eshleman S: **Magnitude and Determinants of Behavioral Risk Reduction: Longitudinal Analysis of a Cohort at Risk for AIDS.** *Psychology & Health* 2007, **1**:73-95.
45. Harman JJ, Amico KR: **The Relationship-Oriented Information-Motivation-Behavioral Skills Model: A Multilevel Structural Equation Model among Dyads.** *AIDS and Behavior* 2009, **13**(2):173-184.

Figures

Figure 1

Figure 1. The Information-Motivation-Behavioral Skills model of condom use frequency among transgender women in China (N = 198) Path coefficients are standardized. Circles represent latent variables and rectangles represent manifest variables. Single arrows represent standardized regression coefficients and multiheaded arrows represent standardized correlations. Solid lines represent statistical significance and dashed lines represent a lack of statistical significance. **P < 0.01, ***P < 0.001.