

The analgesic efficacy of ultrasound-guided transversus abdominis plane block for retroperitoneoscopic renal surgery: A randomized controlled study

Xue Li

peking university first hospital

Zhen-Zhen Xu

Peking University First Hospital

Xue-Ying Li

Peking University First Hospital

Ting-Ting Jiang

Peking University First Hospital

Zeng-Mao Lin (✉ linzengmao@163.com)

<https://orcid.org/0000-0001-6450-8063>

Dong-Xin Wang

Peking University First Hospital

Research article

Keywords: Transversus abdominis plane block; Analgesia; Retroperitoneoscopic renal surgery; Postsurgical recovery.

Posted Date: September 26th, 2019

DOI: <https://doi.org/10.21203/rs.2.9718/v4>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 18th, 2019. See the published version at <https://doi.org/10.1186/s12871-019-0850-3>.

Abstract

Background Ultrasound-guided lateral transversus abdominis plane (TAP) block can provide definite analgesia to the anterior abdominal wall. However, whether this method is useful in renal surgery through the lateral abdominal wall pathway remains unknown. The study aimed to evaluate the analgesic efficacy of lateral TAP block for retroperitoneoscopic partial or radical nephrectomy. **Method** In this prospective, randomized, double-blind, placebo-controlled trial, eligible patients were randomized into two groups. After anaesthesia induction, ultrasound-guided lateral TAP block was performed with either 30 ml of 0.4% ropivacaine (Group T) or an equivalent volume of normal saline (Group C). The primary outcomes were opioid consumption during surgery and in the first 24 hours after surgery. Secondary outcomes included postsurgical pain intensity immediately awakening from anaesthesia and at 0.5, 1, 2, 6, 12, and 24 hours after surgery, as well as recovery variables including the incidence of postoperative nausea and vomiting (PONV), sleep quality, time to first ambulation, drainage and length of hospital stay. **Results** A total of 104 patients were enrolled and randomized (53 in Group T and 51 in Group C). Laparoscopic surgery was converted to open surgery in one patient of Group T; this patient was excluded from the outcome analysis. The opioid consumption during surgery (intravenous morphine equivalent dose: median 35.0 mg [interquartile range 18.0, 49.6] in Group C vs. 40.3 mg [20.9, 59.0] in Group T, $P=0.281$) and in the first 24 hours after surgery (10.8 mg [7.8, 21.7] in Group C vs. 13.2 mg [8.0, 26.6] in Group T, $P=0.311$) did not differ significantly between groups. There were no significant differences between groups regarding the pain intensity at all time points after surgery and the recovery variables (all $P>0.05$). **Conclusions** Our results showed that, in patients undergoing retroperitoneoscopic renal surgery, preoperative lateral TAP did not decrease intra- and postoperative opioid consumption, nor did it relieve pain intensity or promote postoperative recovery in the first 24 hours after surgery. However, the trial might be underpowered. **Trial registration** This study was registered on November 4, 2017, in the Chinese Clinical Trial Registry with the identification number ChiCTR-17013244.

Background

For laparoscopic renal surgery, the retroperitoneal approach is an alternative pathway of the transperitoneal approach. The overall outcomes of both approaches, such as the rates of perioperative complications, positive surgical margin and postoperative recurrence, are similar; whereas the retroperitoneoscopic approach is advantageous in terms of easier hilar control and shorter total operative time, especially in patients with a past history of intraperitoneal procedures or with a posteriorly located renal tumour [1-3]. Therefore, the retroperitoneoscopic approach is the most popular approach for radical or partial nephrectomy in Peking University First Hospital. Multimodal analgesia including nerve block is advocated to improve early recovery after renal surgery [4,5].

Transversus abdominis plane (TAP) block is an effective regional anaesthetic technique that blocks neural afferents of the T6-L1 spinal nerve innervating the anterolateral abdominal wall [6]. Since the original report by Rafi [7], there have been a plethora of studies on this block and variations of the original approach, among which lateral TAP is the most commonly used approach in abdominal surgery, with its

dermatomal sensory block covering T10 to L1 [8]. A recent systematic review demonstrated that TAP block had definite analgesic efficiency for some kinds of lower abdominal surgeries, such as gynaecological surgery, caesarean section and hernia repair, but not urologic surgery [9]. However, it was worth noting that there was a high heterogeneity among the urologic studies included in that review. In addition, some studies focusing on renal surgery were missed.

To our knowledge, only three randomized controlled trials compared the effect of lateral TAP block with placebo in laparoscopic live-donor nephrectomy [10-12]. All of these found that TAP block surely reduced postoperative pain severity and opioid requirements. However, this conclusion could not be extrapolated to retroperitoneal laparoscopic renal surgery (RLRS) of which the main incision was completely different from those of live-donor nephrectomy. Theoretically, the dermatomes of lateral TAP block can only partially cover the incision in RLRS. Therefore, whether it could reduce opioid consumption and subjective pain intensity, and ultimately promote postoperative recovery in patients undergoing RLRS remains unknown. The purpose of this study was to determine whether lateral TAP block could provide effective analgesia and improve recovery in patients undergoing RLRS.

Methods

Study design

This prospective, randomized, double-blinded trial was approved by the Biomedical Research Ethics Committee of Peking University First Hospital (2017-1398). It was registered at <http://www.chictr.org.cn> with an identification number of ChiCTR-INR-17013244. Written informed consent was obtained from each patient.

Participants

Potential participants were screened the day before surgery. Patients aged between 18 and 70 years and scheduled to undergo elective laparoscopic radical or partial nephrectomy through the retroperitoneal approach were included. Patients who met any of the following criteria were excluded: (1) chronic opioid addiction and/or use of other kinds of analgesic drugs for more than 3 months; (2) inability to communicate due to severe dementia, language barrier, or end-stage disease; (3) allergic to local anaesthetics; (4) nerve block contraindication such as an infection in the puncture site or severe coagulation dysfunction; and (5) refusal to participate in the study. Patients who were eligible for this trial were taught how to evaluate pain intensity by using the numeric rating scale (NRS, an 11-point scale where 0 indicates no pain and 10 indicates the worst pain) and how to use a patient-controlled analgesia (PCA) device.

Anaesthesia management and surgical technique

All patients were managed according to a standardized anaesthetic protocol. Anaesthesia was induced with sufentanil, propofol and etomidate. Endotracheal intubation was facilitated with cisatracurium or

rocuronium. Anaesthesia was maintained with continuous infusion of propofol, remifentanyl (and intermittent sufentanyl) or sufentanyl, with or without dexmedetomidine; the aim was to maintain the BIS values between 40 and 60, and the mean arterial pressure and heart rate within 20% of the preoperative values. At 30 minutes before the end of surgery, 50 mg of flurbiprofen axetil and 5 mg of tropisetron were administered intravenously. After emergence from anaesthesia, all patients were monitored in the post-anaesthesia care unit (PACU) for at least 1 hour before transferred to the general ward.

The retroperitoneal laparoscopic procedure was usually performed through three ports. The primary port was invariably placed through the incision made for the creation of the working space, which was just below the tip of the 12th rib on the posterior axillary line. The secondary port was placed 2 cm above the iliac crest on the midaxillary line. The third port was placed under the costal margin on the anterior axillary line. In case of radical nephrectomy, the initial incision was extended ventrally for kidney removal. The pneumoperitoneum was maintained at approximately 12-14 mmHg throughout the procedure (Figure 1).

For all patients, a standard postoperative pain management was provided, i.e., a PCA pump, which was established with 1.25 µg/ml sufentanyl and programmed to administer a background rate of 0.5 ml/h and an on-demand bolus of 4 ml every 10 minutes, together with a rigorous rescue analgesia plan. The target was to maintain the NRS pain score below 4. In the PACU, regular pain evaluation was performed every 30 minutes. If the NRS score was higher than 4, a PCA bolus of 4 ml was administered first, and pain was evaluated 5 min later. If the NRS score remained higher than 4, another 3-5 µg sufentanyl was administered intravenously according to the patient's body weight. No more rescue analgesics were administered if the NRS score decreased to 4 or less. In the general ward, pain evaluation was performed at 2, 6, 12, and 24 hours after surgery; in addition, patients were instructed to request additional analgesia in case of breakthrough pain. Pain control measures were similar to those in the PACU, except that morphine (3-5 mg) was administered instead of sufentanyl. NSAIDs or other analgesics could also be administered according to the surgeons' prescription.

Randomization and intervention

Stratified randomization with a block size of 4 was performed using the SAS statistical package version 9.3 (SAS Institute, Cary, NC, USA) by a biostatistician (Xiao-Lu Nie) who was not involved in the data management and statistical analyses. Stratification was performed according to the planned type of surgery, i.e., radical or partial nephrectomy. The randomization results were then sealed in sequentially numbered envelopes, transferred to a study coordinator (Ting-Ting Jiang) with the Good Clinical Practice (GCP) certification and stored at the site of the investigation until the end of the study.

The day before surgery, an investigator (Zeng-Mao Lin) screened potential participants and recruited patients after obtaining written informed consents. On the day of surgery, the study coordinator opened the envelopes consecutively according to the recruitment sequence and prepared the study drugs for each patient, but did not participate in the rest of the trial. All study drugs were provided as clear aqueous solutions in the same 20 ml syringes for TAP block. In this way patients were randomly assigned into two

groups: patients in Group T received 30 ml of 0.4% ropivacaine, while those in Group C received an equivalent amount of normal saline. Apart from the study drugs used for TAP block, other perioperative management was identical in both groups. All health-care team members, investigators, and patients themselves were fully blinded to the group assignments throughout the study period.

Ultrasound-guided TAP block was performed by two experienced anaesthetists (Da Huang and Hao Kong) immediately after the induction of anaesthesia and approximately 15 minutes before skin incision. With the patient in the supine position, the ultrasound probe was placed at the midaxillary line between the lower costal margin and the iliac crest. At this point, the plane between the internal oblique and transverse abdominal muscles was identified (Figure 2A). A special needle used for nerve block (80 mm or 100 mm, Stimuplex D, Germany) was inserted using an in-plane technique in the anteroposterior direction. After aspiration, to avoid inadvertent intravascular injection and abdominal paracentesis, an injection with 2 ml of normal saline was used to ensure correct positioning of the needle. The prepared study drug was then injected into this plane. Successful study drug injection was defined as the appearance of a hypoechoic ellipsoid with well-defined margins on ultrasonic imaging (Figure 2B).

Follow-up schedule and outcomes

Investigators (Xue Li, Zhen-Zhen Xu, Zeng-Mao Lin) who were blinded to the study group assignment were in charge of the perioperative data collection. Patients were followed-up at several time points in the first 24 hours after surgery. In addition, the electronic medical record was reviewed to obtain necessary data.

The primary outcomes were opioid consumption (intravenous morphine equivalent dose) during surgery and within the first 24 hours after surgery. The secondary outcomes included the following: (1) the NRS pain scores both at rest and with coughing immediately awakening from anaesthesia and at 0.5, 1, 2, 6, 12 and 24 hours after surgery; (2) time to the first bolus demand in the PCA system, as well as the numbers of required and administered bolus; (3) time to the first rescue analgesic, as well as its use and frequency in addition to the PCA system; (4) the incidence of postoperative nausea and vomiting (PONV) within 24 hours after surgery and the use of antiemetics; (5) subjective sleep quality on the night of surgery evaluated by the NRS (an 11-point scale where 0 indicates the best sleep quality and 10 indicates the worst sleep experience); (6) time to the first ambulation after surgery; (7) drainage during the first 24 hours after surgery; and (8) the length of hospital stay after surgery.

Safety outcomes were monitored from the beginning of anaesthesia until 24 hours after surgery. The adverse events associated with TAP block included but not limited to the following: numbness in the lower extremities, haematoma and bleeding in the needle trajectory, visceral organ injury, anaphylaxis, local anaesthetic toxicity. Other perioperative adverse events were also documented.

Statistical analysis

Sample size estimation

According to previous studies [10,12, 13], the use of TAP block decreased opioid consumption by 13.5%-45.3% compared with the placebo during the first 24 hours after surgery. We conservatively assumed that opioid consumption would be reduced by 10% in the TAP block group. Sample size calculation was performed based on the previous data obtained from our clinical follow-up system, which showed that the total consumption of sufentanil (within 24 hours after surgery) in patients who underwent RLRS without TAP block was $36.5 \pm 5.4 \mu\text{g}$. With the significance and power set at 0.05 (two-sided) and 90%, respectively, the sample size required to detect differences was 94 patients. Taking into account a drop-out rate of approximately 10%, we planned to enrol 104 patients. Sample size calculation was performed with the PASS 11.0 software (Stata Corp. LP, College Station, TX).

Outcome analyses

Normally distributed continuous variables are expressed as the mean \pm SD values and were compared using a two-tailed Student's t-test. Non-normally distributed continuous variables and ordinal data are expressed as medians (interquartile range) and were compared using the Mann-Whitney U test. Categorical variables are expressed as numbers (percentages) and were compared with Chi-squared analysis or Fischer's exact test. Time-to-event data were analysed by the Kaplan-Meier estimator, with the difference between groups compared by the log-rank test. Per-protocol analysis was performed. Two-sided P values of less than 0.05 were regarded as statistically significant. All statistical analyses were performed with the SPSS statistical package version 25.0 (IBM SPSS Inc., Chicago, IL, USA).

Results

From January 1, 2018, to March 20, 2018, 166 patients were screened for eligibility; of these, 130 met the inclusion/exclusion criteria, 104 gave consents and were randomized into the study, with 53 patients in Group T and 51 in Group C. During surgery, one patient developed major haemorrhage and was converted to open surgery. This patient was excluded from the per-protocol analysis (Figure 3).

No patient received opioid treatment before surgery. The two groups were comparable with regard to the demographic or baseline characteristics except that the percentage with diabetes mellitus was higher in Group T ($P=0.041$; Table 1). Intraoperative variables, including durations of anaesthesia and surgery, use and dose of dexmedetomidine, and estimated blood loss, did not differ significantly between groups (Table 2).

The opioid consumption during surgery (intravenous morphine equivalent dose: median 35.0 mg [interquartile range 18.0, 49.6] in Group C vs. 40.3 mg [20.9, 59.0] in Group T, $P=0.281$) and in the first 24 hours after surgery (10.8mg [7.8, 21.7] in Group C vs. 13.2 mg [8.0, 26.6] in Group T, $P=0.311$) did not differ significantly between groups. Stratified analysis did not find any significant differences regarding intraoperative and postoperative morphine equivalent dose between the two groups in patients receiving either the partial or radical nephrectomy (Table 3).

There were no significant differences between groups regarding the numbers of required and administered bolus, as well as the time to the first required bolus from the PCA system ($P=0.335$, 0.338 and 0.105 , respectively). There were no significant differences between groups regarding the percentage and frequency of rescue analgesics, as well as the time to first dose rescue analgesics in addition to the PCA system ($P=0.153$, 0.306 , and 0.845 , respectively) (Table 3). Post-surgical pain scores at those abovementioned time points were similar between the two groups both at rest and with coughing (all $P>0.05$) (Figure 4A and 4B).

As for postoperative recovery in the first 24 hours after surgery, the incidence of PONV, percentage of antiemetic therapy, subject sleep quality, time to the first ambulation, volume of drainage, and length of stay in hospital after surgery were not significantly different between the two groups (all $P>0.05$) (Table 4).

No adverse events related to the TAP block technique were observed in either group. One patient in Group T developed major haemorrhage during surgery (estimated blood loss of 6500 ml), one patient in Group C developed emergence delirium. Perioperative management was uneventful in other patients.

Discussion

To our knowledge, this was the first study investigating the efficiency of lateral TAP block in patients undergoing RLRS. Our results showed that preoperative lateral TAP block did not decrease intra- and postoperative opioid consumption, nor did it relieve pain intensity or promote postoperative recovery early after surgery. Our study added new evidence to the current knowledge of analgesic measures for laparoscopic urological surgery.

Our results conflicted with those of previous studies. For example, both Parikh and Guner found that lateral TAP block performed at the end of laparoscopic donor nephrectomy significantly decreased pain score and total opioid consumption in the first 24 hours [11,12]. Similarly, Hosgood also claimed that TAP block reduced the early morphine requirement (within 6 hours after surgery) in a similar patient population [10]. It should be noted that there were two important differences between our study and the above-mentioned others. First, the location of the main surgical incision for kidney retrieval and trocar sites in the present study was completely different from those in the previous studies [10,11]. Second, we performed TAP block before surgery, whereas others performed TAP block after surgery [11,12]. Thus, when comparing results among different trials, it is crucial to take the surgical technique, the block approach and the time of block into consideration. Full clarification of our negative findings requires detailed understanding of the innervation of the abdominal wall.

The anterolateral abdominal wall is mainly innervated by the anterior rami of thoracolumbar spinal nerves (T6-L1), which follow a curvilinear course from the back towards the midline of the body [6]. Generally, as they proceed, after giving off lateral cutaneous branches near the costal angle innervating the lateral areas of the abdominal wall [14], they enter into the TAP with a varied course and finally perforate the rectus abdominis and end as the anterior cutaneous branches innervating the anterior

abdomen (area from the midline to the midclavicular line) [15]. Most of the lateral cutaneous branches arise before the main nerves enter the TAP, and only those of T11 and T12 have a short course within or through the TAP [15]. Thus, it is not surprising that lateral TAP block can reliably provide analgesia for the lower anterior abdomen but not the lateral abdominal wall. Ma et al. further confirmed this by detecting the blocking dermatomes in 19 areas of the abdominal wall after lateral TAP block [16]. This might be an important reason for our negative results. The dermatomes of lateral TAP block we performed only covered part of the wounds in RLRS. For urological surgery, the posterior TAP approach [13,17] or quadratus lumborum block [18] may be the better choice because they can block the lateral cutaneous branches of thoracolumbar spinal nerves and provide better lateral abdominal wall analgesia.

Regarding postoperative recovery, we observed no advantages in Group T in terms of the incidence of PONV. This was understandable since the perioperative opioid consumptions were comparable in the two groups. Similar findings were also reported in other studies [10, 19]. We found rather low sleep quality in both groups with no significant difference. We believed that not only pain intensity but also the surrounding environment in the ward affected sleep quality. Given that none of the above variables were different, it was no wonder that the length of hospital stay was similar in the two groups.

Our study has some limitations. First, we did not assess sensory dermatome blockage to confirm a successful TAP block because the block was performed after anaesthesia induction for blinding. Second, we only collected opioid consumption at a single time point (24 hours after surgery) during the follow-up periods. Since we didn't record the opioid consumption at earlier postsurgical time-points, such as at 6 and 12 hours after surgery, we could not analyse the early effect of TAP block. Third, the range of 95% confidence interval of the median difference in opioid consumption between the two groups were large, which indicated that the present trial might be underpowered.

Conclusion

Results of this prospective, randomized, double-blind trial showed that preoperative single-shot lateral TAP block did not decrease intra- and postoperative opioid consumption, nor did it relieve pain intensity or promote postoperative recovery in the first 24 h after surgery for patients undergoing retroperitoneoscopic renal surgery. However, considering the wide range of confidence interval of median difference in opioid consumption between the two groups, the trial might be underpowered.

List Of Abbreviations

ASA: America Society of Anaesthesiologists; BMI: body mass index; BIS: Bi-spectral index; COPD: chronic obstructive pulmonary disease; NYHA: New York Heart Association; NSAIDs: non-steroidal anti-inflammatory drugs; NRS: numeric rating scale; PONV: postoperative nausea and vomiting; PCA: patient-controlled analgesia; PACU: post-anaesthesia care unit; RLRS: retroperitoneal laparoscopic renal surgery; TAP: transversus abdominis plane.

Declarations

Ethics approval and consent to participate

The research protocol was approved by the Biomedical Research Ethics Committee of Peking University First Hospital (Number: 2017-1398), and all the patients signed written informed consent voluntarily.

Consent for publication

Not applicable.

Availability of data and materials

The dataset supporting the conclusions of this article is included within the article and its additional files.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding.

Authors' contributions

Xue Li: Study design, data collection and analysis, manuscript drafting.

Zhen-Zhen Xu: Data collection.

Xue-Ying Li: Sample size calculation.

Ting-Ting Jiang: Randomization implementation and drug preparation.

Zeng-Mao Lin: Study design, data collection, manuscript revision.

Dong-Xin Wang: Critically revised the manuscript.

Acknowledgements

We thank Xiao-Lu Nie a great deal for performing the randomization sequence, making the covering envelopes and statistical consultation. Furthermore, we appreciate the efforts of Da Huang and Hao Kong for performing the TAP block. We also acknowledge the Department of Anesthesiology team and the Department of Urological Surgery team in Peking University First Hospital for supporting the research.

References

1. Liatsikos E, Kallidonis P, Do M, Dietel A, Al-Aown A, Constantinidis C, Stolzenburg JU. Laparoscopic radical and partial nephrectomy: technical issues and outcome. *Word J Urol* 2013, 31(4):785-791.
2. Ren T, Liu Y, Zhao X, Ni S, Zhang C, Guo C, Ren M: Transperitoneal approach versus retroperitoneal approach: a meta-analysis of laparoscopic partial nephrectomy for renal cell carcinoma. *Plos One* 2014, 9(3):e91978.
3. Fan X, Xu K, Lin T, Liu H, Yin Z, Dong W, Huang H, Huang J: Comparison of transperitoneal and retroperitoneal laparoscopic nephrectomy for renal cell carcinoma: a systematic review and meta-analysis. *Bju Int* 2013, 111(4):611-621.
4. Tan M, Law LS, Gan TJ. Optimizing pain management to facilitate enhanced recovery after surgery pathways. *Can J Anaesth* 2015, 62(2):203-218.
5. Nasrallah G, Souki FG: Peri-anesthetic management of laparoscopic kidney Surgery. *Curr Urol Rep* 2018, 19(1):1.
6. Tsai HC, Yoshida T, Chuang TY, Yang SF, Chang CC, Yao HY, Tai YT, Lin JA, Chen KY. Transversus abdominis plane block: an updated review of anatomy and techniques. *Biomed Res Int* 2017, 2017:8284363.
7. Rafi AN. Abdominal field block via the lumbar triangle revisited. *Anaesthesia* 2012, 67(12):1399-1401.
8. Tran TM, Ivanusic JJ, Hebbard P, Barrington MJ. Determination of spread of injectate after ultrasound-guided transversus abdominis plane block: a cadaveric study. *Br J Anaesth* 2009, 102(1):123-127.
9. Brogi E, Kazan R, Cyr S, Giunta F, Hemmerling TM: Transversus abdominal plane block for postoperative analgesia: a systematic review and meta-analysis of randomized-controlled trials. *Can J Anaesth* 2016, 63(10):1184-1196.
10. Hosgood SA, Thiyagarajan UM, Nicholson HF, Jeyapalan I, Nicholson ML. Randomized clinical trial of transversus abdominis plane block versus placebo control in live-donor nephrectomy. *Transplantation* 2012, 94(5):520-525.
11. Parikh BK, Waghmare VT, Shah VR, Mehta T, Butala BP, Parikh GP, Vora KS. The analgesic efficacy of ultrasound-guided transversus abdominis plane block for retroperitoneoscopic donor nephrectomy: A randomized controlled study. *Saudi J Anaesth* 2013, 7(1):43-47.
12. Guner CM, Goz R, Berber I, Kaspar C, Cakir U. Ultrasound/ laparoscopic camera-guided transversus abdominis plane block for renal transplant donors: a randomized controlled trial. *Ann Transplant* 2015, 20:418-423.
13. Aniskevich S, Taner CB, Perry DK, Robards CB, Porter SB, Thomas CS, Logvinov II, Clendenen SR. Ultrasound-guided transversus abdominis plane blocks for patients undergoing laparoscopic hand-assisted nephrectomy: a randomized, placebo-controlled trial. *Local Reg Anesth* 2014, 7:11-16.
14. Sakamoto H, Akita K, Sato T. An anatomical analysis of the relationships between the intercostal nerves and the thoracic and abdominal muscles in man. I. Ramification of the intercostal nerves. *Acta Anat (Basel)* 1996, 156(2):132-142.

15. Hebbard PD, Barrington MJ, Vasey C. Ultrasound-guided continuous oblique subcostal transversus abdominis plane blockade: description of anatomy and clinical technique. *Reg Anesth Pain Med* 2010, 35(5):436-441.
16. Ma J, Pan Y, Yang J, Xie Z, Li J: Efficacy of mid-axillary approach to transversus abdominis plane block in each abdominal and back region. *Chin J Anesthesiol* 2016, 10(01):1247-1249 (in chinese).
17. Qu G, Cui XL, Liu HJ, Ji ZG, Huang YG. Ultrasound-guided Transversus Abdominis Plane Block Improves Postoperative Analgesia and Early Recovery in Patients Undergoing Retroperitoneoscopic Urologic Surgeries: A Randomized Controlled Double-blinded Trial. *Chin Med Sci J* 2016, 31(3):137-141.
18. Elsharkawy H, Ahuja S, DeGrande S, Maheshwari K, Chan V: Subcostal approach to anterior quadratus lumborum block for pain control following open urological procedures. *Journal of Anesthesia* 2019, 33(1):148-154.
19. Ma N, Duncan JK, Scarfe AJ, Schuhmann S, Cameron AL: Clinical safety and effectiveness of transversus abdominis plane (TAP) block in post-operative analgesia: a systematic review and meta-analysis. *J Anesth* 2017, 31(3):432-452

Tables

Table 1. Demographic and baseline characteristics

	Group C (n=51)	Group T (n=52)	P value
Age, year	51.1±11.1	51.9±10.3	0.717
Body mass index, kg/ m ²	24.8±3.3	24.8±3.8	0.929
Male	31 (60.8%)	32 (61.5%)	0.937
Type of surgery			0.927
Radical nephrectomy	24 (47.1%)	24 (46.2%)	
Partial nephrectomy	27 (52.9%)	28 (53.8%)	
ASA class			
I	27 (52.9%)	24 (46.2%)	0.421
II	24 (47.1%)	27 (51.9%)	
III	0 (0.0%)	1 (1.9%)	
NYHA class			0.495
I	51 (100%)	50 (96.2%)	
II	0 (0.0%)	2 (3.8%)	
Comorbidities			
Stoke	2 (3.9%)	4 (7.7%)	0.692
Hypertension	19 (37.3%)	16 (30.8%)	0.487
Coronary artery disease	1 (2.0%)	3 (5.8%)	0.624
Diabetes Mellitus	3 (5.9%)	10 (19.2%)	0.041
Asthma and/or COPD	2 (3.9%)	0 (0.0%)	0.243
Previous abdominal and back surgery	14 (27.5%)	10 (19.2%)	0.324

Data are presented as mean ± standard deviation or number (%).

ASA: America Society of Anaesthesiologists; NYHA: New York Heart Association; COPD: chronic obstructive pulmonary disease.

Table 2. Intraoperative data

	Group C (n=51)	Group T (n=52)	P value
Duration of anaesthesia, min	144 (122, 168)	140 (126, 165)	0.805
Duration of surgery, min	78 (59, 112)	79 (64, 102)	0.934
Use of dexmedetomidine	20 (39.2%)	13 (25.0%)	0.122
Dose of dexmedetomidine, µg	30 (23, 39) (n=20)	30 (22, 50) (n=13)	0.785
Estimated blood loss, ml	50 (50, 50)	50 (50, 50)	0.387

Data are presented as median (interquartile range) or number (%).

Table 3. Effectiveness outcomes

	Group C (n=51)	Group T (n=52)	Estimated effects (95% CI) ^a	P value
Opioid consumption during surgery				
Sufentanil, mg	20 (15, 38)	23 (20, 30)	Median D=0 (-5, 5)	0.685
Remifentanil, µg	600 (502, 794) (n=33)	607 (428, 818) (n=39)	Median D=0 (-119, 120)	0.977
Morphine equivalent dose, mg	35.0 (18.0, 49.6)	40.3 (20.9, 59.0)	Median D=4.4 (-3.6, 13.4)	0.281
Morphine equivalent dose, mg/kg	0.58 (0.29, 0.77)	0.59 (0.30, 0.87)	Median D=0.05 (-0.06, 0.18)	0.326
Opioid consumption within 24 h after surgery				
Sufentanil, mg	33 (23, 65)	40 (24, 80)	Median D=4 (-4, 14)	0.311
Morphine equivalent dose, mg	10.8 (7.8, 21.7)	13.2 (8.0, 26.6)	Median D=1.2 (-1.3, 4.8)	0.311
Morphine equivalent dose, mg/kg	0.16 (0.12, 0.31) (n=27)	0.19 (0.12, 0.39) (n=28)	Median D=0.02 (-0.02, 0.07)	0.252
Laparoscopic partial nephrectomy				
Intraoperative MED, mg	41.2 (14.0, 49.6)	38.0 (19.0, 60.4)	Median D=4.4 (-10.6, 16.6)	0.480
Postoperative MED within 24 h, mg	10.1 (7.4, 19.8)	17.7 (8.4, 26.1)	Median D=3.1 (-0.3, 10)	0.070
Laparoscopic radical nephrectomy				
Intraoperative MED, mg	30.0 (19.2, 52.7)	40.7 (26.9, 53.8)	Median D=5.5 (-6.8, 18.7)	0.370
Postoperative MED within 24 h, mg	14.5 (8.2, 25.0)	10.6 (8.0, 27.8)	Median D=-0.9 (-6.0, 4.4)	0.773
Data from the PCA system				
Number of required bolus	4 (1, 10)	6 (1, 15)	Median D=1 (-1, 4)	0.335
Number of administered bolus	3 (1, 10)	5 (1, 12)	Median D=1 (-1, 3)	0.338
Time to first required bolus, hour ^b	1.7 (0.4, 3.0)	6.0 (2.8, 9.2)	HR=1.46 (0.93, 2.30)	0.105
Data of rescue analgesia within 24 h after surgery				
Percentage of rescue analgesics	12 (23.5%)	19 (36.5%)	OR=0.53 (0.23, 1.26)	0.153
Frequency of rescue analgesics	2 (1, 3)	1 (1, 4)	Median D=0 (-1, 0)	0.306
Time to first rescue analgesics, hour ^b	21.0 (14.4, 27.6)	25.0 (19.3, 30.7)	HR=1.07 (0.52, 2.24)	0.845

Data are presented as median (interquartile range) or number of patient (%), unless otherwise indicated.

MED: morphine equivalent dose; PCA: patient controlled analgesia.

^a Calculated as Group T vs. or minus Group C.

^b Data were analysed by Kaplan-Meier analysis and compared by log-rank test; the results are presented as the median (95% confidence interval).

Table 4. Comparisons of recovery variables

	Group C (n=51)	Group T (n=52)	P value
PONV within 24 hours	15 (29.4%)	17 (32.7%)	0.719
Use of antiemetics within 24 hours	10 (19.6%)	8 (15.4%)	0.573
NRS subject sleep quality, score ^a	5 (2, 7)	4 (2, 7)	0.717
Time to first ambulation, hour ^b	19.5 (17.8, 21.2)	20.0 (18.2, 21.8)	0.314
Volume of drainage within 24 hours, ml	30 (0, 55)	35 (0, 90)	0.248
LOS in hospital after surgery, day ^b	4.0 (3.7, 4.3)	4.0 (3.6, 4.4)	0.754

Data are presented as number (%) or median (interquartile range).

PONV: postoperative nausea and vomiting; NRS: numeric rating scale; LOS: length of stay.

^a Subjective sleep quality on the night of surgery;

^b Data were analysed by Kaplan-Meier analysis and compared by log-rank test; the results are presented as median (95% confidence interval).

Figures




Figure 1

The three trocar sites in retroperitoneoscopic renal surgery.




Figure 2

Sonography of the lateral TAP block, indicating the ultrasound anatomical structure (A) and the spread of local anaesthetic (B). The green arrow indicates the needle trajectory; EO, external oblique; IO, internal oblique; TA, transversus abdominis; LA, local anaesthetic.




Figure 3

Flow diagram of the study





Figure 4

NRS pain score at rest (A) or with coughing (B) within 24 hours after surgery. There were no significant differences between the two groups. IAFA: immediately awakening from anaesthesia.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementTableS1R.pdf](#)
- [TAPdataset2PPanalysisfinal.xlsx](#)