Statin Use is Associated With The Reduction in Hepatocellular Carcinoma Recurrence After Liver Surgery

Elias Khajeh

University of Heidelberg

Arash Dooghaie Moghadam

University of Heidelberg

Pegah Eslami

University of Heidelberg

Sadeq Ali-Hasan-Al-Saegh

University of Heidelberg

Ali Ramouz

University of Heidelberg

Saeed Shafiei

University of Heidelberg

Omid Ghamamejad

University of Heidelberg

Sepehr Abbasi

University of Heidelberg

Christian Rupp

University of Heidelberg

Christoph Springfield

Liver Cancer Center Heidelberg (LCCH)

Carlos Carvalho

Digestive Unit, Clinical Oncology, Champalimaud Clinical Centre

Pascal Probst

University of Heidelberg

Seyed Mostafa Mousavizadeh

University of Heidelberg

Arianeb Mehrabi (✉ Arianeb.Mehrabi@med.uni-heidelberg.de)

University of Heidelberg

Keywords: hepatocellular carcinoma, liver resection, liver transplantation, statins, meta-analysis.

Posted Date: October 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-950411/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Hepatocellular carcinoma (HCC) is the sixth most common form of cancer worldwide. Although surgical treatments have an acceptable cure rate, tumor recurrence is still a challenging issue. In this meta-analysis, we investigated whether statins prevent HCC recurrence following liver surgery.

Methods

PubMed, Web of Science, EMBASE and Cochrane Central were searched. The Outcome of interest was the HCC recurrence after hepatic surgery. Pooled estimates were represented as hazard ratios (HRs) and odds ratios (ORs) using a random-effects model. Summary effect measures are presented together with their corresponding 95% confidence intervals (CI). The certainty of evidence was evaluated using the Grades of Research, Assessment, Development and Evaluation (GRADE) approach.

Results

The literature search retrieved 1,362 studies excluding duplicates. Nine retrospective studies including 44,219 patients (2,243 in the statin group and 41,976 in the non-statin group) were included in the qualitative analysis. Patients who received statins had a lower rate of recurrence after liver surgery (HR: 0.41; 95% CI: 0.31–0.54; $p < 0.001$). Moreover, Statins decreased the recurrence one year after surgery (OR: 0.27; 95% CI: 0.16–0.47; $P < 0.001$), three years after surgery (OR: 0.22; 95% CI: 0.15–0.33; $P < 0.001$), and five years after surgery (OR: 0.28; 95% CI: 0.19–0.42; $P < 0.001$). The certainty of evidence for the outcomes was moderate.

Conclusion

Statins increase the disease-free survival of patients with HCC after liver surgery. These drugs seem to have chemoprevention effects that decrease the probability of HCC recurrence after liver transplantation or liver resection.

Background

Hepatocellular carcinoma (HCC) is the sixth most common malignancy (1) and is emerging as the fastest-growing fatal cancer in the United States with a rapidly rising mortality rate worldwide (2). Establishing the best treatment option for HCC is difficult and depends on the tumor stage at the time of diagnosis (3). Surgical or interventional curative approaches such as local ablation, surgical resection, and liver transplantation are the treatments of choice for tumors diagnosed in earlier stages (4–6). However, detecting early-stage tumors is not feasible in many cases and only 13% of HCC cases are

diagnosed early enough for curative therapy (7). Although surgical procedures like tumor resection or liver transplantation have acceptable cure rates, they also have high recurrence rates (8, 9), with recurrence in more than 50% of patients after five years of surgery (10, 11). Recurrence rates after transplantation are between 8 and 21% despite the use of new predictive models (12).

Both animal and human studies have shown an independent relationship between cholesterol levels and HCC progression (13). HCC cell lines use cholesterol in their cell membranes and for organelle division (14) and levels of high-density lipoprotein cholesterol have been associated with tumor aggressiveness (13). Statins are hydroxymethyl glutaryl coenzyme A (HMG-CoA) reductase inhibitors and are commonly used to lower cholesterol levels in blood (15). In addition, statins have immunomodulatory effects and can protect against cancer (16–19). Several studies have attempted to investigate the possible role of statins in preventing HCC recurrence (20). Early work showed that statins affect molecular pathways in HCC cell lines to prevent over-proliferation in vivo (21). These anti-cancer effects make statins an interesting candidate for HCC prevention (14).

Recent studies have suggested that statins might increase survival rates in HCC patients and reduce HCC recurrence rates after curative treatment (22, 23). In this systematic review and meta-analysis, we investigate the role of statins in preventing HCC recurrence after hepatic surgery.

Methods

The present study was reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and recommendations of the Study Center of the German Society of Surgery (24, 25).

Eligibility criteria

The research question and eligibility criteria were formulated based on the PICOS strategy (population, intervention, comparison, outcomes, and design of studies).

- *Population:* All patients with HCC who underwent hepatic surgery, including liver resection and liver transplantation
- *Intervention:* Treatment with statins
- *Comparators:* No statin treatment
- *Outcome:* HCC recurrence after surgery
- *Study design:* Any study design except case reports, study protocols, animal studies, conference papers, and letters to the editor.

To eliminate the risk of analyzing the same patients more than once, the studies were thoroughly assessed and double publications and overlapping reports were excluded. The remaining studies were selected for full-text review by reviewing the titles and abstracts for eligibility.

Literature search

The predefined search terms were: ("carcinoma, hepatocellular" OR "adenoma hepatocellular" OR "adenomas hepatocellular" OR "hepatocellular adenoma" OR "HCC" OR "hepatocellular carcinoma") AND ("Statin" OR "Hydroxymethylglutaryl-CoA Reductase Inhibitors" OR "HMG-CoA Reductase Inhibitor"). Our comprehensive literature search was conducted in Medline/PubMed, EMBASE, Web of Science and Cochrane Central databases from their inception until February 2021. We also searched PubMed/Medline and Cochrane Central for systematic reviews of randomized clinical trials on surgical interventions. All studies comparing HCC recurrence in adult patients who underwent liver surgery were included.

Study selection

Two authors (ADM and PE) independently screened all titles and abstracts and made their selections according to PICOS eligibility criteria. The full texts of appropriate studies were evaluated and their data were extracted by two authors (SAHS and AR) independently. Discrepancies were resolved through discussions with the first and senior authors (EK and AM). For each treatment group, the following data were extracted: study characteristics, patient characteristics, study quality, and outcome measures.

Critical appraisal

The quality of each study was assessed by two independent reviewers (SAHA and AR) using the methodological index for non-randomized studies (MINORS). Quality was determined based on 12 MINORS items and was scored as follows: 0 (not reported), 1 (reported but inadequate) or 2 (reported and adequate). The best score was 24 for comparative studies. Studies with 20 points or lower were deemed high risk of bias, 21–23 points intermediate risk of bias, and 24 points low risk of bias. The overall quality of the evidence for each outcome was also assessed using Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.

Statistical analysis

All data were analyzed by RevMan version 5.3 (Nordic Cochrane Centre, Cochrane Collaboration, Copenhagen, Denmark). The effect size for dichotomous outcomes were presented as odds ratios (OR) or hazard ratios (HR) with their corresponding 95% confidence intervals (CI). For analyzing ORs and HRs of included studies, random effects model was used. Statistical heterogeneity was evaluated with the I^2 statistic. I^2 values of 0–25% indicate insignificant heterogeneity, 26–50% indicate low heterogeneity, 51–75% indicate moderate heterogeneity, and 76–100% indicate high heterogeneity. The fixed-effects model was used when the I^2 value was <50% and the random-effects model was used when the I^2 value was >50%. A p-value less than 0.05 was considered statistically significant in all analyses.

Results

Literature search

The literature search retrieved 1,362 studies after duplicates were excluded. Evaluation of titles and abstracts excluded a further 1,297 articles. Of the 65 full-text articles assessed for eligibility, 56 were excluded for various reasons, including insufficient data on survival. Finally, 9 articles were included in the qualitative and five articles in the quantitative meta-analysis (Figure 1). All included studies were retrospective and reported on 44,219 patients (2,243 in the statin group and 41,976 in the non-statin group). All studies were published between 2012 and 2021. The certainty of evidence for the outcomes, assessed using the GRADE approach, was moderate.

Qualitative report

Of the nine included studies (Table 1), four studies (20, 23, 26, 27) had sufficient information regarding HRs of disease recurrence as well as recurrence free survival in statin and no-statin groups. The study of Young et al (28) only reported HR of using statin in the recurrence of HCC after liver surgeries. Lee (29), Yeh (30), and Wu Cy (31) studies reported neither crude HRs of recurrence nor disease free survival data in statin and non-statin groups and therefore were not included in our meta-analyses. The HRs reported in these three studies were results of multivariate analysis or were adjusted for some other variables. The Wu LL study (22) was also excluded from meta-analyses, because statin use was not reported separately in patients who underwent surgery. According to available data in included studies, we performed two separate meta-analyses of HRs and ORs.

Table 1
Included studies in qualitative analysis.

Author	Year	Country	Type	Statin group	Non-statin group	LRx or LTx	Remarks and main findings
Yang (26)	2021	Taiwan	Retrospective	46	774	LRx	<ul style="list-style-type: none"> - Statins included atorvastatin, fluvastatin, pitavastatin, and rosuvastatin. - Statin use significantly reduced HCC recurrence (HR: 0.354; p value < 0.001). - The statin group had higher RFS than the non-statin group after propensity score matching.
Young (28)	2020	Taiwan	Retrospective	30	400	LRx	<ul style="list-style-type: none"> - Statins included lovastatin, fluvastatin, rosuvastatin, atorvastatin, and pravastatin. - Statin use (HR = 0.50; 95% CI = 0.27–0.94, p = 0.031) was significantly associated with decreased recurrence in univariate analysis.

Abbreviations: HR, Hazard Ratio; LRx, liver resection; LTx, liver transplantation; RFS, recurrence-free survival

Author	Year	Country	Type	Statin group	Non-statin group	LRx or LTx	Remarks and main findings
Cho (23)	2019	Korea	Retrospective	112	235	LTx	<p>- The types of statins used in the study are not specified.</p> <p>- Statin therapy was associated with a reduced risk of HCC recurrence (OR = 0.38, 95% CI = 0.16–0.91).</p>
Nishio (27)	2018	Japan	Retrospective	43	600	LRx	<p>- 17 patients used pravastatin, 15 patients atorvastatin, 7 patients rosuvastatin, and 4 patients pitavastatin.</p> <p>- Significant improvement of both RFS (5-yr RFS, 55.4% in the statin group versus 25.0% in the non-statin group) and OS (5-yr OS, 73.1% versus 56.7%, respectively) in perioperative statin users.</p>

Abbreviations: HR, Hazard Ratio; LRx, liver resection; LTx, liver transplantation; RFS, recurrence-free survival

Author	Year	Country	Type	Statin group	Non-statin group	LRx or LTx	Remarks and main findings
Kawaguchi (20)	2017	Japan	Retrospective	31	703	LRx	<p>- Statins included pravastatin, simvastatin, fluvastatin, pitavastatin, atorvastatin, and rosuvastatin.</p> <p>- The RFS was significantly higher in the statin than non-statin group ($P < 0.001$): the 1-, 3-, and 5- year RFS were 87.1%, 76.7%, and 76.7%, respectively, in the statin group, and 65.3%, 40.6%, and 32.9%, respectively, in the non-statin group.</p> <p>- The OS was not significantly different between the groups.</p>
Wu LL (22)	2016	Taiwan	Retrospective	934	17958	LRx, and other treatments	<p>- The types of statins used in the study are not specified.</p> <p>- Better OS with surgery and statin use compared with RFA/PEI and statin use ($p = 0.0003$ and $p = 0.019$, respectively, for stages I and III).</p>

Abbreviations: HR, Hazard Ratio; LRx, liver resection; LTx, liver transplantation; RFS, recurrence-free survival

Author	Year	Country	Type	Statin group	Non-statin group	LRx or LTx	Remarks and main findings
Lee (29)	2016	Taiwan	Retrospective	132	2078	LRx	<ul style="list-style-type: none"> - The types of statins used in the study are not specified. - Use of statins and NSAIDs also can reduce the risk of recurrence of HCC and mortality after surgery.
Yeh (30)	2015	Taiwan	Retrospective	740	14834	LRx	<ul style="list-style-type: none"> - The types of statins used in the study are not specified. - The use of statin can significantly reduce risk of recurrent HCC (HR, 0.51; 95% CI, 0.42–0.61; P < 0.001).
Wu Cy(31)	2012	Taiwan	Retrospective	175	4394	LRx	<ul style="list-style-type: none"> - The use of statin was significantly associated with lower risk of tumor recurrence (HR: 0.68; 95% CI: 0.53-0.87; p = 0.002).
Abbreviations: HR, Hazard Ratio; LRx, liver resection; LTx, liver transplantation; RFS, recurrence-free survival							

The quality assessment of included studies is shown in Table 2. All included studies had an overall MINORS score of less than 20, indicating a considerable risk of bias. All included studies had a high risk of biased assessment of the study endpoint because they were non-randomized and non-blinded. In addition, all included studies were retrospective so did not prospectively calculate the sample size.

Table 2
Assessment of the quality of studies included in qualitative and quantitative analyses.

Authors	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Score¹
Yang (26)	2	1	0	2	2	2	2	0	2	0	2	2	17
Young (28)	2	2	0	2	1	2	2	0	2	0	2	2	17
Cho (23)	2	2	0	2	1	2	2	0	2	0	2	2	17
Nishio (27)	2	1	0	2	1	2	2	0	2	0	2	2	16
Kawaguchi (20)	2	2	2	2	1	2	2	0	2	0	2	2	19
Wu LL(22)	2	1	1	2	1	2	1	0	2	0	2	2	16
Lee (29)	2	1	0	2	1	2	1	0	2	0	2	2	15
Yeh (30)	2	1	0	2	1	2	2	0	2	0	1	2	15
Wu Cy(31)	2	1	1	2	1	2	1	0	2	0	2	2	16
<p>Q1. A clearly stated aim Q2. Inclusion of consecutive patients Q3. Prospective collection of data Q4. Endpoints appropriate to the aim of the study Q5. Unbiased assessment of the study endpoint Q6. Follow-up period appropriate to the aim of the study Q7. Loss to follow up less than 5% Q8. Prospective calculation of the study size Q9. An adequate control group Q10. Contemporary groups Q11. Baseline equivalence of groups Q12. Adequate statistical analyses</p> <p>¹ The items are scored 0 (not reported), 1 (reported but inadequate) or 2 (reported and adequate). The best total score is 16 for non-comparative studies and 24 for comparative studies.</p>													

Quantitative analysis

In 5 studies with totally 2,374 participants, association between statin use and HCC recurrence was reported as HRs. After pooling of HRs of HCC recurrence using a random-effect model, the rate of recurrence was significantly lower in patients who received statins before their surgery (HR: 0.41; 95% CI: 0.31– 0.54; $p < 0.001$; Figure 2). No significant heterogeneity was seen between the studies in this regard ($I^2 = 0\%$; $P = 0.64$).

HCC recurrence after surgery was reported in 2,544 patients from four studies (232 patients in the statin group and 2,312 patients in the non-statin group). HCC recurrence one year after surgery occurred in 16 patients (6.89%) in the statin group and in 663 patients (28.67%) in the non-statin group. Meta-analysis indicated that statin use decreased HCC recurrence one year after surgery (OR: 0.27; 95% CI: 0.16–0.47; $p < 0.001$; Figure 3) using a random-effects model. There was no considerable heterogeneity among the studies ($I^2 = 0\%$; $p = 0.39$).

HCC recurrence three years after surgery was reported in 31 patients (13.3%) in the statin group and in 1,159 patients (50.1%) in the non-statin group. Meta-analysis indicated that statin use decreased HCC recurrence three years after surgery (OR: 0.22; 95% CI: 0.15–0.33; $p < 0.001$; Figure 4) using a random-effects model. There was no considerable heterogeneity among the studies ($I^2 = 0\%$; $p = 0.56$).

HCC recurrence five years after surgery was reported in 49 patients (20.9%) in the statin group and in 1360 patients (58.8%) in the non-statin group. Meta-analysis indicated that statin use decreased HCC recurrence five years after surgery (OR: 0.28; 95% CI: 0.19–0.42; $p < 0.001$; Figure 5) using a random-effects model. There was no considerable heterogeneity among the studies ($I^2 = 18\%$; $p = 0.30$).

Discussion

HCC comprises more than 80% of primary liver cancers (32) and is the third most deadly cancer worldwide. In Europe, it is the seventh leading cause of death (33, 34). OS is higher if the diagnosis is made early (9), but high HCC recurrence rates after surgical treatment still remain a big challenge (12, 35). New methods are urgently needed to reduce recurrence, thereby improving long-term surgical outcomes and reducing healthcare-related costs in the future.

Because of the virulent character of HCC, prevention plays a significant role in treatment (36). Different studies have investigated the protective effect of several drugs against HCC recurrence, including statins, aspirin, and anti-diabetic agents (37). Statins are classified into lipophilic and hydrophilic statins and have been used extensively to prevent and treat cardiovascular diseases (38, 39). Studies have recently demonstrated that statins can reduce the risk of many cancers, including liver cancers (38, 40). Furthermore, statins can inhibit progression of liver fibrosis and cirrhosis in HCC patients (41) and can reduce the risk of HCC in patients with Hepatitis C Virus (HCV) infection and also with nonalcoholic fatty liver disease (NAFLD) (42, 43). Lipophilic statins seem to be more effective at preventing HCC than hydrophilic statins are (38). Fluvastatin has been revealed as the most effective drug in reducing HCC risk (37).

In this systematic review and meta-analysis, we investigated whether statins can reduce HCC recurrence following liver surgery. We showed that the recurrence was lower at one, three, and five years after surgery in patients with HCC who underwent liver surgery in combination with statin treatment than in patients who underwent liver surgery without statin treatment. These results indicate that statins should be considered effective at reducing the recurrence of HCC tumors.

Statins may reduce the risk of cancer via several mechanisms, including inhibiting oncogenic pathways, promoting tumor-specific apoptosis, inhibiting the proteasome pathway, inhibiting hepatitis virus replication, and reducing cholesterol synthesis (44, 45). Statins can also decrease endothelial dysfunction, intrahepatic vasoconstriction, inflammation, and fibrosis (46–48). The anti-inflammatory and immunomodulatory effects of statins allow them to inhibit harmful inflammatory and immunologic responses that may promote cancer (46–48). The effect of statins on liver regeneration and ischemia-

reperfusion injury after extensive hepatectomy has been investigated in animal models. These studies showed that these drugs can improve outcomes by facilitating regeneration and by inhibiting the harmful inflammatory response (49, 50). In a pilot clinical study, preoperative oral atorvastatin therapy for 3 days prior to liver resection reduced the harmful immunologic and inflammatory responses due to ischemia-reperfusion injury (51).

Patients who are not eligible for liver resection can be treated with more conservative options (22). Wu et al. (22) reported that combining statin treatment with these conservative methods improves the survival of patients with advanced HCC. In patients with a contraindication for these conservative therapies, especially those with HBV/HCV, palliative treatment with statins can reduce the mortality rate (22). It seems that combining statins and conservative therapies in patients who cannot undergo surgical therapies can improve the HCC prognosis (22, 52). According to Aaron et al., statins can improve the survival of HCC patients when administered both before or after HCC is diagnosed (53).

HCC recurrence is detected in half of patients three years after liver surgery (54). Recurrent disease is not easily treatable, so it is important to prevent recurrence after resection (20). Studies have shown that statins can reduce HCC recurrence rate by reducing viremia in patients with HBV and HCV. Possible mechanisms include reducing pro-inflammatory cytokines in serum (37, 55, 56), reducing the virulent potency of viral infections (20, 56), or inhibiting cirrhotic progression (56).

Preventing HCC recurrence after liver transplantation is also an important issue (57) but has not been well investigated. Statins have several side-effects (including myalgia and myotoxicity that may lead to rhabdomyolysis), especially when administered at high doses. These side-effects need to be properly investigated in post-transplant patients (58)(59). The prevalence of post-liver transplant dyslipidemia is 16–66% worldwide (46). Statins decrease lipidemia in patients after liver transplantation, thereby preventing cardiovascular events (58).

There are some limitations to the present systematic review and meta-analysis. We found no RCT that compared the clinical outcomes of liver surgery between statin and non-statin groups. Furthermore, the timing, dosage, and type of statin is important to evaluate the outcomes; however, this information was not provided in every study. Further well-designed, large-scale RCTs are needed to determine whether statin therapy prevents HCC recurrence after liver resection or transplantation.

In conclusion, statins increased disease-free survival of patients with HCC after liver surgery. They may reduce HCC recurrence after liver surgery by chemoprevention effects. Unfortunately, the existing evidence is still too limited (small study populations, retrospective study designs, and single-center studies) to confirm a role for statins in reducing disease recurrence in HCC patients. Further randomized clinical trials should confirm the effectiveness of statins in preventing HCC recurrence after liver surgery, and should determine the importance of different types of surgery and types of statins.

List Of Abbreviations

HCC

Hepatocellular Carcinoma

HR

hazard Ratio

OR

Odds Ratio

GRADE

Grades of Research, Assessment, Development, and Evaluation

HMG-CoA

Hydroxy-Methyl Glutaryl Coenzyme A

PRISMA

Preferred Reporting Items for Systematic Reviews and Meta-Analyses

MINORS

Methodological Index for Non-Randomized Studies

HCV

Hepatitis C Virus

NAFLD

Nonalcoholic Fatty Liver Disease.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Only publicly available data were used in this study, and data sources and handling of these data are described in the Materials and Methods and in the additional files. Further information is available from the corresponding author upon request.

Competing interests

All authors declare no conflicts of interests.

Funding

This study was funded by the Champalimaud Clinical Centre of the Champalimaud Foundation.

Authors' contributions

A.M. and E.K. contributed to the conception of study. A.D.M., P.E., S.A.H.S., and A.R. conducted the literature search. E.K., S.A.H.S. and A.R. analyzed the data and interpreted data. A.D.M., P.E., O.G., S.S., and S.M.M. drafted the manuscript, which E.K. A.M., C.R., C.S., C.C. and P.P. critically revised. All authors gave final approval of the article to be published.

Acknowledgements

None.

References

1. El-Serag HB. Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology*. 2012;142(6):1264-73.e1.
2. Mokdad AH, Dwyer-Lindgren L, Fitzmaurice C, Stubbs RW, Bertozzi-Villa A, Morozoff C, et al. Trends and Patterns of Disparities in Cancer Mortality Among US Counties, 1980-2014. *Jama*. 2017;317(4):388–406.
3. Akamatsu N, Cillo U, Cucchetti A, Donadon M, Pinna AD, Torzilli G, et al. Surgery and Hepatocellular Carcinoma. *Liver Cancer*. 2017;6(1):44–50.
4. Belghiti J, Kianmanesh R. Surgical treatment of hepatocellular carcinoma. *HPB: the official journal of the International Hepato Pancreato Biliary Association*. 2005;7(1):42–9.
5. Ho MC, Hasegawa K, Chen XP, Nagano H, Lee YJ, Chau GY, et al. Surgery for Intermediate and Advanced Hepatocellular Carcinoma: A Consensus Report from the 5th Asia-Pacific Primary Liver Cancer Expert Meeting (APPLE 2014). *Liver Cancer*. 2016;5(4):245–56.
6. Rahbari NN, Mehrabi A, Mollberg NM, Müller SA, Koch M, Büchler MW, et al. Hepatocellular carcinoma: current management and perspectives for the future. *Annals of surgery*. 2011;253(3):453–69.
7. El-Serag HB, Siegel AB, Davila JA, Shaib YH, Cayton-Woody M, McBride R, et al. Treatment and outcomes of treating of hepatocellular carcinoma among Medicare recipients in the United States: a population-based study. *Journal of hepatology*. 2006;44(1):158–66.
8. Fujiwara N, Friedman SL, Goossens N, Hoshida Y. Risk factors and prevention of hepatocellular carcinoma in the era of precision medicine. *Journal of hepatology*. 2018;68(3):526–49.
9. Wang JH, Wang CC, Hung CH, Chen CL, Lu SN. Survival comparison between surgical resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *Journal of hepatology*. 2012;56(2):412–8.
10. Al-Saeedi M, Ghamarnejad O, Khajeh E, Shafiei S, Salehpour R, Golriz M, et al. Pringle Maneuver in Extended Liver Resection: A propensity score analysis. *Scientific reports*. 2020;10(1):8847.
11. Khajeh E, Shafiei S, Al-Saegh SA, Ramouz A, Hammad A, Ghamarnejad O, et al. Meta-analysis of the effect of the pringle maneuver on long-term oncological outcomes following liver resection. *Scientific reports*. 2021;11(1):3279.

12. Bodzin AS. Hepatocellular carcinoma (HCC) recurrence and what to do when it happens. *Hepatobiliary Surg Nutr.* 2016;5(6):503–5.
13. Carr BI, Giannelli G, Guerra V, Giannini EG, Farinati F, Rapaccini GL, et al. Plasma cholesterol and lipoprotein levels in relation to tumor aggressiveness and survival in HCC patients. *The International journal of biological markers.* 2018;33(4):423–31.
14. Kaplan DE. Statins and Hepatocellular Carcinoma Protection. *Gastroenterol Hepatol (N Y).* 2019;15(4):190–3.
15. Ramkumar S, Raghunath A, Raghunath S. Statin Therapy: Review of Safety and Potential Side Effects. *Acta Cardiol Sin.* 2016;32(6):631–9.
16. Van Wyhe RD, Rahal OM, Woodward WA. Effect of statins on breast cancer recurrence and mortality: a review. *Breast Cancer (Dove Med Press).* 2017;9:559–65.
17. Strykowska-Gora A, Karczmarek-Borowska B, Gora T, Krawczak K. Statins and cancers. *Contemporary oncology (Poznan, Poland).* 2015;19(3):167–75.
18. Nielsen SF, Nordestgaard BG, Bojesen SE. Statin use and reduced cancer-related mortality. *The New England journal of medicine.* 2012;367(19):1792–802.
19. Demierre MF, Higgins PD, Gruber SB, Hawk E, Lippman SM. Statins and cancer prevention. *Nature reviews Cancer.* 2005;5(12):930–42.
20. Kawaguchi Y, Sakamoto Y, Ito D, Ito K, Arita J, Akamatsu N, et al. Statin use is associated with a reduced risk of hepatocellular carcinoma recurrence after initial liver resection. *Bioscience trends.* 2017;11(5):574–80.
21. Mullen PJ, Yu R, Longo J, Archer MC, Penn LZ. The interplay between cell signalling and the mevalonate pathway in cancer. *Nature reviews Cancer.* 2016;16(11):718–31.
22. Wu LL, Hsieh MC, Chow JM, Liu SH, Chang CL, Wu SY. Statins improve outcomes of nonsurgical curative treatments in hepatocellular carcinoma patients. *Medicine.* 2016;95(36):e4639.
23. Cho Y, Kim MS, Nam CM, Kang ES. Statin Use is Associated with Decreased Hepatocellular Carcinoma Recurrence in Liver Transplant Patients. *Scientific reports.* 2019;9(1):1467.
24. Moher D, Liberati A, Tetzlaff J, Altman DG, JIS. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. 2010;8(5):336–41.
25. Kalkum E, Klotz R, Seide S, Hüttner FJ, Kowalewski KF, Nickel F, et al. Systematic reviews in surgery-recommendations from the Study Center of the German Society of Surgery. *Langenbecks Arch Surg.* 2021.
26. Yang SY, Wang CC, Chen KD, Liu YW, Lin CC, Chuang CH, et al. Statin use is associated with a lower risk of recurrence after curative resection in BCLC stage 0-A hepatocellular carcinoma. *BMC cancer.* 2021;21(1):70.
27. Nishio T, Taura K, Nakamura N, Seo S, Yasuchika K, Kaido T, et al. Impact of statin use on the prognosis of patients with hepatocellular carcinoma undergoing liver resection: a subgroup analysis of patients without chronic hepatitis viral infection. *Surgery.* 2018;163(2):264–9.

28. Young SH, Chau GY, Lee IC, Yeh YC, Chao Y, Huo TI, et al. Aspirin is associated with low recurrent risk in hepatitis B virus-related hepatocellular carcinoma patients after curative resection. *Journal of the Formosan Medical Association = Taiwan yi zhi*. 2020;119(1 Pt 2):218–29.
29. Lee PC, Yeh CM, Hu YW, Chen CC, Liu CJ, Su CW, et al. Antiplatelet Therapy is Associated with a Better Prognosis for Patients with Hepatitis B Virus-Related Hepatocellular Carcinoma after Liver Resection. *Ann Surg Oncol*. 2016;23(Suppl 5):874–83.
30. Yeh CC, Lin JT, Jeng LB, Ho HJ, Yang HR, Wu MS, et al. Nonsteroidal anti-inflammatory drugs are associated with reduced risk of early hepatocellular carcinoma recurrence after curative liver resection: a nationwide cohort study. *Annals of surgery*. 2015;261(3):521–6.
31. Wu CY, Chen YJ, Ho HJ, Hsu YC, Kuo KN, Wu MS, et al. Association between nucleoside analogues and risk of hepatitis B virus–related hepatocellular carcinoma recurrence following liver resection. *Jama*. 2012;308(18):1906–14.
32. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*. 2018;68(6):394–424.
33. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA: a cancer journal for clinicians*. 2011;61(2):69–90.
34. Yuen MF, Hou JL, Chutaputti A. Hepatocellular carcinoma in the Asia pacific region. *Journal of gastroenterology and hepatology*. 2009;24(3):346–53.
35. Yang Z, Miao R, Li G, Wu Y, Robson SC, Yang X, et al. Identification of recurrence related microRNAs in hepatocellular carcinoma after surgical resection. *International journal of molecular sciences*. 2013;14(1):1105–18.
36. Kim G, Jang SY, Nam CM, Kang ES. Statin use and the risk of hepatocellular carcinoma in patients at high risk: A nationwide nested case-control study. *Journal of hepatology*. 2018;68(3):476–84.
37. Fujiwara N, Friedman SL, Goossens N, Hoshida Y. Risk factors and prevention of hepatocellular carcinoma in the era of precision medicine. *Journal of hepatology*. 2018;68(3):526–49.
38. Shi M, Zheng H, Nie B, Gong W, Cui X. Statin use and risk of liver cancer: an update meta-analysis. *BMJ Open*. 2014;4(9):e005399.
39. Björkhem-Bergman L, Backheden M, Söderberg Löfdal K. Statin treatment reduces the risk of hepatocellular carcinoma but not colon cancer—results from a nationwide case-control study in Sweden. *Pharmacoepidemiology and Drug Safety*. 2014;23(10):1101–6.
40. Li Z, Li Y, Li X, Zhang L, Zhao N, Du H, et al. Statins on hepatocellular carcinoma risk in hepatitis B or C patients protocol for a systematic review and meta-analysis. *Medicine*. 2018;97(34):e11950.
41. Kaplan DE, Serper MA, Mehta R, Fox R, John B, Aytaman A, et al. Effects of Hypercholesterolemia and Statin Exposure on Survival in a Large National Cohort of Patients With Cirrhosis. *Gastroenterology*. 2019;156(6):1693-706.e12.
42. Tsan YT, Lee CH, Ho WC, Lin MH, Wang JD, Chen PC. Statins and the risk of hepatocellular carcinoma in patients with hepatitis C virus infection. *Journal of clinical oncology: official journal of*

- the American Society of Clinical Oncology. 2013;31(12):1514–21.
43. Hajifathalian K, Tafesh Z, Rosenblatt R, Kumar S, Homan EA, Sharaiha RZ, et al. Effect of Statin Use on Cancer-related Mortality in Nonalcoholic Fatty Liver Disease: A Prospective United States Cohort Study. *Journal of clinical gastroenterology*. 2021.
 44. Higashi T, Hayashi H, Kitano Y, Yamamura K, Kaida T, Arima K, et al. Statin attenuates cell proliferative ability via TAZ (WWTR1) in hepatocellular carcinoma. *Medical oncology (Northwood, London, England)*. 2016;33(11):123.
 45. Chiu HF, Ho SC, Chen CC, Yang CY. Statin use and the risk of liver cancer: a population-based case-control study. *The American journal of gastroenterology*. 2011;106(5):894–8.
 46. Gazi IF, Liberopoulos EN, Athyros VG, Elisaf M, Mikhailidis DP. Statins and solid organ transplantation. *Current pharmaceutical design*. 2006;12(36):4771–83.
 47. Greenwood J, Mason JC. Statins and the vascular endothelial inflammatory response. *Trends in immunology*. 2007;28(2):88–98.
 48. Dulak J, Józkowicz A. Anti-angiogenic and anti-inflammatory effects of statins: relevance to anti-cancer therapy. *Current cancer drug targets*. 2005;5(8):579–94.
 49. Colakoglu T, Nursal TZ, Ezer A, Kayaselcuk F, Parlakgumus A, Belli S, et al. Effects of different doses of statins on liver regeneration through angiogenesis and possible relation between these effects and acute phase responses. *Transplantation proceedings*. 2010;42(9):3823-7.
 50. Tokunaga T, Ikegami T, Yoshizumi T, Imura S, Morine Y, Shinohara H, et al. Beneficial effects of fluvastatin on liver microcirculation and regeneration after massive hepatectomy in rats. *Digestive diseases and sciences*. 2008;53(11):2989–94.
 51. Sarin S, Kaman L, Dahiya D, Behera A, Medhi B, Chawla Y. Effects of preoperative statin on liver reperfusion injury in major hepatic resection: a pilot study. *Updates in surgery*. 2016;68(2):191–7.
 52. Shao JY, Lee FP, Chang CL, Wu SY. Statin-Based Palliative Therapy for Hepatocellular Carcinoma. *Medicine*. 2015;94(42):e1801.
 53. Thrift AP, Natarajan Y, Liu Y, El-Serag HB. Statin Use After Diagnosis of Hepatocellular Carcinoma Is Associated With Decreased Mortality. *Clinical gastroenterology and hepatology: the official clinical practice journal of the American Gastroenterological Association*. 2019;17(10):2117-25.e3.
 54. Poon RT, Fan ST, Lo CM, Liu CL, Wong J. Long-term survival and pattern of recurrence after resection of small hepatocellular carcinoma in patients with preserved liver function: implications for a strategy of salvage transplantation. *Annals of surgery*. 2002;235(3):373–82.
 55. Singh S, Singh PP. Statins for prevention of hepatocellular cancer: one step closer? *Hepatology (Baltimore, Md)*. 2014;59(2):724–6.
 56. Janicko M, Drazilova S, Pella D, Fedacko J, Jarcuska P. Pleiotropic effects of statins in the diseases of the liver. *World J Gastroenterol*. 2016;22(27):6201–13.
 57. Siegel AB, El-Serag HB. Statins for chemoprevention of hepatocellular carcinoma: assessing the evidence. *Expert review of gastroenterology & hepatology*. 2013;7(6):493–5.

58. Bjorkhem-Bergman L. Is There a Role for Statins in Palliative Care for Patients Suffering from Hepatocellular Carcinoma? *Journal of palliative care*. 2015;31(3):172–6.

59. Muck W, Neal DA, Boix O, Voith B, Hasan R, Alexander GJ. Tacrolimus/cerivastatin interaction study in liver transplant recipients. *British journal of clinical pharmacology*. 2001;52(2):213–5.

Figures

Figure 1

Flow chart of study.

Figure 2

Pooled analysis of hazard ratios for recurrence of HCC

Figure 3

Forest plot for HCC recurrence one year after liver surgery.

Figure 4

Forest plot for HCC recurrence three years after liver surgery.

Figure 5

Forest plot for HCC recurrence five years after liver surgery.