Health care use and treatment-seeking for depression symptoms in rural India: A descriptive cross-sectional analysis

Tessa Naomi Roberts (✉ tessa.roberts@kcl.ac.uk)

Institute of Psychiatry Psychology and Neuroscience <https://orcid.org/0000-0001-8584-4162>

Rahul Shidhaye

Pravara Institute of Medical Sciences

Vikram Patel

Harvard Medical School Department of Global Health and Social Medicine

Sujit D Rathod

London School of Hygiene and Tropical Medicine

Research article

Keywords: Depression, help seeking, health service utilisation, India

Posted Date: December 13th, 2019

DOI: <https://doi.org/10.21203/rs.2.18888/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on April 6th, 2020. See the published version at <https://doi.org/10.1186/s12913-020-05162-0>.

Abstract

Background

There is a large “treatment gap” for depression worldwide. This study aimed to better understand the treatment gap in rural India by describing health care use and treatment-seeking for depression.

Methods

Data were analysed from a two round cross-sectional community survey carried out in rural Madhya Pradesh between May 2013 and December 2016. We examined the proportion of individuals with probable depression who sought treatment in different sectors, for depression symptoms and for any reason, and compared the latter with health service use by non-depressed individuals. We showed the frequency with which barriers to healthcare utilisation were reported by adults with probable depression, and examined differences in the proportion of adults who sought treatment for depression by predisposing, enabling and need factors.

Results

86% of adults with probable depression reported seeking no depression treatment. However, 66% had used health services for any reason in the past 3 months, compared to 46% of those without depression ($p < 0.0001$). Private providers were most frequently consulted by adults with probable depression (32%), while only 19% consulted traditional providers. Structural barriers to healthcare use such as cost and distance to services were frequently reported (54% and 52%, respectively) but were not associated with treatment-seeking for depression. The following factors were found to be positively associated with treatment-seeking for depression: higher symptom severity; reporting lack of energy, lack of interest/pleasure, low self-esteem, or slow movements/restlessness on more than 7 days in the past 2 weeks; being married; having discussed depression symptoms; and reporting problems with medication availability and supply as a barrier to healthcare. No evidence was found for an association between treatment-seeking for depression and most socio-economic, demographic or attitudinal factors.

Discussion and conclusions

The majority of adults with probable depression actively seek healthcare, but not specifically for depression symptoms, indicating the need to improve detection of depression during consultations about other complaints. Private providers should be included in programmes to improve the detection and treatment of depression. Unmarried individuals may experience greater difficulties in accessing care in this setting. Further research should test the hypotheses generated in this descriptive study.

Background

Depressive disorders are largely untreated despite accounting for an enormous burden of disease. The 2010 Global Burden of Disease study found that depression was the second leading cause of disability

worldwide (1). However, in developed countries only 54.3% of people with a 12 month major depressive episode report visiting any service provider for mental health reasons in the past year, and just 25.2% in low- and middle-income countries (LMIC) (2). Fewer than half of those who sought help received minimally adequate treatment according to evidence-based guidelines (3).

The reasons for low demand for services in LMIC are poorly understood. The World Health Organization (WHO) advocates integrating evidence-based interventions into primary care to increase the availability and accessibility of services (4) as a strategy to reduce the gap. Yet in the World Mental Health Surveys, only 34.6% of people with depression in LMIC regarded themselves as needing treatment (3), suggesting that the treatment gap cannot be explained solely in terms of limited availability of mental health services.

Few data are available to inform strategies to promote treatment-seeking in LMIC, such as India, where the treatment gap for depression is over 85% (5). Two recent systematic reviews on treatment-seeking for common mental disorders showed that “need factors”, such as greater symptom severity, chronicity, and disability, are positively associated with the likelihood of seeking treatment, and that women, the middle-aged, those with higher levels of education, and people of Caucasian ethnicity are more likely to seek treatment in high-income countries (6, 7). They also showed that factors such as income, employment, and place of residence were generally not associated with treatment-seeking. However, there was a relative lack of evidence from LMIC, and few data were available with which to evaluate factors such as beliefs, attitudes, social support or health systems characteristics, which are hypothesised to be important to treatment decisions (7).

In India, data on treatment-seeking for depression are scarce. In 2016, a systematic review of “contact coverage” (8) (i.e. the proportion of adults with depression who sought treatment for depression (9)) found only one population-based study of treatment-seeking from India. This study reported that rural residents were less likely to seek treatment than urban residents, with no clear association with wealth (10). However, the researchers used receipt of a depression diagnosis as its outcome measure, which conflates treatment-seeking with health care providers’ ability to detect and diagnose depression. Other Indian studies have reported on the use of general health services by people with depression, but without distinguishing between treatment sought for depression and for other health problems (11–13). As such, very little evidence is available from India to inform efforts to reduce the treatment gap for depression.

Evidence-based strategies for reducing the treatment gap can only be devised if service planners have access to information on who seeks treatment, under what circumstances, and from where, some of which may differ between settings. This study is a descriptive analysis of treatment-seeking for depression by adults in Sehore sub-district, Madhya Pradesh, with the following specific objectives:

1. To estimate the proportion of adults with probable depression who consult different types of treatment providers, (a) for depression symptoms and (b) for any reason, and to compare the latter with general health care use by people without probable depression;

2. To measure the prevalence of self-reported barriers to using health services among adults with probable depression;
3. To estimate the change in probability of treatment-seeking for symptoms of depression associated with need, predisposing and enabling factors.

Methods

Setting

Sehore sub-district is a predominantly rural area in Madhya Pradesh, with a population of 427,432 (14). 31.7% live below the poverty line and agriculture is the mainstay of the local economy (15). General health indicators are below the national average (16), literacy rates are 81% for males and 58% for females (14), and 88% of residents have completed only primary education or less (17).

The PRIME programme (Programme for Improving Mental Health Care) aimed to implement and evaluate district-level Mental Health Care Plans (MHCP) (18). The MHCP for Sehore focussed on depression, psychosis and alcohol use disorders and was implemented through community health centres between August 2014 and October 2016 (19).

Prior to implementation of the MHCP, outpatient and inpatient services were provided through Sehore District Hospital, by one psychiatrist and one clinical psychologist who are employed under the District Mental Health Programme and provide their services on alternate days, with periodic “outreach camps” (20). No psychotropic medication or psychosocial interventions were available in primary care facilities, there were no psychiatric social workers or psychiatric nurses, and primary care workers were largely untrained in identifying and treating mental disorders. After the plan was implemented, depression treatment was available at three Community Health Centres with psychological interventions delivered by case managers and pharmacological treatments prescribed for severe cases by medical officers. Community awareness activities were conducted to encourage service uptake, such as community meetings and proactive case finding in the community by the case managers. They also screened patients in Community Health Centres. The study area (20), Mental Health Care Plan (21), and PRIME evaluation plan (22) have been described in more detail elsewhere. The term “implementation area” will be used to refer to those villages where MHCP activities were fully implemented.

Sample

This report is a secondary analysis of data from a population-based, cross-sectional community survey carried out with the primary aim of estimating the change in treatment-seeking among adults with probable depression, before and after implementation of the MHCP. This secondary analysis focuses on characterising treatment-seeking patterns for adults with probable depression in both rounds.

The study design, sampling plan, and data collection have been described in detail elsewhere (21, 22). Briefly, data collection for the first round took place prior to Mental Health Care Plan implementation, in two waves (May-June 2013 and January-March 2014), and the second round after implementation of the plan (October-December 2016). The target population was adults (aged 18 and above) residing within the implementation area, with participants selected from voter lists through systematic random sampling. Inclusion criteria were fluency in spoken Hindi, residency in the selected household, willingness to provide informed consent, and absence of cognitive impairments that would preclude informed consent or ability to participate.

Across both rounds, 6,203 adults were recruited, 6,134 (98.9%) consented to participate, and 4,297 resided within catchment areas of the de facto implementation area, where treatment was made available in the Community Health Centres. Of the 4,297, 568 adults (289 in round 1, 279 in round 2) screened positive for depression and comprise the primary sub-sample for this secondary analysis. No difference in the probability of treatment-seeking for depression was observed between rounds, so for the purposes of the current analyses, data from both rounds were pooled to increase statistical power (23). In order to compare use of health services by adults with and without depression, for this analysis we also included the 3,531 community survey participants who resided within the implementation, did not screen positive for depression, and who did not report equivalent symptoms within the past 12 months. The sample size was calculated for the parent study based on the numbers required to detect a difference in contact coverage between rounds (the proportion of people with depression and alcohol use disorders who sought treatment for their condition), as described elsewhere (21).

Data collection

Interviews were administered orally, in Hindi, by trained local fieldworkers who recorded participant responses using a questionnaire application programmed on Android tablets. The structured questionnaire included sections on socio-demographic details, health care use, barriers to using health services, depression symptoms, treatment-seeking for depression, alcohol use and related treatment, disability, internalised stigma related to depression and alcohol use, suicidal ideation and behaviours, and mental health knowledge and attitudes.

Study measures

The Patient Health Questionnaire (PHQ-9) consists of 9 items on depression symptoms which are summed to generate a symptom score (24). We used a cut-off point of ≥ 10 to indicate probable depression (25, 26) which has previously been validated in India (27, 28). Participants were also asked if they had experienced equivalent symptoms for any 2 week period in the past 12 months.

Barriers to the use of health services were based on the Study on Global Ageing and Adult Health (SAGE) (29). We added one question in round 2 on distance to health services. These barriers were not specific to

depression.

We chose factors to investigate based the Andersen socio-behavioural model (30, 31), which groups factors associated with health service utilisation into; (a) need factors, which include both objective and subjective assessments of health status, (b) predisposing factors, covering both demographic characteristics and attitudinal factors such as health beliefs, and (c) enabling factors, which refers to structural determinants such as financial situation, transport and social support.

Predisposing factors included gender, religion, education, age, caste, marital status and internalised stigma (measured using questions from the Internalized Stigma of Mental Illness (ISMI) scale (32)). Enabling factors included land ownership, housing type, employment status, discussing depression symptoms with someone, and reporting cost and travel barriers to health care. Need factors included symptom severity, disability (measured using the 12-item World Health Organization Disability Assessment Schedule (WHODAS 2.0)(33)), perceived need for health care, probable alcohol use disorder (measured using the Alcohol Use Disorders Identification Test (AUDIT) with a cut-off of ≥ 8 (34–37)), suicidal thoughts (measured using the Composite International Diagnostic Interview (CIDI) suicidality module (38)), and PHQ-9 item-specific symptoms of depression.

Treatment-seeking was measured after completing the PHQ-9 questionnaire by asking “Did you seek any treatment for these problems at any time in the past 12 months?”. Thus, in this report, “treatment-seeking for depression” refers to seeking treatment for the symptoms listed in the PHQ-9. Participants who answered affirmatively were asked to specify the type of provider consulted. In the section on health care utilisation, participants were asked “In the last three months, have you visited any health facility or provider for any health problem?”, and in which sector. Details of all measures used, and how these were treated in the analysis, are presented in the supplementary material.

Analysis

First, we describe the sociodemographic and clinical characteristics of the sub-sample of adults with probable depression, using unweighted counts and weighted percentages to account for the sampling design.

To estimate proportion of adults with probable depression who consult different types of treatment providers for depression symptoms and for general healthcare, we present the frequency of self-reported treatment-seeking for depression symptoms and general health care use, using weighted percentages and unweighted counts. We also present the frequency of general health care use by adults without depression (excluding those who reported depression symptoms over the past 12 months) and compare these proportions using Chi squared tests.

We next measure the prevalence of self-reported barriers to health service use by adults with probable depression, by presenting percentages on the frequency with which each barrier was reported, again

using weighted percentages and unweighted counts.

To assess the association between perceived need, predisposing and enabling factors and treatment-seeking for depression, we present the proportion of adults with probable depression who sought treatment for depression by each characteristic, along with prevalence ratios and 95% confidence intervals, and tested the association between each variable with the outcome of treatment-seeking for depression using univariable log-linear regression analyses. For brevity, we present only the results for factors where this association reached a significance level of $p < 0.05$, but a full table is included in the supplementary material. Since these analyses were intended to be descriptive and hypothesis-generating, rather than causal and hypothesis-testing, we did not conduct multivariable analyses to control for potential confounders. In order to interpret the findings on the effect of discussing depression symptoms (presumed to be a proxy measure for social support), we also examined participants' self-reports on who they discussed symptoms with, but the numbers in each group were too small to treat as separate variables.

All analyses were conducted using Stata/IC 15.1 (39). Frequencies are reported as observed, while percentages, regression coefficients, 95% confidence intervals, and P-values are design adjusted.

Ethics

Researchers explained the purpose of the survey to potential participants, read out the contents of study information sheets, and answered potential participants' questions. Informed consent was indicated with either a signature or a thumbprint. All screen-positive participants who were not receiving treatment were referred to the nearest public health facility where depression treatment was available.

Ethical approval was provided by the World Health Organization Research Ethics Review Committee (Geneva, Switzerland), the Sangath Institutional Review Board (Goa, India), and the London School of Hygiene & Tropical Medicine Observational Ethics Committee (London, United Kingdom) (ref: 10439).

Results

Sample characteristics

The socio-demographic and mental health characteristics of participants with and without probable depression are described in Table 1. Among those with probable depression, the mean age was 45.4 years, there were approximately equal proportions of men and women (53.8% female), and most participants were Hindu (92.1%), married (81.7%), and had not completed primary education (74.1%). The majority of participants with probable depression had moderate symptoms (77.9%). Tiredness or lack of energy was the most frequently reported symptom (reported by 79.2% on more than 7 days in the past 2 weeks), followed by feeling depressed or hopeless (63.3%).

Table 1

Socio demographic and mental health characteristics of adults with and without probable depression in
Sehore sub-district, India, 2013–2016

Characteristic	Adults with probable depression, N (%)	Adults without probable depression, N (%)
Gender		
Female	321 (53.8)	1,589 (43.9)
Male	247 (46.2)	1,942 (56.1)
Age group, years		
18–29	98 (17.5)	1,138 (32.8)
30–49	248 (44.1)	1,514 (43.0)
50–90	222 (38.4)	879 (24.2)
Education level completed		
Less than primary	419 (74.1)	2,017 (56.8)
Primary	129 (22.4)	1,124 (32.8)
Secondary or more	20 (3.5)	390 (10.5)
Employment status		
Unemployed	20 (4.2)	53 (1.7)
Productive non-income	241 (38.5)	1,401 (34.0)
Low income	277 (51.9)	1,785 (52.4)
High income	30 (5.4)	289 (7.9)
Religion		
Hindu	525 (92.1)	3,180 (89.8)
Muslim	43 (7.9)	350 (10.2)
Christian	0 (0)	1 (0.0)
Caste		
Scheduled Caste	101 (15.8)	516 (14.2)
Scheduled Tribe	25 (4.2)	140 (4.0)

Counts reported as observed, percentages are design adjusted.

Characteristic	Adults with probable depression, N (%)	Adults without probable depression, N (%)
Other Backwards Caste	393 (71.0)	2,503 (71.1)
General	49 (9.1)	372 (10.7)
Marital status		
Single	32 (6.4)	375 (10.9)
Married	461 (81.7)	2,953 (83.9)
Widowed / Separated / Divorced	75 (11.9)	213 (5.3)
Current depression severity (PHQ-9 score)		
Moderate (10–14)	450 (77.9)	0 (0.0)
Moderately severe (15–19)	107 (20.1)	0 (0.0)
Severe (≥ 20)	11 (2.0)	0 (0.0)
Depression-related symptoms reported on more than 7 days in past 2 weeks		
Tiredness / lack of energy	450 (79.2)	842 (23.3)
Feeling depressed or hopeless	371 (63.3)	255 (7.5)
Sleep problems	333 (58.1)	347 (9.8)
Lack of interest or pleasure	289 (53.3)	175 (5.5)
Appetite problems	293 (49.9)	282 (8.2)
Lack of concentration	229 (40.3)	155 (4.5)
Low self-esteem / feeling like a failure	123 (22.5)	38 (1.2)
Slow movements / restlessness	119 (22.2)	47 (1.3)
Thoughts of death / self-harm	37 (7.4)	3 (0.0)
Counts reported as observed, percentages are design adjusted.		

The non-depressed group included more males, more people with secondary education, fewer unemployed people, and the mean age was lower (39.6 years).

Objective 1: Use of health services and treatment-seeking for depression

Table 2 shows the health care used for any reason in the past 3 months by adults with and without probable depression.

65.6% of adults with probable depression had used health services for some reason in the past three months. Of these, 48.4% consulted the private sector while 29.8% consulted public providers and 29.3% consulted traditional providers. Those with probable depression were more likely to have used health services in the past 3 months than those without depression (65.6% vs. 45.7%, $p < 0.0001$).

Table 2. Health care used in the past 3 months for any reason by adults with and without probable depression in Sehore sub-district, 2013–2016

	By those with probable depression, N (%) (n = 568)	By those without probable depression, N (%) (n = 3,531)	P-value
Private health care provider	165 (32.0)	638 (19.0)	< 0.0001
Public health care provider	108 (19.6)	408 (11.5)	< 0.0001
Traditional service provider	119 (19.2)	675 (18.2)	0.60
Mental health specialist	1 (0.3)	0 (0)	0.02
Other	3 (0.6)	5 (0.2)	0.06
None	205 (34.4)	1,909 (54.3)	< 0.0001
Counts reported as observed, percentages are design adjusted.			

Table 3 shows treatment sought specifically for depression symptoms in the past 12 months by adults with probable depression.

13.9% of adults with probable depression sought treatment for depression symptoms, and of these, 61.3% did so from generalist providers, compared to 22.1% who consulted specialists and 16.7% who consulted traditional service providers.

Table 3. Health care used in the past 12 months for depression symptoms by adults with probable depression in Sehore sub-district, 2013–2016.

	By those with probable depression, N (%) (n = 568)
Generalist health worker (including case managers employed under the mental health care plan)	48 (8.5)
Specialist mental health worker	13 (3.1)
Traditional service provider	14 (2.3)
None	493 (86.1)
Counts reported as observed, percentages are design adjusted.	
Total exceeds 100% because some participants visited more than one sector.	

Objective 2: Barriers to health care use

Table 4 presents self-reported barriers to health care use by adults with probable depression. Cost and distance barriers were the most commonly reported barriers, with each reported by more than half of the sample who were asked about these (54.3% and 52.3%, respectively). The third most commonly reported barrier was the belief that health services were not needed (31.3%).

Table 4. Self-reported barriers to health care use among adults with probable depression in Sehere sub-district, 2013–2016

Barrier	Number of adults with probable depression who reported barrier (%)
Fees are not affordable	302 (54.3)
Services are too far away	145*(52.3)
Services not currently needed	172 (31.3)
Dislike taking medications	178 (30.9)
Care received is not good enough	148 (26.2)
Care providers do not understand my health problems	135 (23.4)
They don't have medicines I need	95 (17.3)
They frequently run out of medicines	81 (15.9)
Other reason	56 (11.2)
Seeking some kinds of treatment can make me or my family feel embarrassed	38 (8.1)
All percentages are adjusted for the complex sampling strategy.	
*only measured in follow up round, so denominator was 279.	

Objective 3: Factors associated with treatment-seeking for depression symptoms

Table 5 shows those associations between need, predisposing and enabling factors and treatment-seeking among all adults with probable depression that evidence suggestive of an association ($P < 0.05$). See the supplementary material for the full set of results.

Table 5

Association between need, predisposing and enabling factors and treatment-seeking for depression among adults with probable depression in Sehore sub-district, 2013–2016

	Total seeking treatment (n)	Prevalence of treatment-seeking, % (95% CI)	Prevalence ratio (95% CI)	p- value
Need factors				
Symptom severity (total current PHQ score)				
10–14	50/450	11.5 (8.5–15.5)	1	< 0.01
15–19	20/107	20.7 (13.2–30.8)	1.79 (1.11– 2.88)	
≥ 20	5/11	39.5 (12.8–74.5)	3.42 (1.33– 8.81)	
Tiredness/lack of energy				
< 7 days in past 2 weeks	10/118	7.3 (3.8–13.5)	1	0.03
≥ 7 days in past 2 weeks	65/450	15.7 (11.7–20.6)	2.14 (1.08– 4.24)	
Lack of interest or pleasure				
< 7 days in past 2 weeks	26/279	9.7 (6.3–14.7)	1	0.01
≥ 7 days in past 2 weeks	49/289	17.6 (13.2–23.2)	1.82 (1.16– 2.85)	
Low self-esteem / feeling like a failure				
< 7 days in past 2 weeks	51/445	11.5 (8.5–15.3)	1	< 0.01
≥ 7 days in past 2 weeks	24/123	22.4 (15.2–31.9)	1.96 (1.28-3.00)	
Slow movements / restlessness				
< 7 days in past 2 weeks	51/449	12.2 (8.9–16.5)	1	0.01
≥ 7 days in past 2 weeks	24/119	20.1 (14.4–29.3)	1.65 (1.13– 2.39)	

Counts reported as observed. Prevalence ratios, percentages and P-values are design adjusted.

This table presents data for only those factors for which there was evidence suggestive of an association with treatment-seeking for depression ($P < 0.05$). See the supplementary material for full set of findings.

	Total seeking treatment (n)	Prevalence of treatment-seeking, % (95% CI)	Prevalence ratio (95% CI)	p- value
Predisposing factors				
Marital status				
Single / separated / widowed	7/107	5.9 (2.7–12.2)	1	0.02
Married	68/461	15.7 (11.9–20.6)	2.67 (1.19– 5.99)	
Enabling factors				
Spoken to someone about these problems				
No	13/352	3.9 (2.2-7.0)	1	< 0.001
Yes	62/216	29.4 (23.1–36.5)	7.50 (4.11– 13.68)	
Services don't have medications I need				
No	55/473	11.9 (8.7–16.2)	1	0.01
Yes	20/95	24.4 (15.9–35.6)	1.99 (1.19– 3.32)	
Services frequently run out of medications				
No	56/487	11.9 (8.4–16.7)	1	0.01
Yes	19/81	23.6 (16.3–33.0)	2.05 (1.23– 3.39)	
Counts reported as observed. Prevalence ratios, percentages and P-values are design adjusted.				
This table presents data for only those factors for which there was evidence suggestive of an association with treatment-seeking for depression (P < 0.05). See the supplementary material for full set of findings.				

Among the “need factors”, the following were positively associated with treatment-seeking: symptom severity (39.5% of those with severe symptoms sought treatment compared to 11.5% of those with moderate symptoms), and reporting four specific symptoms on the PHQ-9 on 7 or more days in the past 2 weeks; tiredness or lack of energy, lack of interest or pleasure, low self-esteem or feeling like a failure, and slow movements or restlessness.

Under “predisposing factors”, 5.9% of unmarried people (single, separated or widowed) sought help for depression compared to 15.7% of those who were married.

Among “enabling factors”, 29.4% of those who discussed symptoms sought help compared to 3.9% of those who did not. Spouses were the most common person who symptoms were discussed with (67.5%; data not presented). There was a positive association between treatment-seeking for depression and reporting that “services frequently run out of medications” and “services don’t have the medications I need” as barriers.

Discussion

Principal findings

Although few people sought treatment specifically for depression symptoms, almost two thirds of adults with probable depression had recent contact with health services, which was significantly higher than by adults without probable depression. The private sector was most frequently consulted, while traditional services were used least, indicating that private health services are an important platform through which individuals with depression could theoretically be identified and treated. Structural barriers to using health services such as cost and distance are felt to be major barriers to the use of health care, but the current evidence suggests that reporting these barriers is unrelated to treatment-seeking for depression. These findings suggest the potential importance of social support and marriage in seeking treatment for depression in this context.

Implications for service planning and future research

Use of health services for non-depression reasons

Adults with high levels of depression symptoms are likely to be in contact with health services, but their primary complaints are rarely the depression symptoms listed in the PHQ-9. This echoes previous findings from India that depressed individuals frequently present to health services with somatic symptoms (11, 12, 40–42). Therefore, the most important challenge from a public health perspective appears not to be to persuade depressed individuals to visit services, but rather to enable health workers to recognise their mental health needs during consultations about other complaints. In other words, the relevant “treatment gap” is not between those who do and do not consult health services, but between those who receive effective treatment and those who do not. Health workers should be trained and supervised to distinguish psychosomatic symptoms from other health problems that are comorbid with depression, and provide appropriate care.

Use of the private sector

Importantly, however, we found that adults with depression are more likely to consult private than public health care providers, highlighting the importance of engaging private providers in initiatives to improve depression care. In the state of Madhya Pradesh, 76% of qualified medics and 72% of qualified paramedical staff are employed in the private sector (43). India has one of the most privatised health systems in the world (44), with around 80% of outpatient care provided in the private sector (45, 46). High rates of private health care use have been linked to the underfunding and poor performance of the public health sector (44), and public perceptions that public services are of poor quality (47).

Interventions delivered through the public health system have little chance of reducing the treatment gap in a context where the majority of health care consultations take place elsewhere. The current landscape of the Indian health system is not reflected in the Global Mental Health literature, where traditional services are often discussed (48–54), but private providers are rarely mentioned, despite evidence that they frequently dispense psychotropic medications in India (55). The MANAS trial in Goa demonstrated the feasibility of training and supervising private providers to strengthen their ability to detect and treat depressive disorders (56); this strategy should be evaluated in other regions of India.

Use of traditional services

We also found that only a small proportion of treatment sought by people with probable depression was in the traditional sector. The report of the 2015-16 National Mental Health Survey of India posits preference for traditional services as a major barrier to the use of formal treatment (5), based on qualitative interviews with health professionals and community leaders, but presents no quantitative data on service use. Common mental disorders were not distinguished from severe mental illness in these interviews, so it is possible that the difference between our results and the perceptions of these stakeholders arose because traditional providers play an important role in treating people with psychotic disorders but not depression. Our estimates are backed up by a recent national survey showing that the use of traditional healers is low relative to the use of allopathic care for all health conditions, even in rural areas (57). This suggests that engaging with or influencing the use of traditional services should not be a major policy focus in improving care for depression in this context.

Detecting depression: Symptomatology and help-seeking

In terms of improving detection of depression in health services, health workers should be aware that tiredness or lack of energy is the most common symptom reported by depressed people in this population, followed by depressed mood or hopelessness. Those experiencing lack of energy are more likely to present to health services with depression symptoms than those with depressed mood, potentially because the former symptom is seen as a more legitimate medical complaint than emotional symptoms (58). Future research should test the predictive value of brief questions using local idioms of distress, as in recent research in Nepal (59), to find the most efficient way of detecting depression among primary care attendees who present with somatic symptoms, during short consultations (60).

Who seeks health care for depression symptoms?

An important unanswered question remains about why people with probable depression do not seek health care for depression symptoms specifically. Our findings on factors associated with treatment-seeking for depression symptoms should be interpreted with caution, since these were descriptive rather than hypothesis-testing: The results show which groups seek treatment rather than establishing causal relationships between these factors. However, some intriguing hypotheses were generated that deserve further investigation, to avoid wasting resources on ineffective strategies to reduce the treatment gap.

Firstly, we found no evidence that those with lower levels of self-stigma, exposure to mental health communications, or indicators of higher mental health literacy were more likely to seek treatment for depression. This contrasts with the conclusions of previous Indian studies, which have implicated these factors as barriers to treatment-seeking (13, 61). This may be because in the current study research workers referred to specific symptoms of depression, rather than to mental illness or psychiatric treatment, and some evidence suggests that symptoms of common mental disorders are not associated with mental illness in India (41). Service planners should exercise caution before investing resources in anti-stigma or awareness campaigns without further evidence of their effectiveness in stimulating treatment-seeking for depression (62).

Secondly, while a majority of participants felt that cost and distance barriers are important, in line with previous research (13, 61, 63), those who reported these barriers were no less likely to seek treatment for depression. We also found little evidence to support differences in treatment-seeking by socio-economic status. While structural factors may operate as barriers to general health care use, these findings suggest that lack of treatment-seeking for depression symptoms is not linked to economic and practical constraints, but rather to the nature and severity of symptoms. This requires further investigation, both to explicitly test these hypotheses using quantitative data and to qualitatively explore why enabling factors are not associated with treatment-seeking.

Thirdly, the evidence did not support an association between treatment-seeking and disability or perceived need for health care, which is at odds with international evidence (7). In light of the high rates of general health service use, this may be because people with depression consider themselves to have other health problems, and attribute their disability and associated need for health care to these non-depression symptoms. Future research should assess the overall health needs of adults with depression and investigate the effect of comorbid conditions on help-seeking behaviour.

Intriguingly, the gender differences often reported in studies from high-income countries were not replicated in this setting, and we found the opposite association between marital status and treatment-seeking that which is typically reported elsewhere (6, 7, 64). This demonstrates the importance of local data in identifying vulnerable groups for service planning, and provides suggestive evidence that processes believed to inhibit treatment-seeking in other cultural contexts – such as masculine ideals of self-sufficiency (65, 66) – may not apply in the same way to Indian populations.

Finally, participants who reported limited availability or irregular supply of medications were counter-intuitively more likely to seek treatment for depression than those who did not report these barriers. This

may result from retrospective measurement of these factors, since negative experiences of health care affect attitudes towards services (67). Longitudinal studies are needed to establish causal relationships between attitudes to services and help-seeking behaviour, and test the impact of negative experiences of health care on subsequent attitudes and treatment-seeking behaviour.

Future research should test the hypotheses generated here while controlling for confounding factors, and investigate factors for which data were not available including contextual influences such as social norms. Qualitative research is important to identify factors that the community perceives to be important, and to better understand why so few adults with probable depression consider treatment to be necessary for these symptoms specifically.

Strengths and limitations

To our knowledge, this study is the most comprehensive population-based study to explore patterns of treatment-seeking for depression in India. The current study used a large, representative community-based sample, to show which groups should be targeted in order to reduce the treatment gap for depression. Given the dearth of research on this topic from LMIC, the current findings may provide useful insights for service planning and policy, and generate hypotheses about barriers to treatment-seeking for further testing.

Since this was a secondary analysis of data collected for another primary purpose, however, we were limited by the measures used. More detailed, mental health-specific measures of barriers to care exist that were not employed due to interview length considerations, such as the Barriers to Access to Care Evaluation scale (BACE) (68). This limits the extent to which our results can be compared to recent studies from other settings (e.g. (69)), which could help to distinguish context-specific from more universal barriers. The measures used to indicate economic status are also imperfect proxies, meaning that we cannot be sure from our findings that poverty does not inhibit treatment-seeking for depression, despite the lack of association found here. Equally, the sample size was determined with reference to the primary aim of the parent study, and as such some of the current analyses may have been under-powered to detect an association, particularly for rare characteristics such as unemployment and suicidal thoughts.

Participants' mental health status was determined using a screening tool, not full diagnostic interviews, so the sample is likely to include some false positives, especially given the low positive predictive value of the PHQ-9 reported in Goa (27). Furthermore, since these data were generated through a cross-sectional survey, symptom severity, level of disability and attitudes towards health services were measured only at the time of the interview despite being subject to change over time, whereas treatment-seeking was measured retrospectively over the past 12 months.

There is also the potential for non-response bias, since only 62.5% of selected adults were located at baseline, and 76.2% at endline, due to death or migration. If those who were not located differ

systematically from those who were, this would result in biased estimates. We have no data on those who were not located, although our sample characteristics are generally comparable to the most recent census data (14).

Finally, self-reported data are always potentially open to social desirability bias, especially when using face-to-face interviews, and it is possible that this led to under-reporting of traditional service use. However, our estimates are in line with a recent national survey showing that the use of traditional healers is low relative to the use of allopathic care for both minor and major morbidity, even in rural areas (57).

Conclusion

Although most participants had not sought help specifically for depression symptoms, almost two thirds reported recent contact with health services, most frequently in the private sector. Private health care providers are an important group to engage in efforts to improve detection and treatment of depression in this area, and should be included in programmes of training and supervision to reduce the treatment gap for depression. Future research should investigate why adults with probable depression seek help for other symptoms rather than for depression, and replicate the current findings on factors associated with treatment-seeking for depression symptoms.

References

1. Ferrari AJ, Charlson FJ, Norman RE, Patten SB, Freedman G, Murray CJ, et al. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS medicine*. 2013;10(11):e1001547.
2. Kessler RC, Birnbaum HG, Shahly V, Bromet E, Hwang I, McLaughlin KA, et al. Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO World Mental Health Survey Initiative. *Depression and anxiety*. 2010;27(4):351-64.
3. Thornicroft G, Chatterji S, Evans-Lacko S, Gruber M, Sampson N, Aguilar-Gaxiola S, et al. Undertreatment of people with major depressive disorder in 21 countries. *The British Journal of Psychiatry*. 2017;210(2):119-24.
4. WHO, WONCA. Integrating mental health into primary care: A global perspective. World Health Organization and World Organization of Family Doctors 2008.
5. Gururaj G, Varghese M, Benegal V, Rao G, Pathak K, Singh L, et al. National Mental Health Survey of India, 2015-16: Summary. Bengaluru: National Institute of Mental Health and Neuro Sciences, NIMHANS Publication. 2016(128).
6. Magaard JL, Seeralan T, Schulz H, Brütt AL. Factors associated with help-seeking behaviour among individuals with major depression: A systematic review. *PloS one*. 2017;12(5):e0176730.
7. Roberts T, Miguel Esponda G, Krupchanka D, Shidhaye R, Patel V, Rathod S. Factors associated with health service utilisation for common mental disorders: a systematic review. *BMC Psychiatry*. 2018;18(1):262.

8. Patel V, Xiao S, Chen H, Hanna F, Jotheeswaran AT, Luo D, et al. The magnitude of and health system responses to the mental health treatment gap in adults in India and China. *The Lancet*. 2016.
9. De Silva MJ, Lee L, Fuhr DC, Rathod S, Chisholm D, Schellenberg J, et al. Estimating the coverage of mental health programmes: a systematic review. *Int J Epidemiol*. 2014;43(2):341-53.
10. Raban MZ, Dandona R, Kumar GA, Dandona L. Inequitable coverage of non-communicable diseases and injury interventions in India. *National medical journal of India*. 2010;23(5):267.
11. Mathias K, Goicolea I, Kermode M, Singh L, Shidhaye R, San Sebastian M. Cross-sectional study of depression and help-seeking in Uttarakhand, North India. *BMJ open*. 2015;5(11):e008992.
12. Soni A, Fahey N, Byatt N, Prabhakaran A, Simas TAM, Vankar J, et al. Association of common mental disorder symptoms with health and healthcare factors among women in rural western India: results of a cross-sectional survey. *BMJ open*. 2016;6(7):e010834.
13. James S, Chisholm D, Murthy RS, Kumar KK, Sekar K, Saeed K, et al. Demand for, access to and use of community mental health care: lessons from a demonstration project in India and Pakistan. *International Journal of Social Psychiatry*. 2002;48(3):163-76.
14. Office of the Registrar General and Census Commissioner I. Census 2011 2011 [Available from: <http://www.census2011.co.in/census/district/312-sehore.html>].
15. Horwitz AV, Wakefield JC. The loss of sadness: How psychiatry transformed normal sorrow into depressive disorder: Oxford University Press; 2007.
16. International Institute for Population Sciences. 2009. District Fact Sheet: Sehore, Madhya Pradesh. National Family Health Survey - 4, 2015-16. Available from: http://rchiips.org/NFHS/FCTS/MP/MP_FactSheet_445_Sehore.pdf [Accessed 24/09/18] [
17. Rathod SD, Nadkarni A, Bhana A, Shidhaye R. Epidemiological features of alcohol use in rural India: a population-based cross-sectional study. *BMJ open*. 2015;5(12):e009802.
18. Lund C, Tomlinson M, De Silva M, Fekadu A, Shidhaye R, Jordans M, et al. PRIME: a programme to reduce the treatment gap for mental disorders in five low- and middle-income countries. *PLoS medicine*. 2012;9(12):e1001359.
19. Shidhaye R, Shrivastava S, Murhar V, Samudre S, Ahuja S, Ramaswamy R, et al. Development and piloting of a plan for integrating mental health in primary care in Sehore district, Madhya Pradesh, India. *The British journal of psychiatry : the journal of mental science*. 2016;208 Suppl 56:s13-20.
20. Shidhaye R, Raja A, Shrivastava S, Murhar V, Ramaswamy R, Patel V. Challenges for Transformation: A Situational Analysis of Mental Health Care Services in Sehore District, Madhya Pradesh. *Community Ment Health J*. 2015;51(8):903-12.
21. Rathod SD, De Silva MJ, Ssebunnya J, Breuer E, Murhar V, Luitel NP, et al. Treatment contact coverage for probable depressive and probable alcohol use disorders in four low-and middle-income country districts: the PRIME cross-sectional community surveys. *PLoS One*. 2016;11(9):e0162038.
22. De Silva MJ, Rathod SD, Hanlon C, Breuer E, Chisholm D, Fekadu A, et al. Evaluation of district mental healthcare plans: the PRIME consortium methodology. *The British journal of psychiatry : the journal of mental science*. 2016;208 Suppl 56:s63-70.

23. Shidhaye R, Baron E, Murhar V, Rathod S, Khan A, Singh A, et al. Community, facility and individual level impact of integrating mental health screening and treatment into the primary healthcare system in Sehore district, Madhya Pradesh, India. *BMJ global health*. 2019;4(3):e001344.
24. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*. 2001;16(9):606-13.
25. Moriarty AS, Gilbody S, McMillan D, Manea L. Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): a meta-analysis. *General hospital psychiatry*. 2015;37(6):567-76.
26. Manea L, Gilbody S, McMillan D. Optimal cut-off score for diagnosing depression with the Patient Health Questionnaire (PHQ-9): a meta-analysis. *Canadian Medical Association Journal*. 2012;184(3):E191-E6.
27. Patel V, Araya R, Chowdhary N, King M, Kirkwood B, Nayak S, et al. Detecting common mental disorders in primary care in India: a comparison of five screening questionnaires. *Psychological medicine*. 2008;38(2):221-8.
28. Ganguly S, Samanta M, Roy P, Chatterjee S, Kaplan DW, Basu B. Patient health questionnaire-9 as an effective tool for screening of depression among Indian adolescents. *Journal of Adolescent Health*. 2013;52(5):546-51.
29. Kowal P, Chatterji S, Naidoo N, Biritwum R, Fan W, Lopez Ridaura R, et al. Data resource profile: the World Health Organization Study on global AGEing and adult health (SAGE). *International journal of epidemiology*. 2012;41(6):1639-49.
30. Andersen RM. Revisiting the behavioral model and access to medical care: does it matter? *Journal of health and social behavior*. 1995:1-10.
31. Andersen R. A behavioral model of families' use of health services. *A behavioral model of families' use of health services*. 1968(25).
32. Boyd JE, Adler EP, Otilingam PG, Peters T. Internalized Stigma of Mental Illness (ISMI) scale: a multinational review. *Comprehensive Psychiatry*. 2014;55(1):221-31.
33. Üstün TB, Chatterji S, Kostanjsek N, Rehm J, Kennedy C, Epping-Jordan J, et al. Developing the World Health Organization Disability Assessment Schedule 2.0. 2010.
34. Pal HR, Jena R, Yadav D. Validation of the Alcohol Use Disorders Identification Test (AUDIT) in urban community outreach and de-addiction center samples in north India. *Journal of Studies on Alcohol*. 2004;65(6):794-800.
35. Carey KB, Carey MP, Chandra PS. Psychometric evaluation of the alcohol use disorders identification test and short drug abuse screening test with psychiatric patients in India. *The Journal of clinical psychiatry*. 2003;64(7):767.
36. de Meneses-Gaya C, Zuardi AW, Loureiro SR, Crippa JAS. Alcohol Use Disorders Identification Test (AUDIT): An updated systematic review of psychometric properties. *Psychology & Neuroscience*. 2009;2(1):83.

37. Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG, Organization WH. AUDIT: The alcohol use disorders identification test: Guidelines for use in primary health care. 2001.
38. Kessler RC, Üstün TB. The world mental health (WMH) survey initiative version of the world health organization (WHO) composite international diagnostic interview (CIDI). *International journal of methods in psychiatric research*. 2004;13(2):93-121.
39. StataCorp. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP; 2015.
40. Rathod SD, Roberts T, Medhin G, Murhar V, Samudre S, Luitel NP, et al. Detection and treatment initiation for depression and alcohol use disorders: facility-based cross-sectional studies in five low- and middle-income country districts. Manuscript submitted for publication. 2018.
41. Andrew G, Cohen A, Salgaonkar S, Patel V. The explanatory models of depression and anxiety in primary care: a qualitative study from India. *BMC research notes*. 2012;5(1):499.
42. Patel V, Pereira J, Countinho L, Fernandes R, Fernandes J, Mann A. Poverty, psychological disorder and disability in primary care attenders in Goa, India. *The British Journal of Psychiatry*. 1998;172(6):533-6.
43. De Costa A, Diwan V. 'Where is the public health sector?': Public and private sector healthcare provision in Madhya Pradesh, India. *Health Policy*. 2007;84(2):269-76.
44. De Costa A, Johannson E. By 'default or design'? The expansion of the private health care sector in Madhya Pradesh, India. *Health policy*. 2011;103(2):283-9.
45. Selvaraj S, Karan AK. Deepening health insecurity in India: evidence from national sample surveys since 1980s. *Economic and Political Weekly*. 2009:55-60.
46. Kumar AS, Chen LC, Choudhury M, Ganju S, Mahajan V, Sinha A, et al. Financing health care for all: challenges and opportunities. *The Lancet*. 2011;377(9766):668-79.
47. Levesque J-F, Haddad S, Narayana D, Fournier P, editors. Access to health care in urban Kerala: emerging inequalities across the public-private divide. Presentation made at Global Forum for Health Research, Forum; 2005: Citeseer.
48. Gureje O, Nortje G, Makanjuola V, Oladeji BD, Seedat S, Jenkins R. The role of global traditional and complementary systems of medicine in the treatment of mental health disorders. *The Lancet Psychiatry*. 2015;2(2):168-77.
49. Incayawar M. Future partnerships in global mental health: Foreseeing the encounter of psychiatrists and traditional healers. *Psychiatrists and traditional healers: Unwitting partners in global mental health*. 2009:251-60.
50. Patel V, Saxena S. Transforming lives, enhancing communities—innovations in global mental health. *New England Journal of Medicine*. 2014;370(6):498-501.
51. Patel V. Traditional healers for mental health care in Africa. *Global health action*. 2011;4.
52. Raguram R, Venkateswaran A, Ramakrishna J, Weiss MG. Traditional community resources for mental health: a report of temple healing from India. *Bmj*. 2002;325(7354):38-40.

53. Campbell-Hall V, Petersen I, Bhana A, Mjadu S, Hosegood V, Flisher AJ, et al. Collaboration between traditional practitioners and primary health care staff in South Africa: developing a workable partnership for community mental health services. *Transcultural psychiatry*. 2010;47(4):610-28.
54. Quack J. Ignorance and utilization: mental health care outside the purview of the Indian state. *Anthropology & Medicine*. 2012;19(3):277-90.
55. Ecks S. Commentary: Ethnographic critiques of global mental health. *Transcultural psychiatry*. 2016;53(6):804-8.
56. Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, et al. Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *The Lancet*. 2010;376(9758):2086-95.
57. Singh A, Madhavan H. Traditional vs. non-traditional healing for minor and major morbidities in India: uses, cost and quality comparisons. *Tropical Medicine & International Health*. 2015;20(9):1223-38.
58. Patel V, Prince M. Ageing and mental health in a developing country: who cares? Qualitative studies from Goa, India. *Psychological medicine*. 2001;31(1):29-38.
59. Kohrt BA, Luitel NP, Acharya P, Jordans MJ. Detection of depression in low resource settings: validation of the Patient Health Questionnaire (PHQ-9) and cultural concepts of distress in Nepal. *BMC psychiatry*. 2016;16(1):58.
60. Irving G, Neves AL, Dambha-Miller H, Oishi A, Tagashira H, Verho A, et al. International variations in primary care physician consultation time: a systematic review of 67 countries. *BMJ open*. 2017;7(10):e017902.
61. Tewari A, Kallakuri S, Devarapalli S, Jha V, Patel A, Maulik PK. Process evaluation of the systematic medical appraisal, referral and treatment (SMART) mental health project in rural India. *BMC psychiatry*. 2017;17(1):385.
62. Gulliver A, Griffiths KM, Christensen H, Brewer JL. A systematic review of help-seeking interventions for depression, anxiety and general psychological distress. *BMC psychiatry*. 2012;12(1):81.
63. Maulik PK, Tewari A, Devarapalli S, Kallakuri S, Patel A. The Systematic Medical Appraisal, Referral and Treatment (SMART) Mental Health Project: development and testing of electronic decision support system and formative research to understand perceptions about mental health in rural India. *PloS one*. 2016;11(10):e0164404.
64. Demyttenaere K, Bruffaerts R, Posada-Villa J, Gasquet I, Kovess V, Lepine J, et al. Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *Jama*. 2004;291(21):2581-90.
65. Galdas PM, Cheater F, Marshall P. Men and health help-seeking behaviour: literature review. *Journal of advanced nursing*. 2005;49(6):616-23.
66. Addis ME, Mahalik JR. Men, masculinity, and the contexts of help seeking. *American psychologist*. 2003;58(1):5.

67. Naidu A. Factors affecting patient satisfaction and healthcare quality. International journal of health care quality assurance. 2009;22(4):366-81.
68. Clement S, Brohan E, Jeffery D, Henderson C, Hatch SL, Thornicroft G. Development and psychometric properties the Barriers to Access to Care Evaluation scale (BACE) related to people with mental ill health. BMC psychiatry. 2012;12(1):36.
69. Luitel NP, Jordans MJ, Kohrt BA, Rathod SD, Komproe IH. Treatment gap and barriers for mental health care: A cross-sectional community survey in Nepal. PloS one. 2017;12(8):e0183223.

Abbreviations

AUDIT	Alcohol Use Disorders Identification Test
BACE	Barriers to Access to Care Evaluation scale
CIDI	Composite International Diagnostic Interview
ISMI	Internalized Stigma of Mental Illness scale
LMIC	Low- and middle-income countries
MHCP	Mental Health Care Plan
PHQ-9	Patient Health Questionnaire – 9 item version
PRIME	Programme for Improving Mental Health Care
SAGE	Study on Global Ageing and Adult Health
WHO	World Health Organization
WHODAS	World Health Organization Disability Assessment Schedule

Declarations

Ethics approval and consent to participate

The study reported here was approved by the appropriate ethics committees and performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments, as well as with all relevant laws and regulations. Ethical approval was provided by the World Health Organization Research Ethics Review Committee (Geneva, Switzerland), the Sangath Institutional Review Board (Goa, India), and the London School of Hygiene & Tropical Medicine Observational Ethics Committee (London, United Kingdom) (ref: 10439).

Informed consent was indicated with either a signature or – in cases of low literacy – a thumbprint. This procedure was approved by all of the ethics committees above.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analysed during the current study are available from the PRIME consortium on reasonable request. See PRIME website for details and to request access: <http://www.prime.uct.ac.za/>. Queries regarding access to the data can be directed to the PRIME project manager, Erica Breuer: erica.breuer@uct.ac.za.

The PRIME community survey questionnaire that was used here has been previously published elsewhere (see e.g. Rathod et al, 2016, (21)).

Competing interests

Rahul Shidhaye is a member of the editorial board of BMC Health Services Research. The authors declare that other than this they have no competing interests.

Funding

This study used data from the PRogramme for Improving Mental health care (PRIME) (<http://www.prime.uct.ac.za>). This work was supported by the United Kingdom Department for International Development (Grant number 201446).

The first author was funded by a PhD studentship from the Bloomsbury Colleges (University of London) and subsequently by an ESRC Post Doctoral Fellowship (ES/T007125/1). The funders had no role in the design or execution of this study, nor in the decision to submit the results for publication.

Authors' contributions

TR was the lead author, and was responsible for the study design, data analysis, and the writing of the manuscript. RS and VP advised on the design of the study and the presentation of the results, and offered feedback on the draft manuscript. SR provided guidance throughout the process of study design, analysis, and writing, and also gave detailed comments and feedback on earlier drafts. RS, VP and SR were all involved in the design of the parent study and the collection of the data used in the current analyses. All authors have read and approved the manuscript.

Acknowledgements

This study used data from the PRogramme for Improving Mental health care (PRIME), which is a Research Programme Consortium led by the Centre for Public Mental Health at the University of Cape Town (South Africa), and funded by the UK government's Department for International Development (DFID). Note that the views expressed in this article do not necessarily reflect the UK Government's official policies. We wish to thank the PRIME field team in Bhopal, India, for their work in collecting the study data. Partners and collaborators in the consortium include Sangath, the Public Health Foundation of India and Madhya Pradesh State Ministry of Health (India), Addis Ababa University and Ministry of Health (Ethiopia), Health Net TPO and Ministry of Health (Nepal), University of Kwazulu-Natal, Human Sciences Research Council, Perinatal Mental Health Project and Department of Health (South Africa), Makerere University and Ministry of Health (Uganda), BasicNeeds, Centre for Global Mental Health (London School of Hygiene & Tropical Medicine and Kings Health Partners, UK) and the World Health Organisation (WHO).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementarymaterialBMCHSR.docx](#)