Intracranial aneurysms associated with middle cerebral artery anomalies: the clinical characteristics and treatment considerations: a systematic review

Kun Hou

Jilin University First Hospital

Guichen Li

Jilin University First Hospital

Hongping Liu

Jilin University First Hospital

Kan Xu

Jilin University First Hospital

Jinlu Yu (✉ jlyu@jlu.edu.cn)

Jilin University First Hospital <https://orcid.org/0000-0003-2329-7946>

Research article

Keywords: middle cerebral artery anomalies, accessory middle cerebral artery, duplicate middle cerebral artery, middle cerebral artery fenestration, intracranial aneurysm

Posted Date: December 13th, 2019

DOI: <https://doi.org/10.21203/rs.2.18957/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Frontiers in Neurology on October 27th, 2020. See the published version at <https://doi.org/10.3389/fneur.2020.564797>.

Abstract

Background: As a result of its low occurrence, most of the studies on intracranial aneurysms associated with MCA anomalies were presented as case reports or small case series. In this study, a systematic review on this specific entity was conducted.

Methods: A PubMed search of the published studies was performed on April 6th, 2019 for patients who had intracranial aneurysms associated with MCA anomalies. The languages included in this study were English, Chinese, and Japanese.

Results: Finally, 58 articles reporting of 67 patients including 1 case in our center were included. The identified patients (37 females, 55.2%) aged from 4 to 81 (49.85 ± 15.22) years old. Fifty (50/67, 74.6%) patients presented with hemorrhagic stroke either because of rupture of the aneurysms associated with MCA anomalies or other sources. Sixty-three aneurysms (63/67, 94.0%) were saccular, 3 (4.5%) were dissecting or fusiform, and 1 (1.5%) was pseudoaneurysm. Thirty-two (32/65, 49.2%) patients had other concurrent cerebrovascular anomalies. Fifty-six (83.6%) patients underwent open surgeries, 8 (11.9%) patients underwent endovascular treatment, and 3 (4.5%) patients were conservatively managed. Fifty-six (56/61, 91.8%) patients achieved a good recovery.

Conclusions: The pathophysiological genesis of MCA anomalies associated aneurysms is still obscure to us. The inflicted patients tend to have concurrent other cerebrovascular anomalies, which denotes that congenital defect in cerebrovascular development might play a role in this process. Open surgery is, hitherto, the mainstay of treatment for this specific entity. Most of the affected patients could experience a good recovery.

Background

Middle cerebral artery (MCA) is the largest and most important artery of the anterior circulation. Compared to its counterparts of the posterior circulation, MCA has a lower incidence of vascular anomalies [1]. Generally speaking, MCA anomalies includes accessory MCA (ac-MCA), duplicate MCA (d-MCA), d-MCA origin, MCA fenestration, and twig-like MCA. In rare circumstances, the MCA anomalies can be associated with intracranial aneurysms [2–4]. As a result of its low occurrence, most of the studies on intracranial aneurysms associated with MCA anomalies were presented as case reports or small case series. Hence, large-scale investigation on this rare entity in a single center is unrealistic. In this study, we would like to perform a systematic review on this specific entity to further elucidate its demographic, clinical, therapeutic, and prognostic characteristics.

Methods

A PubMed search of the published studies was performed on April 6th, 2019 for patients who had intracranial aneurysms associated with MCA anomalies. The languages included in this study were English, Chinese, and Japanese. The algorithm used in this search was (((((((accessory middle cerebral artery[Title/Abstract]) OR duplicate middle cerebral artery[Title/Abstract]) OR duplicated middle cerebral artery[Title/Abstract]) OR duplicate middle cerebral artery origin[Title/Abstract]) OR duplicated middle cerebral artery origin[Title/Abstract]) OR fenestration of middle cerebral artery[Title/Abstract]) OR fenestrated middle cerebral artery[Title/Abstract])AND aneurysm[Title/Abstract]. Only articles of which the full text or enough information could be obtained were included in this study. Reference lists of the identified articles were also manually searched for additional studies. Glasgow Outcome Scale was used for the outcome assessment. A Glasgow Outcome Scale score ≥ 4 was defined as good recovery. An aneurysm < 10 mm was defined as small aneurysm.

Definition of intracranial aneurysm associated with MCA anomalies

Intracranial aneurysms located at the beginning or on the trunk of the abnormal MCAs were considered as in association with MCA anomalies. Aneurysms having no direct anatomical neighborhood with the MCA anomalies were excluded in the final analysis.

Location of ac-MCA and the associated aneurysm

Based on their sites of origin along the anterior cerebral artery (ACA), the ac-MCA were divided into 3 types: 1) originating from A1 segment of ACA, 2) originating from the anterior communicating artery (AComA) or A1-A2 junction, 3) originating from A2 segment. The locations of aneurysms were at the beginning or on the trunk of ac-MCA.

Location of d-MCA associated aneurysm

The locations of aneurysms were at the beginning or on the trunk of d-MCA.

Location of MCA fenestration associated aneurysm

The locations of aneurysms were proximal to fenestration, in fenestration, or distal to fenestration.

D-MCA origin aneurysm

The locations of aneurysms were at the beginning of any branch of duplicate origins or on the common trunk.

Results

General information

The PubMed search identified 113 records. Fifty-nine records were excluded based on titles and abstracts screening. After assessing the full text of the remaining 54 articles, 5 were further excluded. A manual searching of the reference lists of the remaining 49 articles was performed, which yielded 9 additional articles. Finally, 58 articles reporting of 67 patients including 1 case in our center were included for the analysis (Fig. 1).

The identified patients (37 females, 55.2%) aged from 4 to 81 (49.85 ± 15.22) years old. Fifty (50/67, 74.6%) patients presented with hemorrhagic stroke either because of rupture of the aneurysms associated with MCA anomalies or other sources. The MCA anomalies associated aneurysms were located on the left side in 32 (32/66, 48.5%) patients. Sixty-three aneurysms (63/67, 94.0%) were saccular, 3 (4.5%) were dissecting or fusiform, and 1 (1.5%) was pseudoaneurysm. Thirty-two (32/65, 49.2%) patients had other concurrent cerebrovascular anomalies in addition to MCA anomalies associated aneurysms. Fifty-six (83.6%) patients underwent open surgeries, 8 (11.9%) patients underwent endovascular treatment, and 3 (4.5%) patients were conservatively managed. Ten (10/62, 16.1%) patients experienced procedure-related complications. Fifty-six (56/61, 91.8%) patients achieved a good recovery.

Accessory MCA aneurysm

General information

The PubMed search identified 113 records. Fifty-nine records were excluded based on titles and abstracts screening. After assessing the full text of the remaining 54 articles, 5 were further excluded. A manual searching of the reference lists of the remaining 49 articles was performed, which yielded 9 additional articles. Finally, 58 articles reporting of 67 patients including 1 case in our center were included for the analysis (Figure 1).

The identified patients (37 females, 55.2%) aged from 4 to 81 (49.85 ± 15.22) years old. Fifty (50/67, 74.6%) patients presented with hemorrhagic stroke either because of rupture of the aneurysms associated with MCA anomalies or other sources. The MCA anomalies associated aneurysms were located on the left side in 32 (32/66, 48.5%) patients. Sixty-three aneurysms (63/67, 94.0%) were saccular, 3 (4.5%) were dissecting or fusiform, and 1 (1.5%) was pseudoaneurysm. Thirty-two (32/65, 49.2%) patients had other concurrent cerebrovascular anomalies in addition to MCA anomalies associated aneurysms. Fifty-six (83.6%) patients underwent open surgeries, 8 (11.9%) patients underwent endovascular treatment, and 3 (4.5%) patients were conservatively managed. Ten (10/62, 16.1%) patients experienced procedure-related complications. Fifty-six (56/61, 91.8%) patients achieved a good recovery.

Accessory MCA aneurysm

Nineteen studies reporting of 20 patients including 1 case in our center were identified (Table 1) [4-22]. The patients aged from 4 to 73 (48.65 ± 16.70) years old, with a male to female ratio of 1:1. Eighteen (18/20, 90%) patients presented with intracranial bleeding from ac-MCA aneurysms or other sources. The sizes of aneurysms were below and above 10 mm in 18 (90%) and 2 (10%) patients, respectively. Seventeen (85%) aneurysms were saccular, 2 (10%) were dissecting, 1 (5%) was pseudoaneurysm. The left to right ratio of aneurysm allocation was 1:1. The locations of ac-MCAs were A1, A1-A2 junction, and A2 in 16 (16/19, 84.2%), 2 (2/19, 10.5%), and 1 (1/19, 5.3%) patient, respectively. Of the 20 aneurysms, 14 (70%) were located at the origin of ac-MCA, 6 (30%) were on the trunk. Nine (9/18, 50%) patients had other concurrent cerebrovascular anomalies. With respect to the treatment, 14 (70%) patients underwent microsurgical clipping of the aneurysms, 4 (20%) (3 coiling, 1 glue embolization) underwent endovascular treatment, 1 (5%) underwent resection of the pseudoaneurysm and distal ac-MCA, and 1 (5%) underwent aneurysm wrapping. Procedure-related complications occurred in 1 (1/19, 5.3%) patient. Eighteen (18/20, 90%) patients experienced good recovery.

Duplicate MCA aneurysm

Twenty-seven studies reporting of 34 patients were finally included (Table 2) [3, 7, 23-47]. The patients aged from 20 to 76 (50.79 ± 13.72) years old, with a male to female ratio of 0.62:1 (13:21). Twenty (20/34, 58.8%) patients presented with intracranial bleeding from d-MCA aneurysms or other sources. The sizes of aneurysms were below and above 10 mm in 33 and 1 patients, respectively. All of the aneurysms were saccular except a fusiform one. The left to right ratio of aneurysm allocation was 1.2:1 (18:15). Of the 34 aneurysms, 32 (94.1%) were located at the origin of d-MCA, 2 were on the trunk. Nine (9/18, 50%) patients had other concurrent cerebrovascular anomalies. Of the 34 patients, 18 (52.9%) have concurrent cerebrovascular anomalies. Twenty-seven (79.4%) patients underwent microsurgical clipping of the aneurysms, 3 (8.8%) patients underwent endovascular coiling, 1 (2.9%) underwent trapping of the aneurysm and simultaneous superficial temporal artery-d-MCA anastomosis, and 3 (8.8%) patients were conservatively followed up. Procedure-related complications occurred in 7 (22.6%) patients. Twenty-six (89.7%, 26/29) patients experienced good recovery.

MCA fenestration aneurysm

Twelve studies reporting of 12 patients were identified, aging from 14 to 81 (49.3 ± 13.1) years old (Table 3) [2, 48-58]. The male to female ratio was 1:1. All of the patients were admitted for intracranial bleeding. All of the aneurysms were smaller than 10 mm except for 1 the size which could not be determined. The morphology of all of the aneurysms were saccular. Nine (9/12, 75%) of the aneurysms were located at the right side. All of the fenestrations were located on M1 segment of the MCAs. The aneurysms were located proximal to, in, and distal to the fenestration in 5 (41.7%), 4 (33.3%), and 3 (25%) patients, respectively. Concurrent cerebrovascular anomalies were identified in 5 (41.7%) patients. With respect to the treatment, 10 patients underwent surgical clipping, 1 underwent aneurysm wrapping, and 1 underwent coiling. Two (2/11, 18.2%) patients experienced procedure-related complications. All of the patients experienced good recovery except for 1 patient without mention of outcome.

Duplicate MCA origin aneurysm

d-MCA origin aneurysm was only identified in a 49-year man incidentally, who was admitted for vertigo [59]. No other cerebrovascular anomaly was reported. The saccular unruptured d-MCA origin aneurysm was microsurgically clipped. The postoperative course was uneventful and no neurological deficit was reported.

Illustrative case

A 59-year old man was admitted for sudden onset of headache 2 days ago. He was a smoker and denied history of any chronic diseases. He was alert on admission. Physical examination was unremarkable except for neck rigidity. Head computed tomography (CT) revealed subarachnoid hemorrhage of modified Fisher grade 2 (Figure 2 A-B). Further CT angiography showed two ac-MCAs originated from the A1 segment of bilateral ACAs (Figure 2 C). A saccular aneurysm was also noted at the origin of the left ac-MCA (Figure 2 C-D). No other cerebrovascular anomaly was noticed. After discussion between the neurosurgical and neuro-interventional members and sufficient negotiation with the patient's legal relatives, endovascular coiling of the aneurysm was planned.

Preprocedural digital subtraction angiography also confirmed the findings on CT angiography (Figure 3 A-B). An Echelon-10 (Medtronic, Irvine, CA) microcatheter was advanced into the left ACA directed by a 0.010-in guidewire. The tip of the microcatheter was introduced into the aneurysm. The aneurysm was satisfactorily coiled using 3 detachable coils with preservation of the distal ACA and ac-MCA (Figure 3 C-D). He experienced an uneventful postprocedural recovery and was discharged the next day without neurological deficit. Follow-up CT angiography 1 year later revealed no recurrence of the aneurysm.

Discussion

According to Padgett's description, at 34–36 days of the embryonal stage (12–14 mm), multiple plexiform arterial twigs develop just distally to the anterior choroidal artery [60]. The plexiform arterial twigs would evolve into the MCA and lateral striate arteries through subsequent fusion and regression. Hypothetically, failure of this process can lead to diverse variations of the MCA (e.g. ac-MCA, d-MCA, MCA fenestration, d-MCA origin, and twig-like MCA) [1]. Generally speaking, the incidence of any MCA anomalies is very low [1, 61–63]. However, in even rarer circumstances, the MCA anomalies could be associated with intracranial aneurysms [2–4]. As a result of the low incidence of MCA anomalies, the reported cases of MCA anomalies associated intracranial aneurysms were all presented as case reports. Hence, the true incidence of MCA anomalies associated aneurysms in patients with MCA anomalies is still unknown. According to this study, 49.2% of the patients had other concurrent cerebrovascular anomalies in addition to MCA anomalies associated aneurysms, which implies that this specific subset of patients might have congenital defect in cerebrovascular development and be prone to cerebrovascular anomalies. Modern investigations using whole genome sequencing and epigenetics in this subset of patients may shed light on this issue.

According to Teal et al., anomalous arteries originating from the ACAs and coursing in parallel to or in close relationship with MCA were defined as ac-MCAs. And those arising from ICAs were considered d-MCAs [64]. The ac-MCA could be subdivided into 3 types based on the site of origin. Type Ⅰ denotes those originating from the A1 segment of ACA, type Ⅱ denotes those originating from the A1-A2 junction (including AComA), and type Ⅲ denotes those originating from the A2 segment. According to Kai et al., the d-MCA was further divided into 2 types (type A and type B) [34]. Those d-MCAs arising from the top of ICA bifurcation were defined as type A, and those between the anterior choroidal artery and ICA top were type B. However, an exceptional case had been reported by Tutar et al., of which the d-MCA originated from the petrous portion of ICA [65]. In another case, a d-MCA originated from the ICA about 10 mm proximal to the ICA bifurcation and an ipsilateral fetal-type posterior cerebral artery originated from the d-MCA [27]. In our systematic review of the literature, ac-MCAs originated from the A1 segment in 84.2% (16/19) of the patients with ac-MCA associated aneurysms. And the origin of d-MCA lied between the anterior choroidal artery and ICA bifurcation in all but 1 patient.

In contrast to the tendency for endovascular treatment for other intracranial aneurysms, 83.6% of the reported patients with MCA anomalies associated aneurysms underwent open surgeries, and only 11.9% of the patients underwent endovascular treatment. This phenomenon is still obscure to us. In our opinion, the reasons are multi-factorial. Firstly, as a result of the technical constraint, earlier cases had to undergo open surgeries. And then due to the unconventional locations and low incidence, medical practitioners are prone to the seemingly safer open surgical approaches. Thirdly, due to the specific characteristics of local vascular configuration, endovascular treatment might be more difficult. However, according to our study of the past reports, the outcome in the patients undergoing endovascular treatment was not inferior to those undergoing open surgeries. And most of the cases with endovascular treatment were reported recently [20, 39, 46, 47, 56].

Due to rarity of the studied issue, the data of this review was extracted from retrospective case reports or small case series. The results would be biased by many factors. Some anatomical, clinical, therapeutic, and prognostic details might be missed due to the different reporting customs. Statistical analysis is inappropriate for this kind of study. No comparative study between endovascular treatment and open surgery could be performed at present.

Conclusions

MCA anomalies have a low incidence of occurrence. Their associated intracranial aneurysms are even rarer than only sporadic reports were presented. The pathophysiological genesis of this specific entity is still obscure to us. The patients with MCA anomalies associated aneurysms tend to have concurrent other cerebrovascular anomalies, which denotes that congenital defect in cerebrovascular development might play a role in this process. Open surgery is, hitherto, the mainstay of treatment for this specific entity. Most of the affected patients could experience a good recovery.

Abbreviations

ACA, anterior cerebral artery

AComA, anterior communicating artery

ac-MCA, accessory middle cerebral artery

CT, computed tomography

d-MCA, duplicate middle cerebral artery

GOS, Glasgow Outcome Scale

ICA, internal carotid artery

MCA, middle cerebral artery

Declarations

Ethics approval

This study was approved by the institutional review board of The First Hospital of Jilin University, and the participant gave his/her informed consent before inclusion in the study.

Acknowledgements: None

Funding: This research received funding support from the Ninth Youth Scientific Research Funding of The First Hospital of Jilin University (jdyy92018035).

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Authors' Contributions

Conception and design: JY, KH. Acquisition of data: KH, GL. Analysis and interpretation of data: HL, KX. Drafting of the article: KH, GL. Critical revision of the article: JY, KX. All of the authors have read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Written informed consent was obtained from the patient for publication of this manuscript and any accompanying images. A copy of the written consent is available for review by the editor of this journal.

References

1. Uchiyama N. Anomalies of the Middle Cerebral Artery. *Neurologia medico-chirurgica*. 2017;57 6:261-6; doi: 10.2176/nmc.ra.2017-0043.
2. Xue X, Li Y, Yin Y, Yuan X, Xiao H, Wei L. Aneurysms Arising from the Fenestrated Middle Cerebral Artery. *The Journal of craniofacial surgery*. 2019; doi: 10.1097/scs.0000000000005357.
3. Mori K, Tamase A, Seki S, Iida Y, Kawabata Y, Nakano T, et al. Duplicated middle cerebral artery associated with aneurysm at M1/M2 bifurcation: a case report. *Journal of medical case reports*. 2018;12 1:283; doi: 10.1186/s13256-018-1824-7.
4. Nomura M, Tamase A, Kamide T, Mori K, Seki S, Iida Y. Accessory middle cerebral artery associated with an unruptured aneurysm at its origin. *Surgical neurology international*. 2015;6 Suppl 16:S421-3; doi: 10.4103/2152-7806.166179.
5. Waga S, Kojima T, Morooka Y, Sakakura M. Aneurysm of the accessory middle cerebral artery. *Surgical neurology*. 1977;8 5:359-60.
6. Handa J, Nakasu Y, Matsuda M, Kyoshima K. Aneurysms of the proximal anterior cerebral artery. *Surgical neurology*. 1984;22 5:486-90.
7. Fuwa I, Matsukado Y, Wada H. [Intracranial aneurysms associated with the accessory middle cerebral artery and duplication of the middle cerebral artery. Report of two cases]. *Neurologia medico-chirurgica*. 1984;24 3:207-11.
8. Miyazaki S, Ito K, Ishii S. Aneurysm at the origin of the accessory middle cerebral artery. *Surgical neurology*. 1984;22 3:292-4.
9. Kuwabara S, Naitoh H. Ruptured aneurysm at the origin of the accessory middle cerebral artery: case report. *Neurosurgery*. 1990;26 2:320-2.
10. Han DH, Gwak HS, Chung CK. Aneurysm at the origin of accessory middle cerebral artery associated with middle cerebral artery aplasia: case report. *Surgical neurology*. 1994;42 5:388-91.
11. Sugita S, Yuge T, Miyagi J, Fujimura N, Shigemori M. Giant aneurysm at the origin of the accessory middle cerebral artery. *Surgical neurology*. 1995;44 2:128-30.
12. Otawara Y, Suzuki M, Abe M, Tomizuka N, Ogawa A. Dissecting aneurysms of the anterior cerebral artery and accessory middle cerebral artery. Case report. *Neurosurgical review*. 1997;20 2:145-8.
13. Georgopoulos CE, Papanikolaou PG, Vlachos KI, Atzemi-Moldow N, Hatzidakis GI. Ruptured aneurysm at the trunk of the accessory middle cerebral artery. *Acta neurochirurgica*. 1999;141 11:1233-5.
14. Fujiwara K, Saito K, Ebina T. Saccular aneurysm of the accessory middle cerebral artery—case report. *Neurologia medico-chirurgica*. 2003;43 1:31-4.

15. Kang DH, Park J, Park SH, Hamm IS. Saccular aneurysm at the anterior communicating artery complex associated with an accessory middle cerebral artery : report of two cases and review of the literature. *Journal of Korean Neurosurgical Society*. 2009;46 6:568-71; doi: 10.3340/jkns.2009.46.6.568.
16. Lee IH, Jeon P, Kim KH, Byun HS, Kim HJ, Kim ST, et al. Endovascular treatment of a ruptured accessory middle cerebral artery aneurysm. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2010;17 3:383-4; doi: 10.1016/j.jocn.2009.02.023.
17. Teramoto S, Tokugawa J, Nakao Y, Yamamoto T. Caudate haemorrhage caused by pseudoaneurysm of accessory middle cerebral artery. *BMJ case reports*. 2015;2015; doi: 10.1136/bcr-2015-213335.
18. Parthasarathy R, Goel G, Gupta V, Narang KS, Anand S, Jha AN. Endovascular glue embolization of dissecting aneurysm of type-3 accessory middle cerebral artery: A contralateral approach. *Interventional neuroradiology : journal of peritherapeutic neuroradiology, surgical procedures and related neurosciences*. 2015;21 6:664-8; doi: 10.1177/1591019915609780.
19. Kheyreddin AS, Filatov YM, Kaftanov AN, Sazonov IA, Bukharin EY, Dmitriev AV. [Rupture of a giant aneurysm of the accessory middle cerebral artery: a case report and a literature review]. *Zhurnal voprosy neirokhirurgii imeni N N Burdenko*. 2017;81 6:103-7; doi: 10.17116/neiro2017816103-107.
20. Ren H, Ma L, Wei M, Li J, Yu M, Yin L. Duplicated middle cerebral artery origin with an aneurysm. *Medicine*. 2018;97 9:e9947; doi: 10.1097/md.00000000000009947.
21. Lee CC, Liu ZH, Jung SM, Yang TC. Ruptured aneurysm of the accessory middle cerebral artery associated with moyamoya disease: a case report. *Chang Gung medical journal*. 2011;34 5:541-7.
22. Wakabayashi Y, Hori Y, Kondoh Y, Asano T, Yamada A, Kenai H, et al. Ruptured anterior cerebral artery aneurysm at the origin of the accessory middle cerebral artery. *Neurologia medico-chirurgica*. 2011;51 9:645-8.
23. Stabler J. Two cases of accessory middle cerebral artery, including one with an aneurysm at its origin. *The British journal of radiology*. 1970;43 509:314-8; doi: 10.1259/0007-1285-43-509-314.
24. In S, In K, Kusano N, Mizuki H, Miyagi J, Kuramoto S. [A case of duplication of the middle cerebral artery with ruptured aneurysm on its origin during pregnancy (author's transl)]. *No shinkei geka Neurological surgery*. 1981;9 3:337-41.
25. Takano S, Nose T, Oowada T, Shirai S, Maki Y. [Aneurysm arising from duplicated middle cerebral artery. Case report]. *Neurologia medico-chirurgica*. 1988;28 9:910-4.
26. Dong LW, Yamada K, Ohta T, Takahashi N. [Ruptured intracranial aneurysm combined with multiple cerebral vessel anomalies; a case report]. *No shinkei geka Neurological surgery*. 1991;19 10:975-8.
27. Takahashi T, Suzuki S, Ohkuma H, Iwabuchi T. Aneurysm at a duplication of the middle cerebral artery. *AJNR American journal of neuroradiology*. 1994;15 6:1166-8.
28. Koyama S, Kotani A, Sasaki J, Tazoe M, Tsubokawa T. Ruptured aneurysm at the origin of duplication of the middle cerebral artery—case report. *Neurologia medico-chirurgica*. 1995;35 9:671-3.
29. Nomura M, Yamashita T, Kita D, Kida S, Kajinami K, Yamashita J. Duplication of the middle cerebral artery associated with an unruptured aneurysm. *Acta neurochirurgica*. 2000;142 2:221-2.
30. Tabuse M, Wakamoto H, Miyazaki H, Ishiyama N. [The usefulness of 3D-CTA for the diagnosis of a ruptured aneurysm at the origin of the duplicated middle cerebral artery: case report]. *No shinkei geka Neurological surgery*. 2002;30 3:327-31.
31. Imaizumi S, Onuma T, Motohashi O, Kameyama M, Ishii K. Unruptured carotid-duplicated middle cerebral artery aneurysm: case report. *Surgical neurology*. 2002;58 5:322-4.
32. Uchino M, Kitajima S, Sakata Y, Honda M, Shibata I. Ruptured aneurysm at a duplicated middle cerebral artery with accessory middle cerebral artery. *Acta neurochirurgica*. 2004;146 12:1373-4; discussion 5; doi: 10.1007/s00701-004-0353-x.
33. Hori E, Kurosaki K, Matsumura N, Yamatani K, Kusunose M, Kuwayama N, et al. Multiple aneurysms arising from the origin of a duplication of the middle cerebral artery. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2005;12 7:812-5; doi: 10.1016/j.jocn.2004.08.033.
34. Kai Y, Hamada J, Morioka M, Yano S, Kudo M, Kuratsu J. Treatment of unruptured duplicated middle cerebral artery aneurysm: case report. *Surgical neurology*. 2006;65 2:190-3; discussion 3; doi: 10.1016/j.surneu.2005.05.032.
35. Kaliaperumal C, Jain N, McKinstry CS, Choudhari KA. Carotid "trifurcation" aneurysm: surgical anatomy and management. *Clin Neurol Neurosurg*. 2007;109 6:538-40; doi: 10.1016/j.clineuro.2007.04.006.
36. Miyahara K, Fujitsu K, Ichikawa T, Mukaiyama S, Okada T, Kaku S. [Unruptured saccular aneurysm at the origin of the duplicated middle cerebral artery: reports of two cases and review of the literature]. *No shinkei geka Neurological surgery*. 2009;37 12:1241-5.
37. Otani N, Nawashiro H, Tsuzuki N, Osada H, Suzuki T, Shima K, et al. A ruptured internal carotid artery aneurysm located at the origin of the duplicated middle cerebral artery associated with accessory middle cerebral artery and middle cerebral artery aplasia. *Surgical neurology international*. 2010;1:51; doi: 10.4103/2152-7806.69378.
38. Kimura T, Morita A. Treatment of unruptured aneurysm of duplication of the middle cerebral artery—case report. *Neurologia medico-chirurgica*. 2010;50 2:124-6.
39. Takahashi C, Kubo M, Okamoto S, Matsumura N, Horie Y, Hayashi N, et al. "Kissing" aneurysms of the internal carotid artery treated by coil embolization. *Neurologia medico-chirurgica*. 2011;51 9:653-6.
40. LaBorde DV, Mason AM, Riley J, Dion JE, Barrow DL. Aneurysm of a duplicate middle cerebral artery. *World neurosurgery*. 2012;77 1:201.e1-4; doi: 10.1016/j.wneu.2011.03.038.

41. Rennert J, Ullrich WO, Schuierer G. A rare case of supraclinoid internal carotid artery (ICA) fenestration in combination with duplication of the middle cerebral artery (MCA) originating from the ICA fenestration and an associated aneurysm. *Clinical neuroradiology*. 2013;23 2:133-6; doi: 10.1007/s00062-011-0120-3.
42. Elsharkawy A, Ishii K, Niemela M, Kivisaari R, Lehto H, Hernesniemi J. Management of aneurysms at the origin of duplicated middle cerebral artery: series of four patients with review of the literature. *World neurosurgery*. 2013;80 6:e313-8; doi: 10.1016/j.wneu.2012.10.004.
43. Kim JS, Lee CH, Park H, Han JW. An Unruptured Cerebral Aneurysm at the Origin of the Duplicated Middle Cerebral Artery. *Journal of cerebrovascular and endovascular neurosurgery*. 2015;17 3:223-6; doi: 10.7461/jcen.2015.17.3.223.
44. Iida Y, Tamase A, Kamide T, Mori K, Seki S, Nomura M. Aneurysm at origin of duplicated middle cerebral artery associated with another aneurysm. *Surgical neurology international*. 2015;6 Suppl 21:S549-52; doi: 10.4103/2152-7806.168069.
45. Miyoshi H, Migita K, Kumano K, Hashimoto N, Toyota A. [A Case of Aphasia after Neck Clipping of a Ruptured Aneurysm at the Origin of the Duplicated Middle Cerebral Artery]. *No shinkei geka Neurological surgery*. 2016;44 11:959-64; doi: 10.11477/mf.1436203408.
46. Hayashi K, Matsuo Y, Hayashi Y, Shirakawa K, Kaminogo M. [A Ruptured Aneurysm at the Origin of a Duplicated Middle Cerebral Artery, Treated by Coil Embolization:A Case Report]. *No shinkei geka Neurological surgery*. 2017;45 5:391-6; doi: 10.11477/mf.1436203520.
47. Tsang COA, Smith L, Klostranec J, Orru E, Pereira VM. Ruptured Duplicated Middle Cerebral Artery Aneurysm Successfully Treated by Coil Embolization with Balloon Remodeling. *World neurosurgery*. 2018;120:509-10; doi: 10.1016/j.wneu.2018.09.154.
48. Ueda T, Goya T, Kinoshita K. [Fenestration of the middle cerebral artery associated with an aneurysm]. *Neurologia medico-chirurgica*. 1983;23 4:278-81.
49. Ueda T, Goya T, Kinoshita K, Wakuta Y, Mihara K. [Multiple anomalies of cerebral vessels. A case of multiple aneurysms associated with fenestration of the middle cerebral artery and persistent primitive trigeminal artery]. *No shinkei geka Neurological surgery*. 1984;12 4:531-6.
50. Kalia KK, Ross DA, Gutin PH. Multiple arterial fenestrations, multiple aneurysms, and an arteriovenous malformation in a patient with subarachnoid hemorrhage. *Surgical neurology*. 1991;35 1:45-8.
51. Deruty R, Pelissou-Guyotat I, Mottotese C, Bogner L, Laharotte JC, Turjman F. Fenestration of the middle cerebral artery and aneurysm at the site of the fenestration. *Neurological research*. 1992;14 5:421-4.
52. Nakamura H, Takada A, Hide T, Ushio Y. Fenestration of the middle cerebral artery associated with an aneurysm—case report. *Neurologia medico-chirurgica*. 1994;34 8:555-7.
53. Schmieder K, Hardenack M, Harders A. Proximal long fenestration associated with an internal carotid artery aneurysm. Case illustration. *Journal of neurosurgery*. 1997;86 4:733; doi: 10.3171/jns.1997.86.4.0733.
54. Nussbaum ES, Defillo A, Janjua TM, Nussbaum LA. Fenestration of the middle cerebral artery with an associated ruptured aneurysm. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2009;16 6:845-7; doi: 10.1016/j.jocn.2008.08.039.
55. Sim KB, Lee CS, Park JC, Huh JS. Cerebral aneurysm in the long fenestration at the middle portion of m1 segment. *Journal of Korean Neurosurgical Society*. 2010;48 5:434-7; doi: 10.3340/jkns.2010.48.5.434.
56. Yamaguchi S, Ito O, Suzuki S. Coil embolization of a ruptured aneurysm arising from a middle cerebral artery fenestration—case report. *Neurologia medico-chirurgica*. 2010;50 3:213-6.
57. Tabuchi S, Yoshioka H. Ruptured aneurysm at the fenestration of the middle cerebral artery detected by magnetic resonance angiography in a patient with systemic lupus erythematosus and renal failure: a case report. *Journal of medical case reports*. 2014;8:30; doi: 10.1186/1752-1947-8-30.
58. Sharifi G, Bakhtevari MH, Sabouri S, Rezaei O. An aneurysm at the site of the fenestration of the middle cerebral artery in a patient with multiple aneurysms: A case report. *Surgical neurology international*. 2015;6 Suppl 16:S414-7; doi: 10.4103/2152-7806.166177.
59. Hirayama K, Nakamura H, Yoshimura S, Ozono K, Fukuda Y, Hiu T, et al. [Unruptured Aneurysm Arising from the Proximal End of the Duplicate Origin of the Middle Cerebral Artery]. *No shinkei geka Neurological surgery*. 2018;46 10:901-9; doi: 10.11477/mf.1436203837.
60. Dh P. The development of the cranial arteries in the human embryo. *Contrib Embryol* 1948 212:207–61.
61. Watanabe T, Togo M. Accessory middle cerebral artery. Report of four cases. *Journal of neurosurgery*. 1974;41 2:248-51; doi: 10.3171/jns.1974.41.2.0248.
62. Uchino A, Kato A, Takase Y, Kudo S. Middle cerebral artery variations detected by magnetic resonance angiography. *Eur Radiol*. 2000;10 4:560-3; doi: 10.1007/s003300050960.
63. Crompton MR. The pathology of ruptured middle-cerebral aneurysms with special reference to the differences between the sexes. *Lancet*. 1962;2 7253:421-5.
64. Teal JS, Rumbaugh CL, Bergeron RT, Segall HD. Anomalies of the middle cerebral artery: accessory artery, duplication, and early bifurcation. *Am J Roentgenol Radium Ther Nucl Med*. 1973;118 3:567-75.
65. Tutar NU, Tore HG, Kirbas I, Tarhan NC, Coskun M. Various origins of the duplicated middle cerebral artery. *Journal of neuroimaging : official journal of the American Society of Neuroimaging*. 2008;18 4:448-50; doi: 10.1111/j.1552-6569.2008.00254.x.

Tables

Table 1. Aneurysms associated with accessory MCA

Case	Study	Age/sex	Presentation	Size (mm)	Morphology	Side	Location of ac-MCA and aneurysm	Accompanying intracranial anomalies	Treatment	Procedure-related complication	Outcome (GOS)
1	Waga et al., 1977 ⁵	51/F	SAH	Small	Saccular	L	A1, origin	NA/NM	Clipping	NA/NM	1
2	Handa et al., 1984 ⁶	55/F	SAH	Small	Saccular	R	A1, origin	NA/NM	Clipping	No	5
3	Fuwa et al., 1984 ⁷	57/M	SAH	Small	Saccular	L	A1, origin	Right ICA-PComA aneurysm	Clipping	No	5
4	Miyazaki et al., 1984 ⁸	42/M	SAH	Small	Saccular	L	A1, origin	No	Clipping	No	5
5	Kuwabara et al., 1990 ⁹	73/F	SAH	4×6	Saccular	R	A1, origin	No	Clipping	Hydrocephalus	5
6	Han et al., 1994 ¹⁰	34/F	SAH	4×5	Saccular	L	A1, origin	No	Clipping	No	5
7	Sugita et al., 1995 ¹¹	53/M	Visual disturbance	Giant	Saccular	R	A1, origin	No	Clipping	No	5
8	Otawara et al., 1997 ¹²	66/F	SAH	Small	Dissecting	R	A1, origin	Ipsilateral A1 dissecting aneurysm	Wrapping	No	2
9	Georgopoulos et al., 1999 ¹³	32/F	SAH, IVH, ICH	Small	Saccular	L	A1, trunk	No	Clipping	No	4
10	Fujiwara et al., 2003 ¹⁴	30/M	SAH	3×5	Saccular	R	A1, origin	No	Clipping	No	5
11	Kang et al., 2009 ¹⁵	38/M	SAH	4×5	Saccular	L	A1, origin	No	Clipping	No	5
12	Lee et al., 2010 ¹⁶	59/F	SAH	7.3×4.8	Saccular	L	A1, origin	Contralateral ac-MCA at A1	Coiling	No	5
13	Wakabayashi et al., 2011 ²²	36/F	SAH	3×6	Saccular	L	A1, origin	Ipsilateral ac-MCA at A2	Clipping	No	5
14	Lee et al., 2011 ²¹	56/M	SAH, IVH	Small	Lobular pseudoaneurysm	R	ACA, trunk	Right MMD involving ICA, ACA, and MCA	Resection of aneurysm and distal ac-MCA	No	5
15	Nomura et al., 2015 ⁴	64/M	CI	Small	Saccular	L	A1, origin	No	Clipping	No	5
16	Teramoto et al., 2015 ¹⁷	68/M	ICH	7	Saccular	R	A2, trunk	Ipsilateral M1 stenosis	Clipping	No	5
17	Parthasarathy et al., 2015 ¹⁸	4/F	SAH	8.8×2.1	Fusiform dissecting	R	A1-A2 junction, trunk	Contralateral d-MCA	Aneurysm and parent artery occlusion with glue	No	5
18	Kheyreddin et al., 2017 ¹⁹	37/F	SAH and ICH	25	Saccular	R	A1, trunk	No	Clipping	No	5
19	Ren et al., 2018 ²⁰	59/M	ICH	Small	Saccular	R	A1-A2 junction, trunk	Ipsilateral d-MCA origin	Palliative coiling	No	5
20	Present case	59/M	SAH	6.5 × 3.0	Saccular	L	A1, origin	Contralateral ac-MCA	Coiling	No	5

Abbreviations: ACA, anterior cerebral artery; CI, cerebral infarction; F, female; GOS, Glasgow Outcome Scale; ICA, internal carotid artery; ICH, intracerebral hemorrhage; IVH, intraventricular hemorrhage; L, left; M, male; MCA, middle cerebral artery; MMD, moyamoya disease; NA/NM, not applicable or not mentioned; PComA, posterior communicating artery; R, right; SAH, subarachnoid hemorrhage

Table 2. Aneurysms associated with duplicate MCA

Case	Study	Age/sex	Presentation	Size (mm)	Morphology	Side	Location of aneurysm	Accompanying intracranial anomalies	Treatment	Postoperative complication	Outcome (GOS)
1	Stabler et al., 1970 ²³	31/F	SAH	Small	Saccular	R	Origin	An aneurysm at the bifurcation of the left ICA	Clipping	Hydrocephalus	NA/NM
2	In et al., 1981 ²⁴	29/F	SAH	Small	Saccular	R	Origin	No	Clipping	No	5
3	Fuwa et al., 1984 ⁷	46/F	SAH	Small	Saccular	R	Origin	No	Clipping	No	5
4	Takano et al., 1988 ²⁵	74/M	Head trauma	6	Saccular	R	Origin	No	Clipping	Hydrocephalus	NA/NM
5	Dong et al., 1991 ²⁶	50/M	SAH	Small	Saccular	L	Origin	ACA fenestration, ac-MCA	Clipping	No	5
6	Takahashi et al., 1994 ²⁷	51/F	SAH	Small	Saccular	L	Origin	Contralateral carotid-ophthalmic aneurysm, d-MCA sharing common trunk with fetal PCA	Clipping	No	5
7		54/M	SAH	Small	Saccular	L	Origin	No	Clipping	Vasospasm	3
8	Koyama et al., 1995 ²⁸	28/M	SAH	Small	Saccular	R	Origin	No	Clipping	No	5
9	Nomura et al., 2000 ²⁹	63/F	Incidental	Small	Saccular	L	Origin	No	Clipping	No	5
10	Tabuse et al., 2002 ³⁰	34/F	SAH	Small	Saccular	R	Origin	No	Clipping	No	5
11	Imaizumi et al., 2002 ³¹	52/M	SAH	Small	Saccular	L	Origin	Contralateral ICA-PComA aneurysm	Clipping	No	5
12	Uchino et al., 2004 ³²	45/F	SAH, ICH	Small	Saccular	L	Trunk	Bilateral ac-MCAs	Clipping	No	5
13	Hori et al., 2005 ³³	67/M	SAH, ICH	Small	Saccular	R	Origin	Ipsilateral ICA-PComA aneurysm	Clipping	Aphasia, hydrocephalus	3
14		49/M	Incidental	Small	Saccular	L	Origin	BA tip aneurysm	Clipping	No	5
15	Kai et al., 2006 ³⁴	63/F	Vertigo	Small	Saccular	L	Origin	No	Clipping and STA-d-MCA anastomosis	No	5
16	Kaliaperumal et al., 2007 ³⁵	39/F	SAH	< 10	Saccular	L	Origin	No	Clipping	No	5
17	Miyahara et al., 2009 ³⁶	56/F	Incidental	Small	Saccular	R	Origin	3 aneurysms at other locations	Clipping	No	5
18		58/M	Vertigo	7	Saccular	R	Origin	Ipsilateral ICA-PComA aneurysm	Clipping	No	5
19	Otani et al., 2010 ³⁷	66/F	SAH	6	Saccular	R	Origin	Ipsilateral ac-MCA	Clipping	No	5
20	Kimura et al., 2010 ³⁸	60/F	Incidental	4	Saccular	L	Origin	No	Clipping	No	5
21	Takahashi et al., 2011 ³⁹	62/F	SAH	4.0 × 4.2	Saccular	L	Origin	Ipsilateral AChA aneurysm	Coiling	No	5
22	laBored et al., 2012 ⁴⁰	34/M	Incidental	10	Fusiform	L	Trunk	No	Trapping of the aneurysm and STA-MCA anastomosis	Craniotomy flap infection	5
23	Rennert et al., 2013 ⁴¹	52/F	Recurring headache	2	Saccular	L	Origin	Ipsilateral supraclinoid ICA fenestration, AComA aneurysm	Clipping	No	5
24	Elsharkawy et al., 2013 ⁴²	62/M	Epilepsy	12	Saccular	L	Origin	No	Clipping	No	5
25		55/F	SAH	3	Saccular	L	Origin	Contralateral ICA-PComA aneurysm	Clipping	No	5
26		49/F	Migraine and double vision	1 × 2	Saccular	R	Origin	Contralateral ICA bifurcation aneurysm	Conservative management	NA/NM	NA/NM
27		37/M	Incidental	1	Saccular	L	Origin	No	Conservative management	NA/NM	NA/NM
28	Kim et al., 2015 ⁴³	61/F	Headache	3	Saccular	R	Origin	No	Clipping	No	5
29	Iida et al., 2015 ⁴⁴	41/F	SAH	5.5 × 6.5	Saccular	R	Origin	Contralateral ICA-PComA aneurysm	Clipping	No	5
30		76/F	SAH, ICH	1.8 × 2.5	Saccular	R	Origin	Ipsilateral MCA aneurysm	Clipping	Vasospasm, hydrocephalus	3
31	Miyoshi et	60/F	SAH	Small	Saccular	L	Origin	No	Clipping	Temporary aphasia	5

	al., 2016 ⁴⁵										
32	Hayashi et al., 2017 ⁴⁶	41/M	SAH	Small	Saccular	NA/NM	Origin	No	Coiling	No	5
33	Mori et al., 2018 ³	62/M	Alcohol abuse	< 5	Saccular	L	Origin	Ipsilateral ICA bifurcation aneurysm	Conservative management	NA/NM	NA/NM
34	Tsanget al., 2018 ⁴⁷	20/F	SAH	3.6 × 3.1	Saccular	R	Origin	Ipsilateral AChA aneurysm	Coiling	No	5

Abbreviations: ACA, anterior cerebral artery; AChA, anterior choroidal artery; BA, basilar apex; F, female; GOS, Glasgow Outcome Scale; ICA, internal carotid artery; ICH, intracerebral hemorrhage; L, left; M, male; MCA, middle cerebral artery; NA/NM, not applicable or not mentioned; PCA, posterior cerebral artery; PComA, posterior communicating artery; R, right; SAH, subarachnoid hemorrhage; STA, superficial temporal artery

Table 1. Aneurysms associated with MCA fenestration

Case	Study	Age/sex	Presentation	Size (mm)	Morphology	Side	Location of MCA fenestration and aneurysm	Accompanying intracranial anomalies	Treatment	Procedure-related complication	Outcome (GOS)
1	Ueda et al., 1983 ⁴⁸	65/M	ICH	NA/NM	Saccular	R	M1, ipsilateral MCA bifurcation	No	Clipping	NA/NM	NA/NM
2	Ueda et al., 1984 ⁴⁹	45/F	SAH	Small	Saccular	R	M1, proximal to fenestration	Multiple intracranial aneurysms, PTA	Clipping	No	5
3	Kalia et al., 1991 ⁵⁰	49/M	SAH	Small	Saccular	R	M1, in fenestration	ACoM fenestration, multiple intracranial aneurysms, AVM	Wrapping	No	5
4	Deruty et al., 1992 ⁵¹	52/F	SAH	Small	Saccular	R	M1, in fenestration	ACoM aneurysm	Clipping	No	5
5	Nakamura et al., 1994 ⁵²	36/F	SAH	Small	Saccular	L	M1, proximal to fenestration	No	Clipping	No	5
6	Schmieder et al., 1997 ⁵³	14/M	SAH	Small	Saccular	L	M1, proximal to fenestration	No	Clipping	No	5
7	Nussbaum et al., 2009 ⁵⁴	75/F	SAH	5	Saccular	R	M1, distal to fenestration	No	Clipping	No	5
8	Sim et al., 2010 ⁵⁵	32/M	SAH	6	Saccular	R	M1, in fenestration	Contralateral MCA aneurysm	Clipping	No	5
9	Yamaguchi et al., 2010 ⁵⁶	81/F	SAH	Small	Saccular	R	M1, in fenestration	Contralateral MCA aneurysm	Coiling	No	5
10	Tabuchi et al., 2014 ⁵⁷	47/F	SAH	Small	Saccular	R	M1, proximal to fenestration	No	Clipping	Hydrocephalus	5
11	Sharifi et al., 2015 ⁵⁸	52/M	SAH	Small	Saccular	L	M1, distal to fenestration	Multiple intracranial aneurysms	Clipping	Bacterial meningitis	5
12	Xue et al., 2019 ²	43/M	SAH	2.5	Saccular	R	M1, proximal to fenestration	No	Clipping	No	5

Abbreviations: ACoM, anterior communicating artery; AVM, arteriovenous malformation; F, female; GOS, Glasgow Outcome Scale; ICH, intracerebral hemorrhage; L, left; M, male; MCA, middle cerebral artery; NA/NM, not applicable or not mentioned; PTA, persistent trigeminal artery; R, right; SAH, subarachnoid hemorrhage

Figures

Figure 1

Flow chart of searching strategy

Figure 2

Head CT shows subtle SAH (A-B). CTA reveals two ac-MCAs originate respectively from the A1 segment of the bilateral ACAs (C, black arrow) and an aneurysm originates from the beginning of the left ac-MCA (C-D, asterisk and white arrow). Abbreviations: ACA, anterior cerebral artery; ac-MCA, accessory middle cerebral artery; CT, computed tomography; CTA, CT angiography

Figure 3

Three dimensional (A) and plain (B) angiogram of the left ICA in AP view shows an ac-MCA arises from the A1 segment of ACA and a saccular aneurysm is also note at the origin of ac-MCA. Angiogram of the left ICA in AP view shows the aneurysm is successfully coiled (C-D). Abbreviations: ACA, anterior cerebral artery; ac-MCA, accessory middle cerebral artery; AP, anteroposterior; ICA, internal carotid artery

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [PRISMA2009checklist.doc](#)