Methicillin-resistant *Staphylococcus aureus* nasal colonization among HIV-infected patients in Taiwan: prevalence, molecular characteristics and associated factors with nasal carriage

Yhu-Chering Huang (✉ ychuang@cgmh.org.tw)

Taoyuan Chang Gung Memorial Hospital

Yi-Yu Hsu

Mackay Memorial Hospital

David Wu

Cathay General Hospital

Chien-Ching Hung

National Taiwan University Hospital

Shie-Shian Huang

Keelung Chang Gung Memorial Hospital of the CGMF

Fang-Hsueh Yuan

Chang Gung Memorial Hospital

Ming-Hsun Lee

Chang Gung Memorial Hospital

Ching-Tai Huang

Chang Gung Memorial Hospital

Shian-Sen Shie

Chang Gung Memorial Hospital

Po-Yen Huang

Chang Gung Memorial Hospital

Chien-Chang Yang

Chang Gung Memorial Hospital

Chun-Wen Cheng

Chang Gung Memorial Hospital

Hsieh-Shong Leu

Chang Gung Memorial Hospital

Ting-Shu Wu

Chang Gung Memorial Hospital

Research article

Keywords: nasal carriage, antibiotic use, injection drug use, injection drug user, molecular typing

Posted Date: December 15th, 2019

DOI: <https://doi.org/10.21203/rs.2.18964/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on March 30th, 2020. See the published version at <https://doi.org/10.1186/s12879-020-04979-8>.

Abstract

Objective: To evaluate nasal carriage and molecular characteristics of methicillin-resistant *Staphylococcus aureus* (MRSA), as well as the risk factors of MRSA colonization, in human immunodeficiency virus (HIV)-infected patients in northern Taiwan.

Methods: From September 2014 to November 2015, HIV-infected patients seeking outpatient care at four hospitals were eligible for this study. A nasal specimen was obtained from each subject for the detection of *S. aureus* and a questionnaire was completed by each subject. MRSA isolates once identified were characterized.

Results: Of 553 patients surveyed, methicillin-susceptible *S. aureus* (MSSA) was detected in 119 subjects (21.5%) and MRSA in 19 subjects (3.4%). Female gender, injection drug use, smoking, hepatitis C virus carrier, cancer and antibiotic use within one year were positively associated with MRSA colonization. By multivariate analysis, only cancer (adjusted odds ratio (aOR) 7.78, [95% confidence interval (CI), 1.909–31.731]) and antibiotic use within one year (aOR 3.89, [95% CI, 1.219–12.433]) were significantly associated with MRSA colonization. Ten isolates were characterized as sequence type (ST) 59/staphylococcal chromosome cassette (SCC) IV or V_T, endemic community strains in Taiwan, four isolates as ST 8/SCC*mec* IV (USA 300) and one isolate as ST 239/SCC*mec* IIIA, a hospital strain.

Conclusions: Nasal MRSA carriage in HIV-infected patients seeking outpatient care was low (3.4%) in northern Taiwan, and most of the colonizing isolates were genetically endemic community strains. Cancer and antibiotic use within one year were associated with MRSA colonization.

Background

Methicillin-resistant *Staphylococcus aureus* (MRSA) was first reported in the 1960s [1] and the infections due to MRSA rapidly increased in the 1980s [2–4], most of which were health care-acquired infections. In the late 1990s, MRSA began to be recognized as a major cause of community-onset infections, which was termed as community-associated MRSA (CA-MRSA) later [2–4]. The epidemic of CA-MRSA has spread rapidly both in the community [2–4] and in healthcare settings over the past decade. Carriage of *S. aureus*, including MRSA, has been identified as a predisposing factor for subsequent invasive infections [5, 6], and the anterior nares are the most common site of colonization.

Among human immunodeficiency virus (HIV)-infected patients, *S. aureus* infection is considered to be a major cause of significant morbidity and mortality [7–9]. Previous studies suggest that prevalence of nasal colonization with MRSA is higher in HIV-infected individuals than in the general population [10–12]. HIV infection is also related to persistent colonization [13]. The higher colonization burden may be associated with a higher incidence of subsequent infections [10, 11, 14]. However, risk factors for *S. aureus* colonization appear to be different among different populations, including patients infected with HIV [11, 15–17]. MRSA colonization in HIV-infected patients may vary widely in different geographical

regions [15, 18–21], the time point of survey [14], and the coverage of antiretroviral therapy in the population [22].

MRSA has been a prevalent etiology of infections in Asian countries, including Taiwan, either in healthcare settings or in the community, for decades, but the reports regarding MRSA colonization in HIV-infected population have been scarce [12, 15, 23]. Hence, we conducted this study to evaluate the prevalence of and associated factors for MRSA colonization in HIV-infected population in Taiwan. All collected MRSA isolates were further characterized by molecular methods and antibiotic susceptibility.

Methods

The study was approved by the institutional review board (IRB) of Chang Gung Memorial Hospital and the Research Ethics Committee of National Taiwan University Hospital. A written informed consent was obtained from each subject.

Subject Enrollment

The study was conducted in hospital-based infectious diseases outpatient clinics at Chang Gung Memorial Hospital (CGMH) and National Taiwan University Hospital (NTUH) from September 2014 to November 2015. Both CGMH and NTUH are university-affiliated medical centers situated in northern Taiwan. Data were obtained in multiple CGMH branches, including Linkou branch, Taipei branch and Keelung branch. We also recruited patients from prison and correctional facilities in Keelung, where inmates were regularly followed by infectious diseases specialists in CGMH Keelung branch. All the HIV-infected patients who visited and treated at CGMH and NTUH were eligible and invited to participate in this study. Study subjects were enrolled when they sought HIV care at the infectious diseases outpatient departments.

Data collection

To identify the associated factors with MRSA acquisition, a self-administered questionnaire interview was performed to inquire into the factors for colonization with MRSA in HIV-infected patients. The following information was also collected from medical records of the participants: age, gender, drinking habits and smoking habits, underlying diseases, clinical characteristics, latest hospitalization, community exposure (eg, drug use, and incarceration), previous *S. aureus* infection, antibiotic use, skin disease, plasma HIV RNA load and CD4 count, and antiretroviral therapy. Clinical information regarding hospitalization, residence in a long-term care facility, outpatient department visit, surgery, dialysis, and usage of tubes (nasogastric tube, urine catheter, tracheostomy tube, drainage tube, port-A, and dialysis tube) were also obtained. Substance abuse was defined as current or former use of illicit drugs.

Microbiologic study

Nasal specimens were obtained by swabbing the anterior 1 cm of the nasal vestibule of both anterior nares of the participants. The swabs were preserved in the transport medium (Venturi Transystem, Copan

Innovation Ltd.) immediately. *S. aureus* was identified by colony morphology, Gram stain, positive coagulase tests, and presence of β -hemolysis after subculture to tryptic soy agar plates containing 5% sheep's blood. Methicillin resistance was confirmed by ceftiofur disk-diffusion method according to the recommendation of Clinical and Laboratory Standard Institutes [24].

Antimicrobial susceptibility study

The antimicrobial susceptibility of all MRSA isolates to ten antibiotics, including ciprofloxacin, trimethoprim/sulfamethoxazole (SXT), penicillin, teicoplanin, linezolid, clindamycin, doxycycline, fusidic acid, vancomycin, and erythromycin, was tested in accordance with the guideline of Clinical and Laboratory Standard Institutes by using the disk-diffusion method [24].

Molecular characterization

By pulsed-field gel electrophoresis (PFGE) with *Sma*I digestion [25-27], the genotypes were designated in alphabetical order, as in our previous studies [25-27]; any new genotype, if identified, was designated consecutively. PFGE patterns with fewer than four band differences from an existing genotype were defined as subtypes of that genotype. Staphylococcal chromosome cassette *mec* (*SCCmec*) type, and the presence of Panton-Valentine leukocidin (PVL) genes were determined by a multiplex PCR strategy [28]. Some isolates of representative PFGE patterns were selected and underwent multilocus sequence typing (MLST) [29] and *spa* typing [27]. The allelic profiles were assigned through comparison of the sequences at each locus with those of the known alleles in the *S. aureus* MLST database and were defined as sequence types accordingly. The details of the procedures were described elsewhere previously [25-29].

Statistical Analysis

Statistical analyses were performed with the Statistical Package for the Social Sciences (SPSS software for Windows, version 17.0). The definition of statistical significance was $p < 0.05$. Chi-square analysis or Fisher's exact test was used for examination of categorical variables. Continuous variables were compared between patients with MRSA colonization versus patients without MRSA colonization using two-sample t-test, in which Levene's test was used to determine equality of variances between the two groups.

Results

During the 14-month study period, a total of 810 HIV-infected patients sought outpatient care at CGMH and NTUH (446 from CGMH and 364 from NTUH). Of the 810 eligible patients, 586 patients (275 from CGMH and 311 from NTUH) were interviewed and invited to participate in this study. 553 patients (259 from CGMH and 294 from NTUH) were enrolled in total and were surveyed for nasal carriage of MRSA after giving their written informed consent.

Detailed demographic data are shown in Table 1. 531 were male and 22 female. The majority of the participants were aged 20–60 years (with 47.6% between 20–39 years, and 45.6% between 40–60 years). The main route of HIV acquisition was male-to-male sex contact (68.0%), followed by injection drug use (16.3%). Table 1 shows the comparison among different hospitals and no statistically significant difference was found among different hospitals in terms of nasal MRSA carriage rate among the HIV-infected patients.

Table 1

Comparison of demographics and transmission vehicles between HIV-infected patients with and without methicillin-resistant *Staphylococcus aureus* (MRSA) nasal colonization

Characteristic	No. (%) of subjects			P value
	Total (n = 553)	MRSA carriers (n = 19)	Non-MRSA carriers (n = 534)	
Age in years, mean (SD)	41.2 (11.4)	43.8 (13.0)	41.1 (11.3)	0.296
Gender				
Male	531(96)	15(79)	516(97)	0.005*
Female	22(4)	4(21)	18(3)	
Route of HIV transmission				
Injection drug use	90(16)	9(47)	81((15)	0.001*
Heterosexual activity	62(11)	2(11)	61(11)	1.000
Male-to-male sex	375(68)	6(32)	369(69)	0.002*
Blood transfusion	3(0.5)	0(0)	3(0.6)	1.000
Vertical transmission	1(0.2)	0(0)	1(0.2)	1.000
Unknown	36(6.5)	2(11)	34(6.4)	0.380
Facility				0.203
CGMH, Linko branch	111 (20.1)	3 (15.8)	108 (20.2)	
CGMH, Taipei branch	13 (2.4)	0 (0)	13 (2.4)	
CGMH, Keelung branch	117 (21.2)	8 (42.1)	109 (20.4)	
Keelung jail and prison	18 (3.3)	1 (5.3)	17 (3.2)	
NTUH	294 (53.2)	7 (36.8)	287 (53.8)	
Abbreviations: CGMH, Chang Gung Memorial Hospital; NTUH, National Taiwan University Hospital				

The overall MSSA and MRSA nasal colonization rate was 21.5% and 3.4%, respectively. The comparison of underlying diseases and other medical history between patients with and without MRSA colonization are shown in Tables 1 and 2. In univariate analysis, we found that female gender ($p = 0.005$), injection drug use ($p = 0.001$), smoking ($p = 0.02$), hepatitis C virus (HCV) carrier ($p = 0.003$), cancer ($p = 0.001$) and

antibiotic use within one year ($p = 0.02$) were significant factors associated with MRSA colonization. CD4 count or plasma HIV RNA load was not significantly associated with MRSA colonization.

Table 2

Analysis of risk factors for nasal colonization with methicillin-resistant *Staphylococcus aureus* (MRSA) among HIV-infected Outpatients

Characteristic	No. (%) of subjects			P value
	Total (n = 553)	MRSA carriers (n = 19)	Non-MRSA carriers (n = 534)	
Personal history				
Smoking	290(52)	15(79)	275(51)	0.020*
Alcohol Drinking	139(25)	6((32)	133(25)	0.590
Hospitalization after HIV diagnosis	269(49)	7(37)	262(49)	0.354
Recent hospitalization within one year	80(14)	5(26)	75(14)	0.174
Operation history	16(3)	0(0)	16(3)	1.000
Nurse home	3(0.6)	0(0)	3(0.6)	1.000
Underlying diseases				
Hypertension	59(11)	3(16)	56(10)	0.443
DM	21(4)	1(5)	20(4)	0.527
Arrhythmia history	10(2)	1(5)	9(2)	0.297
Angina history	7(1)	0(0)	7(1)	1.000
Pneumonia history	26(5)	1(5)	25(5)	0.606
COPD	6(1)	0(0)	6(1)	1.000
Asthma	12(2)	0(0)	12(2)	1.000
Tuberculosis	20(4)	1(5)	19(4)	0.509
HBV carrier	62(11)	2(11)	60(11)	1.000
HCV carrier	123(22)	10(53)	113(21)	0.003*
Cirrhosis	3(0.5)	1(5)	2(0.4)	0.100
CKD	10(2)	1(5)	9(2)	0.297

Abbreviations: DM, diabetes mellitus; COPD, chronic obstructive pulmonary disease; HBV, hepatitis B virus; HCV, hepatitis C virus; CKD, chronic kidney disease; URI, upper respiratory infection; SD, standard deviation

Characteristic	No. (%) of subjects			P value
	Total (n = 553)	MRSA carriers (n = 19)	Non-MRSA carriers (n = 534)	
Cancer	27(5)	5(26)	22(4)	0.001*
Recent infectious diseases (within 3 months)				
Recent URI	62(11)	2(11)	60(11)	1.000
Skin infection	29(5)	1(5)	28(5)	1.000
Unhealed wound	16(3)	1(5)	15(3)	0.433
UTI	6(1)	0(0)	6(1)	1.000
Recent treatment (in 3 months) [any information on antiretroviral therapy?]				
On all catheters	10(2)	1(5)	9(2)	0.297
Hemodialysis	2(0.3)	1(5)	1(0.2)	0.068
Antibiotic use in 1 year	88(16)	7(37)	81(15)	0.020*
Laboratory data				
CD4 count (cells/uL), mean $\pm$ SD		469.16 $\pm$ 214.8	540.61 $\pm$ 292.3	0.292
Plasma HIV RNA load (Log ₁₀ of copies/ml), mean $\pm$ SD		1.341 $\pm$ 2.055	0.929 $\pm$ 1.660	0.293
Abbreviations: DM, diabetes mellitus; COPD, chronic obstructive pulmonary disease; HBV, hepatitis B virus; HCV, hepatitis C virus; CKD, chronic kidney disease; URI, upper respiratory infection; SD, standard deviation				

Table 3
Multivariate analysis of factors associated with nasal colonization of MRSA in HIV carriers in Taiwan.

Factor	Adjust odds ratio	95% confidence interval	P value
Female Gender	2.910	0.690–12.276	0.1459
Injection drug user	1.189	0.207–6.837	0.8464
Male-to-male sex	0.241	0.055–1.050	0.0581
Smoking	2.004	0.581–6.911	0.2713
HCV carrier	2.233	0.504–9.885	0.2899
Cancer	7.782	1.909–31.731	0.0042
Antibiotic use within the past one year	3.892	1.219–12.433	0.0218

All 19 MRSA isolates were available for molecular characterization and the results are shown in Fig. 1. Six PFGE patterns were identified. One isolate carried SCCmec type IIIA and the remaining isolates carried either SCCmec type IV or VT. PFGE pattern D/ sequence type (ST) 59 was the relatively common clone, followed by PFGE pattern C/ST 59 and PFGE pattern AI/ST 8.

All the MRSA strains were resistant to penicillin and susceptible to linezolid, teicoplanin, doxycycline, and vancomycin. The susceptibility rates to erythromycin, clindamycin, ciprofloxacin, trimethoprim-sulfamethoxazole (TMP-SMX), and fusidic acid were 15.8%, 42.1%, 84.20%, 94.7%, and 89.5%, respectively.

Discussion

To our knowledge, this is the largest study on this issue in Asian countries [15]. In this cross-section survey among 553 HIV-infected patients seeking care at the outpatient clinics in four hospitals in northern Taiwan, we found that the overall prevalence of MRSA nasal carriage among the HIV-infected individual was 3.4% while the prevalence of MSSA nasal carriage was 21.5%. The nasal carriage of MRSA was associated with cancer and antibiotic exposure within the past 12 months.

The nasal carriage rate of MRSA in this study is slightly lower than that previously reported in Taiwan in 2003 (5.6% of 296 HIV-infected patients) [12] and in 2009–2010 (4.4% of 457 HIV-infected patients) [23]. Compared with other populations in Taiwan, the nasal MRSA carriage rate among HIV-infected patients was significantly lower than that for patients hospitalized in the intensive care units (ICU) (32% of 177 patients) [30], but similar to that for adult patients receiving hemodialysis (3.8% of 296 patients) [31], adult patients visiting emergency room (3.8% of 502 patients) [32] and otherwise healthy adults for health examination (3.8% of 3098 adults) [33]. It could be explained that in the era of highly active antiretroviral therapy (HAART), the majority of the HIV-infected patients recruited in this study came from

community settings, outpatient visiting and these patients were not frequently exposed to the high-risk populations for MRSA acquisition, such as patients admitted to the ICUs.

For patients infected with HIV, the nasal MRSA carriage rate was different, with a huge discrepancy, in different countries and regions. A latest meta-analysis on this issue [15] showed that the estimated pooled worldwide prevalence of MRSA in HIV-infected people is around 7% (5%-9%), with the prevalence of 7%-13% in the region of the Americas and 0–1% in the European region. Compared with other Asian countries, the prevalence rate (3.4%) in this study was lower than that in India (6%-26%), as well as in Singapore (3%-10%) but higher than that in Malaysia (< 1%). This variation may be attributable to different prevailing MRSA clones, study population, the intermittent nature of colonization [10] and different culture of antibiotic use between countries [34].

We observed that MRSA colonization was associated with antibiotic use within 12 months among HIV-infected individuals. This finding supported previous observations that recent receipt of antibiotics was strongly associated with MRSA colonization [12]. Wang et al found that smoking was a protective factor against MRSA colonization in the community setting [33], which was not confirmed in this study. The potential reasons for the disparity might be the difference in questionnaire design. A latest meta-analysis on this issue [15] indicated that the risk factors for MRSA colonization included having a previous MRSA infection, hospitalization in the past year and use of antibiotics in the past six months.

Molecular characterization of MRSA isolates in the present study showed that 10 (52.6%) of the 19 isolates belonged to ST59 lineage, the predominant community-associated MRSA strain in Taiwan. One isolate was characterized by ST239-SCCmec IIIA which had been known as the most dominant healthcare-associated MRSA clone in Taiwan [4]. The patient with the isolate of ST239 did have a medical history of recent hospitalization within the past year and had been admitted four times after being diagnosed with HIV. The remaining eight isolates carried type IV SCCmec. Among these eight isolates, four of them were identified and confirmed as USA 300 later [35], the predominant community-associated MRSA strain in the United States, on the basis of their being sequence type 8 by multilocus sequence typing (MLST), detection of arginine catabolic metabolic element (ACME) gene and harboring SCCmec type IV and genes for PVL. Three of them were characterized by ST30-SCCmec IV strains, known as the Southwest Pacific clone [2, 4]. One isolate characterized as sequence type 45 which had been identified as the predominate strain in nursing homes in Taiwan [36]. In this study, USA300 (ST8) accounted for four isolates and was a second most common clone. All four isolates were from NTUH. This is an emergent issue in Taiwan [35], which needs further surveillance and observation.

There were several limitations to the current study. First, less than 70% of the HIV-infected patients visiting CGMH and NTUH participated in this study which reduced the sample size and indirectly affected the analysis of statistical significance. It may be the reason for the lack of statistically significant association between MRSA colonization and the common risk factors reported in previous studies such as low CD4 T-cell count and hospitalization within the past year [37]. Second, the presence of other unrecognized sites of colonization should be considered. In this study, nasal specimens for MRSA

detection were obtained by swabbing both anterior nares. Patients might be colonized in the inguinal, genital or peri-rectal areas but not in the nasopharynx [11, 37]. Other potential reasons for the disparity between the studies included the intermittent nature of colonization [10] and the difference in questionnaire design.

Conclusions

The prevalence of nasal MRSA colonization in HIV-infected patients seeking outpatient care was low (3.4%) in northern Taiwan and most of the colonizing isolates were genetically endemic community strains. Cancer and antibiotic use within one year were associated with MRSA colonization.

Abbreviations

MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-sensitive *Staphylococcus aureus*; CA, community-associated; PFGE, pulsed-field gel electrophoresis; ST, sequence type; PVL, Panton-Valentine leucocidin; HIV, human immunodeficiency virus

Declarations

Ethics approval and consent to participate

The study was approved by the institutional review board of Chang Gung Memorial Hospital and the Research Ethics Committee of National Taiwan University Hospital. A written informed consent was obtained from each subject.

Consent to publish

Not applicable.

Availability of data and materials

All the data are presented in the paper.

Competing interests

The authors have no conflicts of interest relevant to this article to disclose.

Funding

The study was supported by a grant from the Ministry of Science and Technology, Executive Yuan, Taiwan (MOST 104-2815-C-182A-003B) and a grant from Medical Foundation in Memory of Dr. Deh-Lin Cheng. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contribution

YYH, DW, YCH: Study design, Data collection, Data analysis, Writing.

Others: Data collection, Data analysis

Acknowledgement

The authors thank the research assistants, Ya-Lin Huang and Yu-Chiao Huang for excellent technical assistance.

References

1. Jevons MP. "Celbenin"-resistant staphylococci. *BMJ* 1961; **1**:124-5.
2. Chuang YY, Huang YC: Molecular epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in Asia. *Lancet Infect Dis* 2013; **13**: 698-708.
3. David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev* 2010; **23**:616-87.
4. Chen CJ, Huang YC: New epidemiology of *Staphylococcus aureus* infection in Asia. *Clin Microbiol Infect* 2014; **20**:605-23.
5. von Eiff C, Becker K, Machka K, Stammer H, Peters G. Nasal carriage as a source of *Staphylococcus aureus* bacteremia. *N Engl J Med* 2001;344:11-6.
6. Ellis MW, Hospenthal DR, Dooley DP, Gray PJ, Murray CK. Natural history of community-acquired methicillin-resistant *Staphylococcus aureus* colonization and infection in soldiers. *Clin Infect Dis* 2004;39:971-9.
7. Popovich KJ, Weinstein RA, Aroutcheva A, Rice T, Hota B. Community-associated methicillin-resistant *Staphylococcus aureus* and HIV intersecting epidemics. *Clin Infect Dis* 2010; **50**: 979–87.
8. Jacobson MA, Gellermann H, Chambers H. *Staphylococcus aureus* bacteremia and recurrent staphylococcal infection in patients with acquired immunodeficiency syndrome and AIDS-related complex. *Am J Med* 1988;85:172-6.
9. Witt DJ, Craven DE, McCabe WR. Bacterial infections in adult patients with the acquired immune deficiency syndrome (AIDS) and AIDS-related complex. *Am J Med* 1987; **82**: 900–906.
10. Shet A, Mathema B, Mediavilla JR, Kishii K, Mehndru S, Jeane-Pierre P, et al. Colonization and subsequent skin and soft tissue infection due to methicillin-resistant *Staphylococcus aureus* in a cohort of otherwise healthy adults infected with HIV type 1. *J Infect Dis* 2009; **200**: 88–93.
11. Popovich JK, Hota B, Aroutcheva A, Kurien L, Patel J, Lyles-Banks R, et al. Community-associated methicillin-resistant *Staphylococcus aureus* colonization burden in HIV-infected patients. *Clin Infect Dis* 2013; **56**: 1067–1074.

12. McDonald LC, Lauderdale TL, Lo HJ, Tsai JJ, Hung CC. Colonization of HIV-infected outpatients in Taiwan with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*. *Int J STD AIDS* 2003;**14**: 473-7.
13. Padoveze MC, Pedro RrdJ, Blum-Menezes Da, Bratfich OJ, Moretti ML. *Staphylococcus aureus* nasal colonization in HIV outpatients: persistent or transient? *Am J Infect Control* 2008;**36**:187-91.
14. Delorenze GN, Horberg MA, Silverberg MJ, Tsai A, Quesenberry CP, Baxter R. Trends in annual incidence of methicillin-resistant *Staphylococcus aureus* (MRSA) infection in HIV-infected and HIV-uninfected patients. *Epidemiol Infect* 2013; **141**: 2392-402.
15. Sabbagh P, Riahi SM, Gamble HR, Rostami A. The global and regional prevalence, burden, and risk factors for methicillin-resistant *Staphylococcus aureus* colonization in HIV-infected people: A systematic review and meta-analysis. *Am J Infect Control*. 2019 ;**47**:323-333.
16. Popovich KJ, Smith KY, Khawcharoenporn T, Thurlow CJ, Lough J, Thomas G, et al. Community-associated methicillin-resistant *Staphylococcus aureus* colonization in high-risk groups of HIV-infected patients. *Clin Infect Dis* 2012;**54**:1296-303.
17. Onorato M, Borucki MJ, Baillargeon G, Paa, DP, Freeman DH, Pat Cole C, Mayhall CG. Risk factors for colonization or infection due to methicillin-resistant *Staphylococcus aureus* in HIV-positive patients: a retrospective case-control study. *Infect Control Hosp Epidemiol* 1999;**20**: 26-30.
18. Oliva A, Lichtner M, Mascellino MT, Iannetta M, Ialungo AM, Tadadjeu Mewamba S, et al. Study of methicillin-resistant *Staphylococcus aureus* (MRSA) carriage in a population of HIV-negative migrants and HIV-infected patients attending an outpatient clinic in Rome. *Ann Ig* 2013; **25**: 99-107.
19. Imaz A, Cobos-Trigueros N, Falcó V, Dominguez MA, Manzardo C, Pujol M, et al. Community-associated methicillin-resistant *Staphylococcus aureus* infections in HIV-infected patients in Spain. *J Infect* 2013; **66**: 199-201.
20. Joore IKCW, Rooijen MSV, Loeff MFSVD, Neeling AJD, Dam AV, Vries HJCD. Low prevalence of methicillin-resistant *Staphylococcus aureus* among men who have sex with men attending an STI clinic in Amsterdam: a cross-sectional study. *BMJ* 2013;**3**:002505.
21. Kyaw WM, Lee LK, Siong WC, Ping AC, Ang B, Leo YS. Prevalence of and risk factors for MRSA colonization in HIV-positive outpatients in Singapore. *AIDS Res Ther* 2012;**9**:33.
22. Cenizal MJ, Hardy RD, Anderson M, Katz K, Skiest DJ. Prevalence of and risk factors for methicillin-resistant *Staphylococcus aureus* (MRSA) nasal colonization in HIV-infected ambulatory patients. *J Acquir Immune Defic Syndr* 2008;**48**:567-71.
23. Wu CJ, Ko WC, Ho MW, Lin HH, Yang YL, Lin JN, et al. Prevalence of and risk factors for methicillin-resistant *Staphylococcus aureus* colonization among human immunodeficient virus-infected outpatients in Taiwan: oral *Candida* colonization as a comparator. *J Oral Microbiol* 2017;**9**:1322446.
24. Clinical and Laboratories Standards Institute. Performance standards for antimicrobial susceptibility testing; 21st informational supplement. M100-s21, 2011.
25. Huang YC, Ho CF, Chen CJ, Su LH, Lin TY. Comparative molecular analysis of community-associated and healthcare-associated methicillin-resistant *Staphylococcus aureus* isolates from children in

- northern Taiwan. Clin Microbiol Infect 2008; **14**:1167-72.
26. Huang YC, Su LH, Wu TL, Lin TY. Methicillin-resistant *Staphylococcus aureus* nasal carriage in international medical conference attendees. J Microbiol Immunol Infect 2019;52:242-247.
27. Tsai MS, Chen CJ, Lin TY, Huang YC. Nasal methicillin-resistant *Staphylococcus aureus* colonization among otherwise healthy children aged between 2 months and 5 years in northern Taiwan, 2005-2010. J Microbiol Immunol Infect 2018;51:756-62.
28. Oliveira DC, de Lencastre H. Multiplex PCR strategy for rapid identification of structural types and variants of the mec element in methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 2002; **46**: 2155-61.
29. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. J Clin Microbiol 2000; **38**: 1008-15.
30. Chen CB, Chang HC, Huang YC. Nasal methicillin-resistant *Staphylococcus aureus* carriage among intensive care unit hospitalised adult patients in a Taiwanese medical centre: one time-point prevalence, molecular characteristics and risk factors for carriage. J Hosp Infect 2010; **74**: 238-44.
31. Kang YC, Tai WC, Yu CC, Kang JH, Huang YC. Methicillin-resistant *Staphylococcus aureus* nasal carriage among patients receiving hemodialysis in Taiwan: prevalence rate, molecular characterization and de-colonization. BMC Infect Dis 2012; **12**: 284.
32. Lu SY, Chang FY, Cheng CC, Lee KD, Huang YC. Methicillin-resistant *Staphylococcus aureus* nasal colonization among adult patients visiting emergency department in a medical center in Taiwan. PLoS One 2011; **6**: e18620.
33. Wang JT, Liao CH, Fang CT, Chie WC, Lai MS, Lauderdale TL, et al. Prevalence of and risk factors for colonization by methicillin-resistant *Staphylococcus aureus* among adults in community settings in Taiwan. J Clin Microbiol 2009; **47**: 2957-63.
34. Touboul-Lundgren P, Jensen S, Drai J, Lindbæk M. Identification of cultural determinants of antibiotic use cited in primary care in Europe: a mixed research synthesis study of integrated design “Culture is all around us”. BMC Public Health 2015; **15**: 908.
35. Huang YC, Chen CJ, Kuo CC, Lu MC. Emergence, transmission and phylogeny of methicillin-resistant *Staphylococcus aureus* sequence type 8 (USA300) in Taiwan. J Hosp Infect 2018;100:355-8.
36. Tsao FY, Kou HW, Huang YC. Dissemination of methicillin-resistant *Staphylococcus aureus* sequence type 45 among nursing home residents and staff in Taiwan. Clin Microbiol Infect 2015; **21**: 451-8.
37. Zervou FN, Zacharioudakis IM, Ziakas PD, Rich JD, Mylonakis E. Prevalence of and risk factors for methicillin-resistant *Staphylococcus aureus* colonization in HIV infection: a meta-analysis. Clin Infect Dis 2014; **59**: 1302-11.

Figures

Figure 1

Molecular characteristics of methicillin-resistant *Staphylococcus aureus* isolates from 19 HIV-infected patients. All 19 isolates were resistant to penicillin, and susceptible to vancomycin, teicoplanin, linezolid, and doxycycline. Antimicrobial susceptibility tests (AST): black indicates resistance, and grey indicates susceptibility. Abbreviations are as follows: ciprofloxacin (CIP), erythromycin (E), fusidic acid (FA), clindamycin (CC), trimethoprim-sulfamethoxazole (SXT). PFGE, pulsed-field gel electrophoresis. PVL: black indicates that Pantone-Valentine leucocidin genes were detected. SCCmec, staphylococcal cassette chromosome mec; MLST, multilocus sequence typing. Spa types t2365, t3385 and t3485 are variants of t437, while t2849 is the variant of t008.