Analytical Methods for Assessing Retinal Cell Coupling Using Cut-Loading

William E. Myles (✉ will.myles@newcastle.edu.au)

University of Newcastle Australia

Sally Anne McFadden

University of Newcastle Australia

Research Article

Keywords:

Posted Date: October 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-958415/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Electrical coupling between retinal neurons contributes to the functional complexity of visual circuits. “*Cut-loading*” methods allow simultaneous assessment of cell-coupling between multiple retinal cell-types, but existing analysis impedes direct comparison with gold standard direct dye injection techniques. Therefore cut-loading was used to assess coupling strength in a-type horizontal cells in dark-adapted Guinea pig retinae (n=29) using the standard protocol (Method 1) compared with two non-linear methods. Method 1 describes the distance of dye-diffusion (space constant), while Method 2 extracted the coupling coefficient (k_j) and Method 3 measured the diffusion coefficient (De). Dye transfer was measured after one of five diffusion times (1-20 mins), or with a coupling inhibitor, meclofenamic acid (MFA) (50–500 μ M after 20 mins diffusion). Method 1 includes background fluorescence, producing less accurate coupling estimates than measuring the fluorescence of individual cell-soma ($p<0.001$). The space constant (Method 1) increased with diffusion time ($p<0.01$), whereas Methods 2 ($p=0.54$) and 3 ($p=0.63$) produced consistent results across all diffusion times. Method 1 was less sensitive to detecting changes induced by MFA than Methods 2 ($p<0.01$) and 3 ($p<0.01$). Comparatively, Methods 2 and 3 proved more sensitive and generalisable; allowing for detailed assessment of the coupling between different populations of gap-junction linked cell networks.

Introduction

The cells of the retina are extensively linked by intercellular gap-junctions that allow the intercellular passage of ions and small molecules (typically up to 1000 Da), resulting in correlated spiking between pairs or networks of coupled neurons. In the retina, cell coupling has been observed in all major cell types^{1–4} either as homologous coupling amongst one cell-type or as heterologous coupling between multiple cell-types. These coupled networks facilitate the rapid, lateral spread of signals across the retina that complements slower synaptic transmission. Functionally, cell-coupling between retinal neurons modulates receptive field size, facilitates the transition between photopic and scotopic retinal states and contributes to signal averaging, synchrony and motion detection^{5–7}.

Gap-junctions are comprised of two linked hemichannels (connexons) spanning a 2-4nm intercellular gap, each formed by six connexin (Cx) proteins arranged radially surrounding a central pore^{8,9}. Twenty-one mammalian genes coding connexin proteins have been identified. Connexins are classified base on molecular weight, which ranges from 21 to 70kDa¹⁰. The cells of the retina express several connexin isoforms. Retinal pigment epithelial cells express Cx43, cone and rod photoreceptors express Cx36, horizontal cells express Cx50 and Cx57, bipolar and amacrine cells express Cx36 and Cx45, and retinal ganglion cells express Cx30.2, Cx36, Cx43 and Cx45¹¹.

The conductance of a gap-junction is transiently regulated by the phosphorylation of serine residues in the component connexin proteins¹². In the retina, this occurs in response to the release of light-mediated neuromodulators such as dopamine^{13,14} and nitric oxide¹⁵, facilitating in the switching between light and dark processing pathways. For example, heterotypic Cx36/45 gap-junctions between All amacrine cells

and ON-cone bipolar cells are disinhibited in scotopic illumination, allowing for rod signals to utilise the downstream cone pathway².

Given the fundamental importance of coupled networks in determining the fate of signal processing in the retina, understanding their role relies on measurement of coupling after various experimental manipulations. Methods for assessing cell-cell coupling include simultaneous electrical recordings from cell pairs¹⁶, intracellular microinjection of molecular tracers¹⁷, and cut-loading¹⁸. Patch-clamping and microinjection techniques provide an exceptional level of detail, however the techniques themselves are highly technical, requiring the prior identification and recording from individual cells or cell pairs. Cut-loading is a simple alternative whereby the retina is cut with a scalpel blade dipped in molecular tracer and the relative diffusion through coupled cells is measured at a fixed time point.

The most common technique for analysing cut-loaded tissues is to use the space constant from an exponential decay function fitted to the fluorescent intensity of the tissue measured per unit distance from the cut location^{18–20}. This technique has several limitations in that: 1) it does not account for variation in cell density between tissues or sampling location, 2) the equation is fitted to mean fluorescence data containing non-specific background fluorescence, 3) the technique is limited to studying cell populations isolated in a particular layer of the retina e.g., photoreceptors or horizontal cells and 4) space constant values obtained via this method cannot be directly compared to those obtained via other techniques.

Other techniques such as using compartmental analysis to determine coupling coefficients k_j between coupled cells²¹, or calculating the effective diffusion coefficient of molecular tracers through coupled cell networks²² have been employed to measure dye-diffusion from single cells loaded via microinjection techniques. These techniques have also been applied to microinjection experiments using the intracellular tracer NeurobiotinTM, in which only single time-point data is available²³, however, they have not been integrated as part of the cut-loading protocol.

In the present study, we compared three analytical techniques for assessing cell-coupling in cut-loaded mammalian retinæ. Method 1 was a modified version of the standard exponential decay analysis protocol, Method 2 was Zimmerman and Rose's method of calculating the coupling coefficient k_j^{21} , but also adjusted for the geometric arrangement of retinal cells and location of molecular tracer and Method 3 employed Fick's second law of diffusion to calculate the effective diffusion in one-dimension through a network of coupled cells. We compared these three techniques in retinas across five dye incubation times and in retinas incubated for a set incubation time with increasing concentration of the gap-junction inhibitor meclofenamic acid (MFA). We report that the standard analysis protocol may be improved via some simple procedural modifications, however, Methods 2 and 3 were more versatile and provided greater overall sensitivity.

Methods

Animals

Tri-coloured domestic Guinea pigs (*Cavia porcellus*, n = 29) were reared in plastic boxes with wire tops, illuminated overhead by white light emitting diodes diffused through a 3mm thick Perspex screen located 200 mm above the box lid. Lights operated on a 12-hour light-dark cycle at a constant illuminance of 700lx, measured at eye level within the housing box. Room temperature was maintained at 21(±2)°C and food and water was provided *ad libitum*. All experiments and experimental procedures were approved by the Animal Care and Ethics Committee of the University of Newcastle and were conducted in accordance with the Australian code for the care and use of animals for scientific purposes.

Experimental Design

Two experiments were undertaken on retinae freshly extracted from dark adapted guinea pigs between 14 and 28 days of age (Table 1). Experiment 1 used 17 retinae that were cut-loaded with molecular tracer for different incubation times, and the imaged tissues were analysed using three different methods to yield the coupling space constant, the coupling rate constant, or the diffusion coefficient. In Experiment 2, retinae were cut-loaded with molecular tracer and incubated for a single duration. Cell coupling was inhibited with increasing concentrations of the general gap-junction inhibitor meclofenamic acid and imaged tissues were analysed using the same three methods as in Experiment 1. In both experiments, a-type horizontal cells were selected as a model system for analyses. In Experiment 3, the generalisability to a different coupled network was studied in retinae from Experiment 2, by co-labelling for neuronal nitric oxide synthase (nNOS).

Table 1

Experimental Design showing the primary variables manipulated. The *Experimental Light Exposure* refers to the light level during cut-loading at a wavelength of 850nm and was approximately equivalent to total darkness

	Aim	Incubation Time (mins)	No. of retinae (No. of cuts)	MFA Concentration (μM)	Experimental Light Exposure ($\log R^*.\text{Rod}^{-1}.\text{s}^{-1}$)
Experiment 1	Test analysis methods across different incubation times (Horizontal Cells)	1	2 (5)	N/A	-4.5
		3	4 (10)		
		5	4 (12)		
		10	4 (10)		
		20	3 (7)		
Experiment 2	Test analysis methods for variable coupling at a fixed time-point (Horizontal Cells)	20	2 (4)	50	-4.5
			2 (4)	100	
			2 (5)	150	
			2 (5)	200	
			2 (4)	250	
			2 (4)	500	
Experiment 3	Reanalysing tissues from Exp. 2, for non-horizontal cells	Same as Exp. 2			

Procedures

Light Adaptation and Euthanasia

Guinea pigs were adapted to complete darkness for one hour prior to euthanasia. Animals were anaesthetised using gaseous isoflurane (5% in 1.5 L/min O_2) and euthanised via intracardial injection of Pentobarbitone Sodium (Lethabarb®, Virbac Australia Pty Ltd, NSW, Australia). A dim red head torch (light emitting diode (LED) spectral peak at 635nm, max $20\text{uW}.\text{cm}^{-2}$ at 20cm away from tissue, luminance at tissue was $0.5 \log R^*.\text{Rod}^{-1}.\text{s}^{-1}$) was used during euthanasia and dissection in an otherwise dark room. During cut-loading, infrared LEDs (spectral peak 850nm, $-4.5 \log R^*.\text{Rod}^{-1}.\text{s}^{-1}$) and

night vision goggles (Sionyx Aurora Pro) were used to maintain dark-adaptation of the retina during experimentation.

Cut-loading Procedure

The eye was rapidly enucleated and submerged in Ames solution (catalogue: A1420-10X1L, Sigma-Aldrich, MO, USA) at room-temperature (18 to 21°C). Whilst submerged, the cornea and limbus were removed with a circular cut extending along the posterior pars plana (Figure. 1). A small incision was then made along the long ciliary artery to mark the nasal axis using a No.11 scalpel blade. The crystalline lens and vitreous humour were gently removed using No.5 forceps. The remaining eye cup containing the retina, choroid and sclera were transferred into a well plate containing 10mL of Ames solution (36°C) bubbled with carbonox (95% O₂ 5% CO₂).

The retina was separated from the underlying choroid using a blunt dental spatula. The optic nerve joining the retina and sclera was then cut using 3mm curved scissors and the retina orientated with photoreceptor side down was mounted onto 0.22µm pore size Millipore membrane filter paper (catalogue: GSWP04700, Merck & Co. NJ, USA). Tissues were acclimatised for 15 minutes in complete darkness in the bubbled Ames solution. In Experiment 2, the required dilution of meclofenamic acid (MFA) (catalogue: M4531, Merck & Co. NJ, USA) made from stock solution of 100mg/mL MFA sodium salt dissolved in 100% ethanol was added directly to the acclimatisation Ames bath solution and all bath solutions up until fixation.

The retinas were briefly removed from solution and cut along the superior, temporal, and inferior axes with a size 11 scalpel blade that prior to each cut was dipped in 3% the biotin derivative, N-(2-aminoethyl) biotinamide hydrochloride (Neurobiotin™ Tracer, catalogue: SP-1120, Vector Laboratories, CA, USA) diluted in Ames solution. The tissue was returned to the Ames solution and the Neurobiotin™ dye allowed to disperse through the cell network by incubating for either 1, 3, 5, 10 or 20 minutes in Experiment 1, or 20 minutes in Experiment 2 (see Table 1). At the end of each Experiment, tissues were removed from the bubbled bath solution and fixed in 4% paraformaldehyde (4% wt/v, diluted in 0.1M phosphate buffer) at room temperature for 30 minutes. Retinas were washed in 1 x PBS (30 mins) and reacted with Alexa-Fluor 488 conjugated streptavidin (catalogue: S11223, ThermoFisher Scientific, MA, USA, diluted 1:100 in 0.5% Triton-x in PBS) overnight at 4°C. Retinas were washed in 1 x PBS and mounted, photoreceptor side down, onto microscope slides using anti-fade Vectashield aqueous mounting medium (Vector Laboratories, CA, USA, catalogue: H-1000-10).

Immunofluorescence

Retinae from Experiments 1 and 2 were removed from glass slides and placed in PBS, incubated in Triton-x for 30 minutes (1% Triton-x in PBS) and blocked using normal Donkey serum (10% NDS in 0.5% Triton-x, PBS) for one hour. Retinae from Experiments 1 and 2 were then counter labelled with antibodies for Calbindin and Neuronal Nitric Oxide Synthase (nNOS) respectively (See Table 2 for details). Retinae were

washed in PBS (3 x 10 mins) and incubated in secondary antibody solution (Table 2). Retinas were washed in 1 x PBS and mounted, photoreceptor side down, onto microscope slides using anti-fade Vectashield aqueous mounting medium.

Table 2
Antibody reagent list. Both primary and secondary antibodies were diluted in 1% NDS and 0.5% Triton-x in PBS. RT, room temperature.

Antibody	Catalogue	Host Species	Company	Dilution	Time (Temperature)
Neuronal Nitric Oxide Synthase (nNOS)	N7280	Rabbit	Sigma-Aldrich	1:1000	Overnight (4°C)
Calbindin		Rabbit		1:400	Overnight (4°C)
Donkey anti-Rabbit (CY3)	715-165-150	Donkey	Jackson ImmunoResearch Laboratories, Inc.	1:400	1 hour (RT)

Image Collection and Preparation

Images were collected 3mm from the optic nerve head along the superior, temporal and inferior cuts using an Olympus IX91 scanning laser confocal microscope. Each image comprised of 50 optical slices (1µm apart), taken between the outer plexiform later and the retinal ganglion cell layer. Confocal settings were kept consistent across all image acquisition. Composite images from 10 slices were made using the sum-slices z-project function in Fiji (open-source distribution based on ImageJ2 released by National Institutes of Health)²⁴.

Light Intensity Calculations

Reported values of spectral irradiance ($E(\lambda)$, $\mu\text{W}.\text{cm}^{-2}.\text{nm}^{-1}$) of the home-box during rearing, the red LED head torch used during euthanasia and dissection and the infrared LED lighting used during cut-loading experimentation were measured using an Ocean Optics spectrophotometer (USB-4000, Ocean Optics). Illuminance (lx) was calculated based on the Guinea pig photopic spectral sensitivity curve²⁵. Photometric units were converted to total effective photons. $\text{cm}^{-2}.\text{s}^{-1}$ based on the rod spectral sensitivity curve as determined using Govardovskii nomograms²⁶ with $\lambda_{\text{max}} = 496 \text{ nm}$ (Jacobs & Deegan, 1994). This was then converted into photoisomerisations. $\text{rod}^{-1}.\text{s}^{-1}$ ($R^*.\text{rod}^{-1}.\text{s}^{-1}$) assuming a $1.0 \mu\text{m}^2$ effective collection area for each rod²⁷. At low light levels, irradiance ($\mu\text{W}.\text{cm}^{-2}$) was measured using a Newport optical power meter centred at the spectral peak (main unit: 2936-R, sensor: 818-ST2-UV/DB, Newport).

Modelling Neurobiotin™Dye-Transfer

Standard protocol: The mean fluorescence of the image spanning perpendicular from the cut location was measured using the plot profile function in Fiji²⁴.

Modified protocol: As the true cellular concentration of dye was not known, the relative mean fluorescence of each cell was used to gauge the relative concentration of tracer in cells. The mean fluorescence and position of each cell soma was measured in Fiji using the oval tool. The background fluorescence was measured and subtracted from absolute fluorescence measurements. All measurements were then normalised to the maximum fluorescence measured at the cut location. The distance between adjacent horizontal cells (Experiments 1 and 2) and amacrine cells (Experiment 3) were measured from soma centre to soma centre. Forty measurements for each cell-type per retinal image were used to obtain an average.

Data were analysed as both cell fluorescence per absolute distance (µm) from the cut site and as cell fluorescence per cell-separation from the cut site (calculated by dividing absolute distance by the mean soma-soma distance).

Method 1: The decline in fluorescence intensity per distance was fitted with the exponential decay curve below

$$C = C_0 e^{\left(\frac{-x}{\lambda}\right)}$$

1

Where C is the concentration of the cell at distance x from the cut (either in µm or cells), C_0 is the maximum concentration and λ is the space constant (either in µm or cells).

Method 2: Zimmerman and Rose²¹ described the diffusion kinetics of molecular tracers through a chain of 5-7 coupled giant cells of the Chironomus salivary gland using compartment model analysis. For a series of three coupled cells (C_0, C_1, C_2) the movement of tracer into (q_i) and out of the cell (q_s) C_1 can be approximated using the following equations

$$q_R = -k_j(C_1 - C_0)$$

2

$$q_s = -k_j(C_2 - C_1)$$

3

Where k_j is a rate constant of units (distance².time⁻¹). The net movement of tracer through cell C_1 can then be approximated by the combination of these two equations to yield:

$$q_R - q_S = -k_j(C_1 - C_0 - C_2 + C_1) = k_j(C_0 + C_2 - 2C_1)$$

4

The geometric arrangement of HCs differs to that of giant cells used in Zimmerman and Rose's original paper. The arrangement of HCs can be modelled using a triangular lattice, with each cell connecting to the six neighbouring cells (Fig. 2). Therefore, in cut-loading a line of cells will be initially filled with tracer. If we assume no tracer transfer occurs between cells of equal internal dye concentration, then the number of cells C_0 would feed into depends on the angle of the cut with respect to the lattice (Fig. 2). In situation one, each cell at C_1 would receive tracer from two cells at C_0 and would feed into two cells at C_2 (Fig. 2a). In situation two, the cells at C_1 would either receive tracer from one or three cells at C_0 and would feed into either one or three cells at C_2 (Fig. 2b). As these situations alternate in the second example, each layer receives and exports tracer to an average of two cells.

This was incorporated into equation (4) to yield:

$$2q_R - 2q_S = -k_j(2C_1 - 2C_0 - 2C_2 + 2C_1) = 2k_j(C_0 + C_2 - 2C_1)$$

5

Which yields the series:

$\frac{dC_1}{dx} = 2k_j(C_2 - C_1) - k_s(C_1 V_1)$	(6)
$\frac{dC_n}{dx} = 2k_j(C_{n+1} + C_{n-1} - 2C_n) - k_s(C_n V_n)$	etc.

Where k_j is a rate constant describing dye transfer between coupled cells within a network (cells²/s), k_s is sequestration or loss of dye as it passes through the tissue (cells²/s) and V is the relative volume of the cell (set to 1 as all HC assumed to have equal volume). These equations were solved in MATLAB (mathworks) by first fitting a 2-parameter Gaussian curve to the original data to calculate the mean relative fluorescence at discrete intervals (C_n). A solution for k_j was obtained by fitting the concentration series to the ode45 solver, which solved the above equation series (expanded to $n = 1:40$) at a defined time point based on the 4th and 5th Runge-Kutta method.

When k_j was calculated in terms of absolute distance (cm².s⁻¹), the geometric path of molecular tracer was not accounted for and the non-normalised cell-cut distances were used for calculation via Zimmerman and Rose's²¹ original equation written below:

$\frac{dC_1}{dx} = k_j(C_2 - C_1) - k_s(C_1 V_1) \quad (7)$
$\frac{dC_n}{dx} = k_j(C_{n+1} + C_{n-1} - 2C_n) - k_s(C_n V_n) \quad \text{etc}$

Method 3: The diffusion of dye through a homologously coupled cell network (such as horizontal cells) following cut-loading can be described as the diffusion of a substance along one axis using Fick's second law of diffusion²⁸

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2}$$

8

Where C is concentration of the diffusing substance, t is time in seconds, D is the diffusion coefficient ($\text{cm}^2.\text{s}^{-1}$) and x is distance (cm).

One method for modelling dye-diffusion during cut-loading is to consider that at $t = 0$, all molecular tracer is located within the region $-l < x < +l$ describing the boundary of the initial cut made through the retina. Dye transfer then occurs in one-dimension through the coupled cells at the cut boundary, which can be treated as the sum of an infinite number of line sources with diffusion occurring along the x -axis and modelled using the following equation:

$$C = \frac{C_0}{2} \left\{ \text{erf} \left(\frac{l - |x|}{2\sqrt{Dt}} \right) + \text{erf} \left(\frac{l + |x|}{2\sqrt{Dt}} \right) \right\}$$

9

Where:

$$\text{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$$

10

And C_0 is the initial concentration, with initial boundary conditions at $t = 0$:

$$C = C_0, x < l \text{ and } C = 0, x > l$$

Figure 3 shows an example solution for equation (9). with $l = 0.04 \text{ cm}$, $D = 1.6 \times 10^{-6} \text{ cm}^2.\text{s}^{-1}$ and $C_0 = 6.15$.

Statistical Analysis

Data from the five incubation conditions in Experiment 1 were analysed using a 1x5 way ANOVA in SPSS (IBM Statistics 25). In cases where data failed the assumption of homogeneity of variance (Levene's test,

$p < 0.05$) a Brown-Forsythe ANOVA was used instead. Independent samples t-tests were used when comparing fits (Experiment 1) and when comparing coupling rates between cell types (Experiment 3). In Experiment 2, the reduction in cell-coupling due to MFA (100 μM – 500 μM) was expressed as a percentage reduction from the low-dose (50 μM MFA) experimental condition. Paired samples t-tests were then used to compare the normalised effective response calculated by the three analysis methods (Experiment 2). Wilcoxon signed-rank test was used when data was not normally distributed. All tests were conducted as two-tailed tests. Data from Experiment 2 was fitted with the dose-response Hill curve²⁹:

$$\frac{E}{E_{max}} = \frac{1}{1 + \left(\frac{EC_{50}}{[A]} \right)^{-n}} \quad (11)$$

Where E is the magnitude of the change in coupling, E_{max} is the maximum change in coupling, $[A]$ is the concentration of meclofenamic acid, n is the Hill coefficient and EC_{50} is the concentration at which 50% of the maximum response is observed. Fitting was performed in Matlab (Mathworks) using a modified script by Ritchie Smith³⁰. Power analyses were performed in G*Power (Version 3.1, Universität Kiel and Universität Dusseldorf, Germany) for an independent samples t-test (2-tailed).

Results

Modification of the standard analysis protocol

An example cut-loaded retinal image of NeurobiotinTM filled a-type horizontal cells (aHC) spanning 1.5 mm from the cut is presented in Fig. 4a, the associated fluorescence data in Fig. 4b, and the same data translated onto a log scale in Fig. 4c. The exponential decay function, equation (1), produced a poorer fit to the data in regions closest to and furthest away from the cut (red dashed line in Fig. 4c) but overall was well fitted (mean R^2 of fit $\pm$ S.D = 0.89 ± 0.05 , $n = 44$ cuts). An alternative method was based on the mean fluorescence for each aHC soma (Fig. 4c, closed markers) and had the advantage of excluding background fluorescence that was not relevant to the cells of interest and hence gave a more accurate estimate of the dye transfer within the cell network with increasing distance from the cut. The median decay of the exponential fit was significantly less rapid, corresponding to a larger calculated space constant than that produced by fitting the same equation to the total mean image fluorescence (270.3 vs. 212.7 μm , Fig. 4c, $Z = 3.74$, $p < 0.001$). Importantly, data can also be meaningfully normalised for mean soma-soma distance between adjacent horizontal cells (Fig. 4d) to adjust for variations in cell density between retinæ and sampled areas. This alternative technique was employed in Experiments 1 and 2 as Method 1.

Experiment 1. Time-dependant dye transfer through coupled a-type horizontal cells

The extent to which Neurobiotin™ travelled through coupled aHCs increased with incubation duration (Fig. 5a-e). A-type horizontal cells were easily distinguishable from B-type cells based on morphology (Fig. 5f) and co-labelled with calbindin (Fig. 5g).

The fluorescence of cells situated immediately adjacent to the cut was less than that observed deeper into the tissue for all incubation conditions (Fig. 5h), resulting in non-linear dye-transfer characteristics (compare open and closed markers in Fig. 6a, b, c). Additionally, the distance from the cut where reduced fluorescence occurred, increased with longer incubation times (Fig. 5h), suggesting that the cut in the HC network acts as a dye-sink, allowing dye to be lost into the bath solution. To track true coupling coefficients, each equation was only fitted to the curve following the peak in cell fluorescence (closed markers Fig. 6).

Methods 2 and 3 produced good fits of the cell fluorescence data at all timepoints (Fig. 6b and c). Comparatively, Method 1 did not follow the curvature of the concentration series, resulting in a poorer fits at longer incubation times (Fig. 6a, comparison of regression coefficients at one minute and ten minutes, $R^2 = 0.85 \pm 0.02$ vs. 0.93 ± 0.01 , $T(13) = 2.57$, $p = 0.023$). As expected, the space constant (μm) of dye-transfer in aHCs increased significantly with the duration of diffusion (Fig. 6d, $F(4, 38) = 12.17$, $p < 0.001$). This remained true when data were adjusted for cell-cell spacing (Fig. 6g, Brown-Forsythe, $F(4, 20.6) = 40.80$, $p < 0.001$).

Estimation of the coupling coefficient kj of aHCs via Method 2 produced consistent results across all incubation times, when calculated both in terms of absolute distance (Fig. 6e, $F(4,38) = 0.65$, $p = 0.63$) and normalised cell-spaces (Fig. 6h, $F(4,38) = 0.78$, $p = 0.54$). The mean coupling coefficient kj of aHCs across all conditions was $1.9 \times 10^{-6} \text{ cm}^2 \cdot \text{s}^{-1}$ or $0.038 \text{ cells}^2 \cdot \text{s}^{-1}$. As with Method 2, Method 3 produced consistent estimation of the effective diffusion coefficient De across all incubation times, when calculated both in terms of absolute distance (Fig. 6f, $F(4,38) = 0.65$, $p = 0.63$) and normalised cell-spaces (Fig. 6i, $F(4,38) = 0.45$, $p = 0.77$). The mean effective diffusion coefficient De of aHCs across all conditions was $1.86 \times 10^{-6} \text{ cm}^2 \cdot \text{s}^{-1}$ or $0.076 \text{ cells}^2 \cdot \text{s}^{-1}$. Across all methods, adjusting for the mean cell-cell spacing resulted in fewer outliers than if units were presented in terms of absolute distance (compare Fig. 6d,e,f and g,h,i).

Experiment 2. Analysis of variable tracer movement at a fixed time point

All three methods produced adequate dose-response curves, but with deviations in the calculated EC_{50} between approaches (Fig. 6c-h). A large disparity existed between Method 1 and Methods 2 and 3 (all normalised to cell-cell separation) at $150\mu\text{M}$ MFA, with Method 1 displaying a significantly smaller mean reduction in cell-coupling compared to both Method 2 ($41.1 \pm 10.3\%$ vs. 77.1 ± 16.8 , $T(3) = 7.68$, $p = 0.005$) and Method 3 ($41.1 \pm 10.3\%$ vs. $78.1 \pm 12.6\%$, $T(3) = 8.41$, $p = 0.004$). This was consistent regardless of whether or not distance was normalised to cell-spacing (Table 3, Method 1 in μm). When the extent of coupling with $150\mu\text{M}$ MFA concentration was compared to the baseline MFA concentration of $50\mu\text{M}$, Method 1 displayed a slightly lower statistical power than Methods 2 and 3 (Table 3). At high

concentrations of MFA (250µM), the reduction in coupling compared to the baseline condition of 50 µM MFA when estimated using Method 1 (normalised to cell-cell separation) produced a significantly lower maximum reduction in cell-coupling than Methods 2 (60.6 ± 3.9% vs. 93.3 ± 1.1%, $T(3) = 8.56$, $p = 0.003$) and Method 3 (60.6 ± 3.9% vs. 94.3 ± 1.4, $T(3) = 8.40$, $p = 0.004$). This difference was also evident when data was presented as absolute distance (Table 3, Method 1 in µm).

Table 3

Reduction in cell coupling induced by the coupling inhibitor MFA, presented as a percentage reduction from a low-dose baseline concentration. p -values indicate the statistical difference in calculated normalised effect sizes compared to Method 1.

		150µM Vs. 50 µM MFA			250µM Vs. 50 µM MFA		
	Units	Normalised Effective Reduction (%)	Statistical Power	p -value	Normalised Effective Reduction (%)	Statistical Power	p -value
Method 1 (λ)	µm	36.7 ± 9.5	0.893		48.4 ± 5.0	1.000	
	cells	41.1 ± 10.3	0.911		60.6 ± 3.9	1.000	
Method 2 (kj)	cm ² .s ⁻¹	74.5 ± 10.1	>0.999	0.004	88.9 ± 1.3	1.000	0.011
	cells ² .s ⁻¹	77.1 ± 16.8	>0.999	0.005	93.3 ± 1.1	1.000	0.003
Method 3 (De)	cm ² .s ⁻¹	77.5 ± 12.6	0.983	0.003	90.1 ± 1.8	>0.999	0.012
	cells ² .s ⁻¹	78.1 ± 12.6	0.966	0.004	94.3 ± 1.4	>0.999	0.004

Experiment 3. Identification of coupling rates in different retinal cells subtypes

Amacrine cells expressing the enzyme neuronal nitric oxide synthase (nNOS) were also identified in tissues from Experiment 2 (Fig. 8a). Method 2 and Method 3 were used to measure the coupling coefficient kj (cells².s⁻¹) and non-normalised diffusion coefficient De (cm².s⁻¹) for both type-1 and displaced nNOS amacrine cells (Fig. 8a). The rate of dye-transfer across each cell-cell connection (kj) was significantly faster between displaced nNOS amacrine cells compared to type-1 nNOS amacrine cells (Fig. 8g, $7.91 \pm 1.37 \times 10^{-4}$ vs. $2.69 \pm 0.44 \times 10^{-4}$ cells².s⁻¹, $T(21) = 3.773$, $p = 0.001$). However, the mean distance between type-1 nNOS cells was significantly larger than displaced nNOS cells (Fig. 8f, 177.4 ± 7.8 vs. 74.5 ± 4.5 µm, $T(21) = 10.72$, $p < 0.001$). As such, the overall rate of dye diffusion through the retina (De) was significantly faster in the type-1 nNOS cell network than through the displaced nNOS amacrine cell network (Fig. 8h, $3.11 \pm 0.41 \times 10^{-4}$ vs. $1.73 \pm 0.18 \times 10^{-4}$ cm².s⁻¹, $T(21) = 3.00$, $p = 0.007$).

Discussion

In the present study, we demonstrate the versatility of the cut-loading method and highlight several important considerations that should be taken into account when analysing cut-loading data.

Although the standard analysis protocol has several limitations, many of these can be addressed with some simple modifications to the analysis protocol. The variation arising from changes in cell-density between tissues/sampling area (limitation 1) can be addressed by dividing the absolute distance by the mean cell-cell distance, resulting in the units of the space constant being cells as opposed to μm . Limitation 2 related to non-specific fluorescence can be improved by measuring the mean soma fluorescence of each cell, rather than the mean fluorescence of an area. This modification also addresses the disproportionately high fluorescence measured at the cut by the standard technique, resulting from cells other than HCs filling with dye. By measuring the soma fluorescence, the data may still be normalised based on the fluorescence of cells located at the cut, but this data can then be excluded from the fit. Although these simple modifications improve the validity of using the space constant as a means of describing cell-coupling in cut-loading, they do not allow for direct comparison with studies involving dye-injection.

Methods 2 and 3 described the decay in cell fluorescence more accurately than Method 1 and produced consistent estimations of kj and De respectively when applied across a range of incubation times. Methods 2 and 3 were also more sensitive to smaller changes in cell-coupling at a fixed time point than Method 1, producing approximately twice the effect size without affecting high statistical power in the 150 μM MFA group and approximately 1.5 times the maximum effect size at 250 μM MFA (Table 3).

Comparison of coupling (kj) and diffusion (De) coefficients between techniques

Our estimation of the effective diffusion coefficient De of NeurobiotinTM through coupled HCs was similar to that previously measured in Rabbit retinae ($1.86 \times 10^{-6} \text{ cm}^2.\text{s}^{-1}$ vs. $2.56 \times 10^{-6} \text{ cm}^2.\text{s}^{-1}$)²³. However, our measured value for the coupling coefficient kj was substantially higher ($0.038 \text{ cells}^2.\text{s}^{-1}$ vs. $0.003 \text{ cells}^2.\text{s}^{-1}$). One explanation is that the mathematical model used in equation (6), presumes no rise to maximum fluorescence in the first cell of the coupled network, as this duration could not be known from fixed time-point analysis such as that performed in cut-loading. This is opposed to the method used by Mills and Massey, where a continuous injection of a known rate and duration for c_1 is subsequently incorporated into the analysis²³. However, when one of their example Rabbit traces loaded via microinjection (30 minute incubation) was reanalysed using equation (6) (presuming no rise to maximum, with no adjustment for cell geometry), kj values remained four times higher in Guinea pig HCs ($0.038 \text{ cells}^2.\text{s}^{-1}$ vs. $0.0095 \text{ cells}^2.\text{s}^{-1}$). However, by adjusting for the geometric path of dye-travel through a triangular lattice from a single cell-source, equation (12)³¹, the coupling coefficient kj from cut-loaded Guinea pig HCs was similar to estimates from Rabbit HCs investigated using single cell micro-injection ($0.038 \text{ cells}^2.\text{s}^{-1}$ vs. $0.019 \text{ cells}^2.\text{s}^{-1}$).

$\frac{dC_1}{dx} = 6k_j(C_2 - C_1) - k_s(C_1 V_1) \quad (12)$
$\frac{dC_n}{dx} = k_j(2.5C_{n+1} + 1.5C_{n-1} - 4C_n) - k_s(C_n V_n)$

Use cases of each methodological approach

As both Methods 2 and 3 performed well across our testing, the application of either method would depend on the desired information. When the cell-density and tiling geometry are accounted for, as in equation (6) of Method 2, the rate of dye-transfer across each cell-cell connection (k_j) may be estimated. This method therefore allows for the simultaneous comparison of the relative permeability of gap-junctions present between different cell-types. Comparatively, Method 3 calculates the rate of passive diffusion of tracer through a cell network, away from the initial cut. This is of use when the absolute spread of a signal/tracer across the retina through a specific cell network is of interest. This may be normalised to the cell-cell separation if the relative decay of a signal across adjoining cells is of interest, otherwise, the absolute distance of the signal/tracer spread across the retina may be obtained from non-normalised data. For example, the rate of dye-transfer between individual displaced nNOS amacrine cells (k_j) is faster than type-1 nNOS amacrine cells, however as displaced cells are more densely distributed in the retina, the absolute distance that the signal will reach in a given time (De) is greater in the less dense, type-1 nNOS cell network.

Measuring specific cell-types in highly heterogeneous cell layers

A limitation of the cut-loading method, is that it cannot be used to delineate the path of dye-travel, as information detailing whether cells are homologously or heterologously coupled is not attainable. However, the advantage of the cut-loading technique is that it may be expanded to measure cells interspersed in the inner nuclear and retinal ganglion cell layers by counter labelling cut-loaded tissues with antibodies of specific cell markers and measuring the fluorescence of individual cell soma. This technique is limited by the number of compatible fluorophores for secondary antibodies but offers considerably advantage over dye injected into an individual cell as coupling coefficients can be simultaneously assessed for a range of different cell types in the same tissue after a particular experimental manipulation.

Conclusion

Cell-coupling is important to many visual circuits within the retina. However, the functional significance of cell-coupling between many retinal cell-types as well as how these circuits are modulated or otherwise altered in disease, remains to be established. In spite of its simplicity, cut-loading enables cell-coupling data to be obtained simultaneously through different coupled networks across all layers of the retina. By employing the techniques described here, this analysis can be expanded to include assessments of the

relative coupling coefficient (k_j) and effective diffusion coefficient (De) of molecular tracer through the retina.

Declarations

Acknowledgments

This research was funded by the Hunter Medical Research Institute (Grant ID: G1801044) and supported by the Lions Club, Newcastle.

Competing interests

The authors declare no competing interests.

Author Contributions

W.M and S.M conceived the study, W.M collected and analysed data, W.M and S.M wrote the main manuscript, W.M prepared all figures. All authors reviewed the manuscript.

Data Availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

1. Roy, K., Kumar, S. & Bloomfield, S. A. Gap junctional coupling between retinal amacrine and ganglion cells underlies coherent activity integral to global object perception. *Proceedings of the National Academy of Sciences* **114**, E10484–E10493, doi:10.1073/pnas.1708261114 (2017).
2. O'Brien, J. J. & Massey, S. C. Connexin36 Forms Gap Junctions Between All Amacrine Cells and Diffuse ON Cone Bipolar Cells in the Primate Retina. *Invest Ophthalmol Vis Sci*, **46**, 2333–2333 (2005).
3. Dong, C. J. & McReynolds, J. S. The relationship between light, dopamine release and horizontal cell coupling in the mudpuppy retina. *J Physiol*, **440**, 291–309 <https://doi.org/10.1113/jphysiol.1991.sp018709> (1991).
4. O'Brien, J. J., Chen, X., MacLeish, P. R., O'Brien, J. & Massey, S. C. Photoreceptor Coupling Mediated by Connexin36 in the Primate Retina. *The Journal of Neuroscience*, **32**, 4675–4687 <https://doi.org/10.1523/jneurosci.4749-11.2012> (2012).
5. Trenholm, S. & Awatramani, G. B. in *Webvision: The Organization of the Retina and Visual System* (eds H. Kolb, E. Fernandez, & R. Nelson) (University of Utah Health Sciences Center, Copyright: © 2021 Webvision. 2019).

6. Bloomfield, S. A. & Völgyi, B. The diverse functional roles and regulation of neuronal gap junctions in the retina. *Nature Reviews Neuroscience*, **10**, 495–506 <https://doi.org/10.1038/nrn2636> (2009).
7. Murphy-Baum, B. L. & Awatramani, G. B. An Old Neuron Learns New Tricks: Redefining Motion Processing in the Primate Retina., **97**, 1205–1207 <https://doi.org/10.1016/j.neuron.2018.03.007> (2018).
8. Goodenough, D. A. & Revel, J. P. A fine structural analysis of intercellular junctions in the mouse liver. *J Cell Biol*, **45**, 272–290 <https://doi.org/10.1083/jcb.45.2.272> (1970).
9. Perkins, G., Goodenough, D. & Sosinsky, G. Three-dimensional structure of the gap junction connexon. *Biophys J*, **72**, 533–544 [https://doi.org/10.1016/s0006-3495\(97\)78693-4](https://doi.org/10.1016/s0006-3495(97)78693-4) (1997).
10. Mikalsen, S. O., Kongsstovu, S. & Tausen, M. í & Connexins during 500 Million Years—From Cyclostomes to Mammals. *International Journal of Molecular Sciences* **22**, 1584(2021).
11. Roy, S., Kim, D. & Lim, R. Cell-cell communication in diabetic retinopathy. *Vision. Res*, **139**, 115–122 <https://doi.org/10.1016/j.visres.2017.04.014> (2017).
12. Urschel, S. *et al.* Protein kinase A-mediated phosphorylation of connexin36 in mouse retina results in decreased gap junctional communication between All amacrine cells. *J Biol Chem*, **281**, 33163–33171 <https://doi.org/10.1074/jbc.M606396200> (2006).
13. Kothmann, W. W., Massey, S. C. & O'Brien, J. Dopamine-stimulated dephosphorylation of connexin 36 mediates All amacrine cell uncoupling. *J Neurosci*, **29**, 14903–14911 <https://doi.org/10.1523/jneurosci.3436-09.2009> (2009).
14. Baldrige, W. H., Vaney, D. I. & Weiler, R. The modulation of intercellular coupling in the retina. *Seminars in Cell & Developmental Biology*, **9**, 311–318 <https://doi.org/10.1006/scdb.1998.0235> (1998).
15. Miyachi, E., Murakami, M. & Nakaki, T. Arginine blocks gap junctions between retinal horizontal cells., **1**, 107–110 <https://doi.org/10.1097/00001756-199010000-00006> (1990).
16. MacVicar, B. A. & Dudek, F. E. Electrotonic coupling between pyramidal cells: a direct demonstration in rat hippocampal slices., **213**, 782–785 <https://doi.org/10.1126/science.6266013> (1981).
17. Pollack, G. H. Intercellular coupling in the atrioventricular node and other tissues of the rabbit heart. *The Journal of physiology*, **255**, 275–298 <https://doi.org/10.1113/jphysiol.1976.sp011280> (1976).
18. Ribelayga, C., Cao, Y. & Mangel, S. C. The circadian clock in the retina controls rod-cone coupling., **59**, 790–801 <https://doi.org/10.1016/j.neuron.2008.07.017> (2008).
19. Choi, H. J., Ribelayga, C. P. & Mangel, S. C. Cut-loading: a useful tool for examining the extent of gap junction tracer coupling between retinal neurons. *J Vis Exp*, **3180**, <https://doi.org/10.3791/3180> (2012).
20. Hou, B. *et al.* Homeostatic Plasticity Mediated by Rod-Cone Gap Junction Coupling in Retinal Degenerative Dystrophic RCS Rats. *Front Cell Neurosci*, **11**, 98–98 <https://doi.org/10.3389/fncel.2017.00098> (2017).

21. Zimmerman, A. L. & Rose, B. Permeability properties of cell-to-cell channels: Kinetics of fluorescent tracer diffusion through a cell junction. *The Journal of Membrane Biology*, **84**, 269–283 <https://doi.org/10.1007/BF01871390> (1985).
22. Safranyos, R. G. & Caveney, S. Rates of diffusion of fluorescent molecules via cell-to-cell membrane channels in a developing tissue. *J Cell Biol*, **100**, 736–747 <https://doi.org/10.1083/jcb.100.3.736> (1985).
23. Mills, S. & Massey, S. The kinetics of tracer movement through homologous gap junctions in the rabbit retina. *Visual neuroscience*, **15**, 765–777 <https://doi.org/10.1017/S0952523898154159> (1998).
24. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods*, **9**, 676–682 <https://doi.org/10.1038/nmeth.2019> (2012).
25. Jacobs, G. H. & Deegan, J. F. 2 Spectral sensitivity, photopigments, and color vision in the guinea pig (*Cavia porcellus*). *Behav Neurosci*, **108**, 993–1004 <https://doi.org/10.1037//0735-7044.108.5.993> (1994).
26. Govardovskii, V. I., Fyhrquist, N., Reuter, T. O. M., Kuzmin, D. G. & Donner, K. In search of the visual pigment template. *Vis. Neurosci*, **17**, 509–528 <https://doi.org/10.1017/S0952523800174036> (2000).
27. Li, P. H., Verweij, J., Long, J. H. & Schnapf, J. L. Gap-junctional coupling of mammalian rod photoreceptors and its effect on visual detection. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, **32**, 3552–3562 <https://doi.org/10.1523/JNEUROSCI.2144-11.2012> (2012).
28. Crank, J. & Crank, E. P. J. *The Mathematics of Diffusion* (Clarendon Press, 1979).
29. Di Veroli, G. Y. *et al.* An automated fitting procedure and software for dose-response curves with multiphasic features. *Sci. Rep*, **5**, 14701 <https://doi.org/10.1038/srep14701> (2015).
30. doseResponse v. 1.2.0.0 (MATLAB Central File Exchange, <https://au.mathworks.com/matlabcentral/fileexchange/33604-doseresponse>, 2021).
31. Xia, X. B. & Mills, S. L. Gap junctional regulatory mechanisms in the All amacrine cell of the rabbit retina. *Visual neuroscience*, **21**, 791–805 <https://doi.org/10.1017/S0952523804215127> (2004).

Figures

Figure 1

Flow chart illustrating the key steps in the cut-loading procedure. Figure made using the online application BioRender.com.

Figure 2

Geometric model of dye-transfer through coupled horizontal cells. (a) Cells located in C1 (*) receive dye (q_r) from two cells in C0 and individually feed two cells (q_s) in C2. (b) Cells located in C1 either receive input from three cells in C0 and feed one in C2 (cell a), or they receive input from one cell in C0 and feed three cells in C2 (cell b).

Figure 3

The diffusion of molecular tracer from the cut location, modelled according to equation (9). At $t = 0$, all molecular tracer is located within the region $-l < x < l$ surrounding the x-axis. Concentration of NeurobiotinTM is represented as relative fluorescent intensity along the z-axis.

Figure 4

(a) Example image taken in outer portion of the inner nuclear layer of a Guinea pig retina cut-loaded with NeurobiotinTM. (b) The fluorescent intensity of the whole image measured along the horizontal axis extended away from the cut. (c) Semi-log plot of fluorescent intensity data from (b) and fluorescent intensity of individual horizontal cells both fitted with an exponential decay function, equation (1). (d) Fluorescent intensity of individual horizontal cells adjusted for the mean soma-soma cell distance.

Figure 5

(a-e) The extent of dye-transfer through coupled a-type horizontal cells (aHCs) after different incubation times between 1 and 20 min. Images represent the average of 5, 1 μ m thick optical slices through the outer inner nuclear layer created using the z-project function in Fiji (see methods). The cut-edge is located on the left-hand side of each image. (f) Close up of morphologically distinct horizontal cell subtypes: axon-bearing ('B-type', open arrow) and axon-less ('A-type', closed arrow) HCs. Image created from cut-loaded retina using the temporal colour-code function in Fiji for optical slices spanning the inner nuclear layer (see methods). (g) Cut-loaded HCs labelled with NeurobiotinTM (green) and co-labelled with Calbindin (red). (h) Relative mean cell fluorescence of aHCs per cell from the cut. Curves produced by

fitting 3-parameter polynomial fits to mean aHC data for each incubation time, fluorescence at the cut was excluded when creating the fit. The peak cell fluorescence for each curve is indicated by black stars. Scale bars signify 100 μ m.

Figure 6

Comparison of three analytical techniques for assessing the extent of a-type horizontal (aHCs) cell coupling in cut-loaded Guinea pig retinas with increasing incubation times. Column 1 (a, d, g) contains cell fluorescence data analysed using Method 1. Column 2 (b, e, h) shows the same data analysed using Method 2 and column 3 (c, f, i) shows the same data analysed using Method 3. The first row (a, b, c) contains example fits for cell fluorescence data collected per each cell separation following four incubation times, fitted with (a) equation (1), (b) equation (6) and (c) equation (9) corresponding to Methods 1, 2 and 3 respectively. Row 2 (d, e, f) contains the mean $\pm$ SE (black marker, black error bars) and individual measurements (grey markers) for each measure of cell-coupling obtained via: (d) Method 1 (e) Method 2 and (f) Method 3. Row 3 (g, h, i) contains the mean $\pm$ SE (black marker, black error bars) and individual measurements (grey markers) for each measure of cell-coupling normalised to the mean

cell-cell spacing between adjacent aHCs, obtained via: (g) Method 1 (h) Method 2 and (i) Method 3. Crosses indicate outliers defined as data points greater than 1.5 standard deviations from the mean and are not included in calculated mean.

Figure 7

Comparison of three analytical methods for describing dose-response data for a-type horizontal cell (aHC) coupling and meclofenamic acid (MFA) concentration in vitro. The relative diffusion of NeurobiotinTM through HCs in tissue incubated in (a) 50 μM MFA and (b) 250 μM MFA. Column 1 (c, f) contains data analysed using Method 1, column 2 (d, g) shows the same data analysed using Method 2 and column 3 (c, f, i) shows the same data analysed using Method 3. The first row of plots (c, d, e) contains the mean $\pm$ SE (black marker, black error bars) and individual measurements (grey markers) for each measure of cell-coupling obtained via Methods: (c) 1, (d) 2 and (e) 3. Row 2 (f, g, h) contains the

mean $\pm$ SE (black marker, black error bars) and individual measurements (grey markers) for each measure of cell-coupling normalised to the mean cell-cell spacing between adjacent aHCs, obtained via Methods (f)1, (g) 2 and (h) 3.

Figure 8

Identification of amacrine cells expressing neuronal nitric oxide synthase loaded with molecular tracer via cut-loading. (a) Image created from z-stack slices spanning from the inner nuclear layer to the ganglion

cell layer, flattened using the z-project function in Fiji. Open arrows, type-1 nNOS amacrine cells; solid arrows, displaced nNOS amacrine cells. (b, c) Vertical retinal sections labelled for nNOS (red) and DAPI (blue). (b) Open arrows highlight type-1 nNOS amacrine cells located in the inner nuclear layer (INL), (c) solid white arrows highlight displaced nNOS amacrine cells located in the ganglion cell layer (GCL). (d,e) Fluorescent intensity of NeurobiotinTM in cut-loaded type -1 and displaced nNOS amacrine cells analysed using (d) Method 1 (fitted with equation 6); or (e) Method 3 (fitted with equation 9). (f-g) Comparison of nNOS type-1 and nNOS displaced amacrine cells in their mean: (f) cell density, (g) cell-coupling coefficients k_j , and (h) relative diffusion coefficients. Error bars are SEM. * $p < 0.05$.