

“I want to live, but...” The desire to live and its physical, psychological, and social factors among advanced cancer patients: Evidence from the APPROACH study in India

Anjum S Khan Joad

Bhagwan Mahaveer Cancer Hospital and Research Centre

Arati Hota

Bhagwan Mahaveer Cancer Hospital and Research Centre

Pratima Agarwal

Bhagwan Mahaveer Cancer Hospital and Research Centre

Krimal Patel

Bhagwan Mahaveer Cancer Hospital and Research Centre

Kinjal Patel

Bhagwan Mahaveer Cancer Hospital and Research Centre

Jyotika Puri

Duke-NUS Medical School

Soye Shin ([✉ syshin@duke-nus.edu.sg](mailto:syshin@duke-nus.edu.sg))

Duke-NUS Medical School

Research Article

Keywords: Advanced cancer patients, palliative care, desire to live, desire for hastened death, India, socio-economic status, pain severity, psychological distress

Posted Date: October 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-958916/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

The limited access to palliative care resources along with the social stigma around cancer largely explains the poor quality of life (QoL) of Indian advanced cancer patients. As advanced cancer patients with poor QoL often harbour a desire for hastened death (DHD), it is imperative to understand factors affecting DHD, or the desire to live (DTL) among advanced cancer patients in India. We aim to examine the relationship between DTL and physical, psychological, social and spiritual factors measuring patients' QoL alongside their awareness of their late cancer stage.

Methods

We surveyed 200 patients from a tertiary cancer hospital in India to collect their DTL, awareness of cancer stage, demographic characteristics, and standardized measures for patients' QoL. We used a linear probability regression model to quantify the association between these factors and patients' DTL among the final sample of 192 patients with no missing information for the variables of interest.

Results

Among the various domains affecting cancer patients' QoL, we found that the pain severity score (ranging from 0 to 10) and psychological distress score (ranging from 0 to 42) are negatively associated with the DTL. One point increase in each score reduced the DTL by 2.2% ($p < 0.05$) and 0.7% ($p < 0.05$), respectively. Our results also showed that patients whose perceived socio-economic status (SES) is poor have a 16% ($p < 0.05$) lower probability of DTL, compared to those with higher SES (lower middle class, upper middle class, and wealthy). Controlling for caste, religion, gender, age, marital status and years of education, we found psychological distress is statistically higher among patients belonging to this bottom SES.

Conclusions

We found that pain severity, psychological distress and perceived low SES are negatively associated with the desire to live in advanced cancer patients. Future research should focus on developing interventions to improve physical pain and psychological distress, particularly for patients who are socially and economically disadvantaged.

Introduction

He who has a "why" to live can deal with almost any "how". -Nietzsche^{(1)(p.IX)}

According to Global Cancer Data from GLOBOCAN 2018 (2), an interactive web platform that consolidates worldwide cancer statistics, in India, of 2.2 million cancer patients, 784,000 die annually. This number has doubled since 1990 (3). The age-standardised mortality: incidence ratio for India is 0.68, which is higher than that of very high human development index (HDI) countries (0.38) and high HDI countries (0.57) (4). This is a big concern as India has a relatively younger population than those countries and hence is expected to present lower age-adjusted mortality indices. Part of this mismatch can be attributed to better screening practices in high HDI nations, but a large part of it is because of inequitable access to quality cancer care, lack of affordability in care and delayed diagnosis of cancer due to stigma related to the illness and a lack of awareness of its symptoms (5).

Various myths and misconceptions regarding cancer cause delays in patients seeking medical care. A survey of 95 cancer patients in Delhi showed that one third of the patients believed malignancies can be detected in early stages and that they are curable (6). Most of the surveyed patients were reported to hold fatalistic views about the outcome of cancer and 60% felt that their family and society were discriminating against them because of their cancer diagnosis. As a result, the average time taken by patients to report to a doctor after suspecting their disease was 2 years (6). By that time, their cancer has often developed to advanced stages (7–9). Moreover, limited access to palliative care and comprehensive cancer treatment services contributes to the high mortality rate from cancer in India. Less than 4% of patients with serious health-related suffering have access to opioid morphine-equivalents (10). Also, more than 75 percent of cancer expenditures in India are covered out-of-pocket, which indicates that access to affordable and equitable cancer care in India still remains a major challenge (11).

Combined with the prevalence of advanced stage cancer at the first diagnosis, the limited access to cancer and pain treatment resources along with the social stigma around cancer largely explains poor quality of life (QoL) of Indian advanced cancer patients. There is comprehensive literature showing that cancer patients with poor QoL often harbour a desire for hastened death (DHD) (12–15). The frequency of reporting DHD among advanced cancer patients is noticeably higher than that among patients in nonpalliative stages of cancer (16), which is a significant concern for healthcare professionals and stakeholders (17). Despite the urgency of the issue, there is limited research on factors that may increase DHD in advanced cancer patients, particularly in the context of India.

The aim of the study is to fill this knowledge gap by identifying contributors to the desire to live (DTL), the other side of DHD, with advanced cancer patients in a tertiary cancer centre in Jaipur, India. Previous studies, mostly conducted in developed countries, provided evidence that DTL (or, DHD) was associated with various physical, psychological, spiritual, social or interpersonal and spiritual factors. For instance, one study reported that as many as 77% terminally ill patients desired to hasten death because of experiencing moderate to severe physical pain (18). The psychological and emotional distress of cancer patients is also negatively associated with their DTL. Individuals with advanced-stage cancer reported persistent depressive symptoms more frequently (19) and both depression and hopelessness independently predicted DHD of metastatic cancer patients (20). In particular, patients with low socio-economic status (SES) tend to reveal a higher prevalence of depression (21). In addition, experiencing low

social support hinders people's ability to find meaning in their lives, which contributes to high DHD. Lastly, spiritual well-being or belief in a higher power has been established as an important protective factor that keeps depression and hopelessness at bay and builds some tolerance to physical symptoms (22).

Guided by the evidence, we collected patients' awareness of their advanced cancer along with multiple indexes measuring their QoL, in terms of physical (e.g., pain severity), psychological (e.g., self-blame), spiritual (e.g., faith), and social support (e.g., social stigma) dimensions, and examined the association of DTL and these variables. We hypothesize that patients who suffer from severe physical pain, a high level of anxiety & depression, and have low spiritual well-being and social/family support are likely to have a lower DTL. We also hypothesize that patients from a socially and economically disadvantaged group are likely to present a lower DTL, holding other factors constant.

Focusing on the second hypothesis, we further seek to investigate whether there are significant differences in the QoL factors, by patients' perceived socio-economic strata. We hypothesize that patients who reported to have lower SES would experience higher social stigma, psychological distress, and lower interpersonal support. Testing this hypothesis is especially important in a country as diverse as India with multifaceted segmentation based on caste, class, religion, regional economic advancement, and language. For instance, at the regional level, development of palliative care and pain management services is uneven in that there is greater provision of the services in the southern parts than the north (23). The unequal access to these services is consistent with regional economic gaps that five states of south India account for about 30% of India's Gross Domestic Product (GDP) while eight states of north India account for only 2.8% (24). We expect QoL disparities by SES exist at the household/individual level as well.

To the authors' best knowledge, there are no prior studies comprehensively examining the association between DTL and various QoL factors among advanced cancer patients in India. Previous work with cancer patients in India has rather focused on the prevalence of suicidal ideation (14, 15). We expect findings from this study to deepen our understanding of various factors that may affect DTL in a regional cancer centre in India. This would provide insights into advanced cancer patient care not only in India but in many other developing countries where palliative and pain care services are limited.

Methods

Setting and participants

This study was part of a multi-country cross-sectional survey titled "Asian Patient Perspectives Regarding Oncology Awareness, Care and Health (APPROACH)" to assess gaps in care received by advanced cancer patients seeking care at major public hospitals in low- and middle-income countries in Asia. The site in the study is Bhagwan Mahaveer Cancer Hospital and Research Centre (BMCHRC) in Jaipur, India. Established in 1997, BMCHRC is a specialty hospital offering cancer prevention, treatment, research, and education-related services, especially for economically disadvantaged groups of the population by

subsidizing treatments or in some cases, providing it free of cost. Approximately 14,000 cancer patients visit the hospital for treatment from Rajasthan (state where Jaipur is located) and neighbouring states every year. A large proportion of the patients are from low socio-economic strata and present with advanced stages of cancer when they first consult with physicians.

We recruited inpatients and outpatients from the departments of medical oncology and radiotherapy and palliative care from 7th March 2017 to 16th August 2018. Inclusion criteria was patients who were over the age of 21, diagnosed with stage 4 solid cancer, and aware of cancer diagnosis (not necessarily its stage). Only patients with solid cancer were included because those who were diagnosed with cancers like lymphoma or leukemia progress differently and have a different staging system which would create inconsistencies in analysis and interpretation.

The study was approved by the Institutional Ethics Committee of Bhagwan Mahaveer Cancer Hospital & Research Centre (Ref No: BMH/ 2016/89) and the National University of Singapore (Ref No: B-15-319).

Figure 1 presents the patient recruitment diagram. Targeting a sample of 200 patients, the research team screened 210 medical records for the eligibility criteria and identified 209 eligible patients. Among these eligible patients that our trained interviewers approached, 8 patients refused to participate in the study. From 201 patients who consented to the study, 200 completed the interview, which results in a response rate of 99.5 %.

Survey questionnaire

In consultation with oncologists, we developed a questionnaire and validated instruments to measure patients' QoL in physical, psychological & spiritual, and social domains. We also collected patients' demographic information that includes gender, age, education level, marital status, SES, caste (if any), religion, ethnicity as well as their awareness of their advanced cancer stage. All questions in English were translated by professional translators into Hindi, which is commonly spoken at the site, according to a strict translation protocol for interview, and patients' answers were translated back to English. Cross-validation between the original and back translated versions were made to ensure the consistency. More details about the dependent and independent variables (predictors) of interest are discussed below.

a) Outcome variable: DTL

In the survey, we asked them a single question enquiring if they ever wished their life would end sooner. They could answer it with yes, no, or not sure. The source of this question was the first item of the Desire for Death Rating Scale (18), which underwent pilot testing on 10 patients who met the eligibility criteria before being included in the survey. The reason for choosing a single-item measure as opposed to a multi-item validated scale was because questions discerning the desire for an early death were deemed sensitive and a mental health professional was not always available to alleviate any potential distress participants might experience. To create a dummy variable for DTL, we reversed interpretation of the responses for this negatively-phrased question in a way that patients who answered no/not sure were

considered to have DTL. On the contrary, we assigned a value of zero for patients who responded to ever wish to end their life sooner.

b) Physical factors: Pain severity

To get a holistic picture of the severity of pain experienced by participants, we elicited patients' pain experience through four items from Hindi version of the Brief Pain Inventory- Short Form (BPI-SF) (25), at its different levels (worst, least, and on average) and/or in different times (the last 24 hours and the time of responding to the questionnaire). Specifically, participants were asked to rate their pain on a scale of 0 to 10 wherein they can only choose whole numbers (i.e., 0, 1, 2 etc.). We then computed the mean value as a measure of pain severity for patients who responded to at least three out of the four items and left missing otherwise. The number of the latter case is nil. The range of possible values is 0 to 10.

c) Psychological and Spiritual factors

Psychological Distress

We assessed participants' psychological distress using the Hospital Anxiety and Depression Scale (HADS), which consists of two subscales for measuring anxiety and depression with 7 items each. Each item has 3 levels (1 to 3) with a higher value indicating higher levels of distress (i.e., anxiety or depression). We reverse scored positively phrased statements and summed these item responses to obtain HADS-anxiety and HADS-depression, which ranges from 0 to 21, respectively. For each subscale, HADS defines scores from 0-7 as "Normal", 8-10 as "Borderline Abnormal" or "being suggestive of the presence of anxiety/depression", and 11-21 as "Abnormal" or "indicative of the probable presence of the mood disorder" (26). We also combined these two subscales for HADS-total (0 to 42).

Self-Blame

Participants also answered two questions about how much they blamed themselves for their diagnosis due to their behaviour (e.g., smoking, drinking etc.), which was termed behavioural self-blame (BSB), and due to personal characteristics that they perceive as in-built traits which are not subject to change or susceptible to their control, which was termed characterological self-blame (CSB). Possible responses to these questions were 1 (not at all) to 4 (completely), respectively.

Spiritual well-being: Using the 12-item Functional Assessment of Chronic Illness Therapy–Spiritual Well-Being Scale (FACIT–Sp)(27, 28), we measured spiritual well-being of participants in two domains: spiritual meaning/peace (from the first 8 items) and faith (from the last 4 items). The former scale measures the patients' peacefulness, sense of meaning and purpose in life, whereas the latter scale measures their sense of comfort and strength from spiritual beliefs. We reverse scored negatively framed questions and summed item responses so that higher score indicated higher spiritual well-being. Possible scores range from 0 to 32 for the first subscale and from 0 to 16 for the second.

d) Social support

Social Well-Being

The survey instrument included the 7-item Functional Assessment of Cancer Therapy-General questionnaire to assess patients' social/family well-being (FACT- SWB) (29) through evaluating 1) their communication with and the support provided by their friends, family, partner, and other close relations, and 2) the acceptance of illness by the aforementioned support system (perceived by the patients). Each question asked participants to report levels of support on a scale of 0 (not at all) to 4 (very much). The total score ranges from 0 to 28 with a higher score indicating better social/family well-being.

Social Stigma

We used the Sense of Stigma factor questions from the Shame Stigma Scale (30) to evaluate feelings of shame or embarrassment patients may have around others' cognitive and emotional reactions to their cancer diagnosis. Social stigma was scored from 5 (never feels stigma) to 30 (always feels stigma) based on 6 questions which assessed the extent to which patients felt ashamed of, ostracized for, or discriminated against, due to their diagnosis.

e) Awareness of advanced cancer stage

Lastly, we added one categorical-type variable that recorded whether patients were aware of their late cancer stage. Patients were asked if they knew the current stage of their cancer. If they answered with "Advanced Cancer (Stage IV)", then we assigned a value of one to this variable and zero for otherwise (i.e., "Early Stage (Stage I, II or III) or "I don't know").

Statistical analysis

We analyzed the relationship between patients' DTL (binary variable: Yes/No) and physical, psychological & spiritual, social factors as well as the awareness of the late cancer stage, using a linear probability regression model. The demographic variables that we controlled for include gender (male=1, female=0), years of education, age, a dummy for married (married=1, otherwise=0), religion (non-Hindu=1, Hindu=0), type of caste (non-General Caste=1, General Caste=0), and SES (poor=1, otherwise=0). We used the last variable to identify socially and economically disadvantaged groups of patients whose DTL may be significantly different from that of non-disadvantaged groups. In the survey, patients were asked to report their perceived SES among the four pre-classified options: Poor, Low-Middle class, Upper-Middle class, and Wealthy. We created a dummy variable for the lowest SES, by assigning a value of one for patients reporting "Poor" and zero for others. One study reports a high correlation between the subjective SES and self-rated health and health outcomes (31). The self-reported SES may also be a good proxy for financial distress which is often hard to be measured due to large missing data for household income and difficulty in classifying occupational class.

Using linear regression models, we also examined if there are differences in physical, psychological, and social factors between a low socio-economic and a high socio-economic group by controlling for other

demographic characteristics that include gender, age, education, marital status, caste (general vs. non-general caste), and religion (Hindu vs. non-Hindu).

The number of patients for analysis is 192 with no missing information for the variables of interest. An examination of missing data did not reveal any systematic or non-random patterns, and missing scale scores were minimal, ranging across study measure from 1% (on the primary outcome measure, DTL) to 3 % (on the stigma measure). All analyses were conducted at STATA 14.

Results

Table 1 presents the characteristics of the respondents. The mean age of the patients was 52 years and mean years of education was 7 years. About 23% patients responded that they were aware of their late-stage cancer diagnosis. More than half of the patients were males (58%) and fell under disadvantaged caste groups (60%). Also, 23% reported that their perceived SES was “Poor” and 42% reported that they fell under “Lower Middle” class. Most were married (89%) and Hindu (91%). The most common cancer types were breast (16%), Lung (15%), genitourinary (26%), Oral (8%), and Liver (6%). The proportion of outpatients (60%) exceeded the proportion of inpatients (40%).

Table 1
Patient Characteristics (N=192)

	Mean	SD	Min	Max
Desire to Live (DTL)=1	0.859	0.349	0	1
Socio-Economic Status (SES)				
poor	0.229	0.421	0	1
lower middle	0.422	0.495	0	1
upper middle	0.307	0.463	0	1
wealthy	0.036	0.188	0	1
SES: unknown	0.005	0.072	0	1
Caste				
General caste	0.406	0.492	0	1
Disadvantaged caste	0.510	0.501	0	1
Caste: don't know	0.052	0.223	0	1
Caste: unknown	0.031	0.174	0	1
Male	0.583	0.494	0	1
Age	51.563	13.265	18	81
Years of education	7.130	6.320	0	25
Married=1	0.891	0.313	0	1
Non-Hindu	0.094	0.292	0	1
Awareness of late-stage cancer diagnosis	0.234	0.425	0	1
BPI- pain severity	3.439	2.258	0	8.250
Self-blame				
Behavioural Self-Blame (BSB)	0.375	0.485	0	1
Characterological Self-Blame (CSB)	0.568	0.497	0	1
Social/Family support (FACT-SWB)	14.799	6.154	1	28
Stigma	9.026	3.545	6	26
Spiritual well-being				
FACIT-Sp: Total	24.384	9.136	5	48

	Mean	SD	Min	Max
FACIT-Sp: meaning	16.165	6.046	3	32
FACIT-Sp: faith	8.219	4.570	0	16
Psychological distress				
HADS-total	18.703	7.946	0	38
HADS-anxiety	8.563	4.273	0	21
HADS-depression	10.141	4.243	0	20

Patients in our sample reported a mean score of 3.4 (SD: 2.3) for pain severity with the max value of 8.3 out of 10. The mean for HADS-total was 18.7 (SD: 7.9) and the mean scores for HADS-anxiety and HADS-depression were 8.6 (SD: 4.3) and 10.1 (SD: 4.2), respectively. Among the sample patients, 37.5% and 56.8% responded to have behavioral (BSB) and characterological self-blame (CSB), respectively. The mean social/family support score was 14.8 (SD: 6.2) out of 28, and the mean social stigma measurement is 9.0 (SD: 3.5) out of 30 with the max value of 26. Patients also reported a mean score of 24.4 (SD: 9.1) out of 48 for spiritual well-being. The mean score for Meaning/Peace subscale was 16.2 (SD: 6.0) and 8.2 (SD: 4.6) for the Faith subscale.

Table 2, Column 1 presents the main results of the linear probability model estimation, showing the associations between patients DTL and their physical, psychological, and social factors. We find that pain severity reduced DTL, which is statistically significant. Specifically, one point increase in the BPI pain severity score (ranging from 0 to 10) decreased the probability of having DTL by 2.2%. Among psychological factors, the total score of psychological distress (HADS-total; anxiety & depression, ranging from 0 to 42) is negatively associated with DTL, as we expected. One point rise in the total HADS decreases the probability of having DTL by 0.7%, which is smaller than the effect of pain severity, in absolute sense. On the contrary, we did not find a significant association between the outcome variable and self-blame and social support factors that include the measurement for social and family support and perceived social stigma against cancer and cancer patients. Unlike our hypothesis based on previous literature, spiritual well-being does not significantly affect DTL among our sample patients. We also ran the same regression by disaggregating the total psychological distress and spiritual well-being scores into their sub-scales, respectively. The result is in Column 2, Table 2. The sub-scales of psychological distress (HADS-anxiety and HADS-depression) and spiritual well-being (FACIT-Sp: meaning and FACIT-Sp: faith) are insignificant, possibly due to the lack of statistical power.

Table 2
Linear Probability Model Result of Desire on Live

	(1)	(2)
BPI- pain severity	-0.022** (0.011)	-0.022** (0.011)
Behavioural	0.029	0.029
Self- Blame (BSB)	(0.061)	(0.061)
Characterological Self- Blame (CSB)	-0.079 (0.063)	-0.092 (0.064)
HADS-total	-0.007** (0.003)	
HADS-anxiety		-0.006 (0.009)
HADS-depression		-0.010 (0.009)
Social Well Being (FACT-SWB)	0.003 (0.005)	0.003 (0.005)
Stigma	0.006 (0.010)	0.006 (0.010)
FACIT-Sp: Total	0.002 (0.003)	
FACIT-Sp: meaning		-0.004 (0.006)
FACIT-Sp: faith		0.010 (0.007)
Awareness of late-stage cancer diagnosis	-0.054 (0.067)	-0.053 (0.067)

Note. Robust standard errors are in parentheses. *** p < 0.01, ** p < 0.05, * p < 0.1.

	(1)	(2)
SES: Poor	-0.162** (0.071)	-0.158** (0.071)
General caste	-0.061 (0.050)	-0.055 (0.051)
Male	0.005 (0.049)	0.021 (0.051)
Age	-0.001 (0.002)	-0.002 (0.002)
Years of education	-0.007 (0.005)	-0.006 (0.004)
Married=1	-0.034 (0.076)	-0.019 (0.075)
Non-Hindu	-0.076 (0.095)	-0.076 (0.095)
Constant	1.173*** (0.208)	1.238*** (0.208)
Observations	192	192
R-squared	0.156	0.165

Note. Robust standard errors are in parentheses. *** p < 0.01, ** p < 0.05, * p < 0.1.

Only 23% of the 192 sample patients responded that they were aware of their late cancer stage. However, there was no difference in DTL between those who were aware of the late stage and those who were not, when other factors such as pain severity and psychological distress being constant. With regards to demographic information, only SES is significantly associated with DTL. Patients whose perceived SES was poor were likely to have a 16% lower probability of DTL compared to those who had higher SES (lower middle class, upper middle class, and wealthy). By comparing the effects of each variable, we found that SES has the largest effect on DTL.

This finding motivated us to further investigate whether there are significant differences in patients' physical, psychological, and social factors that may affect DTL between these two different socio-economic groups. Table 3 shows result. By controlling demographic variables that include a dummy for General caste, non-Hindu, male, marriage, age, and years of education, we find that only psychological

distress (HADS-total) is significantly higher among patients with the lowest SES. When the aforementioned demographic variables are constant, being poor increased the score of psychological distress by 4.1 points. Considering that its mean value of non-poor patients is 17.5, this change is considered to be a large increase because it is equivalent to a 23% increase.

Table 3
Physical, Psychological, and Social factors by SES

	BPI-Pain severity	BSB	CSB	FACT-SWB	Stigma	FACIT-Sp: Total	HADS-total
SES:Poor	-0.172 (0.427)	0.063 (0.091)	0.087 (0.092)	-0.490 (1.238)	1.176 (0.860)	0.183 (1.813)	4.115*** (1.579)
Mean among SES: Others	3.434	0.358	0.541	15.203	8.642	24.609	17.520
N	192	192	192	192	192	192	192

Note. Robust standard errors are in parentheses. All regressions include a dummy for general caste, male dummy, age, years of education, marriage dummy and non-Hindu dummy. *** p < 0.01, ** p < 0.05, * p < 0.1.

Discussion

Using data collected from 192 advanced cancer patients in a tertiary cancer centre in Jaipur, India, this study examines the relationship between various physical, social, and psychological factors and desire to live (DTL). Consistent with previous literature (32–34), we found that pain severity reduces DTL. One point increase in the BPI pain severity score decreases DTL by 2.2%. Physical symptoms including pain often bring about feelings of powerlessness and helplessness in patients, which is aggravated by the severity of symptoms and chronicity (35). For advanced cancer patients in India, living with chronic pain would be particularly more challenging due to limited access to palliative care services, cost of care, and strict regulations on using opioids and other such pain-relieving medication (36).

As a number of earlier studies showed (20, 37–40), this study also found that psychological distress is negatively correlated with DTL. However, its effect size is not as large as the effect size of the pain severity measurement. One point increase in the HADS-total (ranging from 0 to 42) leads to a reduction of DTL by 0.7%. These results contrast to the findings of a study with metastatic cancer patients in Canada that physical pain no longer contributes to the prediction of the desire to hasten death (DHD) once psychological distress variables are included in the model (41). The authors found that hopelessness and depression accounted for over 34% of the variance in predicting DHD scores. Although our results show a relatively small effect of HADS-total on DTL, the level of psychological distress such as anxiety and depression among the surveyed patients is still worrying. Their average HADS-anxiety and HADS-depression scores were 8.6 and 10.1 respectively, which are not far below the cut-off (11 or higher) for

"probable presence of anxiety/depression" (26). This calls for further attention from healthcare professionals to develop and implement effective strategies to assist patients' psychological distress management.

Despite its importance, even detecting high psychological distress symptoms is challenging in India. This is because there is a dearth of trained mental health professionals in India (42). Additionally, both healthcare providers and patients are reluctant to openly talk about this issue (43, 44). One study shows that healthcare workers are often uncomfortable with managing patients' extreme emotional reactions (45). Another study shows that patients find it difficult to disclose depressive symptoms to healthcare providers because they are fearful of being perceived as "less" than others (46). Patients' limited understanding of the relationship between psychological distress and physical symptoms could further discourage patients' willingness to reveal anxiety/depression (47). More research is needed to clarify what causes patients to hide their psychological distress and to examine the extent to which this behaviour influences the level of actual psychological distress and its effect on DTL.

Holding degrees of pain severity and psychological distress constant, we found socio-economic status had the largest effect on DTL such that patients with perceived socio-economic status "Poor" were likely to have a 16% lower probability of DTL compared to those with higher socio-economic status (i.e., lower middle class, upper middle class and wealthy). We further examined if there are significant differences in patients' physical, psychological, and social factors between the two socio-economic status groups. Controlling for caste, religion, gender, age, marital status and years of education, we found psychological distress to be statistically higher among patients belonging to "Poor" category. Previous work provided similar evidence that cancer patients facing financial difficulties are more likely to feel anxious and depressed, reporting a lower overall quality of life (QoL) (48, 49). Taken together, our findings imply that it is imperative to improve the QoL of advanced cancer patients from disadvantaged groups by addressing their high psychological distress.

Inconsistent with our hypotheses, we found that self-blame, social support, and spiritual well-being factors have no effect on DTL. In particular, the insignificant effects of social support factors (social well-being and social stigma) are surprising given the prevalence of stigma against cancer patients in a northern state of India (50), very close to where this study was conducted. One possible reason to explain this result could be social desirability: patients might feel the obligation to respond to interviewers in a socially acceptable or desirable manner. This tendency could be salient when the survey questions involve third parties and society, as opposed to questions about how patients feel about themselves. In this case, patients with high social desirability are likely to over-report their social support levels, which leads to measurement errors in the relevant variables.

Lastly, our results show that the awareness of the late cancer stage does not decrease DTL, contributing to the growing evidence that the awareness of diagnosis does not negatively impact patients' QoL. In several South-East Asian (51–54) and South Asian countries including India, patients' families often withhold information regarding a cancer diagnosis from the patient, and make treatment decisions on

their behalf, believing this exercise protects the patients (55). Many of them believe that honest and complete disclosure will destroy the patients' hope and the will to live. However, studies suggest their decision may not be the best one for the patients. One study shows no difference in the prevalence of anxiety and depression between patients who were aware of their diagnosis and those who were not, at an oncology centre in South India (56). Another study in north India found that the awareness of cancer diagnosis had no significant effect on the QoL of lung cancer patients (57). In fact, a study with 300 cancer patients at BMCHRC, our study site, found that 94% of them wanted details about their cancer diagnosis and treatment (58). Combined with these findings, our results lend support to the argument that patients need to be better informed about their diagnosis and prognosis. Yet, this should be carefully interpreted and implemented given the importance of the family's role in decision making in many South Asian cultures.

Our findings come with the following limitations. First, the data for this study is collected from a single hospital site in India, and hence our findings cannot be generalized for advanced cancer patients in the entire nation. Second, the sampling strategy we used was convenience sampling as opposed to random sampling, which potentially biases the results as people who volunteered to participate in our survey could be different in unobserved ways from the ones who did not. Third, we cannot rule out the possibility of patients giving socially desirable answers, especially for the questions about how other people and the society treat them.

Conclusions

We examined the relationship between DTL and various physical, psychological, and social factors by using data from 192 advanced cancer patients in a tertiary cancer centre in Jaipur, India. We found that pain severity, psychological distress and low socio-economic status reduced DTL of the patients. The negative impact of low socio-economic status is noticeably large in that perceiving oneself as "poor" reduced the probability of DTL by 16%. Considering the limited access to palliative care and high cost of health care in India and other lower middle-income (LMIC) countries, future research should be focused on developing interventions to improve physical pain and psychological distress, particularly for patients who are socially and economically disadvantaged.

Abbreviations

APPROACH: Asian Patient Perspectives Regarding Oncology Awareness, Care and Health

QoL: Quality of Life

DHD: Desire for Hastened Death

DTL: Desire to Live

SES: Socio-economic Status

HDI: Human Development Index

GDP: Gross Domestic Product

BMCHRC: Bhagwan Mahaveer Cancer Hospital and Research Centre

BPI-SF: Brief Pain Inventory- Short Form

HADS: Hospital Anxiety and Depression Scale

BSB: Behavioural Self-blame

CSB: Characterological Self-blame

FACIT-Sp: Functional Assessment of Chronic Illness Therapy – Spiritual Well-Being Scale

FACT-SWB: Functional Assessment of Cancer Therapy-General questionnaire to assess social/family well-being

LMIC: Lower-Middle Income Countries

Declarations

Copyright and Licenses

The FACIT and all related works are owned and copyrighted by, and the intellectual property of David Cella, Ph.D. Permission for use of the FACT/FACIT system of questionnaire is obtained by contacting information@facit.org.

BPI-SF © Copyright 1991 Charles S. Cleeland, PhD. Pain Research Group – Used by permission. All rights reserved.

Ethics approval and consent to participate

The study was performed in accordance with the Declaration of Helsinki and approved by the Institutional Ethics Committee of Bhagwan Mahaveer Cancer Hospital & Research Centre (Ref No: BMH/2016/89) and the National University of Singapore (Ref No: B-15-319). All participants provided written informed consent.

Consent for publication

Not Applicable.

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Funding for this study was provided by the Lien Centre for Palliative Care.

Authors' contributions

SS, AK, AH, PA, and KiP were involved in the study design. SS conducted the data analysis. SS, JP, AK, AH, PA, KiP, and KrP were involved in interpretation of findings, manuscript writing, and the decision to submit the paper for publication. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Frankl VE. Man's search for meaning: Simon and Schuster; 1985. Attributed to Friedrich Nietzsche. Götzen-Dämmerung; oder, Wie man mit dem Hammer philosophirt. 1889. Sprüche und Pfeile.
2. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: a cancer journal for clinicians. 2018;68(6):394–424.
3. Dhillon PK, Mathur P, Nandakumar A, Fitzmaurice C, Kumar GA, Mehrotra R, et al. The burden of cancers and their variations across the states of India: the Global Burden of Disease Study 1990–2016. The Lancet Oncology. 2018;19(10):1289–306.
4. Mallath MK, Taylor DG, Badwe RA, Rath GK, Shanta V, Pramesh C, et al. The growing burden of cancer in India: epidemiology and social context. The Lancet Oncology. 2014;15(6):e205-e12.
5. Pramesh C, Badwe RA, Sinha RK. The national cancer grid of India. Indian journal of medical and paediatric oncology: official journal of Indian Society of Medical & Paediatric Oncology. 2014;35(3):226.
6. Kishore J, Ahmad I, Kaur R, Mohanta P. Beliefs and perceptions about cancers among patients attending radiotherapy OPD in Delhi, India. Asian Pac J Cancer Prev. 2008;9(1):155–8.
7. IANS. 70% cancer patients in India consult doctor at terminal stage (February 4 is World Cancer Day). Business Standard. 2017 February 3, 2017.
8. Hebert JR, Ghumare SS, Gupta PC. Stage at diagnosis and relative differences in breast and prostate cancer incidence in India: comparison with the United States. Asian Pacific Journal of Cancer

- Prevention. 2006;7(4):547.
9. Pakseresht S, Ingle GK, Garg S, Sarafraz N. Stage at diagnosis and delay in seeking medical care among women with breast cancer, Delhi, India. *Iranian Red Crescent Medical Journal*. 2014;16(12).
 10. Knaul FM, Farmer PE, Krakauer EL, De Lima L, Bhadelia A, Kwete XJ, et al. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *The Lancet*. 2018;391(10128):1391–454.
 11. Pramesh C, Badwe RA, Borthakur BB, Chandra M, Raj EH, Kannan T, et al. Delivery of affordable and equitable cancer care in India. *The Lancet Oncology*. 2014;15(6):e223-e33.
 12. Monforte-Royo C, Villavicencio-Chávez C, Tomás-Sábado J, Balaguer A. The wish to hasten death: a review of clinical studies. *Psycho-oncology* (Chichester, England). 2011;20(8):795–804.
 13. Rodríguez-Prat A, Balaguer A, Booth A, Monforte-Royo C. Understanding patients' experiences of the wish to hasten death: an updated and expanded systematic review and meta-ethnography. *BMJ open*. 2017;7(9):e016659-e.
 14. Mohandoss A, Thavarajah R. A study on suicide among Indians living with cancer during 2001-2014. *Indian journal of cancer*. 2016;53(3):435.
 15. Latha KS, Bhat SM. Suicidal behaviour among terminally ill cancer patients in India. *Indian journal of psychiatry*. 2005;47(2):79–83.
 16. Jones JM, Huggins MA, Rydall AC, Rodin GM. Symptomatic distress, hopelessness, and the desire for hastened death in hospitalized cancer patients. *Journal of psychosomatic research*. 2003;55(5):411–8.
 17. Guy M, Stern TA. The desire for death in the setting of terminal illness: a case discussion. *Primary care companion to the Journal of clinical psychiatry*. 2006;8(5):299–305.
 18. Chochinov HM, Wilson KG, Enns M, Mowchun N, Lander S, Levitt M, et al. Desire for death in the terminally ill. *The American journal of psychiatry*. 1995.
 19. Nordin K, Berglund G, Glimelius B, Sjödén PO. Predicting anxiety and depression among cancer patients: a clinical model. *European journal of cancer (1990)*. 2001;37(3):376-84.
 20. Rodin G, Lo C, Mikulincer M, Donner A, Gagliese L, Zimmermann C. Pathways to distress: the multiple determinants of depression, hopelessness, and the desire for hastened death in metastatic cancer patients. *Social science & medicine*. 2009;68(3):562–9.
 21. Lorant V, Croux C, Weich S, Deliège D, Mackenbach J, Ansseau M. Depression and socio-economic risk factors: 7-year longitudinal population study. *British journal of psychiatry*. 2007;190(4):293–8.
 22. Khan L, Wong R, Li M, Zimmermann C, Lo C, Gagliese L, et al. Maintaining the will to live of patients with advanced cancer. *The cancer journal (Sudbury, Mass)*. 2010;16(5):524–31.
 23. McDermott EP, Selman LM, Wright MP, Clark DP. Hospice and Palliative Care Development in India: A Multimethod Review of Services and Experiences. *Journal of pain and symptom management*. 2008;35(6):583–93.
 24. Implementation' MoSaP. Indian states by GDP. *Statistics Times* 2021.

25. Cleland CS, Ryan K. The brief pain inventory. Pain Research Group. 1991;143–7.
26. Snaith RP. The hospital anxiety and depression scale. Health and quality of life outcomes. 2003;1(1):1–4.
27. Peterman AH, Fitchett G, Brady MJ, Hernandez L, Cella D. Measuring spiritual well-being in people with cancer: The functional assessment of chronic illness therapy—spiritual well-being scale (FACIT-Sp). *Annals of behavioral medicine*. 2002;24(1):49–58.
28. Webster K, Cella D, Yost K. The Functional Assessment of Chronic Illness Therapy (FACIT) Measurement System: properties, applications, and interpretation. *Health and quality of life outcomes*. 2003;1(1):79-.
29. Cella DF, Tulsky DS, Gray G, Sarafian B, Linn E, Bonomi A, et al. The Functional Assessment of Cancer Therapy scale: development and validation of the general measure. *J Clin Oncol*. 1993;11(3):570–9.
30. Kissane DW, Patel SG, Baser RE, Bell R, Farberov M, Ostroff JS, et al. Preliminary evaluation of the reliability and validity of the Shame and Stigma Scale in head and neck cancer. *Head & neck*. 2013;35(2):172–83.
31. Demakakos P, Nazroo J, Breeze E, Marmot M. Socioeconomic status and health: The role of subjective social status. *Social science & medicine* (1982). 2008;67(2):330-40.
32. Schroepfer TA. Critical events in the dying process: the potential for physical and psychosocial suffering. *Journal of palliative medicine*. 2007;10(1):136–47.
33. Emanuel EJ, Fairclough DL, Emanuel LL. Attitudes and Desires Related to Euthanasia and Physician-Assisted Suicide Among Terminally Ill Patients and Their Caregivers. *JAMA: the journal of the American Medical Association*. 2000;284(19):2460–8.
34. Nissim R, Gagliese L, Rodin G. The desire for hastened death in individuals with advanced cancer: A longitudinal qualitative study. *Social science & medicine* (1982). 2009;69(2):165-71.
35. Sand L, Strang P, Milberg A. Dying cancer patients' experiences of powerlessness and helplessness. *Supportive care in cancer*. 2008;16(7):853–62.
36. LeBaron V, Beck SL, Maurer M, Black F, Palat G. An ethnographic study of barriers to cancer pain management and opioid availability in India. *The oncologist*. 2014;19(5):515.
37. Wilson KG, Scott JF, Graham ID, Kozak JF, Chater S, Viola RA, et al. Attitudes of terminally ill patients toward euthanasia and physician-assisted suicide. *Archives of Internal Medicine*. 2000;160(16):2454–60.
38. O'Mahony S, Goulet J, Kornblith A, Abbatiello G, Clarke B, Kless-Siegel S, et al. Desire for hastened death, cancer pain and depression: report of a longitudinal observational study. *Journal of pain and symptom management*. 2005;29(5):446–57.
39. Brown JH, Henteleff P, Barakat S, Rowe CJ. Is it normal for terminally ill patients to desire death? *The American journal of psychiatry*. 1986.

40. Breitbart W, Rosenfeld B, Pessin H, Kaim M, Funesti-Esch J, Galietta M, et al. Depression, hopelessness, and desire for hastened death in terminally ill patients with cancer. *Jama*. 2000;284(22):2907–11.
41. Rodin G, Zimmermann C, Rydall A, Jones J, Shepherd FA, Moore M, et al. The desire for hastened death in patients with metastatic cancer. *Journal of pain and symptom management*. 2007;33(6):661–75.
42. Garg K, Kumar CN, Chandra PS. Number of psychiatrists in India: Baby steps forward, but a long way to go. *Indian journal of psychiatry*. 2019;61(1):104.
43. Mason A, Juyal R, Das SC, Shikha D, Saini S, Semwal J. Prevalence and correlates of psychological distress among cancer patients in a tertiary care hospital in northern India. *Int J Community Med Public Health*. 2019;6:2223–8.
44. Mishra S, Bhatnagar S, Philip FA, Singhal V, Rana SPS, Upadhyay SP, et al. Psychosocial concerns in patients with advanced cancer: An observational study at regional cancer centre, India. *American Journal of Hospice and Palliative Medicine®*. 2010;27(5):316–9.
45. Sengupta J, Chatterjee SC. Dying in intensive care units of India: Commentaries on policies and position papers on palliative and end-of-life care. *Journal of critical care*. 2017;39:11–7.
46. Raguram R, Weiss MG, Channabasavanna S, Devins GM. Stigma, depression, and somatization in South India. *American Journal of Psychiatry*. 1996;153(8):1043–9.
47. Fulton CL. The physical and psychological symptoms experienced by patients with metastatic breast cancer before death. *European journal of cancer care*. 1997;6(4):262–6.
48. Jacob J, Palat G, Verghese N, Chandran P, Rapelli V, Kumari S, et al. Health-related quality of life and its socio-economic and cultural predictors among advanced cancer patients: Evidence from the APPROACH cross-sectional survey in Hyderabad-India. *BMC palliative care*. 2019;18(1):1–12.
49. Gangane N, Khairkar P, Hurtig A-K, San Sebastián M. Quality of life determinants in breast Cancer patients in central rural India. *Asian Pacific journal of cancer prevention: APJCP*. 2017;18(12):3325.
50. Kaur R. Cultural beliefs, cancer and stigma: experiences of patients from Punjab (India). *Studies on Ethno-Medicine*. 2015;9(2):247-54.
51. Tieying Z, Haishan H, Meizhen Z, Yan L, Pengqian F. Health professionals' attitude towards information disclosure to cancer patients in China. *Nursing ethics*. 2011;18(3):356–63.
52. Wang D-c, Guo C-b, Peng X, Su Y-j, Chen F. Is therapeutic non-disclosure still possible? A study on the awareness of cancer diagnosis in China. *Supportive care in cancer*. 2011;19(8):1191–5.
53. Elwyn TS, Fetters MD, Sasaki H, Tsuda T. Responsibility and cancer disclosure in Japan. *Social science & medicine (1982)*. 2002;54(2):281-93.
54. Pang A, Ho S, Lee S-C. Cancer physicians' attitude towards treatment of the elderly cancer patient in a developed Asian country. *BMC geriatrics*. 2013;13(1):35-.
55. Chaturvedi SK. Ethical dilemmas in palliative care in traditional developing societies, with special reference to the Indian setting. *Journal of medical ethics*. 2008;34(8):611–5.

56. Chandra PS, Chaturvedi SK, Kumar A, Kumar S, Subbakrishna D, Channabasavanna S, et al. Awareness of diagnosis and psychiatric morbidity among cancer patients—a study from South India. *Journal of psychosomatic research*. 1998;45(3):257–61.
57. Aggarwal AN, Singh N, Gupta D, Behera D. Does awareness of diagnosis influence health related quality of life in north Indian patients with lung cancer? *The Indian journal of medical research*. 2016;143(Suppl 1):S38.
58. Laxmi S, Khan JA. Does the cancer patient want to know? Results from a study in an Indian tertiary cancer center. *South Asian journal of cancer*. 2013;2(2):57.

Figures




Figure 1

Patient recruitment log