

Metagenomic Analysis of Viral Diversity and a Novel Astrovirus of Forest Rodent

Hai Yin

Qiqihar University

Li Zhao

Jilin University

Hong Chen (✉ 814760285@qq.com)

Harbin Veterinary Research Institute

Research

Keywords: Metagenomics, Forest Rodent, Virome, Novel, Astrovirus

Posted Date: November 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-960263/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Rodents are important virus reservoirs and natural hosts for multiple viruses. They are one of the wild animals that are extremely threatening to the spread of human viruses. Therefore, research on rodents carrying viruses and identifying new viruses that rodents carry is of great significance for preventing and controlling viral diseases.

Methods: In this study, fecal samples from six species of forest rodents in Northeast China were sequenced using metagenomics, and an abundance of virome information was acquired.

Results: Among the top 10 most abundant viral families, RNA virus include Orthomyxoviridae, Picornaviridae, Bunyaviridae and Arenaviridae, DNA virus include Herpesviridae, Insect virus include Nodaviridae and Baculoviridae, Plant virus Tombusviridae and Phage (Myoviridae). Except for Myoviridae, there was no significant difference in the abundance of virus families in the feces of each rodent species. In addition, a new strain of astrovirus was discovered, with an ORF and genome arrangement comparable to other rodent astroviruses. The newly identified astrovirus had the highest similarity with the rodent astrovirus isolate, CHN/100.

Conclusions: The data obtained in this study provided an overview of the viral community present in these rodent fecal samples, revealing some rodent-associated viruses closely related to known human or animal pathogens. Strengthening our understanding of unclassified viruses harbored by rodents present in the natural environment could provide scientific guidance for preventing and controlling new viral outbreaks that can spread via rodents.

Background

Emerging infectious diseases have severely affected global public health [1]. Most of the newly emerging infectious diseases in the world originate from wild animals [2]. The spread of viruses may be affected by many factors, including the density of animals and the degree of contact with humans, among other factors [3]. Rodents are one of the animals in close contact with humans and have a high distribution density. Therefore, rodents, which are important reservoirs of viruses and are natural hosts of multiple viruses, are one of the wild animals that are extremely threatening to humans [4,5]. Many viruses do not cause obvious clinical symptoms in rodents, but when the virus crosses to a human host, it may cause serious disease. For example, Hantavirus and Arenavirus are extremely harmful viruses to humans, but not to rodents.

With the continuous development of DNA sequencing technologies, many new viruses have been discovered in recent years [2,4], including some pathogenic and non-pathogenic viruses that naturally exist in the host. With the continuous deepening of global research into viruses carried by rodents [6-9], a deep understanding of viruses carried by rodents has begun to be developed. Studies have found that most viruses related to public health have similar viruses in rodents. Such studies have also largely enriched the knowledge of virus diversity [6,8-10] and provide a theoretical basis for the possible occurrence of zoonotic diseases in the future and tracking the origins of transmission. Many new viruses are reported continuously, indicating a large number of viruses that remain to be discovered. Therefore, virus composition and new virus mining in rodents from different geographic regions are greatly important for preventing and controlling new infectious diseases.

In this study, fecal pellets were collected from *Myodes rufocanus* (MR), *Apodemus peninsulae* (AP), *Apodemus agrarius* (AA), *Tamias sibiricus* (TS), *Sciurus vulgaris* (SV), and *Cricetulus triton* (CT), which are wild representatives of the mice forest areas of Hengdaohezi Town, Hailin City, and Heilongjiang Province. The feces from those species were used to assess the variety of viruses carried by forest rodents. Metagenomic next-generation sequencing analyses were performed to screen the viromes of the samples. Herein, we outlined the viral spectrum within these mouse samples and

found a novel astrovirus. These data offer new clues for tracing the sources of important viral pathogens that can cause human and animal disease.

Methods

2.1. Sample processing

82 rodents were captured in the forested areas of Hengdaohezi Town (A, N 44°48′44″, E 129°02′04″) Hailin City, Heilongjiang Province from May 21 to August 22 (Summer Festival) in 2020 were selected as the research samples. The collected samples included 22 *Myodes rufocanus* (MR), 16 *Apodemus peninsulae* (AP), 15 *Apodemus agrarius* (AA), 11 *Tamias sibiricus* (TS), 10 *Sciurus vulgaris* (SV), and eight *Cricetulus triton* (CT). A total of six samples from the same species were mixed.

Fecal pellets samples were homogenized, diluted in a ratio of 1:10 using PBS, made into a suspension, and vortexed for thorough mixing. The samples were then centrifuged at 2000 rpm for 10 min at 4°C. Following this, the supernatants were transferred to a fresh tube and centrifuged for 10 min for the complete removal of cell debris, bacterial cells, and other impurities. The supernatants were filtered through a 0.22 µm syringe filter (Jet, Guangzhou, China) and concentrated. The filtrate was centrifuged in an SW55Ti rotor using a Beckman ultracentrifuge at 45,000 rpm for 2 h. The precipitates were re-suspended in PBS and passed through a 0.22 µm syringe filter. The samples were then stored at -80°C until subsequent analyses.

2.2. Viral Metagenomics Analysis

Sequencing libraries were generated using the NEBNext® Ultra™ DNA Library Prep Kit for Illumina (NEB, USA), following the manufacturer's recommendations. Index codes were added to attribute sequences to each sample. Briefly, the DNA samples were fragmented by sonication to a size of 300 bp, and the DNA fragments were then end-polished, A-tailed, and ligated with the full-length adaptor for Illumina sequencing to aid further PCR amplification.

Finally, PCR products were purified (AMPure XP system), and libraries were analyzed for size distribution using an Agilent2100 Bioanalyzer and quantified using real-time PCR. According to the manufacturer's instructions, the clustering of the index-coded samples was performed using a cBot Cluster Generation System. After cluster generation, the library preparations were sequenced on an Illumina HiSeq2500 platform, and paired-end reads were generated.

2.3. Species annotation and analyses of abundance

Prinseq software (version 0.20.4) was used to assess the sample data quality, filter the low-quality and repetitive sequences and the rodent genome was the reference genome. Bowtie2 software was used to remove host DNA sequences, and Mira (v4.0.2) software was used to splice and assemble the sequences. The assembled and unassembled sequences were searched against the local NCBI virus database using the BLASTX and BLASTN tools in the BLAST+ software package to obtain virus annotations. Geneious version 2019.2.1 was used to predict the ORF of the new annotated virus.

2.4. Passage of fecal supernatants in BHK-21 Cells

The BHK-21 was used to propagate the fecal suspensions from wild mice and detect viruses according to previously published methods [11,12]. Briefly, the supernatants of homogenized fecal pellets were filtered through a 0.22 µm syringe filter (Jet, Guangzhou, China) and inoculated onto the BHK21 cells, followed by incubation for 2 h to allow for virus adsorption. After adding a fresh medium, the cells were incubated at 37°C and monitored daily until 7 days postinfection to develop CPE. The infected cells were blindly transmitted three to six times until CPE appeared. The

supernatant harvested from CPE-positive BHK-21 cells was further inoculated on BHK-21 cells to detect the presence of infectious viruses.

2.5. Detection of Astrovirus in CPE-Positive Cells

BHK-21 cells were inoculated with the supernatant collected from CPE-positive cells and incubated for 36 h. Real-time fluorescent quantitative reverse transcription PCR (qRT-PCR), immunofluorescence detection (IFA) were used to analyze the presence of astroviruses in the inoculated cells. For qRT-PCR analysis, primers that specifically amplify the target genes of astrovirus were designed based on the assembled contig sequences.

2.6. Phylogenetic Analysis

The sequences of reference strains with high similarity to the novel astrovirus were downloaded from NCBI. A phylogenetic tree was constructed using MAGA software based on the neighbor-joining maximum composite likelihood method with 1,000 bootstrap replicates to analyze phylogenetic relationships.

2.7. Statistical analysis

Metastate was used to analyze the top 10 abundant taxonomic sequence tags of the three samples. The differences were considered to be statistically significant when the *P*-value was less than 0.05.

Results

3.1. Sequencing and quality control

DNA and RNA, extracted from fecal samples of 82 forest rodent intestinal contents samples collected in the forest area of Hengdaohezi Town Hailin City, Heilongjiang Province, were sequenced. The length of the insert size was 350 bp. Bases showing overlapping information and low mass, and bases that were not measured, were excluded. The total numbers of clean data obtained from the six samples were: 2,384.93 (*MR*), 2,090.33 (*AP*), 1,994.32 (*AA*), 2,391.77 (*TS*), 2,007.30 (*SV*), and 2,148.70 (*CT*). Sequencing data quality was distributed in the quality score Q20 to ensure a normal order of the subsequent advanced analysis. The clean sequence tags were subjected to redundancy processing using the Mothur software to obtain unique sequence tags. The percentages of effective sequences of the six samples were: 95.242% (*MR*), 93.561% (*AP*), 97.509% (*AA*), 97.258% (*TS*), 93.121% (*SV*), and 94.622% (*CT*). (Table 1).

Table 1. Fecal samples used for metagenomic analysis and data generation

Sample	InsertSize (bp)	SeqStrategy	RawData	CleanData	Clean_Q20	Clean_Q30	Clean_GC (%)	Effective (%)
<i>MR</i>	350	(150:150)	2,504.06	2,384.93	76.53	65.45	58.69	95.242
<i>AP</i>	350	(150:150)	2,234.18	2,090.33	75.94	64.76	57.70	93.561
<i>AA</i>	350	(150:150)	2,045.27	1,994.32	87.12	78.68	52.56	97.509
<i>TS</i>	350	(150:150)	2,459.21	2,391.77	83.63	73.78	52.87	97.258
<i>SV</i>	350	(150:150)	2,155.58	2,007.30	73.76	62.15	59.05	93.121
<i>CT</i>	350	(150:150)	2,270.82	2,148.70	78.59	67.93	55.53	94.622

3.2. Viral Communities in fecal samples, based on family-level classifications

Pre-treated clean data for all samples and the assembled sequence were compared to the reference genomes of viruses in the NCBI database using the BLASTX and BLASTN tools in the BLAST+ software package to obtain the virus annotation results. In total, 82 families of mammalian viruses, plant viruses, phages, insect viruses, and fungal viruses were parsed. An overview of the reads of the top 35 families of viruses in each sample is shown in Figure 1. In addition, an overview visualized presentation of the classification of families, genera, and species for each sample is shown in Figure 2A-C.

(1) Single-stranded RNA viruses(Orthomyxoviridae, Picobirnaviridae, Bunyaviridae, and Arenaviridae)

The members of the family Orthomyxoviridae can cause cyclical pandemics throughout the world in various species[13]. In this study, they are assigned to the genus Influenzavirus A and the species influenza A virus. The reads related to the family Orthomyxoviridae occupied the largest proportion of viruses. The percentage of this family of viruses in each sample was: 45.04% (*MR*), 51.57% (*AP*), 41.08% (*AA*), 41.9% (*TS*), 27.59% (*SV*), and 22.1% (*CT*) (Figure 2A).

The family members of Picobirnaviridae cause a wide variety of mucocutaneous, encephalic, cardiac, hepatic, neurological, and respiratory diseases in vertebrate hosts [14]. The Picornaviridae family viruses were found in all six samples. The viruses were assigned to the genus picobirnavirus, the species human picobirnavirus, *Microtus* picobirnavirus V-111_USA_2008, and fox picobirnavirus. It is worth mentioning that the human picobirnavirus occupied the larger proportion of viruses in the *SV* sample (2.64%) (Figure 2C).

The family Bunyaviridae have strong infectivity, wide distribution, a high fatality rate, and can cause serious infectious diseases in humans and animals [15]. Most of the members of this family, such as Rift valley fever virus, Crimean-Congo hemorrhagic fever virus, La Crosse encephalitis virus, and Hantavirus, cause deadly diseases in humans. The natural hosts for Hantavirus are rodents, and it can cause hemorrhagic kidney fever. The virus was present in all samples of the six forest rodents. Among these, the abundance in fecal of *SV* (3.49%) was higher than that in the other samples. The viruses of this family were assigned to the genus, Orthobunyavirus, and the species, Shamonda virus (Figure 2B, C).

Arenaviridae is an enveloped RNA virus found worldwide. The Lassa fever virus, Junin virus, and Machupo virus can cause severe diseases with a high mortality rate[16]. Thus, the prevalence of infectious diseases is closely related to the local dynamic distribution of rodents. In this study, the virus was detected in the fecal of all six species of mice. The viruses in this family are assigned to the genus, Mammarenavirus, and the species, Lassa mammarenavirus.

(2) DNA viruses (Herpesviridae)

Viruses of the Herpesviridae family are enveloped, double-strand DNA viruses, divided into three genera based on phylogenetic clustering: α -herpesvirus, β -herpesvirus, and γ -herpesvirus [17]. This family was detected in fecal samples of all six species of rodent. The viruses in this family were assigned to the genera, Cytomegalovirus, Varicellovirus, Mardivirus, and the species, Cercopithecine herpesvirus 5 and Gallid herpesvirus 2, respectively (Figure 2B, C).

(3) Other rare viruses (Nodaviridae, Baculoviridae, Tombusviridae, Myoviridae)

Insect viruses (Nodaviridae, Baculoviridae), plant viruses (Tombusviridae), and phages (Myoviridae) were identified in the fecal samples. The viruses in the family of Nodaviridae were assigned to the genera, Alphanodavirus and Betanodavirus, and the species, Pariacoto virus and Barfin flounder nervous necrosis virus, respectively. The viruses in the family of Tombusviridae were assigned to the genera, Tombusvirus, and no virus in the family of Tombusviridae was assigned to the species among the top 10 most widely distributed. It is worth mentioning that *AA* and *CT* did not contain any members from Tombusviridae. No virus in the family of Baculoviridae and Myoviridae was assigned to the genera, and species among the top 10 most widely distributed.

(4) Unclassified viruses

Currently, there is little information about unclassified viruses and their evolution in forest rodents. In our data, many reads are classified as "unclassified virus sequences" in all samples, likely to be previously unidentified viruses that have not been studied. The identification and characterization of these unclassified viruses will provide insight into the evolutionary histories of other clinically important viruses, as well as the genetic basis behind their infectivity and virulence in humans and other animals. Such information is important for the development of future treatment options and vaccine research (Figure 2A).

3.3 Genome features of the novel astrovirus

Using mNGS, we identified a novel astrovirus from the feces of *Myodes rufocanus*, which was tentatively named Rodent Astrovirus CHNDB/2019. Rapid Annotation using Subsystem Technology (RAST, <http://rast.nmpdr.org>) was used to annotate the complete genome sequence. The results revealed that the Rodent Astrovirus CHNDB/2019 genome consists of 6192 nt and contains two open reading frames: ORF1 encodes a polyprotein located at nucleotide (nt) positions 35 - 3750, and ORF2 was located at nt positions 3719 -6178 in *Myodes rufocanus*.

3.4 Phylogenetic Analysis

The phylogenetic analysis of the complete genome sequence showed that ORF1 (Figure 3B), ORF2 (Figure 3C), and the complete genome (Figure 3A) had the highest similarity to the rodent astrovirus isolate, CHN/100, which was isolated from Jiaxiang County, Shandong Province, and the genetic distance between them is the closest. The nt sequence similarity between the entire genome and CHN/100 was 84.39%, which the sequence similarity with other rodent astroviruses was 78.08%–67.74%. The nucleotide sequence similarity between ORF1 and orf1b of Rodent Astrovirus isolate CHN/100 was 86.2%, and the amino acid similarity was 74.45%. The greatest similarity between OFR2 and orf2 of the Rodent astrovirus isolates CHN/100 was 81.27%, while the amino acid sequence similarity was 63.61%. Therefore, it was clear that the virus was a rodent astrovirus (GenBank Submission Number, MW927503).

3.5 Infectivity of Astroviruses in BHK Cells

The fecal samples containing the new astroviruses were inoculated into BHK-21 cells to detect the presence of infectious astroviruses. After four to five blind passages, CPE appeared in BHK cells. IFA and qRT-PCR were used to analyze the presence of astroviruses, which could be detected in the inoculated cells. IFA (Figure 4A) confirmed the presence of antibodies specific to the astrovirus capsid spike protein, VP27. In addition, whether there is astrovirus in the inoculated BHK-21 cells. qRT-PCR was performed using astrovirus ORF1 gene-specific primers ORF1-F (5'-CAGTCCTTGGGATTTCTC-3') and ORF1-R (5'-TATTCTTTTCGCACCATTAG-3') (Figure 4B).

Discussion

Currently, there have been some reports about the metagenomic investigations of the viruses carried by rodents. Duan et al. collected the intestinal contents of 99 *Rattus norvegicus* rats and *Apodemus agrarius* mice in Jiaxiang County, Shandong Province, and studied their viral compositions via high-throughput sequencing. The study also found that multiple strains of astrovirus were circulating among rodents in this geographic area at the same time, indicating the extremely rich genetic diversity [18]. Vandegrift et al. analyzed the serum viromes of 978 free-range *P. leucopus* mice captured in Pennsylvania and discovered multiple new viruses from 26 different families, among which there was a highly divergent segmented flavivirus [19]. Williams et al. analyzed the viruses from mouse feces in seven residential locations in Manhattan, Queens, Brooklyn, and Bronx, New York City, New York, within the same year. Unbiased high-throughput sequencing revealed 36 viruses from 18 families and 21 genera, including at least six new viruses and three

new genera[8]. In this study, fecal samples from forest rodents in Northeast China were sequenced using metagenomics, and an abundance of virome information was acquired. 82 families of mammalian viruses, plant viruses, insect viruses, and phages were detected. Among the top 10 most abundant families were the RNA viruses, Orthomyxoviridae, Picornaviridae, Bunyaviridae, and Arenaviridae; the DNA virus, Herpesviridae; the insect viruses, Nodaviridae and Baculoviridae; the plant virus, Tombusviridae; and the phage, Myoviridae.

Astrovirus is a type of virus that infects mammals and poultry and has a wide host range. Astrovirus infection in humans can cause gastrointestinal diseases, and there have been a few cases of neurological symptoms[20-22]. Many new animal astroviruses have been discovered in recent years[9], and such rich genetic diversity among mammalian astroviruses may be caused by the cross-species transmission in wild animals, domestic animals, or humans[20]. This study also identified new viruses. A new astrovirus was identified through sequence splicing, software prediction, evolutionary tree analysis, and experimental verification, and the sequence was submitted to NCBI. There likely are more than the newly identified virus in the samples. Due to the limited sample size in this study, there may be deviations in these rodents that carry the virus. Therefore, further research will be performed to increase the sample size and expand the range of sampling sites.

Conclusions

In this study, fecal samples from 6 species of forest rodents in Northeast China were sequenced using metagenomics, and an abundance of virome information was acquired. The top 10 most abundant families were Orthomyxoviridae, Picornaviridae, Bunyaviridae, Arenaviridae, Herpesviridae, Nodaviridae, Baculoviridae, Tombusviridae, Myoviridae and Unclassified viruses. In addition, a new strain of astrovirus has been discovered, with an ORF genome arrangement comparable to other rodent astroviruses. It has the highest similarity with the rodent astrovirus isolate CHN/100. The metagenomics approach can greatly improve our understanding of the diversity of viruses in mice. This study analyzed the composition and abundance of virus genomes in fecal from six species of rodents using metagenomics technology. This strategy could be extended to other wild or livestock samples worldwide, ultimately increasing our knowledge of the viral population and ecological community, which would help minimize the impact of potential wildlife-associated viruses on public health by providing meaningful basic data.

Abbreviations

ORF: Open reading frame; BHK21: baby-hamster kidney cell line; CPE: cytopathic effects.

Declarations

Ethics approval and consent to participate: The study was approved by the Harbin Veterinary Research Institute and performed following animal ethics guidelines and approved protocols. The animal ethics approval number is Heilongjiang SYXK-2006-032.

Consent for publication:

Not applicable.

Availability of data and materials: The datasets analyzed in the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that there is no conflict of interest regarding the publication of this article.

Funding: This research was funded by Heilongjiang Province Education Department Fundamental Scientific Research Funds, UNPYSCT-2020070.

Author contributions: Haichang Yin and Lili Zhao wrote the paper. Haichang Yin prepared the graphs and table. Hongyan Chen reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements:

Not applicable.

References

1. Luis, A.D.; Hayman, D.T.; O'Shea, T.J.; Cryan, P.M.; Gilbert, A.T.; Pulliam, J.R.; Mills, J.N.; Timonin, M.E.; Willis, C.K.; Cunningham, A.A.; et al. A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? *Proc Biol Sci* 2013, *280*, 20122753, doi:10.1098/rspb.2012.2753.
2. Wolfe, N.D.; Dunavan, C.P.; Diamond, J. Origins of major human infectious diseases. *Nature* 2007, *447*, 279–283, doi:10.1038/nature05775.
3. Woolhouse, M.E.; Gowtage-Sequeria, S. Host range and emerging and reemerging pathogens. *Emerg Infect Dis* 2005, *11*, 1842–1847, doi:10.3201/eid1112.050997.
4. Firth, C.; Bhat, M.; Firth, M.A.; Williams, S.H.; Frye, M.J.; Simmonds, P.; Conte, J.M.; Ng, J.; Garcia, J.; Bhuva, N.P.; et al. Detection of zoonotic pathogens and characterization of novel viruses carried by commensal *Rattus norvegicus* in New York City. *mBio* 2014, *5*, e01933-01914, doi:10.1128/mBio.01933-14.
5. Olival, K.J.; Hosseini, P.R.; Zambrana-Torrel, C.; Ross, N.; Bogich, T.L.; Daszak, P. Host and viral traits predict zoonotic spillover from mammals. *Nature* 2017, *546*, 646–650, doi:10.1038/nature22975.
6. Wu, Z.; Lu, L.; Du, J.; Yang, L.; Ren, X.; Liu, B.; Jiang, J.; Yang, J.; Dong, J.; Sun, L.; et al. Comparative analysis of rodent and small mammal viromes to better understand the wildlife origin of emerging infectious diseases. *Microbiome* 2018, *6*, 178, doi:10.1186/s40168-018-0554-9.
7. Golden, J.W.; Hammerbeck, C.D.; Mucker, E.M.; Brocato, R.L. Animal Models for the Study of Rodent-Borne Hemorrhagic Fever Viruses: Arenaviruses and Hantaviruses. *Biomed Res Int* 2015, *2015*, 793257, doi:10.1155/2015/793257.
8. Williams, S.H.; Che, X.; Garcia, J.A.; Klena, J.D.; Lee, B.; Muller, D.; Ulrich, W.; Corrigan, R.M.; Nichol, S.; Jain, K.; et al. Viral Diversity of House Mice in New York City. *mBio* 2018, *9*, doi:10.1128/mBio.01354-17.
9. Phan, T.G.; Kapusinszky, B.; Wang, C.; Rose, R.K.; Lipton, H.L.; Delwart, E.L. The fecal viral flora of wild rodents. *PLoS Pathog* 2011, *7*, e1002218, doi:10.1371/journal.ppat.1002218.
10. Chen, L.; Liu, B.; Wu, Z.; Jin, Q.; Yang, J. DRodVir: A resource for exploring the virome diversity in rodents. *J Genet Genomics* 2017, *44*, 259–264, doi:10.1016/j.jgg.2017.04.004.
11. Xiao, P.; Han, J.; Zhang, Y.; Li, C.; Guo, X.; Wen, S.; Tian, M.; Li, Y.; Wang, M.; Liu, H.; et al. Metagenomic Analysis of Flaviviridae in Mosquito Viromes Isolated From Yunnan Province in China Reveals Genes From Dengue and Zika Viruses. *Front Cell Infect Microbiol* 2018, *8*, 359, doi:10.3389/fcimb.2018.00359.
12. Xiao, P.; Li, C.; Zhang, Y.; Han, J.; Guo, X.; Xie, L.; Tian, M.; Li, Y.; Wang, M.; Liu, H.; et al. Metagenomic Sequencing From Mosquitoes in China Reveals a Variety of Insect and Human Viruses. *Front Cell Infect Microbiol* 2018, *8*, 364, doi:10.3389/fcimb.2018.00364.
13. Noda, T. [Orthomyxoviruses]. *Uirusu* 2012, *62*, 219–228, doi:10.2222/jsv.62.219.
14. Wang, X.; Ren, J.; Gao, Q.; Hu, Z.; Sun, Y.; Li, X.; Rowlands, D.J.; Yin, W.; Wang, J.; Stuart, D.I.; et al. Hepatitis A virus and the origins of picornaviruses. *Nature* 2015, *517*, 85–88, doi:10.1038/nature13806.

15. Gerlach, P.; Malet, H.; Cusack, S.; Reguera, J. Structural Insights into Bunyavirus Replication and Its Regulation by the vRNA Promoter. *Cell* 2015, *161*, 1267–1279, doi:10.1016/j.cell.2015.05.006.
16. Sarute, N.; Ross, S.R. New World Arenavirus Biology. *Annu Rev Virol* 2017, *4*, 141–158, doi:10.1146/annurev-virology-101416-042001.
17. Lefkowitz, E.J.; Dempsey, D.M.; Hendrickson, R.C.; Orton, R.J.; Siddell, S.G.; Smith, D.B. Virus taxonomy: the database of the International Committee on Taxonomy of Viruses (ICTV). *Nucleic Acids Res* 2018, *46*, D708-D717, doi:10.1093/nar/gkx932.
18. Jing, Z.; Zhang, X.; Shi, H.; Chen, J.; Shi, D.; Dong, H.; Feng, L. A G3P[13] porcine group A rotavirus emerging in China is a reassortant and a natural recombinant in the VP4 gene. *Transbound Emerg Dis* 2018, *65*, e317-e328, doi:10.1111/tbed.12756.
19. Vandegrift, K.J.; Kumar, A.; Sharma, H.; Murthy, S.; Kramer, L.D.; Ostfeld, S.; Hudson, P.J.; Kapoor, A. Presence of Segmented Flavivirus Infections in North America. *Emerg Infect Dis* 2020, *26*, 1810–1817, doi:10.3201/eid2608.190986.
20. Donato, C.; Vijaykrishna, D. The Broad Host Range and Genetic Diversity of Mammalian and Avian Astroviruses. *Viruses* 2017, *9*, doi:10.3390/v9050102.
21. To, K.K.W.; Chan, W.M.; Li, K.S.M.; Lam, C.S.F.; Chen, Z.; Tse, H.; Lau, S.K.P.; Woo, P.C.Y.; Yuen, K.Y. High prevalence of four novel astrovirus genotype species identified from rodents in China. *J Gen Virol* 2017, *98*, 1004–1015, doi:10.1099/jgv.0.000766.
22. Cortez, V.; Meliopoulos, V.A.; Karlsson, E.A.; Hargest, V.; Johnson, C.; Schultz-Cherry, S. Astrovirus Biology and Pathogenesis. *Annu Rev Virol* 2017, *4*, 327–348, doi:10.1146/annurev-virology-101416-041742.

Figures




Figure 1

Heatmap based on the normalized sequence reads of 35 families of viruses in each sample. The horizontal axis was regarded as the sample name. Portrait-axis is species information. The diagram on the left side of the clustering tree is a species of tree. The below the clustering tree is a sample tree. The boxes colored from blue to red represent the metagenomic sequencing reads observed.




Figure 2

A, B, and C represent the relative abundance of families, genera, and species, respectively, identified in each sample. The Y-axis represents the ratio annotation to a certain species, while the X-axis lists the sample names. The corresponding color blocks show the species category (legend on the right side).




Figure 3

Phylogenetic relationships between the novel rodent astrovirus described in this study and other Rodent astrovirus isolates (A), concatenated ORFs, (B) RF1ab, and (C) ORF2 sequences. All three trees were constructed using the neighbor-joining maximum composite likelihood method with 1,000 bootstrap replicates. The viruses found in this study are marked with ●; the scale bar indicates genetic distance.




Figure 4

The new rodent astrovirus was detected in BHK-21 cells. BHK-21 cells were inoculated with supernatants collected from CPE-positive cells inoculated with *Myodes rufocanus* faecal samples for 36 h. The astrovirus in the samples was detected by IFA (A) and qRT-PCR (B). The nucleus was stained with DAPI.