Prognostic Values of the Gross Volume of Metastatic Lymph Nodes in Patients with Esophageal Squamous Cell Carcinoma Treated with Definitive Concurrent Chemoradiotherapy

Yang Li

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

<https://orcid.org/0000-0002-6215-1272>

Yanqi Li

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Fuyi Zhu

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Xiaoying Wei

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Zhoubo Guo

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Tian Zhang

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Xi Chen

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Hui Wei

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Wencheng Zhang

Tianjin Medical University Cancer Institute and Hospital: Tianjin Tumor Hospital

Qingsong Pang

Tianjin Medical University Cancer Institute and Hospital :Tianjin Tumor Hospital

Ping Wang (✉ wangping@tjmuch.com)

Tianjin Medical University Cancer Institute and Hospital :Tianjin Tumor Hospital

Research

Keywords: esophageal cancer, chemoradiotherapy, gross tumor volume of metastatic lymph nodes, prognosis

Posted Date: October 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-960346/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Purpose We aim to explore whether the gross volume of metastatic lymph nodes (GTVnd) and the gross volume of primary tumor (GTVp) could be prognostic factors for esophageal squamous cell carcinoma (ESCC) patients treated with definitive concurrent chemoradiotherapy (dCCRT). **Methods** We retrospectively analyzed 252 ESCC patients treated with dCCRT in the era of intensity-modulated radiation therapy (IMRT) at our institution. The cut-off value for the GTVnd derived from the restricted cubic splines (RCS) was determined. Univariate and multivariate Cox proportional hazard models were performed to determine the association between GTVnd and prognosis. we performed recursive partitioning analysis (RPA) method using GTVnd to develop a new risk stratification (TGTVndM). Moreover, the linear trend χ^2 , likelihood ratio χ^2 , and akaike information criterion (AIC) were used to determine the prognostic value between the TNM and TGTVndM staging systems. **Results** The five-year overall survival (OS) rate was 30.6%, with a median follow-up of 38 months. The cut-off value of GTVnd determined by the RCS was 4.35 cm³. GTVnd $\geq$ 4.35 cm³ was an independent and significant negative prognostic factor for OS (HR=1.949, P<0.001), progression free survival (PFS) (HR=1.425, P=0.048), and distance metastasis free survival (DMFS) (HR=2.548, P=0.001). In multivariable analysis, gender, clinical T stage, and GTVnd were independently associated with OS. RPA segregated patients into 3 prognostic groups: high risk (T1-4 GTVnd $\geq$ 4.35, n=126, III stage), intermediate risk (T4 GTVnd<4.35, n=38, II stage), and low risk (T1-3 GTVnd<4.35, n=88, I stage). The 5-year OS (P<0.001), PFS (P=0.002), and DMFS (P=0.001) were significantly worse in high-risk group in comparison with the intermediate and low risk groups. Compared with the TNM staging system, the clinical T stage combined with GTVnd (TGTVndM) had a higher linear trend χ^2 (26.38 versus 25.77), higher likelihood ratio χ^2 (24.39 versus 20.69), and lower AIC (1255.07 versus 1260.06). **Conclusions** GTVnd may serve as a good prognostic factor in predicting distant metastasis and death for ESCC patients treated with dCCRT. The TGTVndM staging system demonstrated superior accuracy for predicting OS and could serve as a more effective prognostic guidance for unresectable ESCC patients.

Introduction

Esophageal cancer (EC) is the sixth most common cancer and the fourth most common cause of cancer-related death in China, with 324,422 new cases and 301,135 deaths estimated based on the Global Cancer Incidence, Mortality and Prevalence 2020 (GLOBOCAN <https://gco.iarc.fr/>). Histologically, esophageal squamous cell carcinoma (ESCC) is the predominant subtype of EC, accounts for nearly 90% of EC worldwide^{1,2}. Concurrent chemoradiotherapy has been widely accepted as standard regimen for unresectable locally advanced EC patients according to RTOG-85-01 trial³. Despite some advances in past decades, the prognosis remains poor with 5-year OS rate about 20% for all stages combined⁴. Thus, finding effective prognostic factors enable to identify patients with poor prognosis after therapy and optimally define risk adapted treatment strategies to further improve the survival rate in ESCC patients.

The TNM staging established by the American Joint Committee on Cancer (AJCC) is considered as a useful staging system to predict outcomes of EC patients underwent surgical resection^{5,6}. However, TNM staging is a tool based on pathological anatomy. It is difficult to obtain accurate TNM staging for ESCC patient with dCCRT. And survival of ESCC patients receiving dCCRT with the same clinical TNM stage varies widely. Therefore, it is far from accurate and sensitive enough to use only TNM staging to predict the prognosis of these patients. At present, there is still a lack of sufficient indicators to predict the effect of definitive chemoradiotherapy on ESCC patients.

With the advancement of intensity-modulated radiotherapy (IMRT) technique, more and more precise data on volume of primary tumor and metastatic lymph nodes are available to be collected, making it possible to study the influence of volumetric parameters on the outcomes of patients with ESCC. In this study, we propose that the GTVp and GTVnd may have an impact on the survival of patients based on our clinical experience. Therefore, we investigated ESCC patients treated with dCCRT to identify the effect of GTVp and GTVnd on OS, PFS, DMFS, local recurrence-free survival (LRFS), and regional recurrence-free survival (RRFS).

Patients And Methods

Patients

We performed a retrospective study in patients with pathologically confirmed inoperable ESCC who were treated with dCCRT in the era of IMRT at Tianjin Medical University Cancer Institute and Hospital from 2010 to 2019. The main inclusion criteria were as follows: 1) pathologically confirmed ESCC; 2) Karnofsky performance status (KPS) score ≥ 70 ; 3) no distant organ metastasis; 4) no history of a concomitant or previous malignancy; 5) underwent IMRT-based definitive concurrent chemoradiotherapy; and 6) had unresectable EC or refused surgery. A total of 252 patients with complete clinical and treatment information met the above criteria and were selected for analysis. The study was approved by the Ethics Committee of Tianjin Medical University Cancer Institute and Hospital. The patients were not required to sign an informed consent form for this retrospective study.

Clinical Stage

All patients underwent complete evaluations before treatment including detailed medical history, physical examination, upper gastrointestinal radiography, neck, chest and abdominal contrast enhanced computed tomography (CT), endoscopy with biopsy, endoscopic ultrasound, and external ultrasonography of the neck. Patients were re-staged according to the 8th staging system of the AJCC based on all information provided by imaging examination.

Treatment

All ESCC patients received dCCRT and radiotherapy was performed as IMRT. All treatments were planned based on CT simulation planning system with and without contrast and a slice thickness of 2.5 to 5 mm

throughout the entire neck, thorax and upper abdominal under shallow breathing. Primary tumor and metastatic lymph nodes >1 cm (≥ 5 mm in tracheoesophageal groove) were contoured as primary gross tumor volume (GTVp) and gross tumor volume of metastatic lymph nodes (GTVnd) based on both physical examination and all available diagnostic images. The clinical target volume (CTV) was defined as the visible GTVp, GTVnd and subclinical regions at risk for involvement. The planning gross target volume (PGTV) was obtained by adding an isotropic margin of 0.5 cm to the GTVp combined with GTVnd. The planning target volume (PTV) was defined as the CTV plus a 0.5 cm margin in all directions. Two radiation oncologists reviewed all contoured structures to ensure accuracy and consistency. The dosimetric parameters for GTVp and GTVnd were calculated from the Pinnacle system for every patient.

Patients received concurrent chemotherapy with a weekly or three-weekly schedule of paclitaxel and platinum-based drugs. For subsequent consolidation chemotherapy, chemotherapy regimens were selected based on patient age, general physical condition, and physician judgment.

Endpoints

The end points included the OS, PFS, DMFS, LRFS, and RRFS. We defined OS as the time from the first treatment to the date of death from any cause or the date of last follow-up. The PFS was calculated from the time of first treatment to disease progression, including local recurrence (LR), regional recurrence (RR), and distant metastasis (DM). The DMFS was set for the period from the date of treatment to distant metastasis. LRFS and RRFS were defined as the intervals between the beginning of treatment and the date of recurrence that occurred in the esophageal lumen, and between the beginning of treatment and the date of regional lymph nodes. Patients who did not experience an event of interest were censored at their last follow-up or the date of death.

Follow-up

Patients were follow-up at least every 3 months during the first 2 years, every 6 months for 2 to 5 years, and then annually thereafter. The imageological examination including chest CT scans, upper gastrointestinal radiography, and neck and upper abdomen ultrasonography were routinely performed at each follow-up study. In addition, physical examinations, routine blood tests and liver and kidney function analyses were also necessary. The results of the re-examination were confirmed by electronic medical records, paper medical documents, and telephone. If the patients died by the time of contact, available family members provided the needed information.

Statistical analysis

Categorical variables were summarized using frequencies and percentages, and the χ^2 test and Fisher's exact test were used to compare the differences among different groups. Continuous variables were presented as the median values and interquartile range (IQR). Age, KPS, radiation dose and primary tumor length were categorized with the median value as the cut-off. With the utilization of RCS curves, the association between the GTVnd (as a continuous variable) and OS, PFS, DMFS, RRFS were evaluated

based on Cox regression model⁷. The Kaplan-Meier method was used to plot the survival curves and the log-rank test was used to compare the survival distributions. We used univariate and multivariate Cox proportional hazard models to evaluate the influence of different variables on the OS, PFS, DMFS, RRFS, and LRFS. Adjusted hazard ratios (HR) were obtained with corresponding 95% confidence intervals (95% CIs) from the Cox regression analysis. *P* values were calculated using the Cox regression forward-LR model. Additionally, we use the linear trend χ^2 test to measure the discriminatory ability and monotonicities, the likelihood ratio χ^2 test based on the Cox regression model to the homogeneity, and the AIC to the optimal prognostic stratifications between the two staging systems⁸. A higher linear trend χ^2 score or likelihood ratio χ^2 score and a lower AIC value indicates a better model for predicting prognosis⁹. We use the online web server called autoRPA (available at <http://rpa.renlab.org>) to establish a decision-making tree from survival data based on the RPA algorithm and log-rank test statistics to correctly stratify risk in the target population¹⁰. All statistical tests were two-sided and *P* values < 0.05 were considered statistically significant. Analyses were conducted using SPSS v22.0 (IBM SPSS, New York) and the R 4.1.0 (<http://www.r-project.org/>).

Results

Patient characteristics

We identified 252 patients with pathologically confirmed inoperable ESCC treated with dCCRT from 2010 to 2019 in our study, with complete information of GTVp and GTVnd. The demographic characteristics of the ESCC patients are shown in Table 1. The age of the patients ranged from 30 to 82 years with a median age of 60 years. Median GTVp and GTVnd were 38.77 cm³ (interquartile range (IQR), 24.00-58.02 cm³) and 4.34 cm³ (IQR, 0.18-11.38 cm³), respectively. And the median ratio between the two (GTVnd/GTVp) was 0.11 with an IQR of 0.01-0.30.

Table 1. Characteristics of the patients with esophageal squamous cell carcinoma treated with definitive concurrent chemoradiotherapy (n=252)

Patient characteristics	No of patients (%)
Age	
Median	60
Range	30-82
Gender	
Male	220(87.3)
Female	32(12.7)
KPS	
≥90	173(68.7)
<90	79(31.3)
Weight loss	
Yes	89(35.3)
No	163(64.7)
Pain of chest and back	
Yes	51(20.2)
No	201(79.8)
Tumor location	
Cervical	22(8.7)
Upper thoracic	78(31.0)
Middle thoracic	93(36.9)
Lower thoracic	59(23.4)
Clinical T stage, 8th	
T1	9(3.6)
T2	16(6.3)
T3	139(55.2)
T4	88(34.9)
Clinical N stage, 8th	
N0	59(23.4)
N1	86(34.1)

N2	77(30.6)
N3	30(11.9)
Clinical TNM stage, 8th	
I	7(2.8)
II	42(16.7)
III	102(40.5)
IV	101(40.1)
GTVp (cm ³)	
Median	38.77
IQR	24.00-58.02
GTVnd (cm ³)	
Median	4.34
IQR	0.18-11.38
GTVnd/GTVp	
Median	0.11
IQR	0.01-0.30
Tumor length, cm	
≤6	126(50.0)
>6	126(50.0)
Radiation dose, Gy	
<54	75(29.8)
≥54	177(70.2)
Consolidation chemotherapy	
Yes	106(42.1)
No	146(57.9)

Treatment outcomes

Within the median duration of follow-up for the whole patients of 38 months (ranging from 1 to 89 months), we identified 117 (46.4%) patients developing locoregional recurrence (LRR), 62 (24.6%) DM and

132(52.4%) death. The five-year OS, PFS, DMFS, LRFS and RRFS rates for the entire cohort were 30.6%, 28.4%, 58.0%, 46.9%, and 66.7%, respectively.

Prognostic value of volumetric parameters on survivals

GTVp, GTVnd and GTVnd/GTVp were segregated into two groups according to their median values. Subsequently, we investigated the prognostic role of the volumetric parameters using univariate and multivariate Cox proportional hazard regression. In univariate analysis (Table 2), GTVnd $\geq 4.34 \text{ cm}^3$ and GTVnd/GTVp ≥ 0.11 were associated with poorer OS (both $P < 0.001$), PFS ($P = 0.001$; $P = 0.014$; respectively), DMFS (both $P < 0.001$) and RRFS ($P = 0.023$; $P = 0.031$; respectively). Then the variables with statistical significance in univariate analysis ($P < 0.05$) were included in the multivariate analysis and P values were calculated using the Cox regression forward-LR model. Multivariable analyses validated the independent prognostic role of the GTVnd in OS ($\text{HR} = 1.949$, $P < 0.001$), PFS ($\text{HR} = 1.425$, $P = 0.048$) and DMFS ($\text{HR} = 2.548$, $P = 0.001$). In addition, the result demonstrated advanced clinical T stage ($P = 0.002$) and male ($P = 0.046$) were independently associated with shorter OS, shown as Table 3.

Table 2 Univariate analysis of prognostic factors in patients with ESCC (n=252)

Variable	OS <i>P</i>	PFS <i>P</i>	DMFS <i>P</i>	RRFS <i>P</i>	LRFS <i>P</i>
Age (>60 years vs. ≤ 60 years)	0.796	0.923	0.412	0.116	0.640
Gender	0.034	0.005	0.025	0.016	0.084
KPS (≥ 90 vs. < 90)	0.343	0.043	0.048	0.144	0.105
Weight loss	0.122	0.637	0.510	0.516	0.628
Pain of chest and back	0.973	0.665	0.167	0.602	0.783
Tumor location	0.010	< 0.001	0.620	0.015	0.059
Clinical T stage	0.001	0.011	0.184	0.324	0.103
Clinical N stage	0.001	0.341	0.152	0.639	0.923
GTVp (≥ 38.77 vs. $< 38.77 \text{ cm}^3$)	0.167	0.496	0.945	0.475	0.480
GTVnd (≥ 4.34 vs. $< 4.34 \text{ cm}^3$)	< 0.001	0.001	< 0.001	0.023	0.464
GTVnd/GTVp (≥ 0.11 vs. < 0.11)	< 0.001	0.014	< 0.001	0.031	0.887
Tumor length (>6cm vs. ≤ 6 cm)	0.048	0.290	0.847	0.259	0.342
Radiation dose (≥ 54 Gy vs. < 54 Gy)	0.242	0.127	0.060	0.093	0.110
Consolidation chemotherapy	0.633	0.189	0.423	0.110	0.478

Table 3 Multivariate cox proportional hazard regression analysis of prognostic factors in patients with ESCC (n=252)

	HR (95%CI)	P
OS		
T stage		0.002
T stage (T2 vs. T1)	2.492(0.290-21.405)	0.405
T stage (T3 vs. T1)	3.341(0.472-24.925)	0.223
T stage (T4 vs. T1)	6.134(0.844-44.550)	0.073
GTVnd (≥ 4.34 vs. $< 4.34\text{cm}^3$)	1.949(1.353-2.808)	< 0.001
Gender (male vs. female)	1.939(1.013-3.712)	0.046
PFS		
Gender (male vs. female)	2.373(1.242-4.535)	0.009
GTVnd (≥ 4.34 vs. $< 4.34\text{cm}^3$)	1.425(1.003-2.024)	0.048
Tumor location		0.001
Tumor location (UT vs. Cervical)	0.274(0.146-0.514)	< 0.001
Tumor location (MT vs. Cervical)	0.543(0.310-0.950)	0.032
Tumor location (LT vs. Cervical)	0.475(0.261-0.8)	0.015
DMFS		
GTVnd (≥ 4.34 vs. $< 4.34\text{cm}^3$)	2.548(1.491-4.355)	0.001

To confirm the optimal GTVnd cutoff value, we used the RCS with 3 knots and OS, PFS, DMFS, and RRFS as endpoint events. The cutoff value determined by RCS was 4.35 cm^3 in this analysis (Fig.1).

Multivariable analysis also showed that $\text{GTVnd} \geq 4.35\text{ cm}^3$ was independent and significant negative prognostic factors for OS, PFS and DMFS.

Construction of Risk Grouping Using GTVnd by RPA model

Considering the prognostic value of GTVnd, we then performed RPA algorithm including T stage, GTVnd/GTVp, and gender to develop a new staging. The significant RPA-derived splits were only the T stage and GTVnd (Fig.2). The RPA model divided the 252 ESCC patients into the following three groups:

high risk (T1-4 GTVnd $\geq$ 4.35, n=126, III stage) , intermediate risk (T4 GTVnd<4.35,n=38,II stage), and low risk(T1-3GTVnd $\leq$ 4.35, n=88, I stage).

Clinical and treatment characteristics of the new risk grouping is shown in Table 4. The differences in age, gender, KPS, weight loss, pain of chest and back, tumor length, radiation dose, induction chemotherapy and consolidation chemotherapy were not statistically significant between the two groups (P > 0.05). Compared with those in the high-risk group, patients in the low-risk and intermediate-risk groups whose tumor were more located in the cervical and upper thoracic (48.9%, 52.6% vs.29.3%, P=0.033). And the proportion of N0-1 was higher in low-risk and intermediate-risk groups than that in the high-risk group (87.5%, 81.6% vs. 44.8%, P<0.001).

Table 4. Baseline characteristics of patients stratified by new risk stratification

Characteristic	Low-risk group (n=88)	Intermediate-risk group (n=38)	High-risk group (n=126)	<i>P</i>
Age				0.751
≤60	44(50.0%)	19(50.0%)	69(54.8%)	
>60	44(50.0%)	19(50.0%)	57(45.2%)	
Gender				0.165
Male	76(86.4%)	30(78.9%)	114(90.5%)	
Female	12(13.6%)	8(21.1%)	12(9.5%)	
KPS				0.083
≥90	53(60.2%)	26(68.4%)	94(74.6%)	
<90	35(39.8%)	12(31.6%)	32(25.4%)	
Weight loss				0.127
Yes	27(30.7%)	10(26.3%)	52(41.3%)	
No	61(69.3%)	28(73.7%)	74(58.7%)	
Pain of chest and back				0.441
Yes	14(15.9%)	8(21.1%)	29(23.0%)	
No	74(84.1%)	30(78.9%)	97(77.0%)	
Tumor location				0.033
Cervical	7(8.0%)	4(10.5%)	11(8.7%)	
Upper thoracic	36(40.9%)	16(42.1%)	26(20.6%)	
Middle thoracic	28(31.8%)	12(31.6%)	53(42.1%)	
Lower thoracic	17(19.3%)	6(15.8%)	36(28.6%)	
Clinical T stage, 8th				<0.001
T1	7(8.0%)	0(0.0%)	2(1.6%)	
T2	12(13.6%)	0(0.0%)	4(3.2%)	
T3	69(78.4%)	0(0.0%)	70(55.6%)	
T4	0(0.0%)	38(100.0%)	50(39.7%)	
Clinical N stage, 8th				<0.001
N0	37(42.0%)	21(55.3%)	1(0.8%)	

N1	40(45.5%)	10(26.3%)	36(28.6%)
N2	9(10.2%)	3(7.9%)	65(51.6%)
N3	2(2.3%)	4(10.5%)	24(19.0%)
GTVnd			<0.001
≥4.35	0(0.0%)	0(0.0%)	126(100.0%)
<4.35	88(100.0%)	38(100.0%)	0(0.0%)
Tumor length, cm			0.211
≤6	49(55.7%)	21(55.3%)	56(44.4%)
>6	39(44.3%)	17(44.7%)	70(55.6%)
Radiation dose, Gy			0.397
<54	26(29.5%)	8(21.1%)	41(32.5%)
≥54	62(70.5%)	30(78.9%)	85(67.5%)
Consolidation chemotherapy			0.174
Yes	33(37.5%)	21(55.3%)	52(41.3%)
No	55(62.5%)	17(44.7%)	74(58.7%)

The prognostic significance of the risk group

Then, we performed the Kaplan-Meier analysis to compare OS, PFS, DMFS, RRFS, and LRFS between the three groups derived by RPA (Fig.3). We found highly significant differences in OS among the three groups ($P < 0.001$; Fig. 3A), with corresponding 5-year OS rates of 17.0 % for high-risk group, 26.3% for intermediate-risk group, and 54.0% for low-risk group. The 5-year PFS rates of high-risk group, intermediate-risk group and low-risk group were 16.6%, 27.6% and 39.8%, respectively. By the log-rank test, there were significant differences in PFS among the three groups ($P=0.002$; Fig.3B). And the 5-year DMFS rates of 37.8% for patients with high-risk group and 72.0% for those with low-risk group showed significant differences ($P =0.001$; Fig.3C). Though a prognostic analysis demonstrated the 5-year RRFS rate in the high-risk group (59.1%) was lower than that of the intermediate-risk group (66.0%) and low-risk group (76.6%), the difference was not significant ($P=0.063$; Fig.3D). No significant difference in LRFS rate was observed among the three groups ($P=0.194$; Fig.3E).

Comparison of the performance of the new risk grouping and TNM staging system

The performance between the new risk grouping (TGTVndM) and TNM staging systems assessed by linear trend χ^2 , likelihood ratio χ^2 , and the AIC is presented in Table 5. The GTVndM staging system demonstrated higher linear trend χ^2 (26.38 versus 25.77), high likelihood ratio χ^2 (24.39 versus 20.69),

and lower AIC (1255.07 versus 1260.06) compared with the TNM stage, indicating the optimum prognostic stratification in predicting the survival of ESCC patients treated with dCCRT.

Table 5 Comparison of the prognostic performance of the TGTVndM and TNM staging systems

Classification	Subgroups	Figure	Linear trend χ^2	Likelihood ratio χ^2	AIC
N stage	N0, N1, N2, N3	SA	5.06	11.68	1271.29
GTVnd stage	GTVnd<4.35cm ³ GTVnd≥4.35cm ³	SB	16.23	18.48	1260.23
TNM stage	I, II, III, IV	SC	25.77	20.69	1260.06
TGTVndM stage	I, II, III	3A	26.38	24.39	1255.07

Discussion

Previous studies had reported the significant association between tumor volume and outcomes in EC patients treated with radical surgery (with or without neoadjuvant chemoradiotherapy) or definitive concurrent chemoradiotherapy^{11–19}. In our routine clinical practice, we found that EC patients with the same TNM stage and undergoing the same treatment might show considerably different clinical outcomes. We hypothesized that some patients with small tumors size and large lymph nodes metastases and those with large tumors size and small lymph nodes might had different prognosis. Indeed, the results of our study supported this hypothesis. In our study, we identified the GTVnd as an independent prognostic factor for OS, PFS and DMFS. Additionally, we identified a value of 4.35 as the optimal cut-off as defined by the RCS. The results showed that patients with GTVnd≥4.35 cm³ had shorter OS and PFS time and more often developed DM compared with GTVnd < 4.35 cm³. This is in agreement with previously reported findings. Sakanaka et al. reported 144 patients with thoracic ESCC who underwent definitive chemoradiotherapy with large metastatic lymph nodes were at high risk of DM in 2016²⁰. In addition, Zhao et al retrospectively reviewed 376 ESCC patients treated with definitive (chemo-) radiotherapy and concluded that bulky lymph nodes were associated statistically with distant failure and poorer OS²¹. It was reported that lymph nodes metastases were more closely associated with systemic micro-metastases than primary tumor progression and a better lymph nodes response to neoadjuvant chemotherapy in patients with metastatic EC predicted a better survival and less lymphatic,

distant metastases^{20,22,23}. Our study provides a scientific basis for the close correlation between lymph nodes metastases and tumor micro-metastases.

Then, we constructed a new prognostic model to divide ESCC patients into high-risk, intermediate-risk and low-risk groups using RPA method. The GTVnd was the first split and the second was clinical T stage in our RPA model. Specially, patients who were T1-4 and $GTVnd \geq 4.35 \text{ cm}^3$ were classified as high-risk group, T4 and $GTVnd < 4.35 \text{ cm}^3$ as intermediate-risk group and T1-3 and $GTVnd < 4.35$ as low-risk group. We found that patients with high-risk group exhibited shorter OS, PFS and DMFS compared with the other two groups and we also showed that the new staging system (TGTVnd) was superior to the traditional TNM staging system in predicting OS of ESCC patients treated with dCCRT. A previous study conducted by Chen et.al demonstrated that GTV and maximum diameter of metastatic lymph nodes (MDMLN) predicted survival of nonsurgical EC patients more accurately than the 8th edition of AJCC/UICC clinical staging system¹⁵. Another study proposed a new nonsurgical staging system based on the gross tumor volume of the primary tumor and N to be better predict the outcome of ESCC patients¹⁷. The majority of EC patients receiving dCCRT are diagnosed with advanced disease (stage III/IV), the survival of these patients varies widely and an effective staging system is desperately needed. The accuracy of CT for predicting the N staging was 78% in EC patients²⁴, which made inaccuracy of TNM staging to predict recurrence and prognosis to some extent. Therefore, we built a new risk staging system based on various combinations of T subgroups and GTVnd subgroups to better evaluate the outcomes of inoperable ESCC patients.

Moreover, the multivariate analysis also revealed that clinical T stage and gender were independent prognostic factors affecting OS for ESCC patients. A study from Taiwan analyzed 14,394 ESCC patients and indicated that sex and clinical T were independent prognostic factors for OS²⁵. This large sample cohort study supported the conclusions of our research. The multicenter studies at home and abroad demonstrated that men exhibited a worse prognosis than women in EC^{26,27}. The reason for the discrepancy is not clear. Sex hormone might be one of the reasons to the difference in prognosis, and early preclinical research showed that estrogens significantly inhibited the growth of esophageal carcinoma cell²⁸. Another possible explanation could be differences in long term smoking history, heavy alcohol consumption and nutritional status between men and women^{29,30}.

We found that the location of primary tumor was independently associated with recurrence in our study. It has been reported in previous studies that tumor location was an independent prognostic factor for PFS in EC patients^{31,32}. Patients with cervical ESCC had the worse PFS rate (3-year PFS, 18.7%), compared with patients with thoracic ESCC (3-year PFS, 36.4%, $P=0.002$) in our study. Münch. et al. including 95 ESCC patients treated with either dCCRT or neoadjuvant chemoradiation followed by surgery reported that proximal tumor location was associated with short PFS³², which was consistent with the results of ours. The location of the primary tumor relating to the mode of tumor recurrence might contribute to the difference^{23,32}, to be specific, the local–regional recurrence rate was 72.2% for cervical ESCC while 43.9% for thoracic ESCC, showing a significant difference by the chi-squared test ($P=0.01$) in our study. Yang

et.al reviewed 1220 thoracic ESCC patients who underwent complete resection to conclude that tumor location did not affect survival prognosis³³ and subsequent a study conducted by Shi et.al reported a decreased OS in pathological T2-3N0M0 ESCC patients with proximal tumor location³⁴. However, the different tumor locations have been used different treatment making comparisons across studies difficult. For example, dCCRT is the general recommendation for patients with cervical tumor location^{35,36} and surgery has been the standard treatment for thoracic EC³⁷. Overall, we should not neglect the tumor location as an important prognostic factor.

This study has several limitations, first, with the retrospective design based on case records, which may lead to bias in patient selection. Second, the difficulty in delineating metastatic lymph nodes due to partial lymph nodes conglomerated with primary tumor may cause certain error in a small number of patients. Finally, further studies with larger sample size and external validation are required to confirm our results. However, one strength of our study is that we investigated the effect of volumetric parameters on prognosis for patients with ESCC treated with dCCRT in the era of IMRT, which appears to be simple and accurate prognostic factors to be obtained. Moreover, we also constructed a prognostic risk group for inoperable ESCC patients, which might be a valid indicator for predicting the sensitivity to chemoradiotherapy.

Conclusions

In conclusion, the present study showed that GTVnd was independently prognostically significant for OS, PFS and DMFS. The optimum cut-off point for GTVnd was 4.35 cm³ in predicting distant metastasis and death for ESCC patients treated with dCCRT. The new risk groups stratify these patients into I, II, and III subgroups to better assess the prognosis and guide the treatment. The results need validation from prospective randomized studies.

Abbreviations

GTVnd: gross volume of metastatic lymph nodes

GTVp: gross volume of primary tumor

ESCC: esophageal squamous cell carcinoma

dCCRT: definitive concurrent chemoradiotherapy

IMRT: intensity-modulated radiation therapy

RCS: restricted cubic splines

RPA: recursive partitioning analysis

AIC: akaike information criterion

OS: overall survival

PFS: progression free survival

DMFS: distance metastasis free survival

LRFS: local recurrence-free survival

RRFS: regional recurrence-free survival

CTV: clinical target volume

PGTV: planning gross target volume

PTV: planning target volume

Declarations

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

None

Funding

This work was supported by grants from the Chinese National Key Research and Development Project [No. 2018YFC1315601].

Author information

Affiliation

Department of Radiation Oncology, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin,China;

Yang Li, Yanqi Li, Fuyi Zhu, Xiaoying Wei, Zhoubo Guo, Tian Zhang, Xi Chen, Hui Wei, Wencheng Zhang, Qingsong Pang & Ping Wang

Department of Thoracic Oncology, Lung Cancer Diagnosis and Treatment Center, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy of Tianjin, Tianjin's Clinical Research Center for Cancer, Tianjin, China.

Fuyi Zhu

Contributions

(1)Conception and design: P Wang, Q Pang, W Zhang; (2) Manuscript writing: Y Li; (3) Data analysis and interpretation: Y Li, YQ Li, FY Zhu;(4) Data collection: XY Wei, ZB Guo(5) Final approval of manuscript: All authors.

Corresponding author

Correspondence to Wencheng Zhang, Qingsong Pang & Ping Wang

Ethics declarations

Ethics approval and consent to participate

The study was approved by the Ethics Committee of Tianjin Medical University Cancer Institute and Hospital. The patients were not required to sign an informed consent form for this retrospective study.

Consent for publication

All authors have approved the manuscript and agree with submission to Radiation Oncology.

Competing interests

The authors have no conflicts of interest to declare.

References

1. Arnold M, Soerjomataram I, Ferlay J, et al: Global incidence of oesophageal cancer by histological subtype in 2012. *Gut* 64:381-7, 2015
2. Domper Arnal MJ, Ferrandez Arenas A, Lanas Arbeloa A: Esophageal cancer: Risk factors, screening and endoscopic treatment in Western and Eastern countries. *World J Gastroenterol* 21:7933-43, 2015
3. Cooper JS, Guo MD, Herskovic A, et al: Chemoradiotherapy of locally advanced esophageal cancer: long-term follow-up of a prospective randomized trial (RTOG 85-01). Radiation Therapy Oncology Group. *JAMA* 281:1623-7, 1999
4. Siegel RL, Miller KD, Fuchs HE, et al: Cancer Statistics, 2021. *CA Cancer J Clin* 71:7-33, 2021
5. Sudo N, Ichikawa H, Muneoka Y, et al: Clinical Utility of ypTNM Stage Grouping in the 8th Edition of the American Joint Committee on Cancer TNM Staging System for Esophageal Squamous Cell Carcinoma. *Ann Surg Oncol* 28:650-660, 2021
6. Ajani JA, Correa AM, Hofstetter WL, et al: Clinical parameters model for predicting pathologic complete response following preoperative chemoradiation in patients with esophageal cancer. *Ann Oncol* 23:2638-2642, 2012

7. Johannesen CDL, Langsted A, Mortensen MB, et al: Association between low density lipoprotein and all cause and cause specific mortality in Denmark: prospective cohort study. *Bmj* 371:m4266, 2020
8. Hsu PK, Wu YC, Chou TY, et al: Comparison of the 6th and 7th editions of the American Joint Committee on Cancer tumor-node-metastasis staging system in patients with resected esophageal carcinoma. *Ann Thorac Surg* 89:1024-31, 2010
9. Lu J, Zheng CH, Cao LL, et al: The effectiveness of the 8th American Joint Committee on Cancer TNM classification in the prognosis evaluation of gastric cancer patients: A comparative study between the 7th and 8th editions. *Eur J Surg Oncol* 43:2349-2356, 2017
10. Xie Y, Luo X, Li H, et al: autoRPA: A web server for constructing cancer staging models by recursive partitioning analysis. *Comput Struct Biotechnol J* 18:3361-3367, 2020
11. Xu Q, Liu ZK, Cao YK, et al: [Relationship of gross tumor volume with lymph node metastasis and prognosis of esophageal carcinoma]. *Zhonghua Zhong Liu Za Zhi* 34:684-7, 2012
12. Huang R, Guo H, Chen J, et al: Intratreatment Tumor Volume Change During Definitive Chemoradiotherapy is Predictive for Treatment Outcome of Patients with Esophageal Carcinoma. *Cancer Manag Res* 12:7331-7339, 2020
13. Boggs DH, Hanna A, Burrows W, et al: Primary Gross Tumor Volume is an Important Prognostic Factor in Locally Advanced Esophageal Cancer Patients Treated with Trimodality Therapy. *J Gastrointest Cancer* 46:131-7, 2015
14. Lv J, Gan H, Zhang W, et al: Relative tumor volume is a better independent prognostic factor in esophageal squamous cell carcinoma: Results of a retrospective study. *Medicine (Baltimore)* 98:e14963, 2019
15. Chen J, Lin Y, Cai W, et al: A new clinical staging system for esophageal cancer to predict survival after definitive chemoradiation or radiotherapy. *Dis Esophagus* 31, 2018
16. Chen CZ, Chen JZ, Li DR, et al: Long-term outcomes and prognostic factors for patients with esophageal cancer following radiotherapy. *World J Gastroenterol* 19:1639-44, 2013
17. Chen M, Li X, Chen Y, et al: Proposed revision of the 8th edition AJCC clinical staging system for esophageal squamous cell cancer treated with definitive chemo-IMRT based on CT imaging. *Radiat Oncol* 14:54, 2019
18. Yamashita H, Takenaka R, Okuma K, et al: Prognostic factors in patients after definitive chemoradiation using involved-field radiotherapy for esophageal cancer in a phase II study. *Thorac Cancer* 7:564-569, 2016
19. Chen Y, Zhang Z, Jiang G, et al: Gross tumor volume is the prognostic factor for squamous cell esophageal cancer patients treated with definitive radiotherapy. *J Thorac Dis* 8:1155-61, 2016
20. Sakanaka K, Ishida Y, Itasaka S, et al: Identification of a predictive factor for distant metastasis in esophageal squamous cell carcinoma after definitive chemoradiotherapy. *Int J Clin Oncol* 21:899-908, 2016
21. Zhao Z, Zhang Y, Wang X, et al: The Prognostic Significance of Metastatic Nodal Size in Non-surgical Patients With Esophageal Squamous Cell Carcinoma. *Front Oncol* 10:523, 2020

22. Urakawa S, Makino T, Yamasaki M, et al: Lymph Node Response to Neoadjuvant Chemotherapy as an Independent Prognostic Factor in Metastatic Esophageal Cancer. *Ann Surg* 273:1141-1149, 2021
23. Doki Y, Ishikawa O, Takachi K, et al: Association of the primary tumor location with the site of tumor recurrence after curative resection of thoracic esophageal carcinoma. *World J Surg* 29:700-7, 2005
24. Wu LF, Wang BZ, Feng JL, et al: Preoperative TN staging of esophageal cancer: comparison of miniprobe ultrasonography, spiral CT and MRI. *World J Gastroenterol* 9:219-24, 2003
25. Cheng YF, Chen HS, Wu SC, et al: Esophageal squamous cell carcinoma and prognosis in Taiwan. *Cancer Med* 7:4193-4201, 2018
26. Bohanes P, Yang D, Chhibar RS, et al: Influence of sex on the survival of patients with esophageal cancer. *J Clin Oncol* 30:2265-72, 2012
27. He Y, Liang D, Du L, et al: Clinical characteristics and survival of 5283 esophageal cancer patients: A multicenter study from eighteen hospitals across six regions in China. *Cancer Commun (Lond)* 40:531-544, 2020
28. Ueo H, Matsuoka H, Sugimachi K, et al: Inhibitory effects of estrogen on the growth of a human esophageal carcinoma cell line. *Cancer Res* 50:7212-5, 1990
29. Xie SH, Lagergren J: Social group disparities in the incidence and prognosis of oesophageal cancer. *United European Gastroenterol J* 6:343-348, 2018
30. Ferndale L, Aldous C, Hift R, et al: Gender Differences in Oesophageal Squamous Cell Carcinoma in a South African Tertiary Hospital. *Int J Environ Res Public Health* 17, 2020
31. Guo JC, Huang TC, Lin CC, et al: Postchemoradiotherapy Pathologic Stage Classified by the American Joint Committee on the Cancer Staging System Predicts Prognosis of Patients with Locally Advanced Esophageal Squamous Cell Carcinoma. *J Thorac Oncol* 10:1481-9, 2015
32. Munch S, Pigorsch SU, Devecka M, et al: Neoadjuvant versus definitive chemoradiation in patients with squamous cell carcinoma of the esophagus. *Radiat Oncol* 14:66, 2019
33. Yang HX, Hou X, Liu QW, et al: Tumor location does not impact long-term survival in patients with operable thoracic esophageal squamous cell carcinoma in China. *Ann Thorac Surg* 93:1861-6, 2012
34. Shi H, Zhang K, Niu ZX, et al: Does tumour location influence postoperative long-term survival in patients with oesophageal squamous cell carcinoma? *Eur J Cardiothorac Surg* 48:266-72, 2015
35. Gkika E, Gauler T, Eberhardt W, et al: Long-term results of definitive radiochemotherapy in locally advanced cancers of the cervical esophagus. *Dis Esophagus* 27:678-84, 2014
36. Buckstein M, Liu J: Cervical Esophageal Cancers: Challenges and Opportunities. *Curr Oncol Rep* 21:46, 2019
37. Mariette C, Piessen G, Triboulet JP: Therapeutic strategies in oesophageal carcinoma: role of surgery and other modalities. *Lancet Oncol* 8:545-53, 2007

Figures

Figure 1

Restricted cubic spline analysis (95% CI) for the association of gross volume of metastatic lymph nodes (GTVnd) with overall survival (OS), progression free survival (PFS), distant metastasis free survival (DMFS), and regional recurrence free survival (RRFS) in ESCC patients treated with dCCRT.

Figure 2

Recursive partitioning analysis (RPA) for OS in ESCC patients based on GTVnd and T stage

Figure 3

Comparison of survival outcomes in prognostic grouping by recursive partition analysis: (A) overall survival, (B) progression free survival, (C) distant metastasis free survival, (D) regional recurrence free

survival, (E) local recurrence free survival

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS.tif](#)