

What's the Cut-Point?: A Systematic Investigation of Tau PET Thresholding Methods

Alexandra June Weigand

University of California, San Diego <https://orcid.org/0000-0002-7931-1383>

Anne Maass

German Centre for Neurodegenerative Diseases Site Magdeburg: Deutsches Zentrum für Neurodegenerative Erkrankungen
Standort Magdeburg

Graham Eglit

University of California San Diego

Mark Bondi (✉ mbondi@health.ucsd.edu)

University of California

Research

Keywords: tau, positron emission tomography, methods, Alzheimer's disease, review

Posted Date: October 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-962722/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Alzheimer's Research & Therapy on April 5th, 2022. See the published version at <https://doi.org/10.1186/s13195-022-00986-w>.

Abstract

Background: Tau positron emission tomography (PET) is increasing in popularity for biomarker characterization of Alzheimer's disease (AD), and recent frameworks rely on tau PET cut-points to stage individuals along the AD continuum. Given the lack of standardization in tau PET thresholding methods, this study sought to systematically canvass and characterize existing studies that have derived tau PET cut-points and then directly assess different methods of tau PET thresholding in terms of their concurrent validity.

Methods: First, a literature search was conducted in PubMed to identify studies of AD and related clinical phenotypes that used the Flortaucipir (AV-1451) tau PET tracer to derive a binary cut-point for tau positivity. Of 540 articles screened and 47 full-texts reviewed, 23 cohort studies met inclusion criteria with a total of 6,536 participants. In a 2x2x2 design that systematically varied region (temporal meta-ROI and entorhinal cortex), analytic method (receiver-operating characteristics and 2 standard deviations above comparison group), and criterion/comparison variable (amyloid-beta positivity or cognitive diagnosis), a sample from the Alzheimer's Disease Neuroimaging Initiative yielded eight different tau PET cut-points.

Results: For the systematic review, notable variability in sample characteristics, preprocessing methods, region of interest, and analytic approach were observed, which were accompanied by discrepancy in proposed tau PET cut points. The empirical follow-up indicated the cut-point derived based on 2 standard deviations above a cognitively unimpaired group in the temporal meta-ROI best differentiated tau positive and negative groups on cerebrospinal fluid phosphorylated tau, Mini-Mental State Examination score, and delayed memory performance.

Conclusions: Given the impact of discrepant thresholds on tau positivity rates, biomarker staging, and eligibility for future clinical treatment trials, recommendations are offered to standardize cut-point derivations as additional tau PET tracers are developed and implemented in AD studies.

Introduction

Tau neurofibrillary tangles (NFTs), along with amyloid ($A\beta$) plaques, represent the two defining pathologic hallmarks of Alzheimer's disease (AD; Alzheimer, 1907). Although $A\beta$ has predominated AD research for the past 30 years (Hardy, 2017), the recent emergence of positron emission tomography (PET) imaging of tau NFTs has allowed for *in-vivo* characterization of this pathology and highlights the enduring importance of tau in the development and progression of AD (Jagust, 2018). Biomarker frameworks of AD, such as the amyloid-tau-neurodegeneration (AT[N]) framework, have capitalized on advancements in tau PET imaging to stage participants based on their biomarker profile and motivate research across the AD continuum using these staging efforts (Jack *et al.*, 2016, 2018).

Classification into AT(N) stages requires the designation of participants as $A\beta$, tau, and neurodegeneration positive or negative (A+/A-, T+/T-, [N+/N-]). Whereas cut-points for $A\beta$ positivity have been well-validated and consistently used (Breault *et al.*, 2017; Joshi *et al.*, 2012; Klunk *et al.*, 2015), there has been considerable methodological variance in tau PET thresholding studies, resulting in discrepant cut-points with little consistency across studies, despite evidence suggesting strong reliability of tau PET (Devous Sr *et al.*, 2018). Some of this methodological heterogeneity may be attributable to the inherent complexity of tau accumulation. In contrast to $A\beta$, which is diffusely distributed and commonly quantified as a global cortical mean tracer uptake, tau accumulation shows a hierarchical pattern of spread (Braak & Braak, 1991). Thus, the question of how to define tau positivity requires examination of both the quantity of tracer retention and its location, each of which can be defined in multiple ways. Importantly, there has not yet been a comprehensive characterization of these differing methods nor a compilation of the various tau PET cut-points currently in use. Such heterogeneity in sample characteristics and methods across studies directly influences the rates of tau positivity and, consequently, the proportions of each resultant AT(N) profile.

This study sought to clarify the impact of these methodological differences in two ways. First, we reviewed previous studies that have derived a cut-point for tau in AD and related phenotypes and comprehensively evaluated the thresholding methods used, as well as the influence of these methods on resultant cut-points. Second, to more systematically evaluate the unique influence of these methodological factors, we conducted an empirical follow-up study employing a systematic comparison that

varies only one factor at a time (e.g., varying the analytic approach while holding the sample characteristics and preprocessing methods constant). This was conducted in order to assess the independent effects of sample ROIs, analytic approach, and comparison group/criterion variable (i.e., A β positivity or cognitive diagnosis) on tau cut-points and resultant tau positivity rates. We then evaluated the criterion validity of these varying methodological approaches to better inform future research using tau PET cut-points for AT(N) classifications or other AD staging paradigms as well as clinical treatment trial eligibility.

Methods

Review. A literature search conducted between January 1, 2020 and October 1, 2020 indexed PubMed/MEDLINE databases following PRISMA guidelines (<http://www.prisma-statement.org/>) to identify studies relevant to the current review. The following search terms were used: tau PET positivity, tau PET threshold, tau PET cutoff (including cut-off and cut off), tau PET cut-point (including cut-point and cut point). The search was then repeated with “tau PET” replaced by “florbetapir” in all previous search terms. There was no restriction placed based on study year, and all studies from January 1, 1993 through October 1, 2020 were included. Exclusionary criteria were as follows: use of (1) a cut-point derived by another study rather than applying novel thresholding methods; (2) continuous tau PET measures with no cut-points; (3) a threshold derived for the purposes of non-binary classification (e.g., AD subtypes); (4) any tau PET tracer other than Flortaucipir; (5) cerebrospinal fluid rather than PET tau biomarkers; (6) non-AD neurological conditions as a focus (e.g., Parkinson’s disease); (7) animal models. All resulting articles were screened by checking the abstract for exclusionary criteria. If no exclusionary criteria were met based on the abstract or if it was unclear, the full text was reviewed to determine eligibility. Finally, any articles that passed full-text review were included and verified independently. See Figure 1 for a flow diagram depicting this process. Quality ratings were conducted based on the Oxford Centre for Evidence-based Medicine (<https://www.cebm.net/>) by three independent raters with no discrepancies. Summary measures include descriptive statistics of tau PET cut-points based on different methods.

Empirical examination of the varying factors influencing cut-points.

In order to assess the reliability of different cut-point methods and their concurrent validity, we systematically varied the methodological approaches identified in this review within a sample of participants from the Alzheimer’s Disease Neuroimaging Initiative (ADNI). In this way, we controlled for the effects of sample size/characteristics and preprocessing methods that complicated comparisons across existing studies.

Within the ADNI sample of 785 older adults, resultant cut-points were examined in a 2x2x2 design that varied methodological decision points. The 8 possible cut-points were derived from different combinations of region (entorhinal cortex or temporal meta-ROI), analytic approach (receiver-operating characteristics (ROC) or 2 standard deviations above comparison group), and comparison group/criterion variable (A β positivity or cognitive diagnosis).

Full information on processing methods for ADNI A β PET (¹⁸F-AV-45, florbetapir) and tau PET (¹⁸F-AV-1451, flortaucipir) has been previously described elsewhere (Landau & Jagust, 2015, 2016). Standardized uptake values (SUVs) were intensity normalized using the whole cerebellum (A β PET; Landau et al., 2012; Landau & Jagust, 2015) or inferior cerebellar gray (tau PET; Landau & Jagust, 2016; Maass et al., 2017) to create SUV ratios (SUVRs). Tau PET data were partial volume corrected using the geometric transfer method (Baker et al., 2017).

Two regions of interest (ROIs) were examined for tau PET. A FreeSurfer-derived entorhinal cortex region (averaged across hemispheres) was used to approximate early Braak staging rather than an entorhinal cortex and hippocampus composite per extant recommendations given the high susceptibility of the hippocampus to partial volume effects (Landau & Jagust, 2016). A temporal meta-ROI was also examined using a composite of the following FreeSurfer-derived regions (averaged across hemispheres): amygdala, entorhinal cortex, fusiform gyrus, inferior temporal gyrus, and middle temporal gyrus. Outliers more than 3 standard deviations from the mean were removed from analyses (n = 16).

Two analytic methods were examined. ROC analysis assessed the performance of predictive models based on a binary criterion variable (Zou, O’Malley, & Mauri, 2007). In this case, tau PET SUVR (using either the temporal meta-ROI or entorhinal cortex) was used to predict classification as either A β positive/negative or cognitively unimpaired (CU)/impaired (CI). ROC

curves represent classification accuracy as sensitivity (y-axis) versus 1 – specificity (x-axis). Optimal cut-points were defined using the Youden index (sensitivity + specificity – 1). The second analytic method defined cut-points for the temporal meta-ROI and entorhinal cortex as a tau PET SUVR that was 2 standard deviations above a comparison group (either A β positive/negative or CU/CI).

Two separate criterion variables/comparison groups were used for each analytic approach. A β PET positivity, defined based on the existing cut-point of >1.11 (Joshi et al., 2012; Landau & Jagust, 2015), served as one criterion/comparison for defining tau PET cut-points. Cognitive diagnosis, either CU or CI, served as the other criterion/comparison. Diagnosis of CU or mild cognitive impairment (MCI) was determined using Jak/Bondi comprehensive neuropsychological criteria (Jak et al., 2009). Participants were diagnosed with MCI if they (1) had two impaired scores in one cognitive domain or (2) had one impaired score across all three cognitive domains. Six individual neuropsychological measures across domains of memory, language, and attention/executive function were included. Raw scores were then converted z-scores based on predicted values from regression equations adjusting for age, sex, and education derived within a robust CU group (i.e., CU throughout their duration in ADNI) based on the entire ADNI sample. Diagnosis of dementia was made based on the following criteria utilized by ADNI (<http://adni.loni.usc.edu/>): (1) subjective memory complaint reported by the participant, study partner or clinician; (2) objective memory impairment defined by a score below education-adjusted cut-offs on Logical Memory Delayed Recall, Story A of the Wechsler Memory Scale–Revised; (3) score between 20 and 26 on the Mini–Mental State Examination; (r) score 0.5 or 1.0 on the Clinical Dementia Rating Scale; and (v) met NINCDS/ADRDA criteria (McKhann et al., 1984) for probable Alzheimer’s disease. MCI and dementia groups were combined into one CI group.

For each of the derived 8 cut-points, tau positivity rates were determined. Concurrent validity of these 8 unique cut-points were assessed via three separate outcomes: (1) cerebrospinal fluid (CSF) phosphorylated tau (p-tau), (2) Mini-Mental State Examination (MMSE) total score, and (3) Logical Memory Story A Delayed Recall z-score (relative to robust CU group). Specifically, for each of the 8 cut-points, tau positive (T+) and tau negative (T-) participants were compared on these outcome variables, resulting in a Cohen’s *d* statistic that indicates the degree of discrepancy on these outcomes between T+ and T- participants for a given cut-point. Cohen’s *d* statistics were then used to qualitatively compare results across all 8 cut-points, with a larger Cohen’s *d* indicative of a higher concurrent validity for that cut-point.

Results

Review

Based on the above search methods, 540 articles were screened and 47 had the full-text reviewed to determine eligibility. The inclusion criteria were then met by 23 studies (see Table 1) with a total of 6,536 participants (note that this participant pool may not have been unique due to overlap in datasets utilized across the published studies). The quality of the evidence was equivalent across all studies included in this review with 23 cross-sectional studies each having a quality rating of 4 (on a scale of 1-5 with 1 being the highest quality) according to the Oxford Centre for Evidence-based Medicine (<https://www.cebm.net/>).

Table 1
Systematic review of 23 studies deriving tau PET thresholds.

Study	Sample	Reference Region	PVC	ROI(s)	Analytic Approach	Cut-point
[20] Botha et al. (2018), <i>Brain</i>	24 CI OAs from MCSA and ADRC	Cerebellar crus gray	Didn't say	Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG)	ROC maximizing sensitivity/specificity between YAs and A+ CI OAs	1.33
[21] Cho et al. (2018), <i>Neurobiol Aging</i>	220 OAs (all dx) from memory clinic	Cerebellar cortex	Yes using region-based voxelwise method (and no)	25 individual ROIs	Mean + 2.5 SDs above CU with minimal cortical Abeta (<1.4) and EC tau (<1.2)	None reported
[22] Dodich et al. (2020), <i>Eur J Nucl Med</i>	81 CI OAs from memory clinic	Cerebellar crus	Didn't say	Meta-ROI (medial temporal lobe, lateral occipital, ITG)	ROC between A-/N- vs everyone else	1.24
[23] Guo et al. (2020), <i>Alz Res Ther</i>	341 OAs (all dx) from ADNI	Inferior cerebellar cortex	Yes using GTM	1) Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG) 2) EC ROI	A) ROC with Youden index between A- CU and A+ C B) Mean +2 SDs above A- CU	1A) 1.25 2A) 1.21 1B) 1.34 2B) 1.31
[24] Jack et al. (2017), <i>Alz Dem</i>	49 YAs and 153 OAs (all dx) from MCSA and ADRC	Cerebellar crus gray median	"Most likely CSF" voxels were removed	Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG)	1) Maximize specificity (95%) based on YAs 2) Maximize sensitivity (10%) based on A+ CI OAs 3) ROC maximizing accuracy between YAs and A+ CI OAs 4) ROC maximizing accuracy between A-CU OAs and A+ CI OAs	1) 1.19 2) 1.21 3) 1.21 4) 1.32
[25] Jack et al. (2017), <i>Lancet Neuro</i>	435 OAs from MCSA (YAs not reported)	Cerebellar crus gray median	No	Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG)	ROC maximizing accuracy between YAs and A+ CI OAs	1.23
[26] Jack et al. (2019), <i>JAMA</i>	480 OAs without dementia from MCSA (YAs not reported)	Cerebellar crus gray	No (and yes using two compartment method)	Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG)	1) ROC maximizing accuracy between YAs and A+ CI OAs 2) ROC maximizing accuracy between A-CU and A+ CI OAs	1) 1.25 2) 1.33

*Additional cut-points are reported in the supplementary material of these studies.

Study	Sample	Reference Region	PVC	ROI(s)	Analytic Approach	Cut-point
[27] Jang et al. (2019), <i>Eur J Nucl Med imaging</i>	60 OAs with SVCI from medical center and 49 OAs (all dx) from hospital	Cerebellar gray	Yes	1) Braak V/VI 2) Braak III/IV 3) Braak I/II	CIDT (criterion variable unspecified); T+ considered III/IV or above	1) 1.58 2) 1.33 3) 1.29
[28] Lowe et al. (2018), <i>Alz Dem: DADM</i>	112 YAs from MCSA and 576 OAs from MCSA and ADRC	Cerebellar Crus	No	Temporal meta-ROI (EC, parahippocampal, hippocampus); 26 individual ROIs also assessed	95% above A- YAs	None reported
[29] Lowe et al. (2018), <i>Brain</i>	98 YAs, 601 A- CU OAs, 86 A+ CI OAs from MCSA and ADRC	Cerebellar crus gray	No (and yes using two compartment method)	47 individual ROIs	95% above A- YAs for each ROI	None reported
[30] Lowe et al. (2019), <i>Neurology</i>	117 YAs; 579 CU OAs from MCSA	Cerebellar crus	No (and yes using GTM)	43 individual ROIs	95% percent above A- YAs	1.17 in EC
[31] Lowe et al. (2020), <i>Alz Dem</i>	26 OAs from MCSA and ADRC	Cerebellar Crus	No (and yes using GTM)	1) Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG) 2) EC ROI	ROC with Youden index between AD and non-AD spectrum pathology	* 1) 1.29 2) 1.27
[32] Maass et al. (2017), <i>Neuroimage</i>	Sample 1: 12 YAs, 74 CU OAs from BACS, and 61 CI OAs from ADRC Sample 2: 42 CU and 28 CI OAs from ADNI	Inferior cerebellar gray	1) Yes using GTM 2) No	A) Temporal meta-ROI B) ITG (Also examined whole brain, AD-vulnerable mask, and Braak stage composite regions)	ROC with Youden index between A- CU OAs and A+ CI OAs) (Also used CIDT with age/diagnostic group as input variable for Braak composite staging)	1A) 1.47 (BACS/UCSF) and 1.34 (ADNI) 1B) 1.27 (BACS/UCSF) and 1.20 (ADNI) 2A) 1.46 (BACS/UCSF) and 1.40 (ADNI) 2B) 1.30 (BACS/UCSF) and 1.23 (ADNI) (other cut-points omitted from table but included in results section)

*Additional cut-points are reported in the supplementary material of these studies.

Study	Sample	Reference Region	PVC	ROI(s)	Analytic Approach	Cut-point
[33] Mattsson-Carligen et al. (2020), <i>Sci Adv</i>	131 OAs (all dx) from BioFINDER	Inferior cerebellar gray	No (and yes using GTM in EC)	1) EC 2) ITG 3) Braak V/VI	Mean +2 SDs above independent A- CU group	1) 1.39 2) 1.31 3) 1.28
[34] Meyer et al. (2020), <i>JAMA Neuro</i>	322 OAs (all dx) from ADNI	Inferior cerebellar gray	No (and yes using GTM)	Temporal meta-ROI (amygdala, EC, fusiform, parahippocampal, ITG, MTG)	ROC with maximum percentage correct classification between A- CU OAs and A+ CI OAs	* 1.37
[35] Mishra et al. (2017), <i>Neuroimage</i>	97 OAs (all dx) from ADRC	1) Whole cerebellum 2) Cerebellar cortex	Yes using RSF	Meta-ROI (EC, lateral occipital, ITG, amygdala)	SKM method to cluster into high and low tau groups; midpoint taken	1) 1.25 2) 1.22
[36] Ossenkoppele et al. (2018), <i>JAMA</i>	719 OAs (all dx) from ADRC, BioFINDER, and memory clinic	Inferior cerebellar gray	No (and yes using GTM)	5 ROIs examined: EC, ITG, temporal meta-ROI, temporoparietal cortex, Braak V/VI	1) Mean +2 SDs above CU 2) ROC Youden index between controls and AD	* For temporal meta-ROI: 1)1.34 2)1.27 (other cut-points omitted from table but included in results section)
[37] Rafii et al. (2017), <i>JAD</i>	9 adults with Down syndrome from DSBI	1) Cerebellar cortex (bottom slice removed and edges eroded) 2) subcortical WM	Yes using Muller-Gartner method	Braak I-VI average	Quantitative discrimination of A- vs A+	1) 1.2 2) 1.05
[38] Schöll et al. (2016), <i>Neuron</i>	5 YAs and 33 CU OAs from BACS; 15 CI OAs from ADRC	Cerebellar gray	Yes using GTM	1) Braak V/VI 2) Braak III/IV 3) Braak I/II	CIDT using age/diagnostic group as input variable	1) 2.79 2) 1.73 3) 1.40
[39] Schwarz et al. (2016), <i>Brain</i>	14 YAs and 173 OAs (all dx) from a clinical study	Cerebellar crus	Didn't say	7 ROIs corresponding to Braak histological stages	Mean +2.5 SDs above YAs	Ranged from 1.22 (STG) to 1.36 (fusiform)

*Additional cut-points are reported in the supplementary material of these studies.

Study	Sample	Reference Region	PVC	ROI(s)	Analytic Approach	Cut-point
[40] Schwarz et al. (2018), <i>Alz Dem</i>	14 YA, 21 OAs (all dx) for test-retest, and 98 OAs (all dx) from ADNI	Cerebellar gray	Didn't say	1) Meta-ROI (hippocampus, TEC, fusiform, MTG, STG, extrastriate, striate) 2) Simplified meta-ROI (medial temporal, lateral temporal, STG, primary visual cortex) 3) Lobar (average temporal, parietal, and frontal lobes) (ROIs gray matter masked)	1) Mean +2.5 SDs above T- YAs for each ROI within meta-ROI; T+ is anyone above threshold in hippocampus, TEC, fusiform, MTG, and extrastriate 2) Mean +3 SDs above T- YAs to obtain same average as (1); T+ is anyone above threshold in MTL and lateral temporal 3) Mean +3 SDs above T- YAs to obtain same average as (1); T+ is anyone above threshold in temporal lobe	* 1.28
[41] Wang et al. (2016), <i>JAMA Neuro</i>	59 OAs (all dx) from ADRC	Cerebellar cortex	No (and yes using linear regression)	1) Hippocampus 2) Meta-ROI (medial temporal, ITG, lateral temporal, inferior parietal, PCC, precuneus, SPL)	A) ROC with Youden index between A- CU and A+ AD B) ROC with Youden index between A+ CU and A+ AD	* 1A) 1.36 2A) 1.19 1B) 1.36 2B) 1.33
[42] Weigand et al. (2020), <i>Brain Comms</i>	523 OAs (all dx) from ADNI	Inferior cerebellar gray	Yes using GTM	1) Braak V/VI 2) Braak III/IV 3) Braak I/II	CIDT with MMSE as input variable	1) 1.96 2) 1.51 3) 1.18
*Additional cut-points are reported in the supplementary material of these studies.						

A = amyloid. ADNI = Alzheimer's Disease Neuroimaging Initiative. ADRC = Alzheimer's Disease Research Center. BACS = Berkeley Aging Cohort Study. CIDT = conditional inference decision tree. CU = cognitively unimpaired. CI = cognitively impaired. DSBI = Down Syndrome Biomarker Initiative. Dx = diagnoses. EC = entorhinal cortex. GTM = geometric transfer matrix. ITG = inferior temporal gyrus. MCSA = Mayo Clinic Study of Aging. MTG = middle temporal gyrus. N = neurodegeneration. OAs = older adults. PCC = posterior cingulate cortex. ROC = receiver operating characteristics. ROI = region of interest. SD = standard deviation. SKM = sparse k-means. SPL = superior parietal lobe. STG = superior temporal gyrus. SVCI = subcortical vascular cognitive impairment. T = tau. TEC = transentorhinal cortex. UCSF = University of California San Francisco. YAs = younger adults.

Cut-points. A total of 82 cut-points were reported in the main text across the 23 studies (note additional cut-points reported as supplementary material were excluded for the sake of parsimony, but such studies are marked in Table 1). The cut-points derived in the included studies ranged from 1.13 to 2.79 (the next highest value was considerably lower at 1.96). The mean value across all cut-points was 1.33 with a standard deviation (SD) of 0.21, and a median of 1.29. With the outlier value of 2.79 removed, the mean was 1.31 with a SD of 0.14, with a median of 1.28. Given that this variability may be due to several methodological differences, we explored this further below.

Sample Characteristics. Inclusion criteria yielded a sample of 6,115 cognitively unimpaired and cognitively impaired older adults, as well as 421 younger adults who served as reference groups. In addition to older adults who were cognitively unimpaired or diagnosed with mild cognitive impairment, many studies included individuals diagnosed with dementia. While the majority of studies recruited participants with probable AD, other dementia syndromes included AD-variants (i.e., dysexecutive AD, posterior cortical atrophy; Maass et al., 2017; Scholl et al., 2016), hippocampal sclerosis (Botha et al., 2018), subcortical vascular cognitive impairment (Jang et al., 2019), and non-AD neurodegenerative disorders (e.g., Lewy body dementia, primary progressive aphasia; Ossenkoppele et al., 2018). The current review also included one study investigating AD in individuals with Down syndrome (Rafii et al., 2017).

Several included studies recruited participants from research initiatives such as the Mayo Clinic Study of Aging (MCSA; $n = 8$), Alzheimer's Disease Neuroimaging Initiative (ADNI; $n = 5$), Berkeley Aging Cohort Study (BACS; $n = 2$), BioFINDER ($n = 2$), and Down Syndrome Biomarker Initiative (DBSI; $n = 1$). Other sources included local Alzheimer's disease research centers ($n = 10$), memory/medical clinics ($n = 4$), or a clinical study ($n = 1$). Note that several studies recruited from multiple sources to increase the generalizability of the findings, and therefore the above total (33) exceeds the number of included studies (23). See Table 1 for further details.

Preprocessing Methods. Differences were observed in preprocessing methods including reference region and use of partial volume correction (PVC), which may contribute to inter-study variability in cut-points. Twenty-one studies used a cerebellar region, with the cerebellar gray ($n = 6$) and inferior cerebellar gray ($n = 6$) as the most commonly used reference regions to calculate SUVRs. Other regions included the cerebellar crus ($n = 5$), cerebellar crus gray ($n = 3$), and cerebellar crus gray median ($n = 2$; note that all other reference regions were calculated using mean as the measure of central tendency). The whole cerebellum and subcortical white matter were also used.

Use of PVC was indicated by 19/23 studies, with 6 only using PVC, 3 not using any PVC (note that one of these studies did remove "most likely CSF" voxels), 4 not reporting either way, and 10 directly comparing PVC and non-PVC approaches. On average, use of PVC resulted in a higher cut-point (PVC mean[SD] = 1.41[.32], or 1.37[.20] with outlier of 2.79 removed; non-PVC = 1.29[.07]). See Figure 1 for a distribution of cut-points based on use of PVC. See Table 1 for further details.

Regions of Interest. Several regions of interest (ROIs) were used to derive cut-points in the included studies, including individual and composite ROIs. A temporal meta-ROI that was first introduced by Jack et al. (2017) consisting of various medial and lateral temporal regions was used by the largest proportion of studies (10/25; cut-point mean[sd] = 1.31[.08]). Similarly, 4 studies thresholded on a larger meta-ROI that extended to occipital or parietal regions, with one of these studies additionally assessing lobar ROIs. Braak stage composites were used by 8 studies, typically including stages I/II (cut-point mean[SD] = 1.29[.09], III/IV (1.38[.16]), and V/VI (1.52[.48], or 1.41[.28] with outlier removed). Additionally, 5 studies selected individual ROIs such as the entorhinal or inferior temporal cortex, and 4 other studies assessed a large number of ROIs throughout the brain. See Table 1 for further details and Figure 2 for distributions of cut-points using a temporal meta-ROI or Braak stages composites. Notably, these statistics are averaged across studies using or not using PVC, which will also contribute to variability.

Thresholding Analytic Procedures. Perhaps the most heterogeneity in tau PET thresholding methods was observed in the type of analysis used, with only a few of the 23 studies employing the same particular analytic approach. There were, however, clusters of similar statistical methods employed across studies that varied in aspects such as criterion variable or groups assessed. The most frequently observed of these analytic "clusters" was use of receiver operating characteristics (ROC) curves discriminating between two groups to identify the optimal cut-point for differentiating tau positive and negative individuals, which was used in 11 studies. This method varied, however, in the criterion variable (i.e., groups) predicted by tau PET SUVRs, with various combinations of younger adults, A β negative (A-) or A β positive (A+) older adults, and cognitively unimpaired (CU) or cognitively impaired (CI) older adults. The second most frequently observed analytic "cluster" was identifying a threshold (e.g., 2 standard deviations) above the average tau PET SUVR from a reference group at which the cut-point would be defined ($n = 9$). With this method, there was substantial variability across studies in the reference group used, including younger adults, A-CU older adults, and all CU older adults. See Table 1 for further details.

Empirical examination of the varying factors influencing cut-points.

The cut-point values and resultant tau positivity rates for each of the 8 cut-points are reported in Table 2. The average cut-point when using the “early” (i.e., entorhinal) ROI was 2.06, whereas the average cut-point when using the “late” (i.e., temporal meta) ROI was 1.73. This discrepancy is unsurprising given that, within the same sample, we would expect higher tau deposition within the “early” ROI relative to the “late” ROI as delineated by the spatiotemporal progression of tau based on Braak staging, necessitating a higher threshold for the “early” ROI. This cut-point adjustment for each ROI based on level of tau resulted in nearly equivalent tau positivity rates, on average, with 31.7% tau positivity for the “early” ROI and 29.8% tau positivity for the “late” ROI, indicating no appreciable effect of ROI independent of analytic method and comparison group/criterion variable.

Table 2

Derivation of tau PET cut-points, resultant tau positivity rates, and Cohen’s *d* statistics discriminating tau positivity/negativity for 8 unique thresholding methods.

Method	Cut-point	T+ rate	CSF p-tau Cohen’s <i>d</i>	MMSE Cohen’s <i>d</i>	Logical Memory Cohen’s <i>d</i>
Entorhinal/ROC/A+	1.88	36.5%	-1.06 (95% CI [-1.27, -.85])	.91 (95% CI [.75, 1.06])	.88 (95% CI [.72, 1.03])
Entorhinal/ROC/dx	2.14	24.0%	-.99 (95% CI [-1.22, -.75])	1.29 (95% CI [1.11, 1.47])	1.22 (95% CI [1.05, 1.40])
Entorhinal/2SD/A+	1.71	52.8%	-.76 (95% CI [-.96, -.56])	.65 (95% CI [.50, .79])	.54 (95% CI [.39, .68])
Entorhinal/2SD/dx	2.52	13.3%	-1.05 (95% CI [-1.35, -.75])	1.53 (95% CI [1.31, 1.75])	1.22 (95% CI [1.01, 1.44])
Meta-ROI/ROC/A+	1.65	31.5%	-1.12 (95% CI [-1.34, -.90])	1.03 (95% CI [.87, 1.19])	.86 (95% CI [.70, 1.02])
Meta-ROI/ROC/dx	1.72	23.7%	-1.23 (95% CI [-1.47, -.99])	1.26 (95% CI [1.08, 1.44])	1.05 (95% CI [.87, 1.22])
Meta-ROI/2SD/A+	1.52	52.7%	-.70 (95% CI [-.89, -.50])	.59 (95% CI [.44, .74])	.39 (95% CI [.25, .54])
Meta-ROI/2SD/dx	2.04	11.1%	-1.23 (95% CI [-1.54, -.91])	1.60 (95% CI [1.36, 1.84])	1.25 (95% CI [1.02, 1.49])

CSF = cerebrospinal fluid. Dx = diagnosis. MMSE = Mini-Mental State Examination. ROC = receiver-operating characteristics. SD = standard deviation. T = tau. The largest effect sizes for each outcome variable are presented in bold.

The average cut-point when using ROC analysis was 1.85, whereas the average cut-point when positivity is defined as 2 standard deviations above a comparison group was 1.94. On average, there was a 28.9% tau positivity rate when using ROC analysis and a 32.5% tau positivity rate with an analytic method of 2 standard deviations above a comparison group, indicating no appreciable effect of analytic method independent of ROI and comparison group/criterion variable. ROC figures with area under the curve (AUC) values for each ROI and criterion variable are presented in Figure 3.

The average cut-point when using an Aβ positive comparison group (or Aβ positivity as the criterion variable) was 1.69, whereas the average cut-point when using a CU comparison group (or diagnosis as the criterion variable) was 2.10. On average, there was a 52.6% tau positivity rate when defining groups based on Aβ positivity and a 18.0% tau positivity rate when defining groups based on cognitive diagnosis. Thus, comparison group/criterion variable does confer a notable effect on cut-point and resultant positivity rates independent of ROI and analytic methods.

Each of the 8 unique methods for deriving cut-points were then assessed for concurrent validity by evaluating differences between tau positive and tau negative participants for each cut-point on CSF p-tau, MMSE score, and logical memory delayed recall (see Table 2). When considering differences in CSF p-tau levels, the largest discrepancy between tau positive and tau negative groups, as measured by Cohen’s *d*, was observed both for the cut-point that was derived using ROC analysis with diagnosis as the criterion variable in the meta-temporal ROI (Cohen’s *d* = -1.23), and for the cut-point that was derived based on

2 standard deviations above the cognitively normal group in the meta-temporal ROI (Cohen's $d = -1.23$). When considering differences in MMSE score and Logical Memory delayed recall, the largest discrepancy between tau positive and tau negative groups, as measured by Cohen's d , was observed for the cut-point that was derived using the cut-point that was derived based on 2 standard deviations above the cognitively normal group in the meta-temporal ROI (MMSE Cohen's $d = 1.60$; Logical Memory Cohen's $d = 1.25$). See Figure 4 for a depiction of tau positivity group differences on CSF p-tau, MMSE score, and logical memory delayed recall when using the cut-point that was derived based on 2 standard deviations above the cognitively normal group in the meta-temporal ROI.

Discussion

This review and empirical examination of cut-points based on varying threshold derivation methods and analyses revealed notable variability in tau PET SUVR cut-points across published studies to date, ranging from 1.13 to 2.79, and highlighted some of the possible sources and differences in methods to which this variability may be attributed. A discussion of the various decision points in thresholding methods is detailed below, including sample composition, preprocessing steps (i.e., reference region, PVC), selection of ROIs, and statistical approaches.

Firstly, samples differed in demographic and clinical characteristics. The choice of the reference group was thought to affect cut-points since entorhinal tau is common in cognitively normal older adults, which results in higher cut-points for abnormal tau relative to use of a younger control group. Similarly, age and clinical syndrome of the AD patients have a considerable effect on the cut-points. While higher tau tracer uptake in neocortical regions and thus higher cut-points are associated with younger age-of-onset and non-amnesic clinical presentations of AD patients, greater medial temporal lobe vulnerability is seen in older patients with amnesic symptoms such as in the ADNI sample (e.g., Maass et al, 2017).

As shown in the results, differences in preprocessing methods may also contribute to cut-point variability across studies. Interestingly, two studies directly compared results using different reference regions. Mishra et al. (2017) compared both the whole cerebellum and the cerebellar gray as reference regions, with a marginally higher cut-point of 1.25 derived from the use of the whole cerebellum reference region relative to the cerebellar gray (SUVR = 1.22). Additionally, Rafii et al. (2017) assessed multiple reference regions that resulted in a cut-point of 1.20 using a cerebellar gray reference (with the bottom slice removed and edges eroded) and a notably lower cut-point of 1.05 when using a subcortical white matter reference. This was the only study in this corpus of studies for review that included a non-cerebellar reference region, so further comparisons of cerebellar vs. white matter and their influences on cut-point values are not available at this time. More research is needed directly comparing cut-points derived from SUVRs based on differing reference regions to determine the magnitude of this effect on the resultant cut-points. Based on the data available from this single study, it appears that using subcortical white matter resulted in a substantially lower cut-point relative to the commonly used cerebellar gray reference region, possibly due to differences in off-target binding in these regions with the Flortaucipir tracer.

Use of PVC also varied considerably across studies, and many studies directly compared cut-points based on data that had or had not undergone PVC. For example, one study (Meyer et al., 2020) found a much higher cut-point when using PVC (SUVR = 1.79) relative to non-PVC (SUVR = 1.37). This discrepancy is likely explained by the higher atrophy that is associated with higher tau pathology, which leads to partial voluming and thus underestimation of tau PET signal in subjects with advanced tau pathology. The indecision across (and within) studies on whether to use PVC indicates the importance of determining a standardized preprocessing approach to tau PET thresholding in order to reduce cut-point variability and facilitate interpretation of positivity rates.

In addition to the above differences in methodology, what varied most considerably across studies was (1) the region(s) on which cut-points were based and (2) the analytic choices used to derive cut-points. Thresholding procedures for tau PET necessitate the selection of an ROI that is susceptible to tau pathology. A single brain region may be used for a sensitive and localized analysis, or a composite of several regions may be used to cover multiple key regions, likely with increased reliability. As noted in Table 1, the majority of studies used a composite of some type, whether combining AD vulnerable regions or recapitulating Braak stages (Braak & Braak, 1991). A smaller number of studies utilized single ROIs in the medial temporal lobe

implicated early in the AD pathologic process; although this approach offers increased sensitivity and may be appropriate in very early stages in which tau is confined to the entorhinal cortex (Braak & Braak, 1991), it may not accurately reflect the more widespread distribution of tau in later disease stages. Further, given that medial temporal structures are often subject to partial volume effects due to proximity to the choroid plexus (Lowe *et al.*, 2016), use of these regions as isolated ROIs may not be as advisable. Alternatively, use of a temporal meta-ROI may provide increased reliability of the estimate with only a marginal decrease in the sensitivity needed to detect early-stage tau pathology.

The type of statistical analysis used also varied significantly, although most approaches were based on discrimination against a comparison group (e.g., ROC with A- and A+ older adults, 2.5 SDs above mean value for young adults). In examination of the analytic procedures employed across studies, a theme emerged: studies derived a tau threshold that was either contingent on A β in some manner or was independent of A β . Those with A β -contingent methods were used in more than half of the included studies and are based on the assumption that A- controls are not on the AD pathway (Jack *et al.*, 2018). Alternatively, statistical approaches not contingent on A β instead used a comparison group of younger adults or older adults irrespective of A β status, or used cognitive status as a criterion variable. Notably, the contingency on A β used in some studies, based on the assumption that A β negative controls are not on the AD pathway, may influence resultant tau PET cut-points and was examined in our empirical follow-up.

Variability in tau PET cut-points inevitably leads to variability in tau positivity rates, which impacts subsequent staging efforts based on biomarker positivities and possible inclusion in clinical treatment trials, as well as increasing estimation uncertainty which hinders reproducibility in AD biomarker research. Using the different methodological approaches presented above yielded remarkably different resultant cut-points, making it difficult to evaluate the variable utility of any single method. This served as the basis for our empirical investigation of cut-point methods (i.e., ROI, analytic method, and comparison group) to examine how different methodological decision-points in a sample standardized on size, characteristics, and preprocessing techniques would influence cut-point values, resultant positivity rates, and concurrent validity with other biomarker and cognitive outcomes.

There was no appreciable difference in the tau PET cut-point values or positivity rates, on average, based on ROI or analytic approaches. However, the comparison group/criterion variable used did yield a notable difference in cut-points and positivity rates such that use of cognitive diagnosis yielded, on average, higher cut-point values and lower tau positivity rates. This pattern of results suggests that thresholding methods using cognitive diagnosis rather than A β positivity are generally more conservative in defining tau positivity.

Interestingly, use of cognitive diagnosis as the comparison group/criterion variable predominantly resulted in higher Cohen's *d* effect size values for prediction of CSF p-tau, MMSE score, and memory recall, indicating higher concurrent validity of this method for discriminating tau positive and tau negative groups. Specifically, when considering unique combinations of these methodological factors, determination of a cut-point based on a value 2 standard deviations above the CU group in the temporal meta-ROI appeared to be the most valid approach, with highest effect sizes across all outcome variables.

These results are critical to improve our understanding how these various methodological decisions and different choices influence the derivation of tau PET cut-points and resultant positivity rates, and which approaches may be most appropriate to include in a standardized approach to tau PET biomarker thresholding based on their concurrent validity with CSF p-tau levels, MMSE score, and memory recall. Based on concurrent validity across a range of biomarker and cognitive outcomes, it is recommended that cognitive diagnosis, rather than A β positivity, be used to derive tau PET cut-points. However, the specific research question under investigation ideally will determine the specific methodological approaches used, which may necessitate use of certain ROIs, analyses, and/or comparison variables. Notably, the thresholding methods based on cognitive diagnosis yielded the lowest positivity rate, which suggests that it may have high specificity but low sensitivity, albeit this determination was not possible in this study without having a neuropathologic validation of tau PET positivity rates. Thus, the context of the use of various thresholding methods in terms of balancing sensitivity with specificity should be taken into account.

Our findings have implications for interpretation of existing studies using tau PET positivity categories and highlight the difficulty of comparing these studies without standardization of thresholding methods. In particular, comparison of methods using a two standard deviation above the mean approach with either A β positivity or cognitive diagnosis as the comparison group yielded very different concurrent validity across outcome measures, indicating difficulty in comparing positivity rates and outcomes across studies that differ in this regard. Taking steps toward the standardization of these thresholding procedures has important implications for increasing reliability and reproducibility of our research and advancing our understanding of biomarker dynamics across the AD continuum.

Limitations

The primary limitation of this review is the difficulty in comparing how a specific method influences cut-point values due to the highly discrepant procedures and its various combinations across studies. For example, comparing analytic procedures either contingent or not contingent on A β status proved difficult since any given study with these groups may have used different reference regions, PVC applications, regions of interest, or even the method by which the determination of A β positivity was achieved. Such a comparison is important given the notion that tau may operate earlier than and independent of A β (Arnsten *et al.*, 2020; Groot *et al.*, 2021). However, we sought to address this in our empirical follow-up in which a consistent sample with the same preprocessing methods was used to compare ROIs, analytic methods, and comparison group/criterion variable. Another limitation of the review is indicated by the quality of evidence, which indicated the need for more rigor in standardizing methods and analytic decisions across studies. All studies included in this review (as well as the parent study [ADNI] used for the empirical follow-up) were cross-sectional at the point of cut-off determination, and longitudinal within-subject change will be an important direction for future studies to further improve rigor in the derivation of cut-points. Furthermore, several studies used overlapping samples which can lead to bias in our results. Finally, the use of MMSE and memory recall as outcome variables may have resulted in circularity with cut-points that were derived based on cognitive diagnosis. However, the specific neuropsychological measures used to determine diagnosis were independent from the cognitive outcomes used, and it is noteworthy that effect sizes for cognitive outcomes were commensurate with CSF outcomes. Additionally, the inclusion of cognitive outcomes to assess concurrent validity is critical given this is ultimately what we want to use biomarkers to predict and treat.

Importantly, the studies reviewed are further limited by the significant lack of diversity in race/ethnicity in tau PET research specifically and AD research in general. This is a significant limitation of our empirical follow-up as well, which was conducted in a predominately White, highly educated ADNI sample. The presence and degree of biomarkers such as tau may vary across racial/ethnic identities (Morris *et al.*, 2019) as a result of numerous sociocultural factors including experience of racism and access to resources (Barnes, 2019). Exploring these and other moderating variables is essential to gain a holistic understanding of tau PET and the derivation of cut-points.

Conclusions

This review and empirical examination of tau PET cut-point methodologies demonstrates the significant heterogeneity in methods used to derive tau PET cut-points with its most widely studied tracer to date, Flortaucipir, resulting in a large range of suggested cut-points across studies. Factors influencing this variability may include differences in sample characteristics, reference region, use of PVC, region(s) of interest, or analytic procedures. This variability in tau PET cut-points has significant implications for increasingly used biomarker classifications that rely on these cut-points for determination of biomarker positivity and for potential selection for clinical treatment trials, as well as increasing estimation uncertainty which hinders reproducibility in AD biomarker research. Our empirical follow-up systematically addressed how these decisional and analytic choices in the methods used influenced cut-points and resultant positivity rates in a single sample of participants. Taking this approach, we found that a cut-point derived using an analytic approach of 2 standard deviations above a cognitively normal control group within a meta-temporal ROI was the most predictive of concurrent CSF p-tau, MMSE score, and memory performance. Taken together, this paper highlights the importance of standardization in thresholding methods to create and apply reliable and optimal cut-points that improve our characterization of AD biomarker risk.

Declarations

Ethics Approval and Consent to Participate

This study was approved locally by the University of California San Diego Human Research Protections Program (protocol #190618), which can be reached at 858-246-4777. Written informed consent was waived for this retrospective data analysis.

Consent for Publication

Not applicable

Availability of Data and Materials

Data used in this study are available through the Alzheimer's Disease Neuroimaging Initiative and can be found at <http://adni.loni.usc.edu/>.

Competing Interests

M.W.B. receives royalties from Oxford University Press. All other authors report no disclosures relevant to the manuscript.

Funding

Funding for this study was provided by grants from the National Institute on Aging (R01 AG049810 and R01 AG054049 to M.W.B.) and the National Science Foundation (DGE-1650012 to A.J.W.)

Author's Contributions

A.J.W. and M.W.B. conceptualized the study design. A.J.W. performed the systematic literature review. G.M.L.E. and A.M. validated the content of the literature review. A.J.W. performed all statistical analyses. A.J.W. drafted the manuscript. All authors reviewed the manuscript for intellectual content.

Acknowledgements

We thank the investigators of the Alzheimer's Disease Neuroimaging Initiative, as well as all of the study participants, for use of these data to advance our understanding of Alzheimer's disease.

References

1. Alzheimer A. Über eine eigenartige Erkrankung der Hirnrinde. *Allg Zeitschr Psychiatr Psychiatr-Gerichtl Med.* 1907;;146–148.
2. Hardy J. The discovery of Alzheimer-causing mutations in the APP gene and the formulation of the “amyloid cascade hypothesis. *The FEBS Journal.* 2017;284(7):1040–4.
3. Jagust W. Imaging the evolution and pathophysiology of Alzheimer's disease. *Nat Rev Neurosci.* 2018;19(11):687–700.
4. Jack CR Jr, Bennett DA, Blennow K, et al. A/T/N: An unbiased descriptive classification scheme for Alzheimer disease biomarkers. *Neurology.* 2016;87(5):539–47.
5. Jack CR Jr, Bennett DA, Blennow K, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimer's Dementia.* 2018a;14(4):535–62.
6. Breault C, Piper J, Joshi AD, et al. Correlation between two methods of florbetapir PET quantitative analysis. *Am J Nucl Med Mol Imaging.* 2017;7(3):84–91.
7. Joshi AD, Pontecorvo MJ, Clark CM, et al. Performance characteristics of amyloid PET with florbetapir F 18 in patients with Alzheimer's disease and cognitively normal subjects. *The Journal of Nuclear Medicine.* 2012;53(3):378–84.

8. Klunk WE, Koeppe RA, Price JC, et al. The Centiloid Project: standardizing quantitative amyloid plaque estimation by PET. *Alzheimers Dement*. 2015;11(1):1–15.e154.
9. Devous MD, Sr, Joshi AD, Navitsky M, et al. Test-Retest Reproducibility for the Tau PET Imaging Agent Flortaucipir F 18. *J Nucl Med*. 2018;59(6):937–43. doi:10.2967/jnumed.117.200691.
10. Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol*. 1991;82(4):239–59.
11. Landau S, Jagust W. Flortaucipir processing methods. <http://adni.loni.usc.edu/>. Published December 3, 2015.
12. Landau S, Jagust W. Flortaucipir (AV-1451) processing methods. <http://adni.loni.usc.edu/>. Published March 2, 2016.
13. Landau SM, Mintun MA, Joshi AD, et al. Amyloid deposition, hypometabolism, and longitudinal cognitive decline. *Ann Neurol*. 2012;72(4):578–86.
14. Maass A, Landau S, Baker SL, et al. Comparison of multiple tau-PET measures as biomarkers in aging and Alzheimer's disease. *Neuroimage*. 2017;157:448–63.
15. Baker SL, Maass A, Jagust WJ. Considerations and code for partial volume correcting [18F]-AV-1451 tau PET data. *Data Brief*. 2017;15:648–57.
16. Zou KH, O'Malley AJ, Mauri L. Receiver-operating characteristic analysis for evaluating diagnostic tests and predictive models. *Circulation*. 2007;115(5):654–7.
17. Joshi AD, Pontecorvo MJ, Clark CM, et al. Performance characteristics of amyloid PET with flortaucipir F 18 in patients with Alzheimer's disease and cognitively normal subjects. *J Nucl Med*. 2012;53(3):378–84.
18. Jak AJ, Bondi MW, Delano-Wood L, et al. Quantification of five neuropsychological approaches to defining mild cognitive impairment. *Am J Geriatr Psychiatry*.
19. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984;34(7):939–44.
20. Botha H, Mantyh WG, Murray ME, et al. FDG-PET in tau-negative amnesic dementia resembles that of autopsy-proven hippocampal sclerosis. *Brain*. 2018;141(4):1201–17.
21. Cho H, Lee HS, Choi JY, et al. Predicted sequence of cortical tau and amyloid- β deposition in Alzheimer disease spectrum. *Neurobiol Aging*. 2018;68:76–84.
22. Dodich A, Mendes A, Assal F, et al. The A/T/N model applied through imaging biomarkers in a memory clinic. *Eur J Nucl Med Mol Imaging*. 2020;47(2):247–55.
23. Guo T, Korman D, La Joie R, et al. Normalization of CSF pTau measurement by A β ₄₀ improves its performance as a biomarker of Alzheimer's disease. *Alzheimers Res Ther*. 2020;12(1):97.
24. Jack CR Jr, Wiste HJ, Weigand SD, et al. Defining imaging biomarker cut points for brain aging and Alzheimer's disease. *Alzheimers Dement*. 2017;13(3):205–16.
25. Jack CR Jr, Wiste HJ, Weigand SD, et al. Age-specific and sex-specific prevalence of cerebral β -amyloidosis, tauopathy, and neurodegeneration in cognitively unimpaired individuals aged 50-95 years: a cross-sectional study. *Lancet Neurol*. 2017;16(6):435–44.
26. Jack CR Jr, Wiste HJ, Therneau TM, et al. Associations of Amyloid, Tau, and Neurodegeneration Biomarker Profiles With Rates of Memory Decline Among Individuals Without Dementia. *JAMA*. 2019;321(23):2316–25.
27. Jang H, Kim HJ, Park S, et al. Application of an amyloid and tau classification system in subcortical vascular cognitive impairment patients. *Eur J Nucl Med Mol Imaging*. 2020;47(2):292–303.
28. Lowe VJ, Bruinsma TJ, Min HK, et al. Elevated medial temporal lobe and pervasive brain tau-PET signal in normal participants. *Alzheimers Dement (Amst)*. 2018;10:210–6.
29. Lowe VJ, Wiste HJ, Senjem ML, et al. Widespread brain tau and its association with ageing, Braak stage and Alzheimer's dementia. *Brain*. 2018;141(1):271–87.
30. Lowe VJ, Bruinsma TJ, Wiste HJ, et al. Cross-sectional associations of tau-PET signal with cognition in cognitively unimpaired adults. *Neurology*. 2019;93(1):e29–39.

31. Lowe VJ, Lundt ES, Albertson SM, et al. Tau-positron emission tomography correlates with neuropathology findings. *Alzheimers Dement.* 2020;16(3):561–71.
32. Maass A, Landau S, Baker SL, et al. Comparison of multiple tau-PET measures as biomarkers in aging and Alzheimer's disease. *Neuroimage.* 2017;157:448–63.
33. Mattsson-Carlgren N, Andersson E, Janelidze S, et al. A β deposition is associated with increases in soluble and phosphorylated tau that precede a positive Tau PET in Alzheimer's disease. *Sci Adv.* 2020;6(16):eaaz2387.
34. Meyer PF, Pichet Binette A, Gonneaud J, Breitner JCS, Villeneuve S. Characterization of Alzheimer Disease Biomarker Discrepancies Using Cerebrospinal Fluid Phosphorylated Tau and AV1451 Positron Emission Tomography [published correction appears in *JAMA Neurol.* 2020 Feb 17;]. *JAMA Neurol.* 2020;77(4):508–16.
35. Mishra S, Gordon BA, Su Y, et al. AV-1451 PET imaging of tau pathology in preclinical Alzheimer disease: Defining a summary measure. *Neuroimage.* 2017;161:171–8.
36. Ossenkoppele R, Rabinovici GD, Smith R, et al. Discriminative Accuracy of [18F]flortaucipir Positron Emission Tomography for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA.* 2018;320(11):1151–62.
37. Rafii MS, Lukic AS, Andrews RD, et al. PET Imaging of Tau Pathology and Relationship to Amyloid, Longitudinal MRI, and Cognitive Change in Down Syndrome: Results from the Down Syndrome Biomarker Initiative (DSBI). *J Alzheimers Dis.* 2017;60(2):439–50.
38. Schöll M, Lockhart SN, Schonhaut DR, et al. PET Imaging of Tau Deposition in the Aging Human Brain. *Neuron.* 2016;89(5):971–82.
39. Schwarz AJ, Yu P, Miller BB, et al. Regional profiles of the candidate tau PET ligand 18F-AV-1451 recapitulate key features of Braak histopathological stages. *Brain.* 2016;139(Pt 5):1539–50.
40. Schwarz AJ, Shcherbinin S, Sliker LJ, et al. Topographic staging of tau positron emission tomography images. *Alzheimers Dement (Amst).* 2018;10:221–31.
41. Wang L, Benzinger TL, Su Y, et al. Evaluation of Tau Imaging in Staging Alzheimer Disease and Revealing Interactions Between β -Amyloid and Tauopathy. *JAMA Neurol.* 2016;73(9):1070–7.
42. Weigand AJ, Bangen KJ, Thomas KR, et al. Is tau in the absence of amyloid on the Alzheimer's continuum?: A study of discordant PET positivity. *Brain Commun.* 2020;2(1):fcz046.
43. Rousset OG, Ma Y, Evans AC. Correction for partial volume effects in PET: principle and validation. *J Nucl Med.* 1998;39(5):904–11.
44. Meltzer CC, Kinahan PE, Greer PJ, et al. Comparative evaluation of MR-based partial-volume correction schemes for PET. *J Nucl Med.* 1999;40(12):2053–65.
45. Rousset OG, Ma Y, Evans AC. Correction for partial volume effects in PET: principle and validation. *J Nucl Med.* 1998;39(5):904–11.
46. Müller-Gärtner HW, Links JM, Prince JL, Bryan RN, McVeigh E, Leal JP, Davatzikos C, Frost JJ. Measurement of radiotracer concentration in brain gray matter using positron emission tomography: MRI-based correction for partial volume effects. *J Cereb Blood Flow Metab.* 1992 Jul;12(4):571–83.
47. Thomas BA, Erlandsson K, Modat M, et al. The importance of appropriate partial volume correction for PET quantification in Alzheimer's disease. *Eur J Nucl Med Mol Imaging.* 2011;38(6):1104–19.
48. Lowe VJ, Curran G, Fang P, et al. An autoradiographic evaluation of AV-1451 Tau PET in dementia. *Acta Neuropathol Commun.* 2016;4(1):58.
49. Groot C, Doré V, Robertson J, et al. Mesial temporal tau is related to worse cognitive performance and greater neocortical tau load in amyloid- β -negative cognitively normal individuals. *Neurobiol Aging.* 2021;97:41–8.
50. Arnsten AFT, Datta D, Tredici KD, Braak H. Hypothesis: tau pathology is an initiating factor in sporadic Alzheimer's disease. *Alzheimer's Dement.* 2020.
51. Morris JC, Schindler SE, McCue LM, et al. Assessment of Racial Disparities in Biomarkers for Alzheimer Disease. *JAMA Neurol.* 2019;76(3):264–73.

Supplementary Material

Supplementary Material is not available with this version.

Figures




Figure 1

Distribution of tau PET SUVR cut-points in methods with (A) or without (B) partial volume correction (PVC). PET = positron emission tomography. SUVR = standardized uptake value ratio.




Figure 2

Distribution of tau PET SUVR cut-points in methods using a temporal meta-ROI (A) or Braak stage ROIs (B). PET = positron emission tomography. SUVR = standardized uptake value ratio.




Figure 3

Receiver-operating characteristics (ROC) curves used to derive cut-points for (A) entorhinal cortex with A β positivity as the criterion, (B) temporal meta-ROI with A β positivity as the criterion, (C) entorhinal cortex with cognitive diagnosis as the criterion, and (D) temporal meta-ROI with cognitive diagnosis as the criterion. AUC = area under the curve. A β = amyloid-beta.




Figure 4

Beeswarm plots depicting differences between tau negative and tau positive groups on (A) cerebrospinal fluid (CSF) p-tau level, (B) Mini-Mental State Exam (MMSE) score, and (C) Logical Memory delayed recall, when tau positivity is determined using a cutpoint derived as 2 standard deviations above a cognitively intact comparison group in a temporal meta-region of interest.