

PGC-1 α Regulates Airway Epithelial Barrier Dysfunction Induced by House Dust Mite.

Tsutomu Saito

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Tomohiro Ichikawa (✉ ichikawa@rm.med.tohoku.ac.jp)

Tohoku University Graduate School of Medicine <https://orcid.org/0000-0003-0093-6362>

Tadahisa Numakura

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Mitsuhiro Yamada

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Akira Koarai

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Naoya Fujino

Tohoku University School of Medicine, Department of Respiratory Medicine

Koji Murakami

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Shun Yamanaka

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Yusaku Sasaki

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Yorihiko Kyogoku

Sendai City Hospital, Department of Respiratory Medicine

Koji Itakura

Osaki Citizen Hospital, Department of Respiratory Medicine

Hirohito Sano

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Katsuya Takita

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Rie Tanaka

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Tsutomu Tamada

Tohoku University Graduate School of Medicine, Department of Respiratory Medicine

Masakazu Ichinose

Osaki Citizen Hospital, Department of Respiratory Medicine

Hisatoshi Sugiura

Research

Keywords: asthma, airway barrier dysfunction, PGC-1 α , house dust mite

Posted Date: October 27th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-96584/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 19th, 2021. See the published version at <https://doi.org/10.1186/s12931-021-01663-6>.

Abstract

Background: The airway epithelial barrier function is disrupted in the airways of asthmatic patients. Abnormal mitochondrial biogenesis is reportedly involved in the pathogenesis of asthma. However, the role of mitochondrial biogenesis in the airway barrier dysfunction has not been elucidated yet. This study aimed to clarify whether the peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-1 α), a central regulator of mitochondrial biogenesis, is involved in the disruption of the airway barrier function induced by aeroallergens.

Methods: BEAS-2B cells were exposed to house dust mite (HDM) and the expressions of PGC-1 α and E-cadherin, a junctional protein, were examined by immunoblotting. The effect of SRT1720, a PGC-1 α activator, was investigated by immunoblotting, immunocytochemistry, and measuring the transepithelial electrical resistance (TEER) on the HDM-induced reduction in mitochondrial biogenesis markers and junctional proteins in airway bronchial epithelial cells. Furthermore, the effects of protease activated receptor 2 (PAR2) inhibitor, GB83, Toll-like receptor 4 (TLR4) inhibitor, lipopolysaccharide from *Rhodobacter sphaeroides* (LPS-RS), protease inhibitors including E64 and 4-(2-Aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF) on the HDM-induced barrier dysfunction were investigated.

Results: The amounts of PGC-1 α and E-cadherin in the HDM-treated cells were significantly decreased compared to the vehicle-treated cells. SRT1720 restored the expressions of PGC-1 α and E-cadherin reduced by HDM in BEAS-2B cells. Treatment with SRT1720 also significantly ameliorated the HDM-induced reduction in TEER. In addition, GB83, LPS-RS, E64 and AEBSF prevented the HDM-induced reduction in the expression of PGC1 α and E-cadherin.

Conclusions: The current study demonstrated that HDM disrupted the airway barrier function through the PAR2/TLR4/PGC-1 α -dependent pathway. The modulation of this pathway could be a new approach for the treatment of asthma.

Background

Asthma is a chronic airway inflammatory disease that is characterized by variable symptoms of wheeze, shortness of breath, chest tightness and/or cough, and by variable expiratory airflow limitation.¹ These variations are often triggered by miscellaneous external factors including allergen, irritant exposure, climate change, or viral respiratory infections. The bronchial epithelium forms a physical barrier against the external environment through the formation of cell-cell junctions consisting of tight junctions, adherence junctions and desmosomes, referred to as the epithelial junctional complex.^{2 3} Tight junctions and adherence junctions are located at the apical end of the lateral membrane and are comprised of numerous membrane proteins including occludin, zonula occludens (ZO)-1 and E-cadherin.^{2 3} A reduction in the expression of E-cadherin and ZO-1 is reportedly observed in the airways of asthmatic patients.^{4 5} Epithelial damage can be caused by respiratory viruses, cigarette smoke, and allergens such

as *Dermatophagoides pteronyssinus* and fungal allergens.^{6 7} Disruption of the epithelial barrier increases the susceptibility to external stimuli leading to airway hyperresponsiveness. Furthermore, a damaged epithelial barrier increases the accessibility of allergens into the submucosa activating the subsequent immune responses. Thus, regulation of the bronchial epithelial function has been attracting attention as an important immunological checkpoint in asthma. However, the precise mechanisms by which epithelial junctions are disrupted are not fully understood.

In airway epithelial cells and BEAS-2B cells, interleukin (IL)-4 reportedly promotes intracellular asymmetric dimethylarginine (ADMA) accumulation, which causes a reduction in mitochondrial biogenesis.⁸ Though the result of the mitochondrial biogenesis reduction is unknown, since most important role of airway epithelial cells is the airway barrier function, it is probable that the reduction affects airway the barrier disfunction.

Mitochondria play a key role in energy homeostasis and the metabolism of reactive oxygen species (ROS).⁹ Appropriate elimination of damaged mitochondria through mitochondrial autophagy (mitophagy) and the renewal of mitochondria by mitochondrial biogenesis are essential for mitochondrial homeostasis.¹⁰ Mitochondrial biogenesis is regulated mainly at the transcriptional level and requires the coordinated expression of both nuclear-encoded and mitochondrial-encoded proteins, including peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-1 α), mitochondrial transcriptional factor A (TFAM), adenosine 5'-monophosphate-activated protein kinase (AMPK), and nuclear respiratory factors (NRF)-1 and -2.¹⁰ Among these molecules, PGC-1 α is the key regulator of mitochondrial biogenesis.¹¹

Sirtuin 1 (SIRT1) is a powerful deacetylase that has been shown to activate PGC-1 α to drive mitochondrial biogenesis,¹² and SRT1720, the activator of SIRT1, is an effective SIRT1 agonist that enhances PGC-1 α activation.^{13 14 15} In previous reports, SRT1720 alleviated lung injury and improved the lung function in rat with emphysema caused by cigarette smoke through protecting against the apoptosis of type II alveolar epithelial cells.¹⁶ SRT1720 inhibited the differentiation of TGF- β 1-induced myofibroblasts.¹⁷ SRT1720 repressed the LPS-induced release of cytokines such as IL-8, IL-6 and tumor necrosis factor (TNF)- α from cultured peripheral blood mononuclear cells.¹⁸ In a report about asthma, SRT1720 also suppressed inflammatory cell infiltration and cytokine production including TNF- α and IL-6 in the lungs of an ovalbumin (OVA)-induced mouse model.¹⁹ It is probable that the activation of mitochondrial biogenesis by SRT1720 in airway epithelial cells contributes to amelioration of the asthma pathophysiology.

Therefore, the current study aimed to clarify the contribution of regulators of mitochondrial biogenesis to airway barrier dysfunction. We assessed the effect of house dust mite (HDM), a common aeroallergen related to asthma, on the expression of mitochondrial biogenesis markers and junctional proteins in airway epithelial cells. Furthermore, we investigated how an activator of PGC-1 α modulates the constitution of junctional proteins and the airway barrier function to explore novel therapeutic targets for bronchial asthma.

Materials And Methods

Materials

The following reagents were used in this study: purified HDM extract from dermatophagoides pteronyssinus was purchased from LSL (Tokyo, Japan); SRT1720 was from Selleck Chemicals (Houston, TX); GB83 was from Axon Medchem (Groningen, Netherlands); Lipopolysaccharide from *Rhodobacter sphaeroides* (LPS-RS) was from Invivogen (San Diego, CA); Dexamethasone, E64, 4-(2-Aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF), and mouse monoclonal anti- β -actin antibody were from Sigma (St Louis, MO). Protein block, a blocking reagent, was from Dako (Kyoto, Japan); Rabbit polyclonal anti-PGC-1 α antibody, rabbit monoclonal anti-TFAM antibody, rabbit monoclonal anti-E-cadherin antibody, rabbit polyclonal anti-ZO-1 antibody, FITC-conjugated goat anti-rabbit secondary antibody, and Alexa Fluor 647-conjugated goat anti-rabbit secondary antibody were from Abcam (Cambridge, MA); horseradish peroxidase-conjugated secondary antibodies were from Santa Cruz Biotechnology (Dallas, TX); MitoTracker Red probe and Hoechst 33342 were from Invitrogen Life Technologies (Eugene, OR); Keratinocyte-SFM (serum-free medium) and Human Keratinocyte Growth Supplements were from Gibco (Grand Island, NY). Bronchial epithelial growth medium (BEGM) bullet kit including bronchial epithelial basal medium (BEBM) and subculture reagents were purchased from Lonza Japan Bioscience (Tokyo, Japan)

Cell culture

BEAS-2B cells, a human bronchial epithelial cell line, were obtained from American Type Culture Collection (Manassas, VA). The BEAS-2B cells were cultured in Keratinocyte-SFM medium supplemented with Keratinocytes Supplements at 37 °C in a humidified atmosphere of 5 % CO₂ and passaged in 75 cm² flasks. Normal primary human bronchial epithelial cells (pHBEC) were obtained from Lonza (Wokingham, UK) and cultured in BEGM. To prepare BEGM, BEBM was supplemented with human recombinant epidermal growth factor (0.5 ng/ml), insulin (5 μ g/ml), transferrin (10 μ g/ml), hydrocortisone (0.5 μ g/ml), triiodothyronine (6.5 μ g/ml), epinephrine (0.5 μ g/ml), retinoic acid (50 nM), gentamycin and amphotericin-B (50 μ g/ml) and bovine pituitary extract (35 mg/ml).

Animals

8-week-female C57BL/6J mice were purchased from SEIMI (Sendai, Japan). All mice were housed in a specific pathogen-free facility and maintained under constant temperature (24°C), humidity (40%), and light cycle (8:00 A.M. to 8:00 P.M.), with food and water provided ad libitum. The mice were exposed to HDM (30 μ g/body, 50 μ l) intranasally every other day as described previously.²⁰ The control mice received PBS. The mice were euthanized on day 15, and the trachea was cannulated after the pulmonary circulation was perfused and free of blood. The lung was inflated with 10% paraformaldehyde for 10 min

at 25 cm H₂O and subsequently removed and fixed in 10% paraformaldehyde for 24 h at room temperature. Samples were then dehydrated in ethanol and xylene, embedded in paraffin, cut in 5- μ m sections. All experiments were approved by the Tohoku University Animal Experiment Ethics Committee and performed in accordance with the Regulations for Animal Experiments and Related Activities at Tohoku University.

Western blotting

The expression of markers of mitochondrial biogenesis (PGC-1 α and TFAM) and junctional proteins (E-cadherin and ZO-1) in BEAS-2B cells was analyzed by western blotting. Cells (2×10^5 cells/ml) were seeded in 6-well culture plates and grown to 80-90 % confluence. Then the cells were maintained in Keratinocytes Supplements free medium for 24 h and stimulated with or without HDM (100 μ g/ml). 24 h after HDM stimulation, the cells were homogenized in cell lysis buffer (0.05 % TritonX, 3 mM Tris-HCl, pH7.4 0.4 mM EGTA, 10 mM MgCl₂, 1 μ M phenylmethylsulfonyl fluoride, 100 μ g/ml aprotinin, 1 μ g/ml leupeptin) on ice and harvested for western blotting. SRT1720 (1 μ M), a PGC-1 α activator, was added to the culture medium at 6 h before HDM stimulation. Protease activated receptor (PAR) 2 inhibitor GB83, Toll-like receptor (TLR) 4 inhibitor LPS-RS, cysteine peptidase inhibitor E64, and serine peptidase inhibitor AEBSF were added at 30 min before HDM stimulation. Synthetic glucocorticoid dexamethasone was added at 24 h before HDM stimulation. The whole cell lysates were homogenized for several seconds using an ultrasonic homogenizer (BANDELIN, Berlin, Germany). The samples were solubilized in SDS-PAGE sample buffer (Bio-Rad, Hercules, CA). Equal amounts of protein were loaded and separated by electrophoresis on 12% SDS polyacrylamide gels. After electrophoresis, the separated proteins were transferred to polyvinylidene difluoride membranes (Millipore, Darmstadt, Germany). The membranes were blocked with a blocking reagent. Rabbit polyclonal anti-PGC-1 α antibody (1:5000), rabbit monoclonal anti-TFAM antibody (1:5000), rabbit monoclonal anti-E-cadherin antibody (1:10000), rabbit polyclonal anti-ZO-1 antibody (1:2000), and mouse monoclonal anti- β -actin antibody (1:5000) were used as primary antibodies to detect the target proteins. Horseradish peroxidase-conjugated secondary antibodies (1:5000) were detected using ECL-prime Western Blotting Reagent (Amersham Biosciences, Buckinghamshire, UK) and visualized with a chemiluminescence imaging system (LAS-4000 mini, FUJIFILM, Tokyo, Japan). Each band intensity was quantified using ImageJ (National Institutes of Health, Bethesda, MD, USA).

Evaluation of mitochondrial mass and immunochemical localization of E-cadherin and ZO-1 in BEAS-2B cells

BEAS-2B cells were seeded in 8-well chamber slides at a density of 1×10^5 cells/ml and cultured for 24 h, and then the media were replaced with the keratinocyte supplement-free medium for 24 h. After incubation with 1 μ M SRT1720 for 6 h, the cells were stimulated with HDM (100 μ g/ml) for 24 h. The mitochondrial mass was evaluated using MitoTracker Red probe. MitoTracker Red was added to the

existing medium to a final concentration of 200 nM and cells were incubated for 30 min at 37 °C, 5 % CO₂. After staining, the cells were washed once with PBS. Then, the slides were fixed with freshly prepared 4 % paraformaldehyde in PBS for 30 min at room temperature. The slides were permeabilized with 0.1 % Triton X-100 in PBS for 10 min at room temperature. The slides were blocked with a blocking solution for 30 min at room temperature. The slides were then incubated overnight with rabbit monoclonal anti-E-cadherin antibody (1:500), rabbit polyclonal anti-ZO-1 antibody (1:200) at 4°C. The slides were incubated with the appropriate FITC-conjugated secondary antibody (1:3000) for 1 h at room temperature. The nuclei of the cells were stained with Hoechst 33342 (1:200). Fluorescent images were obtained using a confocal laser scanning microscope system (Nikon ECLIPSE Ti-E, C2si; Nikon, Tokyo, Japan). The mean fluorescence intensity of E-cadherin, ZO-1, MitoTracker were quantified by using ImageJ.

Immunochemical localization of PGC-1 α , TFAM and E-Cadherin in lung tissues from mice

Tissue samples were fixed in 10% formalin and embedded in paraffin. 5 μ m thick serial tissue sections were obtained and mounted in Superfrost/Plus glass slides (Fischer Scientific). Deparaffinization was performed by washing three times for 5 minutes in xylene, then washing in 100%, 95%, 80%, 70% ethanol three times for 5 minutes and, finally, rinsing with distilled water. Slides were washed in PBS for 5 min and processed for antigen retrieval using citrate buffer. The sections were permeabilized with Triton-X 100 for 15 min. Sections were then washed in PBS, blocked using blocking reagent (Dako) for 60 min at room temperature. Sections were incubated overnight at 4 °C with rabbit polyclonal anti-PGC-1 α antibody (1:1000), rabbit monoclonal anti-TFAM antibody (1:10000), rabbit monoclonal anti-E-Cadherin antibody (1:10000) or nonspecific polyclonal rabbit IgG as a negative control at 4°C overnight. After being washed, the samples were incubated with Alexa Fluor 647-conjugated goat anti-rabbit secondary antibody (1:3000) for 60 min at room temperature. After washing, the nuclei of the cells were stained with Hoechst 33342 (1:200). Fluorescent images were obtained by microscopy (BX53-33-SDO; Olympus, Tokyo, Japan) and photographed with a digital camera (DP71-SET; Olympus). The mean fluorescence intensities of PGC-1 α , TFAM, and E-cadherin were quantified by using ImageJ.

Transepithelial electrical resistance measurements (TEER)

Primary human bronchial epithelial cells were seeded onto collagen type I coated 12-well transwell inserts (transparent, 0.4 μ m; Greiner) at 1×10^5 cells/cm² with 500 μ l apical and 1500 μ l basolateral volumes, maintained in BEGM, and incubated until the formation of cell monolayers. Then, SRT1720 (1 μ M) was added to the apical medium. After 24 h incubation with SRT1720, the cells were stimulated with HDM (100 ng/ml) or DMSO (vehicle) for 24 h. Then, TEER was monitored using a Millicell ERS-2 voltohmmeter (Millipore, Billerica, MA).

Statistical Analysis

Results are presented as mean \pm SD. Comparisons among groups were performed by One-way analysis of variance followed by the Tukey's multiple comparison test or unpaired t-test. $p < 0.05$ was considered statistically significant.

Results

Expression of mitochondrial biogenesis markers and junctional proteins in HDM-exposed BEAS-2B cells

Because HDM is a common aeroallergen in asthmatic patients, we investigated the effect of HDM on the expression of mitochondrial biogenesis makers and junctional proteins in BEAS-2B cells. The cells were treated with HDM at various concentrations and time points. HDM significantly reduced the expression of TFAM in a concentration-dependent (Figure S1A) and time-dependent manner (Figure S1B). In the HDM-treated cells, the expression of markers of mitochondrial biogenesis was significantly reduced compared to the control cells (PGC-1 α , $p < 0.01$; TFAM, $p < 0.01$) (Fig. 1A, B). The expression of junctional proteins was also significantly reduced compared to the control cells (E-cadherin, $p < 0.01$; ZO-1, $p < 0.01$) (Fig. 1C, D).

Expressions of mitochondrial biogenesis markers and junctional proteins in the lungs of HDM-treated mice.

The expressions of mitochondrial biogenesis markers and junctional proteins were also analyzed by immunofluorescence in the lungs of mice treated with HDM. The immunofluorescence intensities of PGC-1 α , TFAM, and E-Cadherin in the epithelial region were significantly reduced in the lungs from HDM-administered mice compared to the lungs from vehicle-treated mice (PGC-1 α , $p < 0.01$; TFAM, $p < 0.01$; E-cadherin, $p < 0.01$) (Fig. 2A-D).

Effect of SRT1720, a PGC-1 α activator, on the HDM-induced reduction of junctional protein expression and epithelial barrier dysfunction in BEAS-2B cells and pHBEC

To identify whether a PGC-1 α -dependent pathway regulates the expression of junctional proteins in bronchial epithelial cells, we investigated the effects of SRT1720, which is an enhancer of mitochondrial biogenesis by activating PGC-1 α , on the HDM-induced reduction in the expression of junctional proteins. SRT1720 (1 μ M) significantly restored the HDM-induced reduction in the expression of PGC-1 α ($p < 0.05$). Treatment with SRT1720 also significantly restored the HDM-induced reduction in the expression of TFAM ($p < 0.05$) and E-cadherin ($p < 0.05$) in BEAS-2B cells (Fig. 3A-D). Next, we investigated the effect of

SRT1720 on the barrier function by measuring TEER in pHBEC treated with HDM. Treatment with SRT1720 significantly ameliorated the HDM-induced reduction in TEER ($p < 0.05$) (Fig. 3E).

Effect of SRT1720 on the HDM-induced reduction of junctional protein expression in BEAS-2B cells by immunocytochemistry.

We also investigated the effect of SRT1720 on the HDM-induced reduction in the expression of junctional proteins in BEAS-2B cells by immunocytochemistry. Treatment with SRT1720 also restored the HDM-induced reduction in the intensity of junctional proteins expression as confirmed (Fig. 4A-D). Next, the mitochondrial mass in BEAS-2B cells was analyzed by immunofluorescence using a MitoTracker Red probe. HDM reduced the mitochondrial mass in the cells, which was restored by treatment with SRT1720 (Fig. 4A,4B and 4E).

The role of PAR2 and TLR4 signaling in the HDM-induced reduction of PGC-1 α and junctional protein expression in BEAS-2B cells

HDM contains both proteases and endotoxin, which activate PAR2 and TLR4, respectively. To determine which pathway regulates the HDM-induced reduction in PGC-1 α and the subsequent disruption of junctional proteins, we investigated the effects of blockade of PAR2 and TLR4 on the HDM-induced decrease in the expression of PGC-1 α and junctional proteins using GB83, a PAR2 inhibitor, and LPS-RS, a TLR4 inhibitor. GB-83 significantly blocked the HDM-induced reduction in the expression of PGC-1 α ($p < 0.01$), E-cadherin ($p < 0.05$) and ZO-1 ($p < 0.01$) (Fig. 5A-C) compared to vehicle treatment. Interestingly, LPS-RS also blocked the HDM-induced reduction in the expression of PGC-1 α ($p < 0.05$), E-cadherin ($p < 0.05$) and ZO-1 ($p < 0.05$) (Fig. 5D-F).

Effect of protease inhibitors on the HDM-induced decrease of PGC-1 α and E-cadherin expressions in BEAS-2B cells

Serine protease and cysteine protease are present in HDM excrement and cleave junctional proteins directly or indirectly via PAR2 activation. To determine the role of protease activity in the HDM-mediated barrier disruption, we investigated the effects of protease inhibitors on the expression of PGC-1 α and E-cadherin in BEAS-2B cells treated with HDM. E64, a cysteine protease inhibitor, significantly blocked the HDM-induced reduction in the expression of PGC-1 α ($p < 0.05$) (Fig. 6A), and E-cadherin ($p < 0.05$) (Fig. 6B). AEBSF, a serine protease inhibitor, also significantly prevented the HDM-induced reduction in the expression of PGC-1 α ($p < 0.05$) (Fig. 6C) and E-cadherin ($p < 0.01$) (Fig. 6D).

Effect of corticosteroids on the expression of mitochondrial biogenesis markers and junctional proteins in BEAS-2B cells

As inhaled corticosteroids are the central medication for the management of patients with asthma, there is concern about the influence of corticosteroids on the expression of mitochondrial biogenesis markers and the composition of junctional proteins in the airways of asthmatic patients. To address this issue, we investigated whether corticosteroids can affect the expression of markers of mitochondrial biogenesis and junction protein in BEAS-2B cells. Dexamethasone had no significant effect on the baseline expressions of PGC-1 α , TFAM and E-cadherin in the cells (Figure S2A-C). In addition, dexamethasone had no preventive effect on the HDM-induced reduction in the expression of PGC-1 α , TFAM and E-cadherin (Fig. 7A-C).

Discussion

In the current study, HDM, a common aeroallergen to which asthmatic patients are sensitized, decreased the expression levels of the regulators concurrently with the reduction in the amounts of proteins related to the formation of epithelial junctional complex. Activation of PGC-1 α by SIRT1720 restored the HDM-induced mitochondrial loss and reduction in the expression of E-cadherin, leading to recovery of the airway barrier function. These data suggest that a PGC-1 α -dependent pathway may play a central role in the airway barrier dysfunction induced by aeroallergen.

In the context of the pathophysiology of asthma, a previous report showed that the up-regulation of PGC-1 α in bronchial smooth muscle from asthmatic patients²¹ and that non-immune immunoglobulin E (IgE) increased the PGC-1 α mRNA and protein levels in primary human airway smooth muscle cells through the activation of phosphatidylinositol 3-kinase (PI3K)-Akt pathway²², suggesting their involvement in airway wall remodeling. This study suggests that SRT1720 may cause exacerbation of the asthma pathology. Alternatively, the airway epithelial barrier dysfunction propagates the infiltration of inflammation into the submucosa and impairs the process of epithelial repair. Therefore, the airway epithelial barrier dysfunction is profoundly involved in the progress of remodeling.²³ Thus, SRT1720 may contribute to the prevention of remodeling due to the barrier repair effect on airway epithelium.

The innate immune system plays critical roles in the pathogenesis of asthma by recognizing allergens through pattern-recognition receptors (PRRs) including TLRs and PARs, and by providing an early warning system through the production of cytokines and danger signals in bronchial epithelial cells.^{24 25} Many human allergens including HDM, fungi, pollen, and cat contain protease activity and can disrupt tight junction proteins through direct proteolytic cleavage.²⁶ Furthermore, proteases in HDM activate PAR2 resulting in the loss of E-cadherin in human bronchial epithelial cells.²⁷ HDM allergens also contain endotoxin and activate TLR4 in bronchial epithelial cells to induce asthmatic features including the release of epithelium-derived cytokines such as IL-25 and IL-33.²⁸ Therefore, we investigated the effects of antagonists against PAR2 and TLR4 on the HDM-induced reduction in PGC-1 α and tight junction

proteins to determine which signaling is responsible. Interestingly, both antagonists significantly prevented the repression of PGC-1 α and E-cadherin in HDM-treated BEAS-2B cells. A previous report demonstrated that LPS triggered the activation of NF- κ B and the subsequent production of pro-inflammatory cytokines through the cooperative activation of PAR2 and TLR4 in HEK293T cells²⁹, which is consistent with our findings. It has been reported that the inhibition of TLR4 reverses the decreased expression of PGC-1 α in the renal tubules of diabetic mice, suggesting that the TLR4-dependent pathway could be the upstream pathway of PGC-1 α .³⁰ Meanwhile, protease inhibitors were shown to block the HDM-induced reduction in the expression of PGC-1 α in this study. Taken together, both TLR4- and PAR2-mediated signaling pathways might be involved in the HDM-induced airway barrier disruption via the regulation of PGC-1 α . Details of the interactions between TLR4 and PAR2 during mitochondrial biogenesis and maintenance of the airway barrier function remain unclear and further study is needed to clarify such interactions.

In the current study, we showed that corticosteroids administration did not affect the expression of mitochondrial biogenesis markers, E-cadherin expression, and their HDM-induced reduction in BEAS-2B cells. From these results, it may be considered that the effects of corticosteroids on mitochondrial biogenesis in asthmatic airway epithelial cells are negligible.

Conclusion

In conclusion, the current study demonstrated the interaction between innate immunity and mitochondrial biogenesis in the context of inhaled allergen-induced airway barrier dysfunction, an initial feature in the development of asthma. The PAR2/TLR4/PGC-1 α -dependent pathway might be a novel and promising therapeutic target for bronchial asthma as, so far, no effective medication for restoring the airway barrier function has been established. Regulating this pathway could inhibit allergic reaction in airways driven by aeroallergens because restoration of the airway barrier function can prevent the penetration of external stimuli, which would suppress the cytokine release that activates type 2 immune reactions. Although further study is needed to confirm the beneficial effect of PGC-1 α activator on immunological reactions and to determine the molecules involved in the pathway, this approach might be applicable to a broad array of allergic diseases including allergic rhinitis and atopic dermatitis in addition to bronchial asthma.

Abbreviations

AEBSF : 4- (2-aminoethyl) benzenesulfonyl fluoride hydrochloride

HDM : house dust mite

LPS-RS : lipopolysaccharide from *Rhodobacter sphaeroides*

NRF1 : nuclear respiratory factor 1

PAR : protease activated receptor

PGC-1 α : peroxisome proliferator-activated receptor γ coactivator-1 α

PRRs : pattern-recognition receptors

Sirtuin 1 : SIRT1

TEER : transepithelial electrical resistance

TFAM : mitochondria transcription factor A

TLR : Toll-like receptor

ZO-1: zonula occludens-1

Declarations

Ethics approval and consent to participate:

Not applicable.

Consent for publication:

Not applicable.

Availability of data and material:

Not applicable.

Funding:

This study was supported by a grant from the Japan Society for the Promotion of Science (grant number: #18K08166), a grant from the Practical Research Project for Allergic Diseases and Immunology from Japan Agency for Medical Research and Development, AMED (grant number: #16ek0410018h0002, #16ek040036h0001, #17ek0410036h0002).

Competing interests:

Authors have no competing interests.

Authors' contributions:

T. Saito and T. Ichikawa equally contributed to this study.

T. Saito: biochemical studies, data collection, writing of the manuscript.

T. Ichikawa: design of the study, biochemical studies, immunocytochemical analysis, data collection, interpretation of the results, technical advice, writing of the manuscript.

T. Numakura: interpretation of the results, technical advice.

M. Yamada: interpretation of the results, technical advice.

A. Koarai: interpretation of the results, technical advice.

N. Fujino: technical advice.

K. Murakami: measuring TEER.

S. Yamanaka: technical advice.

Y. Sasaki: technical advice.

Y. Kyogoku: technical advice.

K. Itakura: technical advice.

K. Takita: technical advice.

R. Tanaka: interpretation of the results.

T. Tamada: interpretation of the results.

M. Ichinose: design of the study, interpretation of the results.

H. Sugiura: design of the study, interpretation of the results, technical advice.

Acknowledgments:

We thank Mr. Brent Bell for reading the manuscript.

References

1. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention 2019. <https://ginasthma.org/wp-content/uploads/2019/06/GINA-2019-main-report-June-2019-wms.pdf> Accessed July 1, 2019.
2. Georas SN, Rezaee F. Epithelial barrier function: at the front line of asthma immunology and allergic airway inflammation. *J Allergy Clin Immunol.* 2014;134(3):509-520. doi:10.1016/j.jaci.2014.05.049
3. Zihni C, Mills C, Matter K, Balda MS. Tight junctions: from simple barriers to multifunctional molecular gates. *Nat Rev Mol Cell Biol.* 2016;17(9):564-580. doi:10.1038/nrm.2016.80
4. Xiao C, Puddicombe SM, Field S, et al. Defective epithelial barrier function in asthma. *J Allergy Clin Immunol.* 2011;128(3). doi:10.1016/j.jaci.2011.05.038
5. de Boer WI, Sharma HS, Baelemans SM, et al. Altered expression of epithelial junctional proteins in atopic asthma: possible role in inflammation. *Can J Physiol Pharmacol.* 2008;86(3):105-112.
6. Blume C, Swindle EJ, Dennison P, et al. Barrier responses of human bronchial epithelial cells to grass pollen exposure. *Eur Respir J.* 2013;42(1):87-97. doi:10.1183/09031936.00075612

7. Swindle EJ, Collins JE, Davies DE. Breakdown in epithelial barrier function in patients with asthma: Identification of novel therapeutic approaches. *J Allergy Clin Immunol*. 2009;124(1):23-34. doi:10.1016/j.jaci.2009.05.037
8. Pattnaik B, Bodas M, Bhatraju NK, et al. IL-4 promotes asymmetric dimethylarginine accumulation, oxo-nitrative stress, and hypoxic response-induced mitochondrial loss in airway epithelial cells. *J Allergy Clin Immunol*. 2016;138(1):130-141 e9. doi:10.1016/j.jaci.2015.11.036
9. Gureev AP, Shaforostova EA, Popov VN. Regulation of Mitochondrial Biogenesis as a Way for Active Longevity : Interaction Between the Nrf2 and PGC-1 α Signaling Pathways. 2019;10(May):1-12. doi:10.3389/fgene.2019.00435
10. Piantadosi CA, Suliman HB. Mitochondrial Dysfunction in Lung Pathogenesis. *Annu Rev Physiol*. 2017;79:495-515. doi:10.1146/annurev-physiol-022516-034322
11. Wu Z, Puigserver P, Andersson U, et al. Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell*. 1999;98(1):115-124. doi:10.1016/S0092-8674(00)80611-X
12. Sepharose PG, Rodgers JT, Lerin C, et al. Nutrient control of glucose homeostasis through a complex of PGC-1 α and SIRT1. *Nature*. 2005;434(March):3-8. doi:10.1038/nature03314.1.
13. Wan X, Wen JJ, Koo SJ, Liang LY, Garg NJ. SIRT1-PGC1 α -NF κ B Pathway of Oxidative and Inflammatory Stress during *Trypanosoma cruzi* Infection: Benefits of SIRT1-Targeted Therapy in Improving Heart Function in Chagas Disease. *PLoS Pathog*. 2016;12(10):e1005954. doi:10.1371/journal.ppat.1005954
14. Funk JA, Odejinmi S, Schnellmann RG. SRT1720 induces mitochondrial biogenesis and rescues mitochondrial function after oxidant injury in renal proximal tubule cells. *J Pharmacol Exp Ther*. 2010;333(2):593-601. doi:10.1124/jpet.109.161992
15. Svensson K, Schnyder S, Albert V, et al. Resveratrol and SRT1720 Elicit Differential Effects in Metabolic Organs and Modulate Systemic Parameters Independently of Skeletal Muscle Peroxisome Proliferator-activated Receptor γ Co-activator 1 α (PGC-1 α). *J Biol Chem*. 2015;290(26):16059-16076. doi:10.1074/jbc.M114.590653
16. Gu C, Li Y, Xu WL, et al. Sirtuin 1 activator SRT1720 protects against lung injury via reduction of type II alveolar epithelial cells apoptosis in emphysema. *COPD J Chronic Obstr Pulm Dis*. 2015;12(4):444-452. doi:10.3109/15412555.2014.974740
17. Zeng Z, Cheng S, Chen H, et al. Activation and overexpression of Sirt1 attenuates lung fibrosis via P300. *Biochem Biophys Res Commun*. 2017;486(4):1021-1026. doi:10.1016/j.bbrc.2017.03.155
18. Yao H, Sundar IK, Huang Y, et al. Disruption of sirtuin 1-mediated control of circadian molecular clock and inflammation in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol*. 2015;53(6):782-792. doi:10.1165/rcmb.2014-04740C
19. Ichikawa T, Hayashi R, Suzuki K, et al. Sirtuin 1 activator SRT1720 suppresses inflammation in an ovalbumin-induced mouse model of asthma. *Respirology*. 2013;18(2):332-339. doi:10.1111/j.1440-1843.2012.02284.x

20. Ishii T, Niikura Y, Kurata K, et al. Time-dependent distinct roles of Toll-like receptor 4 in a house dust mite-induced asthma mouse model. *Scand J Immunol*. 2018;87(3):1-7. doi:10.1111/sji.12641
21. Trian T, Benard G, Begueret H, et al. Bronchial smooth muscle remodeling involves calcium-dependent enhanced mitochondrial biogenesis in asthma. *J Exp Med*. 2007;204(13):3173-3181. doi:10.1084/jem.20070956
22. Fang L, Wang X, Sun Q, et al. IgE downregulates PTEN through microRNA-21-5p and stimulates airway smooth muscle cell remodeling. *Int J Mol Sci*. 2019;20(4):1-20. doi:10.3390/ijms20040875
23. Davies DE. The role of the epithelium in airway remodeling in asthma. *Proc Am Thorac Soc*. 2009;6(8):678-682. doi:10.1513/pats.200907-067DP
24. Lambrecht BN, Hammad H. Allergens and the airway epithelium response: gateway to allergic sensitization. *J Allergy Clin Immunol*. 2014;134(3):499-507. doi:10.1016/j.jaci.2014.06.036
25. Salazar F, Ghaemmaghami AM. Allergen recognition by innate immune cells: critical role of dendritic and epithelial cells. *Front Immunol*. 2013;4:356. doi:10.3389/fimmu.2013.00356
26. Nawijn MC, Hackett TL, Postma DS, van Oosterhout AJ, Heijink IH. E-cadherin: gatekeeper of airway mucosa and allergic sensitization. *Trends Immunol*. 2011;32(6):248-255. doi:10.1016/j.it.2011.03.004
27. Jacquet A. Interactions of airway epithelium with protease allergens in the allergic response. *Clin Exp Allergy*. 2011;41(3):305-311. doi:10.1111/j.1365-2222.2010.03661.x
28. Hammad H, Chieppa M, Perros F, Willart MA, Germain RN, Lambrecht BN. House dust mite allergen induces asthma via Toll-like receptor 4 triggering of airway structural cells. *Nat Med*. 2009;15(4):410-416. doi:10.1038/nm.1946
29. Rallabhandi P, Nhu QM, Toshchakov VY, et al. Analysis of proteinase-activated receptor 2 and TLR4 signal transduction: a novel paradigm for receptor cooperativity. *J Biol Chem*. 2008;283(36):24314-24325. doi:10.1074/jbc.M804800200
30. Yuan S, Liu X, Zhu X, et al. The Role of TLR4 on PGC-1 α -Mediated Oxidative Stress in Tubular Cell in Diabetic Kidney Disease. *Oxid Med Cell Longev*. 2018;2018:1-14. doi:10.1155/2018/6296802

Figures




Figure 1

Expression of mitochondrial biogenesis markers and junction proteins in bronchial epithelial cells. A-D, BEAS-2B cells were treated with HDM (100 μ g/ml) or vehicle for 24 h and the cells were harvested. Expressions of PGC-1 α , TFAM, E-cadherin and ZO-1 were analyzed by western blotting. Data are expressed as means \pm SD (n = 3). *p < 0.05, **p < 0.01.




Figure 2

Expression of mitochondrial biogenesis markers in the lungs of HDM-treated mice. A, C57BL/6 mice were intranasally exposed to PBS or HDM extract every other day for 2 weeks and the mice were sacrificed. Lung tissues were removed after fixation and immune-fluorescence staining was performed to investigate the expressions of mitochondrial biogenesis markers (PGC-1 α and TFAM) and E-Cadherin in the lungs of the mice. Representative photographs are shown. Bars: 100 μ m. B-D, The mean fluorescence intensity of PGC-1 α , TFAM, and E-cadherin were quantified by using ImageJ. Data are expressed as means \pm SD (n = 4). **p < 0.01.




Figure 3

Effect of SRT1720 on mitochondrial biogenesis and airway barrier function mediated by HDM. A, BEAS-2B cells were treated with various concentrations of SRT-1720 or vehicle for 6 hours. Subsequently, the cells were stimulated with HDM (100 μg/ml) for 24 h and the cells were harvested. The expression of PGC-1α in the cells was analyzed by western blotting. (n = 3). **p < 0.01. B-D, BEAS-2B cells were treated with vehicle or SRT-1720 (1 μM) for 6 hours. Subsequently, the cells were stimulated with HDM (100 μg/ml) for 24 h and the cells were harvested. The expressions of PGC-1α, TFAM, and E-cadherin in the cells were analyzed by western blotting (n = 8). E, Transepithelial electrical resistance (TEER) in primary human bronchial epithelial cells (pHBEC) were measured (n = 4) after the cells were stimulated by HDM (100 μg/ml) or PBS for 24 h with or without SRT-1720 (1 μM) using a Millicell ERS-2 voltohmmeter. All data are expressed as means ± SD. *p < 0.05, **p < 0.01.




Figure 4

Effect of SRT1720 on mitochondrial mass and expressions of junctional proteins mediated by HDM. A and B, Localization of E-cadherin, ZO-1 and the mitochondrial mass in the cells after exposure to HDM were analyzed by immunofluorescence. Representative confocal microscopic images are shown. The cells were stained with E-cadherin (green), ZO-1 (green), MitoTracker (red), and nuclear counterstain Hoechst33342 (blue). Bars: 100 μ m. C-E, The mean fluorescence intensity of E-cadherin, ZO-1, MitoTracker were quantified using ImageJ. Data are expressed as means \pm SD. * p < 0.05, ** p < 0.01.




Figure 5

Effects of PAR2 and TLR4 inhibitors on HDM-mediated downregulation of mitochondrial biogenesis and junctional proteins. A-F, BEAS-2B cells were treated with various concentrations of GB83, a PAR2 inhibitor or LPS-RS, a TLR4 inhibitor for 30 min. Subsequently, the cells were stimulated with HDM (100 $\mu\text{g}/\text{ml}$) for 24 h and the cells were harvested. The expressions of PGC-1 α , E-cadherin, and ZO-1 in cells were analyzed by western blotting. (LPS-RS, n=3; others, n=4). Data are expressed as means \pm SD. * $p < 0.05$, ** $p < 0.01$.




Figure 6

Effects of protease inhibitors on HDM-mediated downregulation of mitochondrial biogenesis and junctional proteins. A-D, BEAS-2B cells were treated with E64 (50 μM) or AEBSF (10 and 100 μM) for 30

min. Subsequently, the cells were stimulated with HDM (100 µg/ml) for 24 h and the cells were harvested. The expressions of PGC-1α and E-cadherin were analyzed by western blotting. (n=4). Data are expressed as means ± SD. *p < 0.05, **p < 0.01.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.pdf](#)
- [FigureS2.pdf](#)