Knowledge, Attitudes and Practices regarding HIV among Women of Childbearing Age in NYANGA Province, Gabon

Augustin Mouinga-Ondémé (✉ ondeme@yahoo.fr)

Centre International de Recherches Medicales de Franceville <https://orcid.org/0000-0002-3143-1060>

Sydney Maghendji-Nzondo

Universite des Sciences de la Sante de Libreville

Julie Buekens

Medical Care Development International

Jéordy-Dimitri Engone-Ondo

Centre International de Recherches Medicales de Franceville

Roméo Karl Imboumy-Limoukou

Centre International de Recherches Medicales de Franceville

Pater Noster Sir-Ondo-Enguier

Centre International de Recherches Medicales de Franceville

Julie Niemczura de Carvalho

Medical Care Development International

Nathalie Pernelle Tsafack-Tegomo

Centre International de Recherches Medicales de Franceville

Jean-Bernard Lékana-Douki

Centre International de Recherches Medicales de Franceville

Research Article

Keywords: HIV, PMTCT, Prevention, traditional beliefs, witchcraft, Nyanga, Gabon

Posted Date: February 25th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-968622/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: The seroprevalence of HIV in Gabon is 4.1%. This rate varies among its nine provinces and is estimated at 4.2% in the province of Nyanga. This variance is believed to be the result of a lack of information and beliefs regarding HIV transmission and prevention. There are currently scarce epidemiological data from this region of Gabon. We assessed the knowledge, attitudes and practices of women of childbearing age from Nyanga Province toward HIV.

Methods: A 13-day cross-sectional descriptive study of 631 women aged 15 to 49 years was conducted within the departments of the province to assess their knowledge and attitudes about HIV, including mother-to-child transmission (MTCT). Household recruitment of participants was conducted, and information was collected by means of a questionnaire. Correct answers were considered to indicate basic knowledge of the topic assessed.

Results: Approximately 90% of women (570/631) were of childbearing age but not pregnant during the survey, with a mean age of 28 years. Additionally, 10% (61/631) were pregnant. Among surveyed women, 99.2% had heard of HIV/AIDS, and 81.8% knew at least one screening site for HIV. More than 83.6% knew that MTCT could occur during pregnancy, and 76.5% knew about transmission during breastfeeding. Moreover, 86.5% knew that HIV could not be transmitted through sharing food with an infected person. Similarly, 68.8% answered that HIV transmission cannot occur through mosquito bites. However, 45.8% of women said that HIV could be transmitted by witchcraft. Approximately 81.6% of participants were aware that HIV could be prevented by having a single monogamous sexual partner and protected sex. Approximately 70% of women were aware of antiretroviral therapy. More than 80% of the participants indicated that they had been screened overall, and 77.3% had been offered HIV screening services during antenatal visits.

Conclusions: Almost all women of childbearing age in Nyanga had heard of HIV, yet approximately half of them had poor knowledge of its modes of transmission. It is therefore imperative to conduct awareness campaigns in conjunction with HIV testing services to better understand the current prevalence of HIV infection in the region.

Background

Human immunodeficiency virus (HIV), which causes the acquired immunodeficiency syndrome (AIDS) pandemic, is still a public health concern. In 2020, 37.7 million people were living with HIV [1], and 1.5 million were newly infected. Two-thirds of these new infections are registered in Africa, the most infected continent, where 25.6 million people are living with HIV/AIDS. In Gabon, the last Demographic and Health Survey (DHS) reported infection in 63,000 people, and the prevalence was estimated at 4.1% [2]. According to the DHS, this prevalence seems to be the consequence of a lack of knowledge about the disease and the various routes of transmission and prevention.

Of the two HIVs, HIV-1 is the most endemic. The most well-known modes of transmission in the world are unprotected sexual intercourse, blood transfusion, and mother-to-child [3, 4]. HIV-1 consists of 4 established groups: Groups M, N, O and P [5]. Group M is responsible for a majority of infections around the world and is regularly described in Gabon under different circulating recombinant forms (CRFs) [6, 7]. HIV infection could be reduced if people used barrier methods such as condoms, having a single sexual partner or choosing abstinence. More specifically, if women have unprotected sexual intercourse with other partners beside their primary partner(s) during pregnancy or if their partners have other partners, they could be infected with HIV and have the chance to transmit the virus to their infants [8, 9]. Additionally, women treated early in their pregnancy could decrease mother-to-child (MTC) transmission of HIV [10]. As recommended by the World Health Organization (WHO) and the Joint United Nations Programme on HIV/AIDS [11], prevention of new HIV transmission should be considered an important action in the strategy of fighting HIV/AIDS [12]. HIV testing and immediate treatment could limit the dissemination of the pandemic [13, 14]. A key approach could be the use of rapid diagnostic tests (RDTs), which facilitate screening for HIV infection due to their many advantages [15]. Indeed, RDTs

have a very low price per test, can be stored at room temperature, do not need complex infrastructure, can be performed in very few steps, and the results can be read visually [16]. Due to the absence of a cure for HIV, antiretroviral therapy (ART) continues to expand in resource-limited countries as the primary care and prevention approach, as well as to contribute to the control of viral replication and decreased HIV transmission, particularly from mother to child. Thus, RDTs could provide an opportunity to achieve the UNAIDS' 95-95-95 goals by 2030: 95% of all people living with HIV will know their status, 95% of all people diagnosed will receive sustained antiretroviral therapy (ART), and 95% of all people receiving ART will have viral suppression [17].

Gabon, a west central african country, is divided into nine provinces, and Nyanga is located in the southwest of the country, home to 52,854 people. In Nyanga Province, HIV seroprevalence is 4.2%, one of the highest rates by province and relatively higher than the national seroprevalence [2]. This variance is hypothesized to be the result of a lack of information regarding HIV transmission and prevention. There are currently scarce epidemiological data from this region of Gabon concerning knowledge, attitudes and practices in the fight against HIV/AIDS.

We conducted a cross-sectional household survey to inform the “Improved Integrated Maternal and Neonatal Survival through Malaria and HIV/AIDS Prevention in Gabon Project.” Our goal was to assess the knowledge, attitudes and practices of women of childbearing age from Nyanga Province toward HIV, with particular emphasis on mother-to-child transmission of HIV. The selected approach was to assess some of the possible misconceptions concerning the modes of transmission of HIV among the study population. Thus, this study was organized to obtain a baseline for planned interventions to improve service delivery for the prevention of mother-to-child transmission of HIV that were undertaken by the nongovernmental organization (NGO) Medical Care Development International (MCDI) in Nyanga Province following the study.

Methods

Study sites and Population

This is a 13-day cross-sectional, descriptive, and analytical epidemiological study conducted in March 2018 among women residing in five of six departmental capitals of the Nyanga province, including Mayumba, Moabi, Mabanda, Moulengui-Binza, and Tchibanga (Figure 1), as described previously [18]. The most important natural resources of Nyanga are oil and gas. Additionally, this province is covered with rainforest and hosts Moukalaba-Doudou National Park with prized species such as elephants, small monkeys, gorillas and chimpanzees. The study population matched the following preselected inclusion criteria: 1) resided within the study site during the study period; 2) signed the consent form; and 3) were women of childbearing age, defined as being 15 to 49 years old. All women aged 15 to 49 years old in the households surveyed were included in the study, and information was collected by means of a questionnaire. The study was approved by the Gabonese National Ethics Committee (N° 001/PR/SG/CNE/2018).

Development of a Questionnaire

The data collection questionnaire was generated based on the Demographic and Health Survey (DHS) and various questionnaires from similar studies conducted previously [19-21]. The questionnaire can be accessed here (Additional file 1). Two sections consisted of sociodemographic characteristics of the study population across questions such as age, pregnancy status and professional status and knowledge about HIV transmission and prevention. Specific questions were asked to women of childbearing age regarding understanding and possible common misconceptions about HIV transmission (“by mosquito bite?”, “by food?”, and “by witchcraft?”) and mother-to-child transmission (“during pregnancy?”, “during delivery?”, and “during breastfeeding?”). Similarly, prevention knowledge was assessed with two other questions: HIV transmission can be prevented “by having a single sexual partner?” and “by using a condom during sexual activity?”. Correct answers from participants were considered to have basic knowledge of HIV.

Data collection

The approach applied in this study was the same as that jointly defined for the malaria study we published previously [18]. Before starting interviews, we selected all households with women of childbearing age. Then, a convenience sample of households was selected for inclusion in the study; all neighborhoods of the departmental capitals were represented. If no member from an identified household was willing to participate, the teams approached the next house on the list until reaching a household willing to participate in the study. The survey was conducted by ten teams of three persons deployed across the five cities. The sample size in each departmental capital was calculated based on probability proportional to size.

Data analysis

Statistical analysis was performed with Epi-info version 3.5.3 (2005, CDC, Atlanta, USA) and STATA version 14 (Stata Corp, College Station, USA). Age was expressed as the mean and standard deviation. The age variable was transformed into a categorical variable. The chi-squared test was used to compare categorical variables. Values of $p < 0.05$ were considered to indicate a statistically significant difference.

Results

Characteristics of the study population

A total of 631 women were interviewed for this study; 90% (570/631) were of childbearing age but not pregnant at the time of the survey, and 10% (61/631) were pregnant; the mean age of all participants was 28 years old. Information about occupation was collected for 620 women, most of whom were unemployed ($n=327$; 52.7%), employed ($n=173$; 27.9%) and students ($n= 120$; 19.4%). Table 1 summarizes the demographic characteristics of the study population.

HIV/AIDS-related knowledge on transmission and prevention

Results about knowledge indicated that 99.2% (621/626) women had heard of HIV/AIDS and 81.8% (508/621) knew of at least one screening site for HIV including hospitals and other health facilities ($n= 222$; 39.6%), and the Ambulatory Treatment Centre (CTA) where HIV-1 patients receive ART in Tchibanga ($n= 325$; 57.8%). Regarding HIV transmission, 83.6% of women knew that mother-to-child HIV transmission can occur during pregnancy; 85.4% (527/617) knew that it could occur during delivery; and 78% agreed that a mother could transmit the virus through breastfeeding. Additionally, more than 85.4% (531/614) of participants knew that HIV could not be transmitted through sharing food with an infected person. Similarly, 68.8% (422/613) answered that HIV transmission cannot occur through mosquito bites. On the topic of HIV prevention, 81.6% of participants indicated that HIV can be prevented by having a single sexual partner and protected sex. However, some misconceptions were highlighted among 45.8% (280/612) of women who said that HIV could be transmitted by witchcraft. All HIV/AIDS-related knowledge, transmission and prevention are summarized in Table 2.

Information and practices against HIV/AIDS

The results showed that more than 70.7% of women were informed about practices against HIV/AIDS and were aware that medicines exist that infected pregnant women can take to reduce the risk of MTCT (antiretroviral therapy). In addition, 91% said it is possible that a healthy-looking person could be infected (Table 3). Additionally, 84.5% answered that using a condom during sexual intercourse can prevent HIV infection (Table 2). More than 80% of the participants indicated that they had ever been screened for HIV, and 77.3% said that they had been offered HIV screening services during antenatal visits (Table 3). Regarding practices, 75% of women declared using good practices to prevent HIV/AIDS. In addition to being tested during antenatal visits (77.3%), they also planned to follow up to obtain the results of their

medical examinations (83.9%). We found that only 63.4% of women received advice from medical staff after receiving the results (Table 4).

Discussion

The current cross-sectional descriptive study was conducted to assess HIV/AIDS-related knowledge, attitudes and practices among 631 women aged 15 to 49 years old, with a particular emphasis on mother-to-child transmission of HIV. We found that most women from Nyanga Province have a good level of knowledge of mother-to-child transmission, prevention and practices against HIV/AIDS. Unfortunately, some participants revealed misconceptions about the AIDS pandemic.

According to the last Gabonese Demographic and Health Survey (GDHS) [2], 99% of women in the general population had heard of HIV/AIDS. This rate was not formally estimated in Nyanga Province during the same survey. In our study, 99.2% of women were informed about HIV/AIDS, suggesting that well-organized awareness campaigns have contributed to the dissemination of proper comprehensive HIV/AIDS knowledge in this province since the last GDHS. Additionally, it would seem that information from different communication channels was delivered to the population. During interviews, participants confirmed obtaining information on HIV/AIDS from medical staff, as well as from media outlets such as television, radio, and social media. The fact that 81.8% of participants knew of at least one screening site for HIV could positively impact their families and society as a whole. We believe that a consequence of increased knowledge could probably be the decrease in HIV seroprevalence in this province; the next GDHS will confirm this hypothesis.

Our results demonstrate a good level of knowledge on the prevention of mother-to-child transmission of HIV (PMTCT) among participants during pregnancy, delivery and while breastfeeding. These findings suggest that awareness campaigns in Nyanga Province have also focused on these modes of transmission. The percentage of knowledge of transmission during breastfeeding is similar (76%) to that found during the 2012 GDHS [2]. The early description of the transmission of HIV from mother-to-child after the onset of the epidemic [22] and the joint action between the provincial multisectoral committee against AIDS and the PMTCT program in Gabon could explain the high rates of good knowledge about MTC transmission. In addition to knowledge of the modes of transmission, the use of antiretroviral therapy (ART) is a solution to prevent MTCT by reducing the viral load and improving the quality of life of HIV-infected persons [23, 24]. Furthermore, ART became available in 2001 in the country and was made available free of cost starting in 2011. In our study, 70.7% of women were aware of ART, a result likely related to several good practices against HIV/AIDS collected from participants, such as HIV counseling and testing (HCT). Nevertheless, higher numbers of respondents declared being tested during antenatal visits (81.7%) and going back to collect the results of their medical examinations (83.9%). Other studies corroborate our findings and have indicated that persons who were aware of ART were more likely to use HCT services [19, 25, 26].

Counseling is related to the prevention of HIV transmission [27]; by answering that HIV could be contracted from a healthy-looking person, 91% of women of childbearing age confirmed that they could identify HIV's real modes of transmission [28–30]. Kwaganzo *et al.* published previously that “HIV counseling and testing services provide information on HIV transmission prevention” [27].

Despite this knowledge, misconceptions were also identified during the study. Through our results, we confirm that HIV/AIDS knowledge can be affected by the overall sociocultural context where for example, some modes of transmission can be known, while more culturally sensitive modes of transmission may not be as widely recognized, which may contribute to false perceptions of risk [31]. During our investigation, some beliefs held about HIV transmission modes included witchcraft and mosquito bites. A large proportion of participants, 45.8% (280/612), claimed that HIV could be transmitted by witchcraft. This idea is widespread in several Sub-Saharan African countries

and constitutes a major cultural feature [32]. Previous studies conducted in Ghana, Congo, Zambia, Zimbabwe, etc. concluded that witchcraft played a key role in the social fabric of most African societies [32, 33] and may influence prevention behaviors. Other beliefs of HIV transmission were noted among 23.8% (146/613) of women who answered that HIV transmission occurred through mosquito bites. This misconception was highlighted in other studies: 13.6% among dental patients at a Nigerian teaching hospital, 59.5% among university students in Xinjiang, China and 48.2% among secondary school adolescents in Nigeria [3, 23, 34]. Additionally, the difference in educational status among participants and the fact that so many of them live in rural areas could have influenced their knowledge. Finally, as Gabon is a hyperendemic zone for malaria [35], some participants could believe that HIV transmission occurs by mosquito bites. The risk is the persistence of such beliefs that may reduce or distort prevention behaviors [32], and beliefs can be resistant to change and entrenched in people's behavior, with the consequence being that they are taught by one generation to the next [36].

Limitations of the study

This survey encountered limitations, mainly due to information bias. A particular methodological limitation may have been the use of a closed survey instrument, which asked participants to agree or disagree with an affirmation about HIV. The response options of Yes, No, and I Don't Know may have led some respondents to answer "Yes" to concepts that they lacked previous familiarity with, in order to seem knowledgeable or to please the investigator. The affirmations used in the survey were based on common beliefs about HIV found in the literature. It might have been more informative from a socioanthropological perspective to use an open-ended survey or other qualitative research methods to take stock of novel beliefs on sources of HIV infection that are unique to Nyanga Province; however, doing so would have rendered quantitative data analysis impossible. Another limitation was the lack of answers to all of the survey questions. A large amount of data were missing, leading to variations in the numbers of women for each parameter studied (transmission by mosquito bite, by food, by witchcraft, awareness of the existence of ART, having ever been tested for HIV/AIDS, etc.). The lack of experience of our investigators during interviews, despite initial training on the survey methodology, may have contributed to this difference.

Conclusion

This study reported that women of childbearing age (15-49 years) from Nyanga Province in Gabon have a good level of knowledge, including of mother-to-child transmission, prevention and attitudes against HIV/AIDS. The misconceptions highlighted raise a challenge to achieve the UNAIDS' 95-95-95 goals set for 2030 in this relatively remote part of Gabon. The "Improved Integrated Maternal and Neonatal Survival through Malaria and HIV/AIDS Prevention Project" in Gabon, which was funded by Noble Energy to begin interventions after this study, was designed with these findings in mind to help improve correct knowledge of HIV/AIDS and limit the spread of the virus, particularly through PMTCT.

Abbreviations

HIV/AIDS: Human immunodeficiency virus/Acquired immunodeficiency syndrome; PMTCT: Prevention of mother-to-child HIV transmission; UNAIDS: United Nations programme on HIV/AIDS; WHO: World Health Organization; CRFs: Circulating recombinant forms; ART: antiretroviral therapy; ATC: Ambulatory Treatment Centre; HCT: HIV counseling and testing; NGO: Nongovernment organization; MCDI: Medical Care Development International; GDHS: Gabonese Demographic and Health Survey; CIRMF: Centre International de Recherches Médicales de Franceville.

Declarations

Ethics approval and consent to participate

The study was approved by the Gabonese National Ethics Committee (N°001/PR/SG/CNE/2018). Written informed consent was obtained from a parent of participants under 16 years old.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

We thank sincerely all participants for this study and all investigators who performed interviews. We would like to acknowledge the MCDI (Medical Care Development International, European and Developing Countries Clinical Trials Partnership) and the CIRMF (*Centre International de Recherches Médicales de Franceville*) for this partnership. Finally, we thank Dr Thierry Tsoumbou for providing the Nyanga map and John Pollock for critiquing this manuscript.

Authors' contributions

All authors read and approved the final manuscript. AMO, JBLD, JNDC and JB participated in the conception and design of the study. AMO, MNS and JDEO participated in the interpretation of the results. AMO drafted and coordinated manuscript writing and wrote the first draft. AMO, JBLD, RKIL, MNS, NPTT, PNSOE contributed to the acquisition of data in the field.

Funding

This work was supported by Noble Energy Gabon Ltd. and Woodside Energy (Gabon) Pty Ltd., through the Improved Integrated Maternal and Neonatal Survival Malaria and HIV/AIDS Prevention project, financed with the Gabonese Ministry of Petroleum and Hydrocarbon's Local Community Development Fund, and implemented by Medical Care Development International from 2017-2019.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

1. UNAIDS, *GLOBAL AIDS MONITORING*. Available from: <https://www.unaids.org/en/resources/documents/2020/unaids-data> 2020.
2. EDSG, *Enquête Démographique et de Santé du Gabon 2012: Rapport de synthèse*. 2012, DGS et ICF International Calverton, MD.
3. Opeodu, O.I. and T.J. Ogunrinde, *MODE OF TRANSMISSION OF HIV/AIDS: PERCEPTION OF DENTAL PATIENTS IN A NIGERIAN TEACHING HOSPITAL*. J West Afr Coll Surg, 2015. **5**(1): p. 1–19.
4. Nelson, C.S., G.G.A. Fouda, and S.R. Permar, *Pediatric HIV-1 Acquisition and Lifelong Consequences of Infant Infection*. Curr Immunol Rev, 2019. **15**(1): p. 131–138.
5. Plantier, J.C., et al., *A new human immunodeficiency virus derived from gorillas*. Nat Med, 2009. **15**(8): p. 871–2.

6. Liégeois, F., et al., *Virological failure rates and HIV-1 drug resistance patterns in patients on first-line antiretroviral treatment in semirural and rural Gabon*. J Int AIDS Soc, 2012. **15**(2): p. 17985.
7. Ndjoyi-Mbiguino, A., et al., *Performance of rapid HIV-1/HIV-2 INSTI on plasma and capillary blood for serological diagnosis of non B subtypes and circulating recombinant forms of HIV-1 in Gabon*. Med Mal Infect, 2015. **45**(8): p. 339–40.
8. Drake, A.L., et al., *Incident HIV during pregnancy and postpartum and risk of mother-to-child HIV transmission: a systematic review and meta-analysis*. PLoS Med, 2014. **11**(2): p. e1001608.
9. Mushamiri, I., et al., *Optimizing PMTCT efforts by repeat HIV testing during antenatal and perinatal care in resource-limited settings: A longitudinal assessment of HIV seroconversion*. PLoS One, 2020. **15**(5): p. e0233396.
10. Blanche, S., et al., *Growth in HIV-1-exposed but uninfected infants treated with lopinavir-ritonavir versus lamivudine: a secondary analysis of the ANRS 12174 trial*. Lancet HIV, 2019. **6**(5): p. e307-e314.
11. UNAIDS, *The gap report*. Geneva: UNAIDS; 2014.
https://www.unaids.org/en/resources/documents/2014/20140716_UNAIDS_gap_report, J.U.N.P.o. HIV/AIDS, Editor. 2014.
12. WHO, *WHO: Global health sector strategy on VIH, 2016-2021*. available from: <https://www.who.int/hiv/strategy2016-2021/ghss-hiv/en/>. 2016, World Health Organization.
13. Granich, R.M., et al., *Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model*. Lancet, 2009. **373**(9657): p. 48–57.
14. Havlir, D., et al., *What do the Universal Test and Treat trials tell us about the path to HIV epidemic control?* J Int AIDS Soc, 2020. **23**(2): p. e25455.
15. Mourez, T., et al., *HIV rapid screening tests and self-tests: Be aware of differences in performance and cautious of vendors*. EBioMedicine, 2018. **37**: p. 382–391.
16. Kosack, C.S., A.L. Page, and P.R. Klatser, *A guide to aid the selection of diagnostic tests*. Bull World Health Organ, 2017. **95**(9): p. 639–645.
17. UNAIDS. *Understanding Fast-Track: Accelerating Action to End the AIDS Epidemic by 2030*. 2015 [cited 2021 02/25/2021]; Available from:
https://www.unaids.org/sites/default/files/media_asset/201506_JC2743_Understanding_FastTrack_en.pdf
18. Imboumy-Limoukou, R.K., et al., *Malaria in children and women of childbearing age: infection prevalence, knowledge and use of malaria prevention tools in the province of Nyanga, Gabon*. Malar J, 2020. **19**(1): p. 387.
19. Chimoyi, L., et al., *HIV-related knowledge, perceptions, attitudes, and utilisation of HIV counselling and testing: a venue-based intercept commuter population survey in the inner city of Johannesburg, South Africa*. Glob Health Action, 2015. **8**: p. 26950.
20. Mall, S., et al., *Changing patterns in HIV/AIDS stigma and uptake of voluntary counselling and testing services: the results of two consecutive community surveys conducted in the Western Cape, South Africa*. AIDS Care, 2013. **25**(2): p. 194–201.
21. Shamu, S., et al., *Knowledge, attitudes and practices of young adults towards HIV prevention: an analysis of baseline data from a community-based HIV prevention intervention study in two high HIV burden districts, South Africa*. BMC Public Health, 2020. **20**(1): p. 1249.
22. Oleske, J.M. and A.B. Minnefor, *Acquired immune deficiency syndrome in children*. Pediatr Infect Dis, 1983. **2**(2): p. 85–6.
23. Agu, I.C., et al., *Misconceptions about transmission, symptoms and prevention of HIV/AIDS among adolescents in Ebonyi state, South-east Nigeria*. BMC Res Notes, 2020. **13**(1): p. 244.

24. Mintsa-Ndong, A., et al., *Mother-to-child HIV-transmission prevention programs in a sub-Saharan African setting: The Gabonese experience*. Int J STD AIDS, 2018. **29**(3): p. 221–226.
25. Venkatesh, K.K., et al., *Who gets tested for HIV in a South African urban township? Implications for test and treat and gender-based prevention interventions*. J Acquir Immune Defic Syndr, 2011. **56**(2): p. 151–65.
26. Tran, B.X., et al., *Depression and Quality of Life among Patients Living with HIV/AIDS in the Era of Universal Treatment Access in Vietnam*. Int J Environ Res Public Health, 2018. **15**(12).
27. Kwagonza, L., et al., *Comprehensive knowledge of HIV prevention among fishing communities of Lake Kyoga, Uganda, 2013*. BMC Public Health, 2020. **20**(1): p. 29.
28. Pisani, E., et al., *Back to basics in HIV prevention: focus on exposure*. Bmj, 2003. **326**(7403): p. 1384–7.
29. Gouws, E., et al., *Short term estimates of adult HIV incidence by mode of transmission: Kenya and Thailand as examples*. Sex Transm Infect, 2006. **82 Suppl 3**(Suppl 3): p. iii51-55.
30. Yavuz, B., et al., *Pharmaceutical Approaches to HIV Treatment and Prevention*. Adv Ther (Weinh), 2018. **1**(6).
31. Mumtaz, G.R., et al., *Characterising HIV/AIDS knowledge and attitudes in the Middle East and North Africa: Systematic review and data synthesis*. Glob Public Health, 2020. **15**(2): p. 275–298.
32. Tenkorang, E.Y., et al., *Superstition, witchcraft and HIV prevention in sub-Saharan Africa: the case of Ghana*. Cult Health Sex, 2011. **13**(9): p. 1001–14.
33. Pearson, S. and P. Makadzange, *Help-seeking behaviour for sexual-health concerns: a qualitative study of men in Zimbabwe*. Cult Health Sex, 2008. **10**(4): p. 361–76.
34. Bamise, O.F., C.T. Bamise, and M.A. Adedigba, *Knowledge of HIV/AIDS among secondary school adolescents in Osun state, Nigeria*. Niger J Clin Pract, 2011. **14**(3): p. 338–44.
35. Boumbanda Koyo, C.S., et al., *High Circulation of Malaria and Low Prevalence of Bacteremia in Febrile and Afebrile Children in Northeastern Gabon*. Am J Trop Med Hyg, 2020. **102**(1): p. 121–129.
36. Amuyunzu-Nyamongo, M., et al., *Putting on a brave face: the experiences of women living with HIV and AIDS in informal settlements of Nairobi, Kenya*. AIDS Care, 2007. **19 Suppl 1**: p. S25-34.

Tables

Table 1: Sociodemographic characteristics of the study population

<i>Parameters</i>	<i>Cities (n/N (%))</i>					
	<i>Mabanda</i>	<i>Mayumba</i>	<i>Moabi</i>	<i>Moulengui-Binza</i>	<i>Tchibanga</i>	Total
<i>Categories</i>						
Women of childbearing age not currently pregnant	100% (49/49)	85% (63/74)	87% (65/75)	92%(34/37)	91% (359/396)	570
Pregant women	0%(0/49)	15% (11/74)	13% (10/75)	8%(3/37)	9%(37/396)	61
Total	49	74	75	37	396	631
Women occupation (N=620)						
Student	9.5% (4/42)	1.4% (1/72)	36.0% (27/75)	16.2% (6/37)	20.8% (82/394)	120
Employed	38.1% (16/42)	30.5% (22/72)	33.3% (25/75)	48.7% (18/37)	23.4% (92/394)	173
Unemployed	52.4% (22/42)	68.1% (49/72)	30.7% (23/75)	35.1% (13/37)	55.8% (220/394)	327

Table 2: Women's knowledge of HIV transmission and prevention

	<i>HIV Prevention (% (n))</i>			<i>HIV transmission (% (n))</i>				
	By mosquito bite	By food	By witchcraft	Mother-to-child during pregnancy	Mother-to-child during breastfeeding	Mother-to-child during delivery	Having a single sexual partner	Using condoms during sexual intercourse
No	69% (422)	86.5% (531)	47.2% (289)	8.3% (51)	11.0% (68)	5.2% (32)	12.8% (77)	11.5% (67)
Unknown	7,3% (45)	2.1% (13)	7.0% (45)	8.1% (50)	12.5% (77)	9.4% (58)	5.6% (34)	4.0% (23)
Yes	23.8% (146)	11.40% (70)	45.8% (280)	83.6% (517)	76.5% (471)	85.4% (527)	81.6% (491)	84.5% (492)
Total	613	614	612	618	616	617	602	582

Table 3: Women's knowledge level about HIV in Nyanga Province

	Existence of ART* for PMTCT	Healthy-looking person	Children infected by mothers	Existence of measures against HIV/AIDS	Agreed to be tested for HIV/AIDS	Invitation to be tested for HIV/AIDS during the last pregnancy
Yes	70.7% (437)	90.9% (558)	54.0% (275)	61.9% (312)	82.5% (414)	77.3% (388)
No	13.6% (84)	7.0% (43)	38.3% (195)	30.2% (152)	16.1% (81)	17.7% (89)
Unknown	15.7% (97)	2.1% (13)	7.7% (39)	7.9% (40)	1.39% (7)	5% (25)
Total	618	614	509	504	502	502

*ART: Antiretroviral Therapy

Table 4: Selected HIV testing practices among women in Nyanga Province

	HIV testing	Results collected by the patient	Advice from medical staff after the results
Yes	81.7% (370)	83.9% (374)	63.4% (279)
No	17.9% (81)	15.5% (69)	35.5% (156)
Unknown	0.4% (2)	0.6% (3)	1.1% (5)
Total	453	446	440

Figures

Figure 1

Map of Gabon's Nyanga province. Shown colors are the six departments with their capitals: Tchibanga (departmental and provincial capital), Moabi, Moulengui-Binza, Mayumba, Mabanda and Ndindi (excluded from the study due to access issues).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1questionnaire.docx](#)