

# Angelica Sinensis Polysaccharides Ameliorate 5-Fluorouracil-Induced Bone Marrow Stromal Cell Proliferation Inhibition Via Regulating Wnt/ $\beta$ -Catenin Signaling

**Hanxianzhi Xiao**

Chongqing Medical University

**Rongjia Qi**

Chongqing Medical University

**Ziling Wang**

Chongqing Medical University

**Minghe Xiao**

Chongqing Medical University

**Yue Xiang**

Chongqing Medical University

**YaPing Wang**

Chongqing Medical University

**Lu Wang** (✉ [luwang@cqmu.edu.cn](mailto:luwang@cqmu.edu.cn))

Chongqing Medical University <https://orcid.org/0000-0002-5056-1047>

---

## Research

**Keywords:** Angelica Sinensis Polysaccharides, 5-fluorouracil, Wnt/ $\beta$ -catenin signaling pathway, oxidative stress, cell proliferation, FoxO1

**Posted Date:** October 29th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-97491/v1>

**License:** (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** Chemotherapy-induced bone marrow hematopoietic microenvironment oxidative damage is closely related to myelosuppression. Angelica Sinensis Polysaccharides (ASP) are major effective ingredients of traditional Chinese medicine Angelica with multi-target anti-oxidative stress features. In this study, we investigated the potential benefits and mechanism of action of ASP on chemotherapy-induced bone marrow stromal cell (BMSC) damage.

**Methods:** The human bone marrow stromal cell line HS-5 cells were divided into control group, 5-FU group, 5-FU + ASP group, and 5-FU+ LiCl group to investigate the mechanism of ASP to alleviate 5-FU-induced BMSC proliferation inhibition.

**Results:** The results showed that 5-FU inhibits the growth of HS-5 cells in a time and dose-dependent manner; however, ASP partially counteracted 5-FU-induced decrease in cell viability, whereas Wnt signaling inhibitor Dkk1 antagonized the effect of ASP on HS-5 cells. ASP reversed the decrease in cytoplasmic total  $\beta$ -catenin, p-GSK-3 $\beta$ , and CyclinD1 following 5-FU treatment, and modulated nuclear expression of  $\beta$ -catenin, Lef-1, and C-myc proteins. Furthermore, ASP also enhanced the antioxidant capacity of cells and reduced 5-FU caused oxidative stress, attenuated FoxO1 expression, thus weakened its downstream apoptosis-related proteins and G<sub>0</sub>/G<sub>1</sub> checkpoint-associated p27<sup>Kip1</sup> expression to alleviate 5-FU-induced apoptosis promote cell cycle progression.

**Conclusion:** The protective role of ASP in BMSCs proliferation for the chemotherapy may be related to its activating Wnt/ $\beta$ -catenin signaling and keeping homeostasis between  $\beta$ -catenin and FoxO1 under oxidative stress. The study provides a potential therapeutic strategy for alleviating chemotherapeutic damage on BMSCs.

## Background

Chemotherapy is a common therapeutic modality for malignant tumors. However, this therapy is not tumor-target and is often accompanied by several side effect[1, 2]. The hematopoietic system is highly sensitive to chemotherapy, even conventional doses of chemotherapeutic drugs may cause myelosuppression, leading to hematopoietic dysfunction, hematopoietic reconstitution disorders, and other adverse reactions[3, 4]. The mechanisms of myelosuppression can be various, including direct cytotoxicity to marrow cells, inhibition of bone marrow precursor or progenitor cell proliferation, or interference with hematopoietic growth factor and receptor signaling subsequently affecting the downstream differentiation processes. Chemotherapy-induced myelosuppression not only damage the proliferating hematopoietic progenitors, but also affect the stromal cells of hematopoietic microenvironment, and this may be the reason for chronic hematopoietic dysfunction[5–10]. As chemotherapy disrupts the steady-state function of hematopoietic and stromal cell, disruptions over time may cause severe bone marrow toxicity and the failure of cancer treatment. 5-FU, widely used in high-proliferative, tissue-derived cancers, particularly for colorectal cancer and breast cancer, exerts its anti-

cancer effects through inhibition of thymidylate synthase (TS) and incorporation of its metabolites into RNA and DNA[10–12]. It was reported that the mechanism of stromal cells proliferation inhibition and apoptosis after 5-FU treatment is oxidative damage[13, 14]. Our previous findings have confirmed that following oxidative damage of BMSCs 5-FU may alter bioactive substance and cause stress-induced premature senescence (SIPS) of hematopoietic cells[15]. However, the specific underlying mechanism of 5-FU-induced BMSC proliferation inhibition remains unclear. Therefore, to explore its related mechanisms to reduce the side effects of chemotherapy drugs and to screen protection drugs during chemotherapy is of clinical guidance significance.

Wnt/ $\beta$ -catenin is an evolutionarily highly conserved signaling pathway that plays a key role in development and is involved in cell proliferation, differentiation, apoptosis, and localization control[16, 17]. Particularly, Wnt pathway involves various signal feedback that maintain the processes of stem cell proliferation, differentiation and self-renewal[18, 19]. The properties of stem cells are conferred by the interaction of stem cells with their local microenvironment. Recent studies have evidenced that Wnt/ $\beta$ -catenin signaling pathway is closely related to hematopoietic microenvironment affecting hematopoietic microenvironment function extensively, participate in BMSC proliferation, alleviate oxidative stress, and regulate hematopoietic stem cell self-renewal through stroma-dependent manner[20–22]. It is increasingly realized that the microenvironment keeps the threshold of Wnt signaling in stem cells at a physiological range. In the current work it was clarified herein the roles of Wnt signaling in chemotherapy-induced stromal suppression and the ameliorative effects of ASP.

The Forkhead transcription factors family including FoxO1 (or Fkhr), FoxO3a (or Fkhr1), FoxO4 (or Afx), and FoxO6 are critically involved in the regulation of apoptosis, proliferation, and the control of oxidative stress[23]. Stress conditions such as high levels of ROS induce FoxO nuclear import and trigger the shifting of  $\beta$ -catenin from TCF/LEF to FoxO-mediated transcription[24, 25]. In hematopoietic system, activation of FoxO factor is sufficient to activate a variety of proapoptotic genes and to trigger apoptosis. Meanwhile, overexpression of FoxO factors cause a strong inhibition of cell proliferation[26, 27]. As playing a critical role in proliferation and apoptosis, it has been aware of that FoxO factors are closely related to chemotherapy-induced cell damage, nevertheless studies are needed to clarify the relationship of FoxO factors and Wnt signaling in myelosuppression.

Angelica of Chinese herb is commonly used medicine to enrich blood, promote blood circulation and treat menstrual disorders[28, 29]. Angelica Sinensis Polysaccharides (ASP) are major effective ingredients of Angelica, with significant bioactivities including anti-oxidation[30, 31], anti-tumor[32, 33], promoting hematopoiesis[34–36], and delaying senescence[37, 38] effects. ASP shows antioxidant activity by suppressing the production of ROS and regulating several chemical substances associated with oxidative stress[39, 40]. Our previous work showed marked antioxidative role of ASP in BMSCs from 5-FU injury in vitro, thus protected hematopoietic cells against SIPS via alleviating oxidative stress, preventing oxidative DNA damage, promoting hematopoietic stimulating factors originated from BMSCs, and enhancing intercellular communication between stromal cells and hematopoietic cells[15]. On this basis, we demonstrated herein that ASP alleviated 5-FU-induced stromal cell proliferation inhibition, apoptosis, and

oxidative stress damage, and the underlying mechanism may be related to ASP activating Wnt/ $\beta$ -catenin signaling and keeping homeostasis between  $\beta$ -catenin and FoxO1 under oxidative stress.

## Materials And Methods

### Reagents

5-fluorouracil was purchased from Sigma-Aldrich Co. (St. Louis, the USA). ASP (Purity  $\geq 95\%$ ) were purchased from Ci Yuan Biotechnology Co. Ltd. (Shanxi, China). LiCl (purity  $>95\%$ ) was purchased from Damao Chemical Reagent Factory (Tianjin, China). Fetal bovine serum (FBS) was purchased from MRC Company (Australia). Dulbecco's modified Eagle medium high-glucose(H-DMEM) was purchased from Gibco Co. (NY, USA). Cell Counting Kit-8 was purchased from Dojindo Laboratories (Japan). EdU Cell Proliferation Assay Kit was purchased from RiboBio Co. Ltd.(Guangzhou, China). $\beta$ -catenin, GSK-3 $\beta$ , p-GSK-3 $\beta$ , Lef-1, Cyclin D1, C-myc, FoxO1, p-FoxO1, P27 Kip1, Bim, Bax, Bcl-2, and caspase-3 antibodies were purchased from Cell Signaling Technology(Danvers, the USA). Dkk1 was purchased from R&D Systems (the USA). Reactive Oxygen Species Assay Kit and Senescence  $\beta$ -Galactosidase Staining Kit were purchased from the Beyotime Institute of Biotechnology (Shanghai, China). Superoxide Dismutase (SOD) assay kit, Malondialdehyde (MDA) assay kit, and Catalase (CAT) assay kit were purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China).

### Cell Culture and Groups of Experiment

Human bone marrow stromal cell line HS-5 was cultured in H-DMEM containing with 10% fetal bovine serum and 100 U/mL penicillin and 100  $\mu$ g/mL streptomycin. Cells were cultured in a humidified atmosphere with 5% CO<sub>2</sub> at 37°C. Cells were divided into control group, 5-FU group, 5-FU + ASP group, and 5-FU+ LiCl group. Control group was routinely cultured; 5-FU group was treated with 5-FU on the concentration of 25 $\mu$ g/mL; 5-FU+ ASP group was pretreated with ASP on the concentration of 100  $\mu$ g/mL, and 25  $\mu$ g/mL 5-FU was added after 6 hours; 5-FU+ LiCl group was pretreated with LiCl on the concentration of 10 mmol/L, and 25  $\mu$ g/mL 5-FU was added after 6 hours, each group was cultured for 48h.

### CCK-8 Cell Viability Assay

Cell viability assay and the screening of drug concentration were performed using the Cell Counting Kit-8. Cells were plated in 96-well plates at a density of  $5 \times 10^3$  cells per well. The optical density(OD) value at 450 nm was measured using a microplate reader(Massachusetts, *the USA*). The cell viability of HS-5 was calculated according to the formula: Cell viability= [(OD experimental group - OD blank group)/ (OD control group- OD blank group)] \*100%. Inhibition rate = [(OD control group- OD experimental group)/ (OD control group- OD blank group)] \*100%

### EdU Proliferation Assay

The HS-5 cells were seeded in 96-well plates at a density of  $5 \times 10^3$  cells per well and treated as described in groups of experiment. After 48 h treatment, cells were exposed to 10  $\mu\text{mol/L}$  EdU solution for 24 h. Cells were washed and fixed in 4% paraformaldehyde at room temperature for 30 min. After washing, cells were permeabilized in PBS containing 0.5% Triton X-100 for 20 min. Then, cells were washed and incubated with 1X Apollo<sup>®</sup> reaction cocktail for 30 min. Subsequently, cells were stained with Hoechst33342 for 30 min and observed under a fluorescence microscope (Olympus, Japan). Counting 200 cells at random, the proliferation rate of HS-5 cells was defined as the ratio of EdU-positive cells (green cells) to Hoechst33342-positive cells (blue cells).

### **Flow Cytometry Analysis**

For cell apoptosis assay, the HS-5 cells were cultured then treated as described in groups of experiment. After 48h treatment, cells were harvested and centrifuged at 1000r/min for 5min. Subsequently, cells were resuspended with 500 $\mu$  L PBS solution for each tube. Cell apoptosis was detected by the flow cytometry. For cell cycle assay, the HS-5 cells were cultured then treated as described in groups of experiment. After 48h treatment, cells were harvested and fixed with pre-cooled 75% ethanol at 4°C for at least 5h. After centrifugation, cells were incubated with propidium iodide (PI) and RNase A at 37°C for 30 min in dark. Cell cycle was detected by the flow cytometry. The apoptosis and cell cycle were analyzed on a FAC-Scan laser flow cytometry (BD Biosciences, New Jersey, the USA). The data were processed by Cell Quest software (BD Biosciences, New Jersey, the USA).

### **Immunofluorescence Staining**

Sterile glass slides were put into 24-well plates, the HS-5 cells were cultured at a density of  $5 \times 10^4$  cells per well in 24-well plates then treated as described in groups of experiment. After 48h treatment, cells were fixed with 4% paraformaldehyde for 30 min at room temperature. After washing, cells were permeabilized in PBS containing 0.5% Triton X-100 for 20 min then blocked with 10% goat serum for 1h. Subsequently, cells were incubated with monoclonal antibody  $\beta$ -catenin (1:150) overnight at 4°C. After being washed thrice with PBS solution, cells were incubated with Cy3-labeled goat-anti-rabbit immunofluorescent secondary antibody (1:300) at 37°C for 2h in the dark. The nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI) for the last 5 min. The images were observed and acquired under the fluorescence microscope.

### **Immunoblot Assay**

The HS-5 cells were treated as described in groups of experiment. After 48h treatment, cells were incubated with PIPA lysis buffer containing 1% protease inhibitor and phosphatase inhibitor for 30 min on ice and proteins were isolated after centrifugation. The concentrations of proteins were detected by the BCA Protein Assay Kit (Beyotime, China). The protein samples were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to polyvinylidene fluoride (PVDF) membranes (Millipore, the USA). The membranes were blocked with 5% skim milk for 1h at room temperature and subsequently incubated overnight at 4°C with  $\beta$ -catenin, Cyclin D1, p-GSK-3 $\beta$ , GSK-3 $\beta$ ,

Lef-1, C-myc, FoxO1, p-FoxO1, p27<sup>Kip1</sup>, Bim, Bcl-2, Bax and caspase-3 primary antibodies (1:1000). After washing three times with Tris-Buffered Saline and Tween-20 (TBST), the membranes were incubated with secondary antibodies for 1 h at room temperature. The enhanced chemiluminescence (ECL) kit (Millipore, the USA) was used for color development and Image Lab 5.2.1 software was used for semi-quantitative analysis. The relative expression levels of the target proteins were determined by the ratio of the target protein gray value to internal reference protein gray value.

### **Oxidation-Associated Biological Indicators Assay**

For the detection of intracellular ROS, the HS-5 cells were seeded in 6-well plates at a density of  $2 \times 10^5$  cells per well then treated as described in groups of experiment. After 48 h treatment, the cells were washed thrice by serum-free medium and then incubated with 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA) at 37°C for 20-30 min in the dark. The content of intracellular ROS was observed and acquired under the fluorescence microscope. The average optical density per unit area was analyzed using ImageJ software. For the detection of MDA content and SOD, CAT activity, the HS-5 cells were cultured and treated as described in groups of experiment. After 48 h treatment, cells were harvested, lysed and centrifuged to collect the supernatant. MDA, SOD, and CAT were measured by the corresponding assay kits according to the manufacturer's instruction.

### **Statistical analysis**

For all assays, the experiments were performed at least three times. All the results were analyzed by One-way analysis of variance (ANOVA) with SPSS 20.0 statistical software. All the data were expressed as mean  $\pm$  standard deviation (SD).  $P < 0.05$  was considered statistical significance.

## **Results**

### **5-FU Inhibits the Growth of HS-5 Cells by down-regulating Wnt/ $\beta$ -catenin Signaling Pathway**

To assess the effect of 5-FU on proliferation, HS-5 cells were treated with 5-FU at different concentrations for 72 hours. As shown in Fig.1A, with the increase of 5-FU concentration and the extension of treatment time, the cell growth was significantly inhibited, suggesting that 5-FU has a dose-dependent and time-dependent inhibiting effect on HS-5 cells proliferation. HS-5 cells were incubated with 5-FU at concentration of 25  $\mu$ g/mL for 48 h, half of the cells were suppressed. Thus, 25  $\mu$ g/mL 5-FU treated cells for 48 h was selected for the subsequent experiments. Interestingly, in the current study we found the relationship between 5-FU inhibition of HS-5 cell proliferation and Wnt/ $\beta$ -catenin signaling pathway. Western blot assay demonstrated that 5-FU downregulated the cytoplasmic levels of p-GSK-3 $\beta$ , total  $\beta$ -catenin and Cyclin D1 in HS-5 cells, followed by nuclear protein expression of  $\beta$ -catenin, Lef-1, and C-myc downregulation (Fig.1B). These results hinted that 5-FU promoted the ubiquitination degradation of  $\beta$ -catenin mediated by GSK-3 $\beta$ -complex, inhibits the nuclear translocation of  $\beta$ -catenin and downregulate the downstream target genes. We hypothesize that the growth inhibition of HS-5 cells caused by 5-FU may be related to the down-regulation of Wnt/ $\beta$ -catenin signaling.

Hence, the activator and antagonist of Wnt/ $\beta$ -catenin signaling pathway were used to illustrate the question above. *In vitro* culture, treated with different concentration of LiCl, which could activate the Wnt/ $\beta$ -catenin signaling pathway, under light microscope, it was shown that the cells in 5, 10, 20mmol/L LiCl groups increased the number, however, the cellularity in 40 mmol/L group significantly dropped concomitant with smaller and loosely dispersed shape (Fig.1C). Also, cell viability was tested by CCK-8 assay. As shown in Fig.1D, cells treated with 5 mmol/L ~ 20 mmol/L LiCl for 48h showed different degrees of proliferation, among which the proliferation rate of the cells peaked in 10 mmol/L LiCl group and was increased to 150% compared with the control group. However, LiCl at a high concentration of 40 mmol/L was presented cytotoxic to HS-5 cells even within 24h. Therefore, the pretreatment with 10mmol/L LiCl was utilized for the subsequent experiment as a positive control. Furthermore, Dkk1, an antagonist for Wnt/ $\beta$ -catenin signaling, was used to get more evidence for Wnt/ $\beta$ -catenin signaling pathway on the cell viability. 50 ng/mL Dkk1 treated for 48h, the cells were dramatically inhibited compared with the control group revealed by the results of CCK-8 (Fig.1E). All the results above indicate that the effect of 5-FU on inhibition of HS-5 cell growth correlates with the suppression of Wnt signaling pathway.

### **Angelica Sinensis Polysaccharides Antagonize Growth Inhibition of 5-FU-treated HS-5 Cells through Wnt/ $\beta$ -catenin Signaling Pathway**

EdU is a thymidine analog that replaces thymidine in the DNA being synthesized during DNA replication. The conjugation reaction of EdU with fluorescent dyes can detect the cell proliferation. As shown in Fig.2A, B, the proportion of proliferating cells in 5-FU group was significantly lower than that of the control group; after pretreatment with ASP and LiCl, the proportion of EdU proliferating cells increased markedly compared with 5-FU group. The results of CCK-8 showed an obvious reduction after a 48h incubation with 5-FU compared with untreated control cells; however, ASP pretreatment partially reversed with ASP, the reduction of cellular viability induced by 5-FU. Moreover, ASP-induced increase in the viability was weakened by Dkk1 (Fig. 2C). The results above suggested that ASP may mitigate the proliferation inhibition of HS-5 cells after 5-FU treatment via regulating Wnt signaling.  $\beta$ -catenin protein is a key effector protein of Wnt/ $\beta$ -catenin signaling pathway, as shown in Fig.3A by immunofluorescence assay, the cytoplasmic and nuclear expression of  $\beta$ -catenin was decreased obviously after 5-FU treatment, however, ASP and LiCl pretreatment respectively rescued the expression of  $\beta$ -catenin and its nuclear translocation. Also, the results of western blot revealed that ASP and LiCl pretreatment significantly reversed the 5-FU-induced decrease in cytoplasmic expression of total  $\beta$ -catenin, p-GSK-3 $\beta$ , and CyclinD1, meanwhile modulated nuclear expression of  $\beta$ -catenin, Lef-1, and C-myc proteins (Fig.3B-I). These data suggested that ASP may activate Wnt signaling, which may be one mechanism that ASP counteract the inhibiting effect of 5-FU on HS-5 cell growth.

### **Angelica Sinensis Polysaccharides Relieve 5-FU-induced Intracellular Oxidative Stress**

To elucidate the mechanism of 5-FU induced damage and ASP mediated protective effect on HS-5 cell growth, we assessed the indexes of oxidative damage. Increases of intracellular ROS and MDA were

found in 5-FU group compared with the control group whereas single pretreatment of ASP or LiCl reversed the increase dramatically (Fig.4A, C). On the contrary, ASP or LiCl administration protected the antioxidant enzymes including SOD and CAT in HS-5 cells (Fig.4D, E). These results demonstrate that 5-FU cause oxidative stress to HS-5 cells, whereas ASP exert a significant anti-oxidative role to alleviate 5-FU-induced oxidative stress, which may be related to the activation of Wnt/ $\beta$ -catenin signaling.

### **Angelica Sinensis Polysaccharides Ameliorate the Activation of FoxO1 Induced by 5-FU**

FoxOs are transcriptional factors closely related to cellular survival and oxidative stress, Notably, activated FoxO1 may impair Wnt signaling via competitive combination with  $\beta$ -catenin in nucleus. To address the question that whether 5-FU-induced intracellular oxidative stress and mitigative effects of ASP is related to FoxO1 transcription, we analyzed the levels of FoxO1 protein and its phosphorylation. Western blot results showed that compared with the control group FoxO1 expression in 5-FU group rose concurrent with decreased p-FoxO1 expression. However, ASP or LiCl pretreatment significantly reduced of FoxO1 expression via degradation of FoxO1 by an increased p-FoxO1 expression (Fig.5A). It was inferred that FoxO1 may be involved in response to 5-FU induced intracellular oxidative stress. Antioxidative properties of ASP may play a role in FoxO1 downregulation, which may be another possible mechanism for ASP upregulation of Wnt/ $\beta$ -catenin signaling. Like p53, FoxOs orchestrate apoptosis through multiple mechanisms, also FoxOs may promote cell cycle arrest by upregulating multiple cell cycle suppressor genes such as CDKI p27<sup>Kip1</sup>. Here, in this context, the proteins correlating to apoptosis and cell cycle modulators including Bcl-2, Bim, Bax, caspase-3 and p27<sup>Kip1</sup> were detected via western blot assay. The results showed that ASP or LiCl counteracted apoptosis and cell cycle arrest by abrogating 5-FU-induced increase in Bim, Bax caspase-3 and p27<sup>Kip1</sup> expression, also by enhancing anti-apoptotic protein Bcl-2 expression (Fig.5B-F, H). The data of flowcytometric analysis revealed that compared with the control group there was a 2.2-fold up-regulation of the apoptosis rate in 5-FU group, ASP or LiCl significantly decreased the percentage of apoptotic cells compared with the 5-FU group (Fig.5G). Meanwhile, ASP or LiCl pretreatment weakened 5-FU induced G0/G1 phase retard concurrent with S and G2/M phase recovery (Fig.5I). All these data above hinted that the effects of 5-FU on HS-5 growth inhibition may relate to the activation of FoxO1 leading to apoptosis or cycle arrest. Anti-oxidative property of ASP exerts a protective effect against cycle arrest and apoptosis.

## **Discussion**

Myelosuppression is one of the common side effects of chemotherapy, characterized by depletion of cells within the bone marrow[40, 41]. In general, myelosuppression is primarily attributable to the direct cytotoxicity to bone marrow cells, inhibition of bone marrow precursor or progenitor cell proliferation, the reduction in HSC reserves and impairment in HSC self-renewal. Notably, because of the reduction of HM cellularity in varying degrees, the damaged-hematopoietic microenvironment may result in diminished or delayed hematopoiesis function, immune-related disorders, as well as long-term damage to the bone marrow recovery[42, 43]. It has been shown that chemotherapeutic treatment damage the hematopoietic microenvironment in vitro and vivo[5, 44, 6, 7, 9, 10]. As chemotherapy disrupts the steady-state function



of hematopoietic and stromal cell, disruptions over time may cause severe bone marrow toxicity and the failure of cancer treatment. To ensure this does not occur, finding appropriate agents to promote the recovery process following discontinuation of chemotherapy and to lessen the bone marrow damage has a profound significance.

Since it was first synthesized in 1957, 5-FU has remained one of the most widely used chemotherapeutic agents with broad-spectrum activity against many solid tumors[45]. 5-FU exerts its anticancer effects through inhibition of thymidylate synthase (TS) and incorporation of its metabolites into RNA and DNA, leading to cytotoxicity and cell death[12]. Recent studies have indicated that 5-FU suppressed the proliferation of HSCs and induced the myelosuppression of mice by down-regulating PI3K-AKT signaling pathway[14, 46]. However, the definite mechanism for 5-FU caused myelosuppression remains unclear. Focused on bone marrow stromal cells, we provided the evidence that 5-FU inhibited stromal cell growth and induced apoptosis, which was related to downregulation of Wnt/ $\beta$ -catenin signaling, also up-regulation of FoxO1 concomitant with an increase of cellular oxidative stress. Furthermore, the current work revealed that anti-oxidative property and role in Wnt signaling regulation might be the key mechanisms of ASP to prevent against 5-FU-induced stromal damage.

Stem cells display the defining capacity to self-renew, and their fate are primarily dictated by extrinsic, short-range signals, which typically emanated from the stem cell niche[47]. The non-hematopoietic cells in hematopoietic microenvironment have a functional role in regulating hematopoiesis and the signaling pathways that regulate HM may be necessary for the development of functional niches that regulate hematopoietic stem cells and their progenitors[48, 49]. The Wnt signaling pathway exerts a variety of effects on target cell developmental processes, including cell proliferation, apoptosis, and differentiation. The canonical Wnt pathway affects cellular functions by accumulating of  $\beta$ -catenin in the cytoplasm and eventually translocating into the nucleus. Within the nucleus,  $\beta$ -catenin binds to T cell factor (TCF) family/lymphoid enhancer factor (LEF) and regulates cell proliferation through Wnt downstream target genes[16, 50, 51]. It was reported that Wnt/ $\beta$ -catenin signaling regulates HSCs function in dosage-dependent manner[52–55]. Various degrees of activation of the pathway may cause different outcomes, leading to either enhanced repopulation capacity or exhaustion of the HSCs. A mild increase in Wnt signaling enhanced HSC function[18, 56]. However, a high Wnt level in HSCs eventually leads to stem cell exhaustion and impairment of reconstitution in irradiated recipients[57–59]. Most importantly, Wnt signaling regulates HSC reconstruction in a stromal-dependent manner. It was found that when hematopoietic cells were co-cultured with BMSCs supplemented with Wnt3a conditioned medium, the cellularity of Lin<sup>-</sup>Sca-1<sup>+</sup>c-kit<sup>+</sup> hematopoietic stem cells were increased, and the hematopoietic transplantation and reconstruction capability were enhanced[60, 20]. Hence, in the current study we focused on the Wnt signaling regulation on BMSCs following chemotherapy. It was found that 5-FU induced a decrease in cytoplasmic expression of total  $\beta$ -catenin, p-GSK-3 $\beta$ , and CyclinD1, meanwhile weakened nuclear expression of  $\beta$ -catenin, LEF-1, and C-myc proteins, causing HS-5 cells proliferation inhibition. The results herein are in line with the other data related to the relationship between canonical

Wnt signaling and cell proliferation, which has confirmed that Wnt/ $\beta$ -catenin signaling positively stimulates cell growth via cell cycle regulation[61, 62].

Reactive oxygen species (ROS) are free radicals and active metabolites of oxygen containing unpaired electrons, which take a significant role in cell signal transduction and regulation[63]. Chemical agents, as well as irradiation can cause a persistent ROS production. This accumulation of ROS may lead to excessive oxidative stress and DNA damage such as DSBs (double-strand breaks), which is considered to be the main potential mechanisms causing cellular damage[64, 65]. Previous study in our group has demonstrated that 5-FU weakened the antioxidant capacity of HS-5 cells and caused high sensitivity of cells to ROS, thus HS-5 cells underwent DSB which eventually resulted in either apoptosis or senescence[15]. Oxidative stress is also related to cell cycle arrest. DSBs initiate DNA damage response through sequential stimulation of ATM, Chk2, and p53[66]. Activation of p53 and its downstream p21 may induce the cell cycle arrest. Meanwhile ROS can activate p38 MAPK pathway[67]. Activation of p53 and p38 pathways can converge at p16 and augment of p16 expression may also lead to permanent cell cycle arrest[68, 69]. Interestingly, it is reported that  $\beta$ -catenin may be critical for antagonizing oxidative stress. Exposing  $\beta$ -catenin knock- down mice to chemotherapeutic agent or radiation caused a decreased expression of the hydrogen peroxide ( $H_2O_2$ ) detoxifying enzyme catalase and led to the accumulation of ROS and superoxide ( $O_2^-$ ) free radicals in cells and an inability to repair DNA damage[70]. On the opposite, effector molecules generated from oxidative DNA damage may also down-regulate the Wnt pathway by inhibiting transcriptional activity or participating in post-translational modifications to enhance ubiquitination degradation[71]. These evidences above hint that Wnt signaling also closely correlated with oxidative stress. Therefore, in the current study, increased oxidative stress may be one of the reasons for down-regulation of wnt signaling induced by 5-FU treatment. Whereas decrease in  $\beta$ -catenin protein accompanying reduction of antioxidant SOD and CAT induced by 5-FU treatment may be another mechanism of cell proliferation inhibition.

Forkhead box O (FOXO) family are transcription factors, which promote cell survival by regulating the cell cycle, apoptosis and the response to oxidative stress[23]. The accumulation of ROS may interrupt 14-3-3 combine to FoxO via JNK (c-Jun N terminal kinase), permit FoxO entrance into nucleus, and induce its transcriptional activation[72, 73]. FoxO can be phosphorylated by phosphatidylinositol 3-kinase-Akt pathway[74, 75]. It is of note that FoxO-mediated transcription requires binding of  $\beta$ -catenin. FoxOs can compete with TCF/LEF by directly binding  $\beta$ -catenin, thereby inhibit Wnt/ $\beta$ -catenin downstream signaling[76, 77, 24, 78]. It was demonstrated herein, compared with the control group FoxO1 expression in 5-FU treated HS-5 cells rose dramatically concurrent with decreased p-FoxO1 expression. The reason for up-regulation of FoxO1 may be related to 5-FU triggered oxidative stress, whereas FoxO1 up-regulation may be another reason for 5-FU induced decrease in Wnt signaling[27, 26, 79]. FoxO transcription factor family regulate the proteins that are crucial for the apoptosis, as well as the proteins involved in proliferative status of a cell. FoxO factors may regulate antiapoptotic and proapoptotic proteins at multiple levels, finally trigger activation of the effector caspases. Bim promotes apoptosis by inhibition of antiapoptotic Bcl-2 family members or through direct activation of Bax-like molecules. FoxO factors may

regulate Bim protein expression to cause cell death due to cytokine deprivation. FoxO factors may also repress transcription of Bcl-XL through the induction of the transcriptional repressor[80–82]. Caspase-3 is an important effector protease, when it is cleaved, it acts as the final executor during apoptosis. In the current study, it was found that in 5-FU treated HS-5 cells, FoxO1 targeted apoptosis-related proteins to cause increase in Bim, Bax, and caspase-3 whereas decrease in Bcl-2. FoxO1 targeted apoptosis to disturb the dynamic balance of the cellularity of HS-5 cells, which may be one of reasons for cell growth inhibition. Moreover, the cyclin kinase inhibitor p27<sup>Kip1</sup>, a downstream target of FoxO1, acting as a potent inhibitor of cyclin/cdk complexes in the S-phase of cell cycle progression was also tested[83–86]. It was found herein that 5-FU increased the expression of p27<sup>Kip1</sup>. In addition, 5-FU simultaneously reduced the expression of Cyclin D1. It is of note that transcriptional repression of D-type cyclins is vital to the FoxO-induced cell-cycle arrest, which is evidenced by transcriptional profiling and mRNA analysis. D-type cyclins are required for phosphorylation and inactivation of the retinoblastoma tumor suppressor protein (pRb), an essential determinant of cell-cycle progression in G1[87, 88]. To sum up, 5-FU-induced HS-5 cell growth inhibition is probably associated with FoxO1 targeted apoptosis or cell cycle arrest.

The traditional Chinese medicine *Angelica sinensis* which is commonly used to enrich blood, promote blood circulation[39]. The active constituents of *Angelica sinensis* include polysaccharides, organic acid and phthalides, among which *Angelica sinensis* polysaccharides (ASP) are regarded as the main biological activity ingredient responsible for pharmacological effects with multi-target property[89]. ASP have attracted more and more attention to its beneficial effects, such as hematopoietic effects[90], immunologic enhancement[91], anti-tumor activity[92, 93], and anti-radiation damage[94]. The antioxidant properties of ASP suppress the production of ROS and protected the endothelial progenitor cells, hepatocytes, myocardial cell and nerve cells from oxidative damage[95, 96, 40]. Moreover, evidences demonstrated that ASP promote cell proliferation, including in total spleen cells, macrophages[91], and gastric epithelial cells[97]. Our previous studies suggested that ASP reduced oxidative stress and oxidative DNA damage, boosted direct cell-cell contact between stromal cells and hematopoietic cells through Cx43 junctions, regulated cytokines, growth factors and chemokines such as CXCL12, SCF, GM-CSF, RANTES and thus provided a homeostatic microenvironment for hematopoietic stem/progenitor cells to regenerate following chemotherapeutic myelosuppression. In the present study, it was further demonstrated that ASP protected HS-5 cells from 5-FU-induced proliferation inhibition and ameliorated cellular oxidative stress via the mechanism of up-regulation of Wnt/ $\beta$ -catenin signaling. Most importantly, it was first evidenced herein that ASP balanced the relationship between FoxO-mediated transcription and Wnt signaling in BMSCs under oxidative stress, which might be promising for clinical therapeutic use of ASP to myelosuppression.

## Conclusions

In conclusion, the present study has reported that ASP protect stromal cells against 5-FU-induced proliferation inhibition and apoptosis via activating the Wnt/ $\beta$ -catenin signaling pathway directly or the indirect effects on Wnt/ $\beta$ -catenin signaling by down-regulation of its antagonizing FoxO1, suggesting a

broad role for ASP as a potential antioxidant protective agent for chemoradiation therapeutic preventive agents.

## Abbreviations

5-FU

5-fluorouracil; LiCl:Lithium chloride; ASP:Angelica sinensis polysaccharides; BMSC:BMSC Bone marrow stromal cell; HSC:HSC Hematopoietic stem cell; SOD:Superoxide dismutase; CAT:Catalase; MDA:Malondialdehyde; FoxO:Forkhead box O; ROS:Reactive oxygen species; HM:Hematopoietic microenvironment.

## Declarations

### Acknowledgements

Not applicable.

**Author's contributions:** For research articles, LW conceptualized and designed the experiments; HXZX performed the experiments; RJQ, ZLW and MHX, YX, YPW contributed reagents/materials/analysis tools; HXZX analyzed the data and wrote the paper; LW revised the paper.

### Funding

This work was supported by the National Natural Science Foundation of China (Grant No.81873103), the Foundation and Frontier Research Project of Chongqing Science and Technology Commission (Grant No. cstc2014jcyjA10001).

### Availability of data and materials

The datasets used in this study are available from the corresponding author upon reasonable request.

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that there are no conflicts of interest.

### Author details

## References

1. Chabner BA, Roberts TG, Jr. (2005) Timeline: Chemotherapy and the war on cancer. *Nat Rev Cancer* 5 (1):65-72. doi:10.1038/nrc1529
2. Papac RJ (2001) Origins of cancer therapy. *Yale J Biol Med* 74 (6):391-398
3. Dritschilo A, Sherman DS (1981) Radiation and chemical injury in the bone marrow. *Environ Health Perspect* 39:59-64. doi:10.1289/ehp.813959
4. Marsh JC (1976) The effects of cancer chemotherapeutic agents on normal hematopoietic precursor cells: a review. *Cancer Res* 36 (6):1853-1882
5. Galotto M, Berisso G, Delfino L, Podesta M, Ottaggio L, Dallorso S, Dufour C, Ferrara GB, Abbondandolo A, Dini G, Bacigalupo A, Cancedda R, Quarto R (1999) Stromal damage as consequence of high-dose chemo/radiotherapy in bone marrow transplant recipients. *Exp Hematol* 27 (9):1460-1466. doi:10.1016/s0301-472x(99)00076-4
6. Kemp K, Morse R, Wexler S, Cox C, Mallam E, Hows J, Donaldson C (2010) Chemotherapy-induced mesenchymal stem cell damage in patients with hematological malignancy. *Ann Hematol* 89 (7):701-713. doi:10.1007/s00277-009-0896-2
7. Li J, Law HK, Lau YL, Chan GC (2004) Differential damage and recovery of human mesenchymal stem cells after exposure to chemotherapeutic agents. *Br J Haematol* 127 (3):326-334. doi:10.1111/j.1365-2141.2004.05200.x
8. Nicolay NH, Ruhle A, Perez RL, Trinh T, Sisombath S, Weber KJ, Ho AD, Debus J, Saffrich R, Huber PE (2016) Mesenchymal stem cells are sensitive to bleomycin treatment. *Sci Rep* 6:26645. doi:10.1038/srep26645
9. Oliveira MS, Carvalho JL, Campos AC, Gomes DA, de Goes AM, Melo MM (2014) Doxorubicin has in vivo toxicological effects on ex vivo cultured mesenchymal stem cells. *Toxicol Lett* 224 (3):380-386. doi:10.1016/j.toxlet.2013.11.023
10. Prata Kde L, Orellana MD, De Santis GC, Kashima S, Fontes AM, Carrara Rde C, Palma PV, Neder L, Covas DT (2010) Effects of high-dose chemotherapy on bone marrow multipotent mesenchymal stromal cells isolated from lymphoma patients. *Exp Hematol* 38 (4):292-300 e294. doi:10.1016/j.exphem.2010.01.006
11. Douillard JY, Cunningham D, Roth AD, Navarro M, James RD, Karasek P, Jandik P, Iveson T, Carmichael J, Alakl M, Gruia G, Awad L, Rougier P (2000) Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *Lancet* 355 (9209):1041-1047. doi:10.1016/s0140-6736(00)02034-1
12. Longley DB, Harkin DP, Johnston PG (2003) 5-fluorouracil: mechanisms of action and clinical strategies. *Nat Rev Cancer* 3 (5):330-338. doi:10.1038/nrc1074

13. Somaiah C, Kumar A, Sharma R, Sharma A, Anand T, Bhattacharyya J, Das D, Deka Talukdar S, Jaganathan BG (2018) Mesenchymal stem cells show functional defect and decreased anti-cancer effect after exposure to chemotherapeutic drugs. *J Biomed Sci* 25 (1):5. doi:10.1186/s12929-018-0407-7
14. Wang S, Zheng G, Tian S, Zhang Y, Shen L, Pak Y, Shen Y, Qian J (2015) Echinacoside improves hematopoietic function in 5-FU-induced myelosuppression mice. *Life Sci* 123:86-92. doi:10.1016/j.lfs.2015.01.002
15. Xiao H, Xiong L, Song X, Jin P, Chen L, Chen X, Yao H, Wang Y, Wang L (2017) Angelica sinensis Polysaccharides Ameliorate Stress-Induced Premature Senescence of Hematopoietic Cell via Protecting Bone Marrow Stromal Cells from Oxidative Injuries Caused by 5-Fluorouracil. *Int J Mol Sci* 18 (11). doi:10.3390/ijms18112265
16. Nusse R, Clevers H (2017) Wnt/beta-Catenin Signaling, Disease, and Emerging Therapeutic Modalities. *Cell* 169 (6):985-999. doi:10.1016/j.cell.2017.05.016
17. Petersen CP, Reddien PW (2009) Wnt signaling and the polarity of the primary body axis. *Cell* 139 (6):1056-1068. doi:10.1016/j.cell.2009.11.035
18. Luis TC, Naber BA, Roozen PP, Brugman MH, de Haas EF, Ghazvini M, Fibbe WE, van Dongen JJ, Fodde R, Staal FJ (2011) Canonical wnt signaling regulates hematopoiesis in a dosage-dependent fashion. *Cell Stem Cell* 9 (4):345-356. doi:10.1016/j.stem.2011.07.017
19. Richter J, Traver D, Willert K (2017) The role of Wnt signaling in hematopoietic stem cell development. *Crit Rev Biochem Mol Biol* 52 (4):414-424. doi:10.1080/10409238.2017.1325828
20. Kim JA, Kang YJ, Park G, Kim M, Park YO, Kim H, Leem SH, Chu IS, Lee JS, Jho EH, Oh IH (2009) Identification of a stroma-mediated Wnt/beta-catenin signal promoting self-renewal of hematopoietic stem cells in the stem cell niche. *Stem Cells* 27 (6):1318-1329. doi:10.1002/stem.52
21. Oh IH (2010) Microenvironmental targeting of Wnt/beta-catenin signals for hematopoietic stem cell regulation. *Expert Opin Biol Ther* 10 (9):1315-1329. doi:10.1517/14712598.2010.504705
22. Schreck C, Bock F, Grziwotk S, Oostendorp RA, Istvanffy R (2014) Regulation of hematopoiesis by activators and inhibitors of Wnt signaling from the niche. *Ann N Y Acad Sci* 1310:32-43. doi:10.1111/nyas.12384
23. Eijkelenboom A, Burgering BM (2013) FOXOs: signalling integrators for homeostasis maintenance. *Nat Rev Mol Cell Biol* 14 (2):83-97. doi:10.1038/nrm3507
24. Behrens J, von Kries JP, Kuhl M, Bruhn L, Wedlich D, Grosschedl R, Birchmeier W (1996) Functional interaction of beta-catenin with the transcription factor LEF-1. *Nature* 382 (6592):638-642. doi:10.1038/382638a0
25. Essers MA, de Vries-Smits LM, Barker N, Polderman PE, Burgering BM, Korswagen HC (2005) Functional interaction between beta-catenin and FOXO in oxidative stress signaling. *Science* 308 (5725):1181-1184. doi:10.1126/science.1109083
26. Burgering BM, Medema RH (2003) Decisions on life and death: FOXO Forkhead transcription factors are in command when PKB/Akt is off duty. *J Leukoc Biol* 73 (6):689-701. doi:10.1189/jlb.1202629

27. Ma Y, Wang H (2012) PI3K/Akt/FoxO: a novel participant in signal transduction in bone cells under mechanical stimulation. *Cell Biol Int* 36 (10):923-926. doi:10.1042/CBI20120078
28. Dietz BM, Hajirahimkhan A, Dunlap TL, Bolton JL (2016) Botanicals and Their Bioactive Phytochemicals for Women's Health. *Pharmacol Rev* 68 (4):1026-1073. doi:10.1124/pr.115.010843
29. Zhao KJ, Dong TT, Tu PF, Song ZH, Lo CK, Tsim KW (2003) Molecular genetic and chemical assessment of radix *Angelica* (*Danggui*) in China. *J Agric Food Chem* 51 (9):2576-2583. doi:10.1021/jf026178h
30. Lei T, Li H, Fang Z, Lin J, Wang S, Xiao L, Yang F, Liu X, Zhang J, Huang Z, Liao W (2014) Polysaccharides from *Angelica sinensis* alleviate neuronal cell injury caused by oxidative stress. *Neural Regen Res* 9 (3):260-267. doi:10.4103/1673-5374.128218
31. Zhuang C, Wang Y, Zhang Y, Xu N (2018) Oxidative stress in osteoarthritis and antioxidant effect of polysaccharide from *angelica sinensis*. *Int J Biol Macromol* 115:281-286. doi:10.1016/j.ijbiomac.2018.04.083
32. Tsai NM, Lin SZ, Lee CC, Chen SP, Su HC, Chang WL, Harn HJ (2005) The antitumor effects of *Angelica sinensis* on malignant brain tumors in vitro and in vivo. *Clin Cancer Res* 11 (9):3475-3484. doi:10.1158/1078-0432.CCR-04-1827
33. Zhang Y, Zhou T, Wang H, Cui Z, Cheng F, Wang KP (2016) Structural characterization and in vitro antitumor activity of an acidic polysaccharide from *Angelica sinensis* (Oliv.) Diels. *Carbohydr Polym* 147:401-408. doi:10.1016/j.carbpol.2016.04.002
34. Bradley RR, Cunniff PJ, Pereira BJ, Jaber BL (1999) Hematopoietic effect of *Radix angelicae sinensis* in a hemodialysis patient. *Am J Kidney Dis* 34 (2):349-354. doi:10.1016/s0272-6386(99)70367-7
35. Liu C, Li J, Meng FY, Liang SX, Deng R, Li CK, Pong NH, Lau CP, Cheng SW, Ye JY, Chen JL, Yang ST, Yan H, Chen S, Chong BH, Yang M (2010) Polysaccharides from the root of *Angelica sinensis* promotes hematopoiesis and thrombopoiesis through the PI3K/AKT pathway. *BMC Complement Altern Med* 10:79. doi:10.1186/1472-6882-10-79
36. Wang K, Wu J, Cheng F, Huang X, Zeng F, Zhang Y (2017) Acidic Polysaccharide from *Angelica sinensis* Reverses Anemia of Chronic Disease Involving the Suppression of Inflammatory Hepcidin and NF-kappaB Activation. *Oxid Med Cell Longev* 2017:7601592. doi:10.1155/2017/7601592
37. Lai P, Liu Y (2015) *Angelica sinensis* polysaccharides inhibit endothelial progenitor cell senescence through the reduction of oxidative stress and activation of the Akt/hTERT pathway. *Pharm Biol* 53 (12):1842-1849. doi:10.3109/13880209.2015.1027779
38. Mu X, Zhang Y, Li J, Xia J, Chen X, Jing P, Song X, Wang L, Wang Y (2017) *Angelica Sinensis* Polysaccharide Prevents Hematopoietic Stem Cells Senescence in D-Galactose-Induced Aging Mouse Model. *Stem Cells Int* 2017:3508907. doi:10.1155/2017/3508907
39. Wei WL, Zeng R, Gu CM, Qu Y, Huang LF (2016) *Angelica sinensis* in China-A review of botanical profile, ethnopharmacology, phytochemistry and chemical analysis. *J Ethnopharmacol* 190:116-141. doi:10.1016/j.jep.2016.05.023

40. Ai S, Fan X, Fan L, Sun Q, Liu Y, Tao X, Dai K (2013) Extraction and chemical characterization of *Angelica sinensis* polysaccharides and its antioxidant activity. *Carbohydr Polym* 94 (2):731-736. doi:10.1016/j.carbpol.2013.02.007
41. Testa NG, Hendry JH, Molineux G (1985) Long-term bone marrow damage in experimental systems and in patients after radiation or chemotherapy. *Anticancer Res* 5 (1):101-110
42. Crawford J, Dale DC, Lyman GH (2004) Chemotherapy-induced neutropenia: risks, consequences, and new directions for its management. *Cancer* 100 (2):228-237. doi:10.1002/cncr.11882
43. Kuter DJ (2015) Managing thrombocytopenia associated with cancer chemotherapy. *Oncology (Williston Park)* 29 (4):282-294
44. Hu W, Sung T, Jessen BA, Thibault S, Finkelstein MB, Khan NK, Sacca AI (2016) Mechanistic Investigation of Bone Marrow Suppression Associated with Palbociclib and its Differentiation from Cytotoxic Chemotherapies. *Clin Cancer Res* 22 (8):2000-2008. doi:10.1158/1078-0432.CCR-15-1421
45. Wilson PM, Danenberg PV, Johnston PG, Lenz HJ, Ladner RD (2014) Standing the test of time: targeting thymidylate biosynthesis in cancer therapy. *Nat Rev Clin Oncol* 11 (5):282-298. doi:10.1038/nrclinonc.2014.51
46. Zhang Y, Ye T, Hong Z, Gong S, Zhou X, Liu H, Qian J, Qu H (2019) Pharmacological and transcriptome profiling analyses of Fufang E'jiao Jiang during chemotherapy-induced myelosuppression in mice. *J Ethnopharmacol* 238:111869. doi:10.1016/j.jep.2019.111869
47. Losick VP, Morris LX, Fox DT, Spradling A (2011) *Drosophila* stem cell niches: a decade of discovery suggests a unified view of stem cell regulation. *Dev Cell* 21 (1):159-171. doi:10.1016/j.devcel.2011.06.018
48. Zhang J, Niu C, Ye L, Huang H, He X, Tong WG, Ross J, Haug J, Johnson T, Feng JQ, Harris S, Wiedemann LM, Mishina Y, Li L (2003) Identification of the haematopoietic stem cell niche and control of the niche size. *Nature* 425 (6960):836-841. doi:10.1038/nature02041
49. Morrison SJ, Spradling AC (2008) Stem cells and niches: mechanisms that promote stem cell maintenance throughout life. *Cell* 132 (4):598-611. doi:10.1016/j.cell.2008.01.038
50. Moon RT, Bowerman B, Boutros M, Perrimon N (2002) The promise and perils of Wnt signaling through beta-catenin. *Science* 296 (5573):1644-1646. doi:10.1126/science.1071549
51. Clevers H (2006) Wnt/beta-catenin signaling in development and disease. *Cell* 127 (3):469-480. doi:10.1016/j.cell.2006.10.018
52. Mohammed MK, Shao C, Wang J, Wei Q, Wang X, Collier Z, Tang S, Liu H, Zhang F, Huang J, Guo D, Lu M, Liu F, Liu J, Ma C, Shi LL, Athiviraham A, He TC, Lee MJ (2016) Wnt/beta-catenin signaling plays an ever-expanding role in stem cell self-renewal, tumorigenesis and cancer chemoresistance. *Genes Dis* 3 (1):11-40. doi:10.1016/j.gendis.2015.12.004
53. Malhotra S, Kincade PW (2009) Wnt-related molecules and signaling pathway equilibrium in hematopoiesis. *Cell Stem Cell* 4 (1):27-36. doi:10.1016/j.stem.2008.12.004
54. Huang J, Nguyen-McCarty M, Hexner EO, Danet-Desnoyers G, Klein PS (2012) Maintenance of hematopoietic stem cells through regulation of Wnt and mTOR pathways. *Nat Med* 18 (12):1778-



1785. doi:10.1038/nm.2984


55. Fleming HE, Janzen V, Lo Celso C, Guo J, Leahy KM, Kronenberg HM, Scadden DT (2008) Wnt signaling in the niche enforces hematopoietic stem cell quiescence and is necessary to preserve self-renewal in vivo. *Cell Stem Cell* 2 (3):274-283. doi:10.1016/j.stem.2008.01.003
56. Famili F, Brugman MH, Taskesen E, Naber BEA, Fodde R, Staal FJT (2016) High Levels of Canonical Wnt Signaling Lead to Loss of Stemness and Increased Differentiation in Hematopoietic Stem Cells. *Stem Cell Reports* 6 (5):652-659. doi:10.1016/j.stemcr.2016.04.009
57. Scheller M, Huelsken J, Rosenbauer F, Taketo MM, Birchmeier W, Tenen DG, Leutz A (2006) Hematopoietic stem cell and multilineage defects generated by constitutive beta-catenin activation. *Nat Immunol* 7 (10):1037-1047. doi:10.1038/ni1387
58. Ming M, Wang S, Wu W, Senyuk V, Le Beau MM, Nucifora G, Qian Z (2012) Activation of Wnt/beta-catenin protein signaling induces mitochondria-mediated apoptosis in hematopoietic progenitor cells. *J Biol Chem* 287 (27):22683-22690. doi:10.1074/jbc.M112.342089
59. Kirstetter P, Anderson K, Porse BT, Jacobsen SE, Nerlov C (2006) Activation of the canonical Wnt pathway leads to loss of hematopoietic stem cell repopulation and multilineage differentiation block. *Nat Immunol* 7 (10):1048-1056. doi:10.1038/ni1381
60. Nemeth MJ, Mak KK, Yang Y, Bodine DM (2009) beta-Catenin expression in the bone marrow microenvironment is required for long-term maintenance of primitive hematopoietic cells. *Stem Cells* 27 (5):1109-1119. doi:10.1002/stem.32
61. Shtutman M, Zhurinsky J, Simcha I, Albanese C, D'Amico M, Pestell R, Ben-Ze'ev A (1999) The cyclin D1 gene is a target of the beta-catenin/LEF-1 pathway. *Proc Natl Acad Sci U S A* 96 (10):5522-5527. doi:10.1073/pnas.96.10.5522
62. Braunschweig L, Meyer AK, Wagenfuhr L, Storch A (2015) Oxygen regulates proliferation of neural stem cells through Wnt/beta-catenin signalling. *Mol Cell Neurosci* 67:84-92. doi:10.1016/j.mcn.2015.06.006
63. Owusu-Ansah E, Banerjee U (2009) Reactive oxygen species prime *Drosophila* haematopoietic progenitors for differentiation. *Nature* 461 (7263):537-541. doi:10.1038/nature08313
64. Wang Y, Liu L, Pazhanisamy SK, Li H, Meng A, Zhou D (2010) Total body irradiation causes residual bone marrow injury by induction of persistent oxidative stress in murine hematopoietic stem cells. *Free Radic Biol Med* 48 (2):348-356. doi:10.1016/j.freeradbiomed.2009.11.005
65. Meng A, Wang Y, Van Zant G, Zhou D (2003) Ionizing radiation and busulfan induce premature senescence in murine bone marrow hematopoietic cells. *Cancer Res* 63 (17):5414-5419
66. Sancar A, Lindsey-Boltz LA, Unsal-Kacmaz K, Linn S (2004) Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints. *Annu Rev Biochem* 73:39-85. doi:10.1146/annurev.biochem.73.011303.073723
67. Ito K, Hirao A, Arai F, Takubo K, Matsuo S, Miyamoto K, Ohmura M, Naka K, Hosokawa K, Ikeda Y, Suda T (2006) Reactive oxygen species act through p38 MAPK to limit the lifespan of hematopoietic stem cells. *Nat Med* 12 (4):446-451. doi:10.1038/nm1388

68. Iwasa H, Han J, Ishikawa F (2003) Mitogen-activated protein kinase p38 defines the common senescence-signalling pathway. *Genes Cells* 8 (2):131-144. doi:10.1046/j.1365-2443.2003.00620.x
69. Beausejour CM, Krtolica A, Galimi F, Narita M, Lowe SW, Yaswen P, Campisi J (2003) Reversal of human cellular senescence: roles of the p53 and p16 pathways. *EMBO J* 22 (16):4212-4222. doi:10.1093/emboj/cdg417
70. Lento W, Ito T, Zhao C, Harris JR, Huang W, Jiang C, Owzar K, Piryani S, Racioppi L, Chao N, Reya T (2014) Loss of beta-catenin triggers oxidative stress and impairs hematopoietic regeneration. *Genes Dev* 28 (9):995-1004. doi:10.1101/gad.231944.113
71. Lin CL, Wang JY, Ko JY, Surendran K, Huang YT, Kuo YH, Wang FS (2008) Superoxide destabilization of beta-catenin augments apoptosis of high-glucose-stressed mesangial cells. *Endocrinology* 149 (6):2934-2942. doi:10.1210/en.2007-1372
72. Nakae J, Oki M, Cao Y (2008) The FoxO transcription factors and metabolic regulation. *FEBS Lett* 582 (1):54-67. doi:10.1016/j.febslet.2007.11.025
73. Morrison DK (2009) The 14-3-3 proteins: integrators of diverse signaling cues that impact cell fate and cancer development. *Trends Cell Biol* 19 (1):16-23. doi:10.1016/j.tcb.2008.10.003
74. Han J, Zhao J, Jiang J, Ma X, Liu X, Wang C, Jiang S, Wan C (2015) Zinc deficiency impairs the renewal of hippocampal neural stem cells in adult rats: involvement of FoxO3a activation and downstream p27(kip1) expression. *J Neurochem* 134 (5):879-891. doi:10.1111/jnc.13199
75. Zhang K, Yang Y, Ge H, Wang J, Chen X, Lei X, Zhong J, Zhang C, Xian J, Lu Y, Tan L, Feng H (2020) Artesunate promotes the proliferation of neural stem/progenitor cells and alleviates Ischemia-reperfusion Injury through PI3K/Akt/FOXO-3a/p27(kip1) signaling pathway. *Aging (Albany NY)* 12 (9):8029-8048. doi:10.18632/aging.103121
76. Iyer S, Ambrogini E, Bartell SM, Han L, Roberson PK, de Cabo R, Jilka RL, Weinstein RS, O'Brien CA, Manolagas SC, Almeida M (2013) FOXOs attenuate bone formation by suppressing Wnt signaling. *J Clin Invest* 123 (8):3409-3419. doi:10.1172/JCI68049
77. Hoogeboom D, Essers MA, Polderman PE, Voets E, Smits LM, Burgering BM (2008) Interaction of FOXO with beta-catenin inhibits beta-catenin/T cell factor activity. *J Biol Chem* 283 (14):9224-9230. doi:10.1074/jbc.M706638200
78. Almeida M, Han L, Martin-Millan M, O'Brien CA, Manolagas SC (2007) Oxidative stress antagonizes Wnt signaling in osteoblast precursors by diverting beta-catenin from T cell factor- to forkhead box O-mediated transcription. *J Biol Chem* 282 (37):27298-27305. doi:10.1074/jbc.M702811200
79. Danciu TE, Gagari E, Adam RM, Damoulis PD, Freeman MR (2004) Mechanical strain delivers anti-apoptotic and proliferative signals to gingival fibroblasts. *J Dent Res* 83 (8):596-601. doi:10.1177/154405910408300803
80. Tang TT, Dowbenko D, Jackson A, Toney L, Lewin DA, Dent AL, Lasky LA (2002) The forkhead transcription factor AFX activates apoptosis by induction of the BCL-6 transcriptional repressor. *J Biol Chem* 277 (16):14255-14265. doi:10.1074/jbc.M110901200

81. Stahl M, Dijkers PF, Kops GJ, Lens SM, Coffe PJ, Burgering BM, Medema RH (2002) The forkhead transcription factor FoxO regulates transcription of p27Kip1 and Bim in response to IL-2. *J Immunol* 168 (10):5024-5031. doi:10.4049/jimmunol.168.10.5024
82. Dijkers PF, Medema RH, Lammers JW, Koenderman L, Coffe PJ (2000) Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the forkhead transcription factor FKHR-L1. *Curr Biol* 10 (19):1201-1204. doi:10.1016/s0960-9822(00)00728-4
83. Nakamura N, Ramaswamy S, Vazquez F, Signoretti S, Loda M, Sellers WR (2000) Forkhead transcription factors are critical effectors of cell death and cell cycle arrest downstream of PTEN. *Mol Cell Biol* 20 (23):8969-8982. doi:10.1128/mcb.20.23.8969-8982.2000
84. Medema RH, Kops GJ, Bos JL, Burgering BM (2000) AFX-like Forkhead transcription factors mediate cell-cycle regulation by Ras and PKB through p27kip1. *Nature* 404 (6779):782-787. doi:10.1038/35008115
85. Kops GJ, Medema RH, Glassford J, Essers MA, Dijkers PF, Coffe PJ, Lam EW, Burgering BM (2002) Control of cell cycle exit and entry by protein kinase B-regulated forkhead transcription factors. *Mol Cell Biol* 22 (7):2025-2036. doi:10.1128/mcb.22.7.2025-2036.2002
86. Collado M, Medema RH, Garcia-Cao I, Dubuisson ML, Barradas M, Glassford J, Rivas C, Burgering BM, Serrano M, Lam EW (2000) Inhibition of the phosphoinositide 3-kinase pathway induces a senescence-like arrest mediated by p27Kip1. *J Biol Chem* 275 (29):21960-21968. doi:10.1074/jbc.M000759200
87. Schmidt M, Fernandez de Mattos S, van der Horst A, Klompaker R, Kops GJ, Lam EW, Burgering BM, Medema RH (2002) Cell cycle inhibition by FoxO forkhead transcription factors involves downregulation of cyclin D. *Mol Cell Biol* 22 (22):7842-7852. doi:10.1128/mcb.22.22.7842-7852.2002
88. Ramaswamy S, Nakamura N, Sansal I, Bergeron L, Sellers WR (2002) A novel mechanism of gene regulation and tumor suppression by the transcription factor FKHR. *Cancer Cell* 2 (1):81-91. doi:10.1016/s1535-6108(02)00086-7
89. Deng S, Chen SN, Yao P, Nikolic D, van Breemen RB, Bolton JL, Fong HH, Farnsworth NR, Pauli GF (2006) Serotonergic activity-guided phytochemical investigation of the roots of *Angelica sinensis*. *J Nat Prod* 69 (4):536-541. doi:10.1021/np050301s
90. Liu PJ, Hsieh WT, Huang SH, Liao HF, Chiang BH (2010) Hematopoietic effect of water-soluble polysaccharides from *Angelica sinensis* on mice with acute blood loss. *Exp Hematol* 38 (6):437-445. doi:10.1016/j.exphem.2010.03.012
91. Yang T, Jia M, Meng J, Wu H, Mei Q (2006) Immunomodulatory activity of polysaccharide isolated from *Angelica sinensis*. *Int J Biol Macromol* 39 (4-5):179-184. doi:10.1016/j.ijbiomac.2006.02.013
92. Shang P, Qian AR, Yang TH, Jia M, Mei QB, Cho CH, Zhao WM, Chen ZN (2003) Experimental study of anti-tumor effects of polysaccharides from *Angelica sinensis*. *World J Gastroenterol* 9 (9):1963-1967. doi:10.3748/wjg.v9.i9.1963

93. Cao W, Li XQ, Wang X, Li T, Chen X, Liu SB, Mei QB (2010) Characterizations and anti-tumor activities of three acidic polysaccharides from *Angelica sinensis* (Oliv.) Diels. *Int J Biol Macromol* 46 (1):115-122. doi:10.1016/j.ijbiomac.2009.11.005
94. Zhao L, Wang Y, Shen HL, Shen XD, Nie Y, Wang Y, Han T, Yin M, Zhang QY (2012) Structural characterization and radioprotection of bone marrow hematopoiesis of two novel polysaccharides from the root of *Angelica sinensis* (Oliv.) Diels. *Fitoterapia* 83 (8):1712-1720. doi:10.1016/j.fitote.2012.09.029
95. Zhang S, He B, Ge J, Li H, Luo X, Zhang H, Li Y, Zhai C, Liu P, Liu X, Fei X (2010) Extraction, chemical analysis of *Angelica sinensis* polysaccharides and antioxidant activity of the polysaccharides in ischemia-reperfusion rats. *Int J Biol Macromol* 47 (4):546-550. doi:10.1016/j.ijbiomac.2010.07.012
96. Ji P, Wei Y, Xue W, Hua Y, Zhang M, Sun H, Song Z, Zhang L, Li J, Zhao H, Zhang W (2014) Characterization and antioxidative activities of polysaccharide in Chinese angelica and its processed products. *Int J Biol Macromol* 67:195-200. doi:10.1016/j.ijbiomac.2014.03.025
97. Xie X, Liu M, Meng Q (2019) *Angelica* polysaccharide promotes proliferation and osteoblast differentiation of mesenchymal stem cells by regulation of long non-coding RNA H19: An animal study. *Bone Joint Res* 8 (7):323-332. doi:10.1302/2046-3758.87.BJR-2018-0223.R2


## Figures



**Figure 1**


5-FU inhibits HS-5 cells proliferation by regulating Wnt/ $\beta$ -catenin signaling pathway. (A) Cell Counting Kit-8 (CCK-8) assay was performed to detect the inhibitory effect of 5-FU. (B) Western blot was performed to detect the effects of 5-FU on the expression of Wnt/ $\beta$ -catenin signaling pathway related proteins in HS-5 cells and the histograms of relative protein expression are presented.  $\beta$ -actin and Lamin B are probed as loading controls. Data are presented as means  $\pm$  SD ( $n=3$ /group) (C) Histologic feature of HS-5 cells treated with Wnt signaling agonist LiCl under inverted microscope (Scale bar=40 $\mu$ m). (D) Cell Counting Kit-8 assay showed the viability HS-5 cell treated with LiCl. HS-5 cell treated with 0 mmol/L LiCl were set

as 1.0, and the results of HS-5 cell viability were normalized to the OD value of 0 mmol/L group (E) HS-5 cells were incubated with 50 ng/mL Wnt signaling inhibitor Dkk1 for 48h and cell viability was detected by CCK-8. HS-5 cell treated without Dkk1 was used as a control, and the results of HS-5 cell viability were normalized to the OD value of control group (\* p < 0.05 versus control)




**Figure 2**

Angelica Sinensis Polysaccharides antagonize the growth inhibitory effect of 5-FU on HS-5 cells via up-regulating Wnt/β-catenin signaling. (A) The proliferative HS-5 cells were labeled by 5-ethynyl-20-deoxyuridine (EdU). The green fluorecence presents proliferative cells, the blue fluorescence presents nuclei (Scale bar =40μm). (B) The percentage of proliferating cells (EdU+) was quantitated using ImageJ software. (C) The viability rate of HS-5 cell was measured by Cell Counting Kit-8. Control group was set as 1.0 (\* p < 0.05 versus control, #p < 0.05 versus 5-FU and \*\* p < 0.05 versus 5-FU+ASP group)



**Figure 3**


Angelica Sinensis Polysaccharides activate Wnt/ $\beta$ -catenin signaling pathway. (A) Expression of  $\beta$ -catenin in HS-5 cells was detected by the immunofluorescence method. The red fluorescence presents  $\beta$ -catenin protein expression, the blue fluorescence presents nuclei (Scale bar=40 $\mu$ m). (B) The Wnt signaling related protein expression in HS-5 cells were detected by the Western blot.  $\beta$ -actin and Lamin B were probed as loading controls. (C-I) The histograms of relative protein expression are presented. Data are presented as means  $\pm$  SD (n=3/group) (\* p < 0.05 versus control and #p < 0.05 versus 5-FU)



**Figure 4**

Angelica Sinensis Polysaccharides reduce 5-FU-induced intracellular oxidative stress. (A) The levels of intracellular reactive oxygen species (ROS) in HS-5 cells were measured by DCFH-DA assay under fluorescence microscope. (B) The mean fluorescence intensity of ROS is quantified and presented by histograms. (C) The results of MDA content in HS-5 cells are presented by histograms. (D) The content of SOD in HS-5 cells are presented by histograms. (E) The results of CAT content in HS-5 cells are presented by histograms. Data were presented as means  $\pm$  SD (n=3/group) (\*  $p < 0.05$  versus control and # $p < 0.05$  versus 5-FU)





**Figure 5**

Angelica Sinensis Polysaccharides ameliorate 5-FU-induced activation of FoxO1. (A) The protein expression levels of FoxO1 and p-FoxO1 in HS-5 cells were detected by the Western blot. The relative protein expression is presented by histograms.  $\beta$ -actin is probed as loading control. (B) The protein expression levels of apoptosis in HS-5 cells were detected by the Western blot. (C-F) The histograms of relative protein expression are presented.  $\beta$ -actin is probed as loading control. (G) Annexin V-FITC/PI

double staining was employed to detect cell apoptosis by flow cytometry and the histogram of apoptosis rate is presented. (H) The protein expression level of p27Kip1 in HS-5 cells was detected by the Western blot and the histogram of p27 Kip1 protein expression is presented. (I) Cell cycle was analyzed by flow cytometry and the histograms of phase distribution are presented. Data were presented as means  $\pm$  SD (n=3/group) (\* p < 0.05 versus control and #p < 0.05 versus 5-FU)




Figure 6

Model of Wnt signaling cascade.