

Biotic resistance or invasional meltdown? Diversity reduces invasibility but not exotic dominance in southern California epibenthic communities

Ryan Amir Beshai (✉ rbeshai@uci.edu)

University of California Irvine <https://orcid.org/0000-0001-5319-2235>

Danny Truong

University of California Irvine

Amy Henry

University of California Irvine

Cascade Sorte

University of California Irvine

Research Article

Keywords: biotic resistance, invasional meltdown, diversity, epibenthic, dominance, community assembly

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-980025/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

High community diversity may either prevent or promote the establishment of exotic species. The biotic resistance hypothesis holds that species-rich communities are more resistant to invasion than species-poor communities due to greater interspecific competition. Conversely, the invasional meltdown hypothesis proposes that greater exotic diversity increases invasibility via facilitative interactions between exotic species. To evaluate the degree to which biotic resistance or invasional meltdown influences marine community structure during the assembly period, we studied the development of marine epibenthic “fouling” communities at two southern California harbors. We found that fewer exotic species established as total and exotic richness increased during community assembly and that this effect remained after accounting for space availability. We also found that changes in exotic abundance decreased over time. Throughout the assembly period, gains in exotic abundance were greatest when space was abundant and richness was low. Altogether, we found greater support for biotic resistance than invasional meltdown, suggesting that both native and exotic species contribute to biotic resistance during early community development. However, this resistance may not reduce the total dominance of exotic species.

Introduction

As the number of established exotic species increases worldwide (Seebens et al. 2020), it is increasingly important to understand the role of community diversity in promoting or hindering future invasions. Two competing invasion hypotheses make diverging predictions about the relationship between community diversity and the likelihood of establishment or spread of exotic species. The biotic resistance hypothesis holds that species-rich communities are more resistant to invasion than species-poor communities due to more efficient resource use, leading to fewer available resources and greater interspecific competition (Elton 1958; Levine and D’Antonio 1999). Conversely, the invasional meltdown hypothesis proposes that greater exotic diversity increases invasibility via facilitative interactions between exotic species, resulting in the increased probability of establishment or spread of exotic species (Simberloff and Von Holle 1999; Simberloff 2006). Here, we evaluated the degree to which biotic resistance or invasional meltdown influence community assembly, particularly the establishment and dominance of exotic species, in a marine epibenthic community.

Invasional meltdown and biotic resistance both have broad support in the literature even though they predict opposing responses of invasive establishment to community diversity; however, relatively few empirical studies have considered both alternate hypotheses. Recent reviews have found that the majority of studies—up to 77%—provide support for invasional meltdown (Jeschke et al. 2012; Braga et al. 2018) and positive relationships between exotic diversity and exotic establishment. Invasional meltdown has been studied across a variety of ecosystems, with empirical evidence coming from terrestrial plant, vertebrate, and invertebrate communities (Relva et al. 2010; Green et al. 2011; Ackerman et al. 2014; Collins et al. 2020), as well as freshwater plants and vertebrates (Ricciardi 2001; Meza-Lopez and Siemann 2015; Crane et al. 2020), and marine invertebrate communities (Grosholz 2005; Freeman et

al. 2016). At the same time, biotic resistance (decreased exotic establishment in more diverse communities) has been demonstrated in terrestrial (Levine et al. 2004; Beaury et al. 2020; Zhang et al. 2020), freshwater (Fitzgerald et al. 2016; Liew et al. 2016), and marine (Stachowicz et al. 1999; Stachowicz and Byrnes 2006; Kimbro et al. 2013; Marraffini and Geller 2015) communities.

Both hypotheses suffer from key knowledge gaps. Evidence for invasional meltdown at the community level remains particularly scarce in marine communities (Braga et al. 2018). For biotic resistance, studies have questioned if negative relationships are in fact driven by diversity and resource availability or are instead driven by spatial heterogeneity (Davis et al. 2005; Peng et al. 2019), productivity (Sandel and Corbin 2010), disturbance regimes (Davis et al. 2000; Sandel and Corbin 2010; Hill and Fischer 2014), or environmental stress gradients (Von Holle 2013). Finally, most studies examining biotic resistance or invasional meltdown have done so in developed or manually-assembled communities. There is a paucity of information about how these processes may influence exotic establishment during community assembly, such as during initial establishment in new habitat or reestablishment following extreme disturbance events.

Marine epibenthic communities are model systems for diversity-invasibility studies (Stachowicz et al. 2002; Stachowicz and Byrnes 2006; Marraffini and Geller 2015) and ideal for testing the role of biotic resistance and invasional meltdown during community assembly. Epibenthic communities occur on hard surfaces, including subtidal rocky reefs and infrastructure such as piers and docks. These communities are typically composed of sessile invertebrates such as ascidians (Phylum Chordata; hereafter referred to as chordates), bryozoans, poriferans, and polychaetes (Lord 2017). Epibenthic communities are ideal model systems for studying invasion dynamics during community assembly due to the high diversity of both native and exotic species (Sellheim et al. 2010; Lord 2017), relatively rapid community development (Sutherland and Karlson 1977), and the presence of a known, strongly-limiting resource: available space (Osman 1977; Sebens 1987). Several studies have highlighted the potential for invasional meltdown to occur in this system (Floerl et al. 2004; Freeman et al. 2016; Gavira-O'Neill et al. 2018). Facilitation within epibenthic communities that could lead to invasional meltdown includes the provisioning of resources such as food (Krüger et al. 2018), shelter (Floerl et al. 2004), additional space (Stachowicz and Byrnes 2006), or settlement cues (Wieczorek and Todd 1997; Dunstan and Johnson 2004). It is also possible that high turnover rates allow for opportunistic colonization by exotic species with life history traits that favor rapid establishment and growth (Sakai et al. 2001; Dunstan and Johnson 2004). Biotic resistance also appears to be important in this ecosystem, with more diverse assemblages preventing invasion due to more consistent occupation of primary space (Osman 1977; Stachowicz et al. 1999, 2002).

We tested the biotic resistance and invasional meltdown hypotheses by examining whether diversity decreased or increased exotic establishment, respectively, during assembly of epibenthic communities at two southern Californian sites. We evaluated: (1) the relationships between richness (total or exotic) and the subsequent establishment of exotic species, (2) the relationships between richness (total and exotic) and subsequent changes in the abundance of exotic species, and (3) trends in exotic dominance over

time. We predicted that if invasional meltdown plays a role in community assembly, we would observe the following: (1) higher exotic richness at time t would lead to greater increases in richness between t and $t + 1$; (2) higher exotic richness would lead to greater relative increases in exotic abundance between times t and $t + 1$; and (3) ratio of exotic to native cover (i.e., exotic dominance) would increase over time. In contrast, if biotic resistance were more important in determining the trajectory of community assembly, we predicted negative relationships between richness of established species and changes in exotic richness and abundance, and a saturation in exotic dominance over time.

Methods

Study locations

We tested the relationship between diversity and exotic species establishment during community assembly in two southern California sites: Huntington Harbor, Huntington Beach, California (33°43'18.90"N, 118° 04'20.60"W) and Newport Harbor, Newport Beach, California (33°37'09.70"N, 117°53'45.90"W). Huntington Harbor (3.63 km²; hereafter referred to as Huntington) is fed by Anaheim Bay and receives runoff from the surrounding urban areas of Long Beach and Huntington Beach. Huntington is composed of five man-made islands and provides numerous hard, artificial settlement spaces. Newport Harbor (3.39 km²; hereafter referred to as Newport) is a natural bay located approx. 20 km south of Huntington; it opens directly into the ocean in Corona del Mar and receives both heavy drainage from San Diego creek and urban runoff (Lambert and Lambert 2003). Both Huntington and Newport experience heavy use from recreational boaters.

Experimental Approach

At each site, we mounted 8 clean, 10 × 10 × 1 cm PVC plastic plates approx. 1 m beneath the surface of the water (2 sites × 8 plates = total of 16 plates). We deployed the plates in August 2018 and allowed communities to assemble over a four-month period until December 2018. During the assembly period, we assessed community development in a nondestructive manner by removing the plates from the water briefly (< 5 min) and photographing the sessile invertebrate cover with a Nikon D90 DSLR camera (with Nikon AF-S DX NIKKOR 18-55mm f/3.5-5.6G VR II lens with the lowest f-stop value and the capture mode set to aperture-priority auto-exposure). Photographs were taken weekly for the first 5 weeks post-deployment and then every 2 weeks through the end of the 18-week assembly period. For our statistical analysis, we removed weeks 1 and 3 and we calculated all of our response variables over two-week intervals for consistency.

We estimated the proportional abundance of each species using point classifications generated by the web-based automated annotation tool, CoralNet (Beijbom et al. 2012, 2015; Beijbom 2015; <https://coralnet.ucsd.edu/>). We began by visually-inspecting our photographs and identifying all species to the lowest possible taxonomic resolution, and we used the species identification list to generate a set of image classification labels. We then submitted our photographs to the software and superimposed a

uniform 30 × 30 point grid (for a total of 900 points) on each of the photographs. We trained the software by allowing it to automatically classify the cover directly underneath each point (as either a species identification, bare primary space, or biofilm) and then confirming or correcting each of the classifications, resulting in increasingly accurate automatic classifications over time. Once we reached 80% classification accuracy, we allowed the tool to run automatically for the remaining photographs. We then conducted a visual inspection of all photos and verified or corrected any misidentified classifications. This semi-automated approach has been demonstrated to reduce classification variability compared to a fully-automated approach (Beijbom et al. 2015). We observed that both native and exotic species frequently settled directly on exposed oyster valves and that oysters never overgrew other organisms, so oyster cover was considered equivalent to available space. We therefore calculated the total available space on each plate as the sum of bare space, biofilm, and oyster cover. Available space averaged $48.07 \pm 2.1\%$ (mean \pm SE) cover across all observations.

For each species (Table 1), we determined its biogeographic status (i.e., native versus exotic) based on its classification in southern California according to the NEMESIS database (Fofonoff et al. 2021; <http://invasions.si.edu/nemesis>). If a species was not present in the database or did not have a record for southern California, we conducted a thorough literature search to find records of the species' native range or consulted expert opinion. When invasion status could not be determined or if the species was of uncertain provenance, we classified the species as either unresolved or cryptogenic *sensu* Carlton (1996). Of 29 total taxa identified on the plates, 1 was classified as unresolved and 1 as cryptogenic; hereafter both groups are jointly referred to as unresolved.

Response Variables

To quantify the effects of diversity on establishment by exotic species, we calculated the change in exotic richness (ΔS) over each of the 2-week intervals as:

$$\Delta S = S_{t+1} - S_t$$

Equation 1)

where S_t represents the richness of exotic species at a given observation, and S_{t+1} represents the exotic richness present in the community at the subsequent observation.

We also calculated proportional changes in abundance between observations as:

$$\Delta A = \frac{A_{t+1} - A_t}{A_t}$$

Equation 2)

where A_t represents the summed proportional abundance of exotic species (i.e., the total proportion of all exotic species out of 900 points) at a given observation, and A_{t+1} represents proportional cover of exotic species at the subsequent observation. For instances in which no cover was observed at A_t but was present in A_{t+1} (i.e., periods in which species' [re]established in the community and for which change in abundance would be undefined), we assigned an initial value of 0.001 (equivalent to coverage of a single point out of our total 900 points). We compared initial values ranging from 0.0001 to 0.001 and found that our choice of value did not affect our conclusions (R. Beshai, unpubl. data).

Finally, we calculated exotic dominance using the ratio of total exotic cover to total native cover at a given observation:

Equation 3)

$$E:N = \frac{A_E}{A_E + A_N}$$

where A_E is the summed proportional cover of all exotic species and A_N is the summed proportional cover of native species. Unresolved species accounted for $3.86 \pm 0.36\%$ cover across all plates and time points, with a maximum of 34.67% at one Huntington observation, and were not included in these calculations.

Statistical analysis

We examined the influence of community diversity metrics (total or exotic richness) on changes in exotic richness using a model averaging approach. We defined a global linear mixed model (LMM) that included explanatory variables of richness (total or exotic, which were assessed separately), available space, site (Huntington or Newport), experimental day (the number of days the community had been assembling), and all two-way interactions between richness, available space, and experimental day; plate was included in the model as a random effect to account for repeated measures. To most accurately estimate both main effects and interactions (Grueber et al. 2011; Harrison et al. 2018) as well as alleviate potential issues arising from multicollinearity (Iacobucci et al. 2016), we centered and standardized our variables. We assessed multicollinearity using variance inflation factors, and found that $VIF < 5.5$ for all global models, which falls within the previously-established standard threshold of 10 (Quinn and Keough 2002; Harrison et al. 2018). We then employed an all-subsets model selection approach on our global model to determine the best supported combination of explanatory variables, ensuring that all candidate models retained the random effect of plate to account for our repeated experimental design. Individual candidate model fits were evaluated using the conditional Akaike information criterion (AICc), and models with $\Delta AICc < 6$ (Symonds and Moussalli 2011; Harrison et al. 2018) were incorporated into a best model set along with the best model; candidate models with $\Delta AICc < 6$ that did not converge were excluded from the best model set. We then averaged our best model set and report the results according to the zero

method (Grueber et al. 2011), where zero is substituted for a variable estimate in models where the parameter is absent.

To assess the relationship between richness metrics and changes in exotic abundance, we again applied our model averaging approach. However, because our change in abundance response had a lower bound of -1 (equal to a 100% loss of exotic cover) but no upper bound, we transformed our abundance response by adding 1.001 to all data points and instead fit a generalized linear mixed model (GLMM) with a Gamma distribution and log link. This model structure was appropriate because our response variable was non-negative, positively skewed, and continuous following the transformation. Our global GLMM was then defined with an explanatory variable structure of richness, available space, site, experimental day, and the two-way interactions between richness, space availability, and experimental day; we again included a random effect of plate.

To determine the effects of time and exotic richness on exotic dominance, we applied a LMM to our calculated E:N ratio values with experimental day, exotic richness, the interaction between day and exotic richness, and site as fixed effects and plate as a random effect. We did not include available space in our model because we were predominantly interested in testing the relationship between exotic dominance and time and how it was influenced by exotic richness. All VIF for this model were < 3 .

Due to the heavy influence of a rainfall event toward the end of the assembly period, all analyses presented here in the main text use data from prior to the rainfall event; all data are shown in our figures, and we provide comparisons of analyses run with the full data set (see Supplementary Appendix). All analyses were conducted in R (R Core Team 2020) using the *lme4* package (Bates et al. 2015) v. 1.1-27 to fit the global models, *MuMIn* v. 1.43.17 (Bartoń 2020) for the model selection and averaging process, and *ggplot2* (Wickham 2016) v. 3.3.5 for data visualization.

Results

Community Structure and Dynamics

We observed a total of 29 species across all plates and sites, with 27 in Huntington and 22 in Newport (Table 1). We identified 19 species as exotic in southern California, 8 as native, and 2 as unresolved. Of the 27 species that recruited in Huntington, 18 (67%) were exotic, 7 were native (26%), and 2 (7%) were unresolved. Biogeographic groups were similarly represented in Newport, with 13 (59%) exotic species, 7 (32%) native species, and 2 (9%) unresolved species recruiting over the assembly period. Across both sites, the majority of exotic cover was the tunicates *Botrylloides diegensis*, *Botryllus schlosseri*, *Ciona robusta*, *Ciona savignyi*, *Diplosoma listerianum*, and *Molgula manhattensis* as well as the bryozoans *Tricellaria inopinata* and *Watersipora subtorquata*, while native cover was driven by the bryozoans *Diaperoecia californica* and *Thalamoporella californica* as well as the bivalve *Ostrea lurida*. The total cover from all other native and exotic species averaged $< 3\%$ across all time points and both sites.

Patterns of community assembly followed a similar pattern between sites. In Huntington, community richness, both total and exotic, increased over time but decreased following a large rainfall event (Fig. 1a, Fig. S1a). A similar pattern was observed in Newport, but soft-bodied chordates comprised a greater proportion of the community and there was a greater reduction in richness following the rainfall event (Fig. 1b, Fig. S1b). In Huntington, exotic cover increased rapidly in the early stages of assembly, primarily driven by recruitment by the early-successional species *D. listerianum* (Edwards and Stachowicz 2010), and declined concomitant with *D. listerianum* die-offs (Fig. 1c, Fig. S1a). The total abundance of exotic species then underwent a second rapid increase following the disappearance of *D. listerianum*, driven by the growth of competitively-dominant chordates (Edwards and Stachowicz 2010), including *B. schlosseri* and *C. savignyi*. Native cover remained relatively stable at low levels throughout the assembly period. In Newport, assembly was similarly characterized by a rapid establishment and spread of exotic species, particularly *B. schlosseri* and *D. listerianum*, but the peak abundance of exotic species occurred approx. 50 days later than in Huntington communities (Fig. 1d, Fig. S1b). Native cover exhibited much greater increases over time in Newport than in Huntington, primarily as a result of greater oyster (*O. lurida*) growth (Fig. 1d; Fig. S1c,d).

Impacts of Richness on Exotic Species

We found that changes in exotic richness were consistently driven by the main effects of richness (at the previous time point) and site over the course of the experiment (Tables S1, S2). Specifically, exotic species established less often or were lost from the communities as richness increased; this pattern was evident considering both total richness (Fig. 2a; $\beta=-1.26$, $p<0.001$) and exotic richness (Fig. 2b; $\beta=-1.27$, $p<0.001$). Additionally, significance of the site parameter in both total richness ($\beta=-1.36$, $p<0.001$) and exotic richness ($\beta=-1.06$, $p=0.003$) models revealed that exotic establishment differed between Newport and Huntington, but that greater richness was always associated with decreased exotic establishment. Space availability did not influence exotic establishment in models considering either total richness (Fig. 3a; $\beta=0.437$, $p=0.072$) or exotic richness ($\beta=0.334$, $p=0.14$).

Changes in the abundance of exotic species were influenced by the main effects of space availability and day, as well as richness \times day and space \times day interactions (Tables S3 & S4). Regardless of whether our model was accounting for total richness ($\beta=0.578$, $p<0.001$) or exotic richness ($\beta=0.577$, $p<0.001$), exotic species exhibited greater growth in communities with ample space than in space-limited communities, where they generally experienced reduced growth or declines in abundance (Fig 3b). Available space fluctuated throughout the assembly period, however, and we observed space \times day interactions in both model sets (Tables S3 & S4), where changes in abundance were greatest early in assembly when space was freely available, and decreased over time as space became more limiting (Fig. S2). We did not find a main effect of richness in either the total richness ($p = 0.07$) or exotic richness ($p=0.14$) model. We did, however, detect a richness \times day interaction in both total richness (Fig 2c; $\beta=-603$, $p<0.001$) and exotic richness (Fig 2d; $\beta=-0.669$, $p<0.001$) models, where changes in abundance were large and positive early in assembly when richness is low and then decreased in magnitude over time as community richness increased, eventually resulting in abundance decreases.

Exotic dominance patterns differed between Huntington and Newport (Fig. 1e,f; Table S5), with greater exotic dominance in Huntington than in Newport throughout the assembly period ($\beta=-0.125$, $p=0.009$). For both sites, exotic dominance generally increased with exotic richness ($\beta=0.104$, $p=0.002$) but did not increase over time ($\beta=0.0106$, $p=0.75$). Rather, in both Huntington and Newport, peak values of exotic dominance occurred synchronous with rapid proliferation of *D. listerianum* and *B. schlosseri* or *W. subtorquata* (Fig. 1e,f; Fig. S1a,b), resulting in a significant day \times exotic richness interaction (Fig. S3; $\beta=-0.159$, $p<0.001$). Though not included in our statistical models, we observed that exotic dominance was impacted by the Day 112 rainfall event at both sites. Generally, soft-bodied chordates declined leading to declines in overall exotic dominance, and in Newport, *D. listerianum* die-offs coincided with a period of dominance by native species.

Discussion

Our results suggest that biotic resistance is more important in structuring communities during the assembly period than invasional meltdown. Increased richness, measured by both total richness and that of exotic species only, was associated with reduced establishment success and population growth of later-arriving exotic species (Fig 2a-d) and did not appear alter community assembly in a way that favored exotic species. Our results are consistent with findings in established and/or manipulated (rather than assembling) communities that suggest that greater community richness reduces invasibility (Stachowicz et al. 1999, 2002; Stachowicz and Byrnes 2006) and that exotic species may also contribute to biotic resistance (Collin and Johnson 2014; Marraffini and Geller 2015). As a result, our study provides evidence that negative interactions between exotic species tend to be more prevalent than positive interactions (Jackson 2015).

Mechanistically, classical theory suggests that that invasibility is heavily determined by resource availability and fluctuations in limiting resources (Elton 1958; Davis et al. 2000; Davis and Pelsor 2001). In epibenthic communities, space is known to be a highly limiting resource, and a more consistent occupation of space has been suggested as a strong deterrent to invasions (Stachowicz et al. 1999, 2002; Stachowicz and Byrnes 2006; Marraffini and Geller 2015). Interestingly, our study found mixed evidence for this, as changes in exotic richness were unrelated to the amount of available space but available space significantly interacted with time to modulate changes in abundance. Epibenthic communities are notably stochastic (Sutherland and Karlson 1977), so it is possible that changes in richness were unrelated to resource availability because the space vacated by transient exotic species was rapidly filled through the arrival of secondary exotic species (Dunstan and Johnson 2004) or because high cover may limit, but not entirely prevent, establishment if exotic species are able to settle on other species (Lord 2017). In contrast, changes in exotic abundance are limited by overgrowth and space occupation by neighboring species (Sebens 1987; Lord and Whitlatch 2015; Lord 2017). The greatest increases in exotic abundance occurred in Huntington as a result of rapid growth by *D. listerianum*. At this point in development, few other exotic species were present in the communities and space was freely available, so all growth by *D. listerianum* contributed to the area covered by exotic species. In comparison, the later spread of *D. listerianum* in Newport contributed less to the total abundance of exotic species,

as *D. listerianum* began to compete with and overgrow other exotic species. Thus, while the total abundance of exotic species generally increased over time, relative increases diminished.

Despite the influence of community richness and limiting resources, our results suggest that biotic resistance may not influence ultimate exotic dominance. Instead, we found that exotic dominance was generally high over the course of our experiment. High exotic dominance is likely driven by several interacting factors. First, exotic species tend to display life-history characteristics that allow for rapid growth, maturation, and the production of large numbers of offspring, allowing them to efficiently establish and spread among recipient communities (Sakai et al. 2001; Theoharides and Dukes 2007). Thus, as exotic species establish, they may be able to consume even scarce resources and preempt native species. Second, a sampling effect might allow one or two most impactful exotic species to disproportionately influence community structure during community assembly. The greatest fluctuations in exotic dominance occurred during boom-bust periods of the tunicates *D. listerianum* and *B. schlosseri*. Such dominance by a few exotic species may be particularly apparent in frequently-disturbed or newly-assembling communities due to the combined release or creation of resources (Davis et al. 2000; Strayer et al. 2017) and the “pioneering” characteristics of exotic species (Sakai et al. 2001). Notably, we observed that in Huntington, high exotic dominance was maintained following *D. listerianum*’s “bust” phase via the proliferation of later-successional exotic species (e.g., *B. schlosseri*) (Edwards and Stachowicz 2010), indicative of “over-invasion” (i.e., the replacement of an established exotic species by a second, competitively-dominant exotic; sensu Russell et al. 2014) limiting the establishment or recovery of native species (Strayer et al. 2017). Third, increased propagule pressure is known to drive invasion success (Lockwood et al. 2005; Simkanin et al. 2017; Cassey et al. 2018), and high exotic abundance in Huntington may be sustained by high regional propagule pressure from within and between adjacent ports. Several studies have connected high shipping intensity with increased rates of exotic establishment (Lord and Whitlatch 2015; Lacoursière-Roussel et al. 2016), and, when strong enough, propagule pressure may allow species to bypass biotic resistances (Hollebone and Hay 2007). Huntington lies < 20 km from the Ports of Los Angeles and Long Beach, two of the busiest ports in the United States (Hu et al. 2021), and is likely receiving high regional propagule dispersal, whereas with a separation of > 30km, it is possible that Newport experiences lower regional propagule pressure; both sites, however, are exposed to high local propagule pressure from established communities on adjacent pilings and docks.

Despite the potential for biotic resistance to be greater in established communities, our study provides evidence for biotic resistance in assembling communities, which may influence ultimate community structure. Previous research has shown that in epibenthic communities, larger organisms are competitively dominant over smaller organisms (Buss 1979), and that more consistent utilization of available space—a product of greater diversity—reduces invasibility (Stachowicz et al. 2002). Together, these suggest that biotic resistance would be more likely to occur in older, more diverse communities containing larger organisms than in early-assembly communities that are relatively low-diversity and where species are likely to be initially similar in size. However, our study demonstrates the possibility for negative interactions to limit increases in exotic abundance during the assembly period, largely through

space competition. The existence of competitive interactions is of particular importance during assembly, as priority effects from early-arrivals can influence ultimate community composition (Sutherland 1974; Connell and Slatyer 1977; Vieira et al. 2018a) and invasibility (Gerla and Mooij 2014). For example, Viera et al. (2018b) found that low-diversity assemblages founded by encrusting bryozoans or colonial tunicates, such as *D. listerianum*, were typically dominated by space monopolizers later in development, whereas more diverse communities founded by arborescent bryozoans tended to result in more diverse communities later in assembly. It is possible that rapid establishment of *D. listerianum* may have set our communities on a trajectory that led to reduced diversity of all species, including exotics.

There are several potential explanations for why we did not observe invasional meltdown occurring in our experiment. First, published examples of invasional meltdown typically document exotic species ameliorating conditions in recipient habitats to make them less hostile to subsequent exotic establishment or acting as disturbances in the recipient communities, and freeing up resources (Simberloff and Von Holle 1999; Simberloff 2006). Facilitation among sessile organisms, however, likely influences exotic and native species equally. For example, *W. subtorquata*, which was abundant in Huntington, has been implicated in invasional meltdown elsewhere because it increases habitat availability (Stachowicz and Byrnes 2006; Scott and terHorst 2020) or acts as a settlement refuge in toxic environments (Floerl et al. 2004), thereby promoting exotic species richness (Floerl et al. 2004; Scott and terHorst 2019; but see Scott and terHorst 2020). Newly created biogenic settlement space may not inherently favor settlement by exotic species over natives, so the presence of foundation species may be more important for increasing overall richness than facilitating exotic settlement per se. The presence of foundation species such as *W. subtorquata* may be more important for increasing the richness of mobile than sessile species (Sellheim et al. 2010; Gavira-O'Neill et al. 2018), and we did not record mobile species in our study. Second, habitat modifications made by a particular species may not fully compensate for the resources that it consumes. Epibenthic species have been documented to settle on top of established adult chordates (*Ascidia ceratodes*), but individuals settling on *Ascidia* tended to exhibit reduced performance (in terms of growth rate) compared to settlers on primary substrate (Claar et al. 2011). Third, allelopathic interactions are rarely documented in marine communities (Granéli and Pavia 2006); thus, overgrowth competition, which consumes primary space but does not actively displace adjacent organisms, acts as the primary antagonistic interaction between epibenthic exotics. This limits the potential for exotic species to alleviate space restrictions for subsequently-arriving exotic species by acting as disturbances and instead serves to increase competitive interactions between exotic species. Finally, several of the exotic species present in our study (e.g., *B. shlosseri*, *C. robusta* [previously identified as *C. intestinalis*], and *S. plicata*) are believed to have been introduced in the early 1900s (Lambert and Lambert 1998) and have likely become naturalized (Richardson et al. 2000). Therefore, it is possible that these species are now “behaving like the natives” during assembly (Pearson et al. 2018), where a lack of “novel” biotic interactions between natives and exotics allows assembly patterns to be determined by recruitment patterns and life history trade-offs (Edwards and Stachowicz 2010, 2012), regardless of whether a species is native or exotic.

Biotic and abiotic factors not examined here are known to influence epibenthic community structure and composition. While studies have demonstrated that predation may have strong impacts on epibenthic community composition (Vieira et al. 2012; Oricchio et al. 2016; Freestone et al. 2021), its role in modulating exotic establishment is debated. In epibenthic communities, predation might serve to limit the establishment (Dumont et al. 2011; Collin and Johnson 2014; Rogers et al. 2016; Kremer and da Rocha 2016; Leclerc et al. 2020) or dominance of exotic organisms, as predators tend to target poorly-defended, soft-bodied chordates (Kremer and da Rocha 2016), which were characteristics of the species that we found to be primary drivers of changes in exotic abundance and dominance. Thus, predation could alter the succession sequence by suppressing rapidly growing initial colonizers (Nydam and Stachowicz 2007), which could allow for greater initial growth of native species, though this effect (which we did not observe) is not consistently documented (e.g., Freeman et al. 2016). Epibenthic communities are also known to be sensitive to changes in abiotic conditions, such as fluctuating salinity (Chang et al. 2018). We observed rapid declines in the abundance of chordates at both harbors following a heavy rainfall event, and these effects were particularly pronounced in Newport (Fig. 1c,d), where chordates constituted a relatively greater proportion of the overall community (Fig. S1). Similar effects have been observed in the past, where large, rainfall-driven fluctuations in salinity have been associated with dramatic changes in chordate populations (MacGinitie 1939; Lambert and Lambert 2003; Nydam and Stachowicz 2007). While still exhibiting a dramatic change in the total abundance of invasive species, the decline in exotic abundance in Huntington was less dramatic due to the presence of the bryozoan *W. subtorquata*, which is known to be widely tolerant to both temperature and salinity fluctuations (Afonso et al. 2020). The heavy impacts of environmental disturbance on epibenthic community structure, combined with naturally high turnover rates (Sutherland and Karlson 1997) and the creation of new habitat structures, mean that natural succession patterns in epibenthic community [re]assembly are frequently reset and suggest that it is particularly important to understand processes influencing the establishment and dominance of exotic species during these [re]assembly periods.

In conclusion, our results are more consistent with our predictions of biotic resistance than invasional meltdown, suggesting that biotic resistance plays a role in limiting the establishment and spread of exotic species during community assembly. In heavily-invaded systems such as those studied here, resistance may not reduce the ultimate dominance of exotic species, but does not necessarily lead to meltdown. Community-level properties such as biotic resistance are known to have scale and context-dependent strengths (Fridley et al. 2007; Kimbro et al. 2013). Therefore, future research should examine whether resistance is able to limit exotic dominance along broader scales and environmental gradients.

Statements And Declarations:

Competing Interests: The authors declare no conflicts of interest.

Availability of data and material: The dataset analyzed during the current study will available in the Dryad repository upon publication of this manuscript.

Code availability: Code will be uploaded to Dryad upon publication of this manuscript.

Authors' contributions: DAT, AKH, and CJBS designed the assembly experiment. DAT collected the data. RAB and CJBS generated the hypotheses. RAB, AKH, and CJBS designed the analyses. RAB conducted the statistical analysis and wrote the manuscript. All authors revised drafts and approved the final manuscript.

Funding: Funding for this study was provided to DAT through University of California, Irvine's Undergraduate Research Opportunities Program (UROP). Funding was also provided to CJBS through the Belmont Forum and BioivERsA joint call for research proposals, under the BiodivScen ERA-Net COFUND program, with support from the National Science Foundation (ICER-1852060).

Ethics approval: Not applicable

Consent to participate: Not applicable

Consent for publication: Not applicable

Acknowledgements:

We would like to thank Roy Pizarek, Susan Emerson, Kim Henry, and Susan Henry for providing site access. We thank Jim Carlton for assistance with species identification and provenance as well as Kailen Mooney and Celia Symons for providing feedback that greatly improved this paper. We would also like to acknowledge that this study was conducted on the unceded ancestral homelands of the Tongva people.

References

Ackerman JD, Falcón W, Molinari J, Vega C, Espino I, Cuevas AA (2014). Biotic resistance and invasional meltdown: consequences of acquired interspecific interactions for an invasive orchid, *Spathoglottis plicata* in Puerto Rico. *Biol Invasions*, 16: 2435-2447. <https://doi.org/10.1007/s10530-014-0676-3>

Afonso I, Bercibar E, Castro N, Costa JL, Frias P, Henriques F, Moreira P, Oliveira PM, Silva G, Chainho P (2020) Assessment of the colonization and dispersal success of non-indigenous species introduced in recreational marinas along the estuarine gradient. *Ecol Indic* 113:106147. <https://doi.org/10.1016/j.ecolind.2020.106147>

Bartoń K (2020) MuMIn: Multi-Model Inference. Version 1.43.17URL <https://CRAN.R-project.org/package=MuMIn>

Bates D, Mäechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models using lme4. *J Stat Softw* 67:1–48. <https://doi.org/10.18637/jss.v067.i01>

Beaury EM, Finn JT, Corbin JD, Barr V, Bradley BB (2020) Biotic resistance to invasion is ubiquitous across ecosystems of the United States. *Ecol Lett* 23:476–482. <https://doi.org/10.1111/ele.13446>

- Beijbom O (2015) Automated Annotation of Coral Reef Survey Images. Dissertation, University of California
- Beijbom O, Edmunds PJ, Kline DI, Mitchell BG, Kriegman D (2012) Automated annotation of coral reef survey images. In: 2012 IEEE Conference on Computer Vision and Pattern Recognition. doi:10.1109/CVPR.2012.6247798
- Beijbom O, Edmunds PJ, Roelfsema C, Smith J, Kline DI, Neal BP, Dunlap MJ, Moriarty V, Fan TY, Tan CJ, Chan S, Treibitz T, Gamst A, Mitchell BG, Kriegman D (2015) Towards Automated Annotation of Benthic Survey Images: Variability of Human Experts and Operational Modes of Automation. PLoS One 10:e0130312. <https://doi.org/10.1371/journal.pone.0130312>
- Braga RR, Gómez-Aparicio L, Heger T, Vitule JRS, Jeschke J (2018) Structuring evidence for invasional meltdown: broad support but with biases and gaps. *Biol Invasions* 20:923–936. <https://doi.org/10.1007/s10530-017-1582-2>
- Buss LW (1979) Bryozoan overgrowth interactions—the interdependence of competition for space and food. *Nat* 281:475–477. <https://doi.org/10.1038/281475a0>
- Cassey P, Delean S, Lockwood JL, Sadowski JS, Blackburn TM (2018) Dissecting the null model for biological invasions: A meta-analysis of the propagule pressure effect. *PLOS Biol* 16:e2005987. <https://doi.org/10.1371/journal.pbio.2005987>
- Chang AL, Brown CW, Crooks JA, Ruiz GM (2018) Dry and wet periods drive rapid shifts in community assembly in an estuarine ecosystem. *Glob Chang Biol* 24:e627–e642. <https://doi.org/10.1111/gcb.13972>
- Claar DC, Edwards KF, Stachowicz JJ (2011) Positive and negative effects of a dominant competitor on the settlement, growth, and survival of competing species in an epibenthic community. *J Exp Mar Biol Ecol* 399:130–134. <https://doi.org/10.1016/j.jembe.2011.02.014>
- Collin SB, Johnson LE (2014) Invasive species contribute to biotic resistance: negative effect of caprellid amphipods on an invasive tunicate. *Biol Invasions* 16:2209–2219. DOI 10.1007/s10530-014-0659-4
- Collins RJ, Copenheaver CA, Barney JN, Radtke PJ (2020) Using Invasional Meltdown Theory to Understand Patterns of Invasive Richness and Abundance in Forests of the Northeastern USA. *Nat Areas J* 40:336–344. <https://doi.org/10.3375/043.040.0406>
- Connell JT, Slatyer RO (1977) Mechanisms of Succession in Natural Communities and Their Role in Community Stability and Organization. *Am Nat* 111:1119–1144. <https://doi.org/10.1086/283241>
- Crane K, Coughlan NE, Cuthbert RN, Dick JTA, Kregting L, Ricciardi A, Maclsaac H, Reid N (2020) Friends of mine: An invasive freshwater mussel facilitates growth of invasive macrophytes and mediates their competitive interactions. *Freshw Biol* 65:1063–1072. <https://doi.org/10.1111/fwb.13489>

- Davis MA, Grime JP, Thompson K (2000) Fluctuating resources in plant communities: a general theory of invasibility. *J Ecol* 88:528–534. <https://doi.org/10.1046/j.1365-2745.2000.00473.x>
- Davis MA, Pelsor M (2001) Experimental support for a resource-based mechanistic model of invasibility. *Ecol Lett* 4:421–428. <https://doi.org/10.1046/j.1461-0248.2001.00246.x>
- Davis MA, Thompson K, Grime JP, Liebhold A (2005) Invasibility: The Local Mechanism Driving Community Assembly and Species Diversity. *Ecography* 28:696–704
- DeRivera CE, Ruiz GM, Hines AH, Jivoff P (2005) Biotic resistance to invasion: Native predator limits abundance and distribution of an introduced crab. *Ecol* 86:3364–3376. <https://doi.org/10.1890/05-0479>
- Dumont CP, Gaymer CF, Thiel M (2011) Predation contributes to invasion resistance of benthic communities against the non-indigenous tunicate *Ciona intestinalis*. *Biol Invasions* 13:2023–2034. <https://doi.org/10.1007/s10530-011-0018-7>
- Dunstan PK, Johnson CR (2004) Invasion rates increase with species richness in a marine epibenthic community by two mechanisms. *Oecologia* 138:285–292. <https://doi.org/10.1007/s00442-003-1400-7>
- Edwards KF, Stachowicz JJ (2010) Multivariate trade-offs, succession, and phenological differentiation in a guild of colonial invertebrates. *Ecol* 91:3146–3152. <https://doi.org/10.1890/10-0440.1>
- Edwards KF, Stachowicz JJ (2012) Temporally varying larval settlement, competition, and coexistence in a sessile invertebrate community. *Mar Ecol Prog Ser* 462:93–102. <https://doi.org/10.3354/meps09828>
- Elton CS (1958) *The Ecology of Invasions by Animals and Plants*. Methuen, London
- Fitzgerald DB, Tobler M, Winemiller KO (2016) From richer to poorer: successful invasion by freshwater fishes depends on species richness of donor and recipient basins. *Glob Chang Biol* 22:2440–2450. <https://doi.org/10.1111/gcb.13165>
- Floerl O, Pool TK, Inglis GJ (2004) Positive Interactions Between Nonindigenous Species Facilitate Transport by Human Vectors. *Ecol Appl* 14:1724–1736. <https://doi.org/10.1890/03-5399>
- Fofonoff PW, Ruiz GM, Simkanin C, Carlton JT (2021) National Exotic Marine and Estuarine Species Information System. <http://invasions.si.edu/nemesis>. Access Date: 2021-04-27
- Freeman AS, Frischeisen A, Blakeslee AMH (2016) Estuarine fouling communities are dominated by nonindigenous species in the presence of an invasive crab. *Biol Invasions* 18:1653–1665. <https://doi.org/10.1007/s10530-016-1108-3>
- Freestone AL, Torchin ME, Jurgens LJ, Bonfim M, López DP, Repetto MF, Schlöder C, Sewall BJ, Ruiz GM (2021) Stronger predation intensity and impact on prey communities in the tropics. *Ecol* 102:e03428. <https://doi.org/10.1002/ecy.3428>

Fridley JD, Stachowicz JJ, Naeem S, Sax DF, Seabloom EW, Smith MD, Stohlgren TJ, Tilman D, Von Holle B (2007) The Invasion Paradox: Reconciling Pattern and Process in Species Invasions. *Ecol* 88:3–17. [https://doi.org/10.1890/0012-9658\(2007\)88\[3:TIPRPA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88[3:TIPRPA]2.0.CO;2)

Gavira-O'Neill K, Guerra-García JM, Moreira J, Ros M (2018) Mobile epifauna of the invasive bryozoan *Tricellaria inopinata*: is there a potential invasional meltdown? *Mar Biodivers* 48:1169–1178. <https://doi.org/10.1007/s12526-016-0563-5>

Gerla DJ, Mooij WM (2014) Alternative stable states and alternative endstates of community assembly through intra- and interspecific positive and negative interactions. *Theor Popul Biol* 96:8–18. <https://doi.org/10.1016/j.tpb.2014.07.001>

Granéli E, Pavia H (2006) Allelopathy in marine ecosystems. In: Reigosa MJ, Pedrol N, González L (eds) *Allelopathy*. Springer, Dordrecht, pp 415–431

Green PT, O'Dowd DJ, Abbott KL, Jeffery M, Retallick K, Mac Nally R (2011) Invasional meltdown: Invader-invader mutualism facilitates a secondary invasion. *Ecol* 92:1758–1768. <https://doi.org/10.1890/11-0050.1>

Grosholz ED (2005) Recent biological invasion may hasten invasional meltdown by accelerating historical introductions. *PNAS* 102:1088–1091. <https://doi.org/10.1073/pnas.0308547102>

Grueber CE, Nakagawa S, Laws RJ, Jamieson IG (2011) Multimodel inference in ecology and evolution: challenges and solutions. *J Evol Biol* 24:699–711. <https://doi.org/10.1111/j.1420-9101.2010.02210.x>

Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED, Robinson BS, Hodgson DJ, Inger R (2018) A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* 6:e4794. <https://doi.org/10.7717/peerj.4794>

Hill KC, Fischer DG (2014) Native–Exotic Species Richness Relationships Across Spatial Scales in a Prairie Restoration Matrix. *Restor Ecol* 22:204–213. <https://doi.org/10.1111/rec.12067>

Hollebone AL, Hay ME (2007) Propagule pressure of an invasive crab overwhelms native biotic resistance. *Mar Ecol Prog Ser* 342:191–196. <https://doi.org/10.3354/meps342191>

Hu PS, et al (2021) U.S. Department of Transportation, Bureau of Transportation Statistics, Port Performance Freight Statistics Annual Report to Congress 2020 (Washington, DC: 2020). <https://doi.org/10.21949/1520450>

Iacobucci D, Schneider MJ, Popovich DL, Bakamitsos GA (2016) Mean centering helps alleviate “micro” but not “macro” multicollinearity. *Behav Res Methods* 48:1308–1317. <https://doi.org/10.3758/s13428-015-0624-x>

- Jackson MC (2015) Interactions among multiple invasive animals. *Ecol* 96:2035–2041. <https://doi.org/10.1890/15-0171.1>
- Jeschke JM, Gómez Aparicio L, Haider S, Heger T, Lortie CJ, Pyšek P, Strayer DL (2012) Support for major hypotheses in invasion biology is uneven and declining. *Neobiota* 14:1-20. <https://doi.org/10.3897/neobiota.14.3435>
- Kimbro DL, Cheng BS, Grosholz ED (2013) Biotic resistance in marine environments. *Ecol Lett* 16:821–833. <https://doi.org/10.1111/ele.12106>
- Kremer LP, da Rocha RM (2016) The biotic resistance role of fish predation in fouling communities. *Biol Invasions* 18:3223–3237. <https://doi.org/10.1007/s10530-016-1210-6>
- Krüger I, Lenz M, Thiel M (2018) The colonial ascidian *Diplosoma listerianum* enhances the occurrence of the hydrozoan *Obelia* sp. during early phases of succession. *Helgol Mar Res* 72:4. <https://doi.org/10.1186/s10152-018-0506-6>
- Lacoursière-Roussel A, Bock DG, Cristescu ME, Guichard F, McKindsey CW (2016) Effect of shipping traffic on biofouling invasion success at population and community levels. *Biol Invasions* 18:3681–3695. <https://doi.org/10.1007/s10530-016-1258-3>
- Lambert C, Lambert G (2003) Persistence and differential distribution of nonindigenous ascidians in harbors of the Southern California Bight. *Mar Ecol Prog Ser* 259:145–161. <https://doi.org/10.3354/meps259145>
- Leclerc J-C, Viard F, Brante A (2020) Experimental and survey-based evidences for effective biotic resistance by predators in ports. *Biol Invasions* 22:339–352. <https://doi.org/10.1007/s10530-019-02092-9>
- Levine JM, Adler PB, Yelenik SG (2004) A meta-analysis of biotic resistance to exotic plant invasions. *Ecol Lett* 7:975–989. <https://doi.org/10.1111/j.1461-0248.2004.00657.x>
- Levine JM, D'Antonio CM (1999) Elton Revisited: A Review of Evidence Linking Diversity and Invasibility. *Oikos* 87:15–26. <https://doi.org/10.2307/3546992>
- Liew JH, Carrasco LR, Tan HH, Yeo DCJ (2016) Native richness and species level trophic traits predict establishment of alien freshwater fishes. *Biol Invasions* 18:3495–3512. <https://doi.org/10.1007/s10530-016-1241-z>
- Lockwood JL, Cassey P, Blackburn T (2005) The role of propagule pressure in explaining species invasions. *TREE* 20:223–228. <https://doi.org/10.1016/j.tree.2005.02.004>
- Lord JP, Whitlatch R (2015) Predicting competitive shifts and responses to climate change based on latitudinal distributions of species assemblages. *Ecol* 96:1264–1274. <https://doi.org/10.1890/14-0403.1>

- Lord JP (2017) Temperature, space availability, and species assemblages impact competition in global fouling communities. *Biol Invasions* 19:43–55. <https://doi.org/10.1007/s10530-016-1262-7>
- MacGinitie GE (1939) Some Effects of Fresh Water on the Fauna of a Marine Harbor. *Am Midl Nat* 21:681–686. <https://doi.org/10.2307/2420525>
- Marraffini ML, Geller JB (2015) Species richness and interacting factors control invasibility of a marine community. *Proc Royal Soc B* 282:20150439. <https://doi.org/10.1098/rspb.2015.0439>
- Meza-Lopez MM, Siemann E (2015) Experimental test of the Invasional Meltdown Hypothesis: an exotic herbivore facilitates an exotic plant, but the plant does not reciprocally facilitate the herbivore. *Freshw Biol* 60:1475–1482. <https://doi.org/10.1111/fwb.12582>
- Nydam M, Stachowicz J (2007) Predator effects on fouling community development. *Mar Ecol Prog Ser* 337:93–101. <https://doi.org/10.3354/meps337093>
- Oricchio FT, Flores AAV, Dias GM (2016) The importance of predation and predator size on the development and structure of a subtropical fouling community. *Hydrobiol* 776:209–219. <https://doi.org/10.1007/s10750-016-2752-4>
- Osman RW (1977) The Establishment and Development of a Marine Epifaunal Community. *Ecol Monogr* 47:37–63. <https://doi.org/10.2307/1942223>
- Pasch B, Bolker BM, Phelps SM (2013) Interspecific Dominance Via Vocal Interactions Mediates Altitudinal Zonation in Neotropical Singing Mice. *Am Nat* 182:E161–E173. <https://doi.org/10.1086/673263>
- Pearson DE, Ortega YK, Eren Ö, Hierro JL (2018) Community Assembly Theory as a Framework for Biological Invasions. *TREE* 33:313–325. <https://doi.org/10.1016/j.tree.2018.03.002>
- Peng S, Kinlock NL, Gurevitch J, Peng S (2019) Correlation of native and exotic species richness: a global meta-analysis finds no invasion paradox across scales. *Ecol* 100:e02552. <https://doi.org/10.1002/ecy.2552>
- Quinn GP, Keough MJ (2002) *Experimental Design and Data Analysis for Biologists*. Cambridge University Press, Cambridge
- R Core Team (2020) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria
- Relva MA, Nuñez MA, Simberloff D (2010) Introduced deer reduce native plant cover and facilitate invasion of non-native tree species: evidence for invasional meltdown. *Biol Invasions* 12:303–311. <https://doi.org/10.1007/s10530-009-9623-0>

Ricciardi A (2001) Facilitative interactions among aquatic invaders: is an “invasional meltdown” occurring

in the Great Lakes? *Can J Fish Aquat* 58:2513-2525. <https://doi.org/10.1139/f01-178>

Richardson DM, Pyšek P, Rejmánek M, Rejmánek M, Barbour MG, Panetta FD, West CJ (2000) Naturalization

and invasion of alien plants: concepts and definitions. *Divers Distrib* 6:93–107.

<https://doi.org/10.1046/j.1472-4642.2000.00083.x>

Rogers T, Byrnes J, Stachowicz J (2016) Native predators limit invasion of benthic invertebrate

communities in Bodega Harbor, California. *Mar Ecol Prog Ser* 545:161-173.

<https://doi.org/10.3354/meps11611>

Russell JC, Sataruddin NS, Heard AD (2014) Over-invasion by functionally equivalent invasive species.

Ecol 95:2268–2276. <https://doi.org/10.1890/13-1672.1>

Sakai AK, Allendorf FW, Holt JS, Lodge DM, Molofsky J, With KA, Baughman S, Cabin RJ, Cohen JE, Ellstrand

NC, McCauley DE, O’Neil P, Parker IM, Thompson JN, Weller SG (2001) The Population Biology of Invasive Species. *Annu Rev Ecol Evol Syst* 32:305–332.

<https://doi.org/10.1146/annurev.ecolsys.32.081501.114037>

Sandel B, Corbin JD (2010) Scale, disturbance and productivity control the native-exotic richness

relationship. *Oikos* 119:1281–1290. <https://doi.org/10.1111/j.1600-0706.2010.18230.x>

Scott ZR, terHorst CP (2019) The effect of an invasive bryozoan on community diversity and structure

varies across two locations. *Community Ecol* 20:258–265. <https://doi.org/10.1556/168.2019.20.3.6>

Scott ZR, terHorst CP (2020) The effect of an invasive foundation species on diversity is due to increased

habitat availability. *J Exp Mar Biol Ecol* 528:151384. <https://doi.org/10.1016/j.jembe.2020.151384>

Sellheim K, Stachowicz JJ, Coates RC (2010) Effects of a nonnative habitat-forming species on mobile and sessile epifaunal communities. *Mar Ecol Prog Ser* 398:69–80.

<https://doi.org/10.3354/meps08341>

Simberloff D (2006) Invasional meltdown 6 years later: important phenomenon, unfortunate metaphor, or

both? *Ecol Lett* 9:912–919. <https://doi.org/10.1111/j.1461-0248.2006.00939.x>

Simberloff D, Von Holle B (1999) Positive interactions of nonindigenous species. *Biol Invasions* 1:21–32

Simkanin C, Davidson IC, Therriault TW, Jamieson G, Dower JF (2017) Manipulating propagule pressure to test

the invasibility of subtidal marine habitats. *Biol Invasions* 19:1565–1575.

<https://doi.org/10.1007/s10530-017-1379-3>

Stachowicz J, Byrnes J (2006) Species diversity, invasion success, and ecosystem functioning:

disentangling the influence of resource competition, facilitation, and extrinsic factors. *Mar Ecol Prog Ser* 311:251–262. <https://doi.org/10.3354/meps311251>

Stachowicz JJ, Fried H, Osman RW, Whitlatch RB (2002) Biodiversity, Invasion Resistance, and Marine Ecosystem Function: Reconciling Pattern and Process. *Ecol* 83:2575–2590.

[https://doi.org/10.1890/0012-9658\(2002\)083\[2575:BIRAME\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[2575:BIRAME]2.0.CO;2)

Stachowicz JJ, Whitlatch RB, Osman RW (1999) Species diversity and invasion resistance in a marine ecosystem. *Sci* 286:1577–1579. <https://doi.org/10.1126/science.286.5444.1577>

Strayer DL, D'Antonio CM, Essl F, Fowler MS, Geist J, Hilt S, Jarić I, Jöhnk K, Jones CG, Lambin X, Latzka AW,

Pergl J, Pyšek P, Robertson P, von Schmalensee M, Stefansson RA, Wright J, Jeschke JM (2017) Boom-bust dynamics in biological invasions: towards an improved application of the concept. *Ecol Lett* 20:1337–1350. <https://doi.org/10.1111/ele.12822>

Sutherland JP (1974) Multiple Stable Points in Natural Communities. *Am Nat* 108:859–873.

<https://doi.org/10.1086/282961>

Sutherland JP, Karlson RH (1977) Development and Stability of the Fouling Community at Beaufort, North Carolina. *Ecol Monogr* 47:425–446. <https://doi.org/10.2307/1942176>

Symonds MRE, Moussalli A (2011) A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behav Ecol Sociobiol* 65:13–21.

<https://doi.org/10.1007/s00265-010-1037-6>

Theoharides KA, Dukes JS (2007) Plant invasion across space and time: Factors affecting nonindigenous species success during four stages of invasion. *New Phytol* 176:256–273.

<https://doi.org/10.1111/j.1469-8137.2007.02207.x>

- Vieira EA, Duarte LFL, Dias GM (2012) How the timing of predation affects composition and diversity of species in a marine sessile community? *J Exp Mar Biol Ecol* 412:126–133.
<https://doi.org/10.1016/j.jembe.2011.11.011>
- Vieira EA, Flores AAV, Dias GM (2018a) Persistence and space preemption explain species-specific founder effects on the organization of marine sessile communities. *Ecol Evol* 8:3430–3442.
<https://doi.org/10.1002/ece3.3853>
- Vieira EA, Dias GM, Flores AAV (2018b) Adding early-stage engineering species affects advanced-stage organization of shallow-water fouling assemblages. *Hydrobiol* 818:211-222.
<https://doi.org/10.1007/s10750-018-3612-1>
- Von Holle B (2013) Environmental stress alters native-nonnative relationships at the community scale. *Biol Invasions* 15:417–427. <https://doi.org/10.1007/s10530-012-0297-7>
- Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*. Version 3.3.5. Springer-Verlag New York
- Wieczorek SK, Todd CD (1997) Inhibition and facilitation of bryozoan and ascidian settlement by natural multi-species biofilms: effects of film age and the roles of active and passive larval attachment. *Mar Biol* 128:463–473. <https://doi.org/10.1007/s002270050113>
- Zhang Z, Liu Y, Brunel C, van Kleunen M (2020) Evidence for Elton’s diversity–invasibility hypothesis from belowground. *Ecol* 101:e03187. <https://doi.org/10.1002/ecy.3187>

Tables

Table 1 Taxa identified on our plates. Taxa were identified to the lowest possible level (species or genus)

Phylum	Name	Biogeographic Status	Presence (Y/N)	
			Huntington	Newport
Porifera	<i>Halichondria bowerbanki</i>	Exotic	Y	Y
Cnidaria	<i>Anthopleura artemisia</i>	Native	Y	Y
Mollusca	<i>Ostrea lurida</i>	Native	Y	Y
Bryozoa	<i>Celleporella hyalina</i>	Unresolved	Y	Y
Bryozoa	<i>Amathia verticillata</i>	Exotic	Y	N
Bryozoa	<i>Cryptosula pallasiana</i>	Exotic	Y	N
Bryozoa	<i>Schizoporella</i> spp.	Exotic	Y	Y
Bryozoa	<i>Tricellaria inopinata</i>	Exotic	N	Y
Bryozoa	<i>Watersipora subtorquata</i>	Exotic	Y	Y
Bryozoa	<i>Bugula californica</i>	Native	Y	Y
Bryozoa	<i>Diaperoecia californica</i>	Native	Y	Y
Bryozoa	<i>Membranipora</i> spp.	Native	N	Y
Bryozoa	<i>Thalamoporella californica</i>	Native	Y	Y
Polychaeta	<i>Demonax</i> sp.	Exotic	Y	N
Polychaeta	<i>Hydroides elegans</i>	Exotic	Y	Y
Polychaeta	<i>Spirorbis</i> sp.	Unresolved	Y	Y
Chordata	<i>Ascidia zara</i>	Exotic	Y	N
Chordata	<i>Botrylloides diegensis</i>	Exotic	Y	Y
Chordata	<i>Botrylloides violaceus</i>	Exotic	Y	Y
Chordata	<i>Botryllus schlosseri</i>	Exotic	Y	Y
Chordata	<i>Ciona robusta</i>	Exotic	Y	N
Chordata	<i>Ciona savignyi</i>	Exotic	Y	Y
Chordata	<i>Didemnum vexillum</i>	Exotic	Y	Y
Chordata	<i>Diplosoma listerianum</i>	Exotic	Y	Y
Chordata	<i>Molgula manhattensis</i>	Exotic	Y	N
Chordata	<i>Styela clava</i>	Exotic	Y	Y
Chordata	<i>Styela plicata</i>	Exotic	Y	Y

Chordata	<i>Aplidium californicum</i>	Native	Y	Y
Chordata	<i>Metandrocarpa taylora</i>	Native	Y	N

Figures




Figure 1

Mean (\pm SE) richness (a. & b.), abundance (c. & d.) and ratios of exotic:native cover (i.e., exotic dominance; e. & f.) observed in fouling communities assembling from August-December 2018 in Huntington (a., c., e.) and Newport (b., d., f.) Harbors, California, USA. Available space is calculated as the sum of bare primary space, biofilm, and oyster (*O. lurida*) cover. Values are averages from $n = 8$ plates sampled every two weeks. Assembling communities were impacted by a heavy rainfall event (indicated by a dashed vertical line) that occurred on Day 112 of the experiment




Figure 2

The relationships between the total richness (a. & c.) and exotic richness (b. & d.) of a community at a given time point and changes in exotic richness (a. & b.) or exotic abundance (c. & d.) at the following time point in assembling fouling communities in Huntington and Newport Harbors, California, USA. In both sites, (a.) fewer exotic species established as total richness increased ($\beta = -1.26$, $p < 0.001$) and (c.)

changes in total abundance of exotic species decreased via a richness \times day interaction ($\beta=0.469$, $p<0.001$). A similar influence of exotic richness was observed when modelling both (b.) changes in exotic richness ($\beta=-1.27$, $p<0.001$) and (d.) changes in exotic abundance ($\beta=0.520$, $p<0.001$). Points represent values from eight individual plates at two sites (Huntington in blue, Newport in green) across 9 time intervals ($n = 144$ total points, 112 of which were used in our statistical models after exclusion of post-rainfall data). Points below $y=0$ indicate either losses in richness (a. & b.) or decreases in abundance (c. & d.). Data are slightly horizontally jittered to increase visibility. The data are presented unstandardized but were standardized for analysis. Black trendlines indicate significant relationships that do not vary by site, while colored trendlines indicate significant site-specific relationships




Figure 3

The relationships between available space at a given time point and changes in (a.) exotic richness and (b.) proportional exotic abundance at the following time point during fouling community assembly in Huntington and Newport Harbors, California, USA. Available space was positively correlated with changes in exotic abundance but uncorrelated with changes in exotic, regardless of the richness metric used in the statistical model (Tables S1-S4). Points represent values from eight individual plates across two sites (Huntington in blue, Newport in green) and all 9 time intervals ($n = 144$ total points, 112 of which were used in our statistical models after exclusion of post-rainfall data). The data are presented unstandardized but were standardized for analysis. Data are slightly horizontally jittered to increase visibility. Colored trendlines indicate significant differences between sites, and plots without trendlines are nonsignificant

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryAppendix.pdf](#)
- [SupplementaryMaterials.pdf](#)