Modeling the Growth of Four Commercial Broiler Genotypes Reared in the Tropics

Bernard Ato Hagan (✉ bernard.hagan@uenr.edu.gh)

University of Energy and Natural Resources <https://orcid.org/0000-0003-2902-5271>

Christian Asumah

University of Energy and Natural Resources

Ernest Darkwah Yeboah

University of Energy and Natural Resources

Vida Korkor Lamptey

CSIR-Animal Research Institute

Research Article

Keywords: Gompertz function, polynomial function, body weight, growth, Ghana

Posted Date: October 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-985098/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Tropical Animal Health and Production on January 25th, 2022. See the published version at <https://doi.org/10.1007/s11250-022-03082-1>.

Abstract

Genetic improvement in commercial broilers worldwide is heavily focused on selection for higher final body weight at a given age. Although commercial broilers are mostly sold by their final body weight, it is important to pay attention to how this weight is attained and at what cost. The cost of feeding broilers, which constitutes about 70% of the total cost of broiler production, varies considerably at different stages of the bird. It is, therefore, important to pay attention to the growth curve of broilers and the parameters of the growth curve to maximise profitability of commercial broiler production. The objective of this study was to model the variations of the growth curves of 4 commercial broiler genotypes reared in Ghana using the Gompertz and polynomial growth functions. Data on body weights at 1, 7, 14, 21, 28, 35 and 42 days for 4 unsexed commercial broiler genotypes were used to model both the Gompertz and polynomial growth functions. The 4 genotypes ranked differently for Gompertz predicted early (1 - 28 days), late growth (28 – 42 days) and body weight at 42 days. Gompertz function predicted growth better for broiler chicken than the polynomial as the parameters of the Gompertz function are biologically meaningful and heritable. Selection of broiler genotypes for production based on their growth curve (slower early growth and faster late growth) could minimize cost of production and thereby increase the profitability of commercial broiler production in the tropics.

Introduction

The rate of genetic improvement in poultry is relatively faster than most livestock species due to their short generation interval and higher reproductive rate. The global broiler breeding industry, over the years, has focused on genetic improvement of growth through selection for higher body weight at a fixed age (Anthony, 2007; Siegel and Dunnington, 1987). The extensive genetic modification of the body weight of broilers has partly contributed to some adverse effects on the birds such as increased feed intake (Barbato et al. 1983), body fat (Griffin, 1996; Katanbaf et al. 1988) and a general decrease in reproductive performance (Barbato, 1999; Dunnington and Siegel, 1985; Pollock, 1999). In addition, there are also problems of ascites (Pakdel et al. 2005) and leg lameness (Kestin et al. 2001) which have been attributed to selection solely for increased body weight or growth rate. Havenstein et al. (2003) reported that fat in 43 days old broilers accounted for about 10-15% of total carcass weight.

In broiler breeding, growth or live body weight at usually 42 days of age are considered traits of interest in selective breeding programmes. The selection of animals on only superior body weight at a fixed age may be partly deceptive as this does not give information on how the weight was attained (Anthony et al. 1991). The growth curve of an animal could give information on how it attained its body weight and at what cost. Variations in the growth curve could have implication on the maintenance cost of a given animal and the eventual profitability of keeping such an animal. For instance, the financial cost of maintaining an animal that has a slower growth early in life and subsequently picks up later in life is quite lower than a rapidly fast growing animal at an earlier age. This is because the metabolic activity of a slower growing animal is not as high as the later hence the cost of feeding the former is relatively low. It

is therefore important not to select animals using only their final body weight without paying attention to their growth curve.

Selection for growth curve parameters can be employed to solve some of the problems associated with selection for final body weight since the mathematical parameters of the growth curve have moderate heritability (Barbato, 1991) and could be used to change the growth trajectory of broilers.

A variety of logistic, polynomial and other sigmoidal curves have been fitted to the growth curves of chickens (Grossman and Bohren, 1982). The Gompertz equation has been reported to have a best fit with its parameters being biologically meaningful (Aggrey, 2002; Ricklefs, 1985). Other importance of mathematical growth functions are prediction of the daily energy, protein and mineral dietary requirements of birds; appropriate slaughter age; and age of sexual maturity (Kaplan and Gürcan, 2018). The objective of this study was to model the variations of the growth of four commercial broiler chicken genotypes reared in Ghana using the Gompertz and polynomial growth functions.

Materials And Methods

Study area

The study was conducted on the Katamanso station of Animal Research Institute of the Council for Scientific and Industrial Research (CSIR-ARI). The area is located at latitude 5° 44' and longitude 0° 8' in the Accra Plains of Ghana. The annual rainfall ranges from 600 to 1000 mm and the mean annual temperature is between 20°C and 34°C.

Genotypes and Husbandry

Four exotic commercial broiler chicken genotypes marketed in Ghana namely Ross 308, Cobb 500, Hybro and Cobb 700 designated as genotypes A, B, C and D, respectively were obtained from hatcheries as unsexed day old chicks. Two hundred (200) chicks of each genotype were wing-banded, weighed and randomly placed into 16 floor pens covered with wood shavings in groups of 50 thus constituting 4 replicates per stock. Genotypes were not intermingled in a pen. Chicks were fed a starter diet of 12.12 MJ/kg Metabolisable energy and 23% crude protein for 3 weeks, followed by a finisher diet 13.2 MJ/kg of Metabolisable energy and 19.98% crude protein until the end of the experiment when broilers were 6 weeks of age. Both feed and water were provided for *ad libitum* access.

Data

The body weights of each of the four unsexed broiler genotypes were recorded at 1, 7, 14, 21, 28, 35 and 42 days of age. The observed body weights of the broiler genotypes were analysed for the mean body weights at 28 and 42 days of age. These traits are important in the broiler industry as animals are usually fed on finisher diet from 28 days onwards and broilers experience their late growth from that period whilst animals are finished at 42 days of age.

Growth functions

The Gompertz and Polynomial growth functions were used to describe the growth patterns of the four commercial broiler genotypes marketed in Ghana. Each growth function was fitted to the body weights at 1, 7, 14, 21, 28 and 42 days of age to estimate the predicted body weights of the birds. The Gompertz predicted mean body weights at 28 and 42 days of age for each genotype were then estimated.

Observed and Gompertz predicted early and late growths for each individual bird of the four broiler chicken genotypes were computed by the formulae below:

$$\text{Observed early growth (grams)} = BW_{28} - BW_1$$

$$\text{Observed late growth (grams)} = BW_{42} - BW_{28}$$

where BW_1 , BW_{28} and BW_{42} are observed body weights of broiler chickens at days 1, 28 and 42, respectively.

$$\text{Gompertz early growth (grams)} = GBW_{28} - GBW_1$$

$$\text{Gompertz late growth (grams)} = GBW_{42} - GBW_{28}$$

where GBW_1 , GBW_{28} and GBW_{42} are Gompertz predicted body weights of broiler chickens at days 1, 28 and 42, respectively.

Broiler genotypes were then ranked based on the observed and Gompertz predicted early and late growths. The Gompertz predicted growth curve parameters for each of the birds were estimated using the Microsoft Excel Macros (Microsoft Office, 2013). The Gompertz function was used to describe the age-weight relationships for each bird. In the function, the body weight of the i^{th} animal at the t^{th} age was computed as

$$BW_{it} = A_i \times \exp(-B_i (\exp^{-Kt})) \text{ (Aggrey, 2002)}$$

where BW_{it} = body weight of the i^{th} broiler at time, t

A = asymptotic body weight of the i^{th} broiler

B = transformed initial body weight of i^{th} bird, $\ln(A/BW_0)$ if BW_0 is estimated hatching weight

K = Maturation rate (exponential rate of decay of the specific growth rate)

The estimated values of the Gompertz parameters (A , B and K) of each of the birds of the genotypes, BW_{28} , BW_{42} , GBW_{28} and GBW_{42} were analysed using General Linear Model procedure of SAS (version 9.4) with Model 1 below:

$$Y_{ij} = \mu + G_i + e_{ij} \text{ Model 1}$$

where Y_{ij} = observation on a bird in the i^{th} genotype (A, B, K, BW₂₈, BW₄₂, GBW₂₈ or GBW₄₂);

G_i = fixed effect of the i^{th} genotype;

e_{ij} = the random residual $\sim N(0, \sigma_e^2)$ where σ_e^2 is the residual variance.

Differences between means were separated using Tukey's test.

Pearson correlations among observed body weights at 28 and 42 days of age and the parameters of the Gompertz growth curve were also estimated. Correlations among early growth, late growth and body weights at 42 days of age for observed and Gompertz predicted body weight were estimated.

The polynomial growth function was also fitted to the body weights and the parameters of the growth function were also estimated for each individual bird in each genotype and the mean parameters for each of the four broiler genotypes were determined.

$$BW_t = B_0 + B_1t + B_2t^2 + B_3I_s + B_4I_s \text{ (Walmsley, 2007)}$$

where BW_t = body weight at age t ;

t = age in days from hatch;

I_s = feed intake of time period;

B_0, B_1, B_2, B_3 and B_4 are unknown coefficients.

The polynomial predicted body weights at 1, 35 and 42 days of age were estimated. These were used to calculate the predicted early and late growths for each bird using the formulae below.

$$\text{Polynomial early growth} = PBW_{35} - PBW_1$$

$$\text{Polynomial late growth} = PBW_{42} - PBW_{35}$$

where PBW_1, PBW_{35} and PBW_{42} are polynomial predicted body weights of broilers at day 1, 35 and 42, respectively.

The polynomial predicted body weight at 35 and 42 days, early and late growths were analysed using GLM procedure of SAS (version 9.4).

Differences between means were separated using the Tukey's test.

Correlations among the parameters of the polynomial function and observed body weights at 42 days of age were estimated. Correlation among early, late and body weight at 42 days were also estimated.

Results

Comparison of observed and Gompertz predicted body weights

The BW_{28} , BW_{42} , GBW_{28} and GBW_{42} of the four commercial broiler genotypes are presented in Table 1. Ranking of genotypes based on body weight at 28 and 42 days of age followed the same trend for both observed and Gompertz predicted body weights. For BW_{28} , genotypes rank in descending order of D, A, C and B. However, for BW_{42} , the genotype ranking was D, C, A and B. Genotype D was superior for both traits in both the observed and Gompertz predicted body weights whilst genotype B was the least in both traits.

Table 1: Mean ($\pm$ standard deviation) observed body weights (in kg) at 28 and 42 days of age (BW_{28} and BW_{42}) and Gompertz predicted body weights at 28 and 42 days of age (GBW_{28} and GBW_{42}) for four commercial broiler genotypes

Genotype	BW_{28}	BW_{42}	GBW_{28}	GBW_{42}
A	1.0323 ± 0.046^{ab}	1.8831 ± 0.063	1.0116 ± 0.040^{ab}	1.8868 ± 0.063
B	0.9368 ± 0.047^b	1.7627 ± 0.065	0.9421 ± 0.041^b	1.7619 ± 0.065
C	1.0000 ± 0.046^b	1.8940 ± 0.063	0.9960 ± 0.040^{ab}	1.8940 ± 0.063
D	1.1947 ± 0.047^a	1.9249 ± 0.065	1.1466 ± 0.041^a	1.9363 ± 0.065
Overall	1.0404 ± 0.036	1.8668 ± 0.058	1.0236 ± 0.032	1.8703 ± 0.032

Comparison of observed and polynomial predicted body weights

The ranking of broiler genotypes by body weights followed a similar pattern for both observed and polynomial predicted body weight at 35 days of age but not for 42 days of age (Table 2). At 35 days, ranking of genotypes on body weight was D, C, A and B in descending order. At 42 days of age, whilst genotype D ranked highest for observed body weight, genotype C ranked highest for polynomial predicted body weight.

Table 2: Mean ($\pm$ standard deviation) observed body weight (in kg) at 35 and 42 days of age (BW_{28} and BW_{42}) and the Polynomial predicted body weights at 35 and 42 days of age (PBW_{35} and PBW_{42}) for four commercial broiler genotypes

Line	BW ₃₅	BW ₄₂	PBW ₃₅	PBW ₄₂
A	1.2006 ± 0.018 ^c	1.8831 ± 0.063	1.4486 ± 0.023 ^b	1.8272 ± 0.023 ^b
B	1.1695 ± 0.019 ^c	1.7627 ± 0.065	1.3391 ± 0.024 ^c	1.7009 ± 0.023 ^c
C	1.2726 ± 0.019 ^b	1.8940 ± 0.063	1.4956 ± 0.024 ^b	1.9301 ± 0.023 ^a
D	1.3776 ± 0.018 ^a	1.9249 ± 0.065	1.6359 ± 0.023 ^a	1.8405 ± 0.023 ^b
Overall	1.2562 ± 0.038	1.8668 ± 0.058	1.4818 ± 0.018	1.8250 ± 0.018

Comparison of stages of growth (Gompertz function)

Table 3 shows the magnitude of growths for the two stages of growth dubbed early and late growths for both observed and predicted growths. Ranking of genotypes based on the growth (early and late growths) followed a similar pattern for both the observed and Gompertz predicted. For early growth, genotypes are ranked D, A, C and B whilst for late growth they ranked C, A, B and D in descending order of magnitude. With the exception of genotype A which retained its rank for both early and late growths, all other genotypes had a different pattern of growth. Genotype D recorded the largest early growth but a slower growth as it approached maturity. Genotype C, on the other hand, was slow growing in its early life and picked up later in life.

Table 3: Mean (± standard deviation) in kg of observed early and late growths (Early growth_(obs) and late growth_(obs)) and the Gompertz predicted early and late growths (Early growth_(Gomp) and late growth_(Gomp)) for four broiler genotypes

Genotype	Early growth _(obs)	Late growth _(obs)	Early growth _(Gomp)	Late growth _(Gomp)
A	0.9078 ± 0.046 ^{ab}	0.8506 ± 0.053	0.9169 ± 0.042 ^b	0.8752 ± 0.046
B	0.8171 ± 0.047 ^b	0.8259 ± 0.055	0.8353 ± 0.043 ^b	0.8198 ± 0.047
C	0.8692 ± 0.046 ^b	0.8940 ± 0.053	0.8642 ± 0.042 ^b	0.8980 ± 0.046
D	1.0718 ± 0.047 ^a	0.7302 ± 0.055	1.0782 ± 0.043 ^a	0.7897 ± 0.047
Overall	0.9157 ± 0.037	0.8264 ± 0.039	0.9228 ± 0.037	0.8467 ± 0.038

Comparison of stages of growth (Polynomial function)

The ranking of the genotypes based on the observed early and late growths and polynomial predicted early and late growths followed similar trends (Table 4). However, the patterns of growth of the genotypes were different. Genotype D recorded the highest growth at the first stage of growth but the slowest growth at later stage of life.

Table 4: Mean ($\pm$ standard deviation) in kg of observed early and late growths (Early growth_(obs) and late growth_(obs)) and the polynomial predicted early and late growths (Early growth_(poly) and late growth_(poly)) for four commercial broiler genotypes

Genotype	Early growth _(obs)	Late growth _(obs)	Early growth _(poly)	Late growth _(poly)
A	1.1553 $\pm$ 0.018 ^c	0.6266 $\pm$ 0.016 ^a	1.4039 $\pm$ 0.023 ^b	0.3786 $\pm$ 0.016 ^{ab}
B	1.1276 $\pm$ 0.019 ^c	0.5540 $\pm$ 0.017 ^b	1.2958 $\pm$ 0.024 ^c	0.3618 $\pm$ 0.017 ^b
C	1.2232 $\pm$ 0.019 ^b	0.6673 $\pm$ 0.017 ^a	1.4433 $\pm$ 0.024 ^b	0.4345 $\pm$ 0.017 ^a
D	1.3369 $\pm$ 0.018 ^a	0.4626 $\pm$ 0.016 ^c	1.5970 $\pm$ 0.023 ^a	0.2046 $\pm$ 0.016 ^c
Overall	1.2120 $\pm$ 0.015	0.5765 $\pm$ 0.014	1.4373 $\pm$ 0.020	0.3431 $\pm$ 0.014

The parameters of the Gompertz function were different for the different broiler genotypes (Table 5). Genotype D had the highest ($P < 0.05$) maturation rate, K compared to the other genotypes even though it came up with the lowest asymptotic body weight, A. There was a strong negative correlation between K and transformed initial body weight, B; and a weak correlation between K and BW₄₂ (Table 7). There was a weak positive correlation between BW₄₂ and parameter A though not significant. The opposite seems to be the case for BW₂₈ and the Gompertz growth curve parameters. There was a significant positive correlation between B and K; and between B and BW₂₈ (Table 7). However, a negative correlation existed between BW₂₈ and A. A significant positive correlation exists between B and K whilst a negative correlation exists between A and B; and between A and K.

The coefficients of the polynomial functions and the observed body weights for the four broiler genotypes are presented in Table 6. For all broiler genotypes, the polynomial coefficients decreased from B₀ to B₃. Parameter B₃ remained negative for all broiler genotypes.

Table 5: Mean ($\pm$ standard deviation) observed body weights at 28 and 42 days of age (in kg) and Gompertz growth parameters for four commercial broiler genotypes

Genotype	BW ₂₈	BW ₄₂	A	B	K
A	1.0323 $\pm$ 0.046 ^{ab}	1.8831 $\pm$ 0.063	11.77 $\pm$ 5.89	8.10 $\pm$ 0.89 ^b	0.365 $\pm$ 0.03 ^b
B	0.9368 $\pm$ 0.047 ^b	1.7627 $\pm$ 0.065	12.70 $\pm$ 6.05	7.35 $\pm$ 0.91 ^b	0.334 $\pm$ 0.03 ^b
C	1.0000 $\pm$ 0.046 ^b	1.8940 $\pm$ 0.063	14.47 $\pm$ 5.89	7.31 $\pm$ 0.89 ^b	0.293 $\pm$ 0.03 ^b
D	1.1947 $\pm$ 0.047 ^a	1.9249 $\pm$ 0.065	3.11 $\pm$ 6.05	12.39 $\pm$ 0.91 ^a	0.515 $\pm$ 0.03 ^a
Overall	1.0404 $\pm$ 0.046	1.8668 $\pm$ 0.058	10.58 $\pm$ 5.72	8.76 $\pm$ 0.87	0.376 $\pm$ 0.02

Table 6: Mean ($\pm$ standard deviation) polynomial growth parameters and the observed body weight at 42 days of age for four commercial broiler genotypes

Genotype	B_0	B_1	B_2	B_3	BW_{42}
A	551.42 ± 7.62^c	50.34 ± 1.16^{bc}	1.51 ± 0.08^b	-0.017 ± 0.002^a	1.8831 ± 0.063
B	569.05 ± 7.84^{bc}	46.46 ± 1.20^c	0.94 ± 0.08^c	-0.015 ± 0.002^a	1.7627 ± 0.065
C	613.05 ± 7.84^a	51.97 ± 1.20^b	1.15 ± 0.08^c	-0.016 ± 0.002^a	1.8940 ± 0.063
D	592.80 ± 7.60^{ab}	62.02 ± 1.16^a	2.06 ± 0.08^a	-0.042 ± 0.002^b	1.9249 ± 0.065
Overall	581.32 ± 7.35	52.81 ± 1.04	1.43 ± 0.07	-0.023 ± 0.04	1.8668 ± 0.058

Table 7: Phenotypic correlation among observed body weights at 42 and 28 days of age (BW_{42}), BW_{28} and Gompertz predicted growth parameters (A, B and K)

	A	B	K
BW_{42}	0.020	-0.117	-0.044
BW_{28}	-0.378*	0.318*	0.701*
A		-0.070	-0.482*
B			0.748*

*Correlation is significant at 0.01 level

Table 8: Phenotypic correlation between observed body weight at 42 days of age (BW_{42}) and polynomial predicted growth parameters

	B_0	B_1	B_2	B_3
BW_{42}	0.635*	0.500*	0.234*	-0.103*
B_0		0.471*	-0.050	-0.229*
B_1			0.777*	-0.891*
B_2				-0.744*

*Correlation is significant at 0.01 level

The observed growth curves of the broiler genotypes looked similar at the early stage of growth (first 28 days) after which genotype D increased slightly (Figure 1). However, at the end of growth, all genotypes attained similar body weights with the exception of genotype B which lacked behind but not statistically significant ($P>0.05$).

The Gompertz predicted growth curves of the four broiler genotypes showed similar pattern as those of the observed growth curves, however the observed growth curves separated at an early stage of 3 weeks of age (21 days) (Figure 2). Genotype B still lacked behind in the Gompertz predicted growth compared to the other genotypes.

Table 9: Correlations between observed body weights and Gompertz predicted (GBW) and Polynomial predicted body weights (PBW) at days 1, 7, 14, 21, 28, 35 and 42

Observed body weight	GBW	PBW
BW1	0.793*	0.552*
BW7	0.402*	
BW14	0.832*	
BW21	0.884*	
BW28	0.982*	
BW35		0.197*
BW42	0.992	0.150*

*Correlation is significant at 0.01 level

The correlations among observed body weight at 42 days, polynomial parameters B_0 , B_1 , and B_2 , were all positive and significant (Table 8). Parameter B_3 was negatively correlated with all other parameters (B_0 , B_1 and B_2). There were strong positive correlations between the observed body weights at different ages and their corresponding Gompertz predicted body weights at different ages (Table 9) with all the correlations being significant. The overall mean observed body weight at 42 days was similar to polynomial predicted body weight at 42 days but not Gompertz predicted 42 days' body weight (Table 10). Gompertz function had a slightly lower variation for body weight at 42 days compared to those of observed and polynomial functions.

Table 10: Mean body weights at 42 days, minimum, maximum body weights and standard deviations (grams) for observed, Gompertz and polynomial predicted using pooled data

	BW_{42Obs}	BW_{42Gomp}	BW_{42Poly}
Overall Mean	1866.83	1870.28	1825.04
Minimum	643.60	1330.00	643.97
Maximum	2652.40	2658.00	2652.64
Standard deviation	296.90	283.24	296.75

Discussion

Ranking on body weight

The similarity in the ranking of broiler genotypes by their observed body weights and Gompertz predicted body weights could be due to the use of only actual body weights by the Gompertz function in predicting the body weights of chicken. However, the differences in ranking observed in broiler genotypes using polynomial predicted body weights could be attributed to the incorporation of feed intake in the function (Walmsley, 2007). Praharaj et al. (1996) reported that ranking of sire families of broilers are significantly different under full and restricted feeding regimes or programme. The appreciable variations within body weights suggest that further genetic improvement could be achieved within the genotypes through selection.

Ranking on early and late growths

The four commercial broiler genotypes had similar rankings for both observed and predicted growths. The pattern of growths, however, differed for the different genotypes. The use of Gompertz predicted growth seems to have a slightly lower variation than the observed growths. Growth of broilers is a highly heritable trait (Siegel and Wilford, 2003), therefore breeding programmes could be designed to select for favourable pattern of growth. Genotype D which had the highest early growth ended up with the lowest late growth.

This is evident in their growth curves for both the observed and Gompertz growth curves (Figures 1 and 2). Genotype C ranked third in magnitude for early growth and first for late growth. Growth of animals is characterized by higher composition of bone and muscle at early growth and higher amount of fat at later stage of growth (Lawrence and Fowler, 2002). Selection in broilers could be designed as high early growth and low late growth. However, the maintenance cost of high early growth is enormous since the metabolic maintenance energy requirement is high. Therefore, although genotype D had the highest BW₄₂, genotype C could be selected based on its slow early growth and high late growth pattern. This might help prevent or minimize some of the challenges associated with selection solely on the basis of body weight (Barbato et al. 1983; Dunnington and Siegel, 1985). The ranking of the broiler genotypes using the polynomial predicted body weights also followed a similar trend. Both the polynomial and

Gompertz functions could be said to give a similar ranking for growth. The pattern of growth is also similar. Slow growing animals at early stage tend to hasten their growth in later stages of life.

Ranking using growth functions (Gompertz and Polynomial)

The use of asymptotic body weight, A, ranks genotypes in descending order of C, B, A and D. The parameter, A estimates the weight of an animal at an infinite age. In commercial broiler chicken production, animals are ready for slaughter between 6 and 8 weeks of age hence hardly reach this matured body weight therefore parameter A is unimportant. Transformed initial body weight, B, and exponential rate of decay of specific growth rate, K, both ranked genotypes in the same order of D, A, B and C. This could be due to the strong positive correlation between the two parameters. Parameter K is however, the most biologically meaningful of the three parameters and has a greater effect on the growth curve. Genotype D which had the highest K value also had the lowest value for late growth. The overall estimated K for the pooled data, 0.376, was higher than that reported by Barbato (1991) but lower than that of Mignon-Grasteau et al. (2000) which were 0.037 and 2.15, respectively. The variations within the estimate for this work were, however, lower. A lower K has a favourable growth curve of low early growth and a high late growth. Barbato (1991) has reported a heritability of 0.28 for this parameter. This trait can therefore be passed on from parents to their progeny. Parameter K has a negative correlation with BW_{42} but a positive correlation with BW_{28} . This is expected since as animals grow, the rate of deterioration of growth increases hence the negative correlation between K and BW_{42} . The phenotypic correlation between BW_{28} and K (0.701) was similar to that of Barbato (1991) who reported 0.74 at 14 days. Rapid progress for a trait can be achieved by selecting on a highly-correlated trait if the trait under consideration is difficult to measure with precision. Parameter K can therefore be selected for by indirectly selecting for BW_{28} . However, there was a big difference between the correlation at 42 days reported by Barbato (1991) and that of this work at the same age. Environmental and genetic variations could have influenced the trait in the present study. There were also negative correlations between the growth curve parameters, A and K. This could be from the rapid decrease in growth rate after inflection point resulting in a lower asymptotic body weight (Mignon-Grasteau et al. 2000).

The polynomial parameters ranked the broiler genotypes differently. The parameter, B_1 however, gave a similar ranking as those of BW_{42} . The parameters of the polynomial function all have a positive correlation with BW_{42} with the exception of B_3 . Aggrey (2002) explained that polynomial function has a better fit for poultry data but the parameters of this function have no biological interpretation because they give no insight into the growth process. The use of this function seems to be more suitable for predicting early body weight rather than late growth in broilers.

Chambers (1990) observed that the correlations between weights at a given age and previous gain were high. Gompertz predicted growth could thus be employed to give a better prediction of body weight than polynomial.

The standard deviation of a given estimation method could be used to predict the robustness of the method for estimation. However, the Gompertz function slightly overestimates the mean body weights. Gompertz function is a better predictor of body weights at later age than the polynomial function.

There was an interesting trend for the correlations between observed body weights at fixed ages and their predicted weights at the same ages. Whilst the correlations between observed and Gompertz predicted body weight generally increased with age the reverse was observed for the polynomial function.

It is possible to predict weight of broiler chickens using the Gompertz and polynomial growth functions. However, the Gompertz function is a better estimator of growth since it has a number of advantages over the polynomial. Among these are: its parameters are biologically meaningful and heritable, lower variation between the parameters it estimates and the high correlations between its predicted weights and those of the observed. Nonetheless, a simple method of slow early growth and fast late growth could also be employed in a broiler breeding programme to rank animals for selection. This will modify the growth curve and minimize the problems related to selection solely on final body weight.

Declarations

Acknowledgement

The authors are grateful to the technical staffs of Animal Research Institute of the Council of Scientific and Industrial Research (CSIR-ARI) for the data collection and management of the experimental flock.

Funding

There was no funding for this work as the information used for the study was data provided by the Poultry Unit of CSIR-Animal Research Institute.

Conflicts of interest

The authors declare that they have no competing interests.

Ethics approval

The manuscript does not contain clinical studies.

Consent to participate

Not applicable

Consent for publication

There are consent from CSIR-Animal Research Institute barring the publication of this work.

Availability of data and material

The data for this work is available upon request.

Code availability

Not applicable.

Authors' contribution

Bernard Ato Hagan – conceived the idea for this work; analysed that data; wrote the manuscript.

Christian Asumah – Wrote the manuscript.

Ernest Darkwah Yeboah – Wrote the manuscript.

Vida Korkor Lamptey – Data collection; wrote the manuscript.

References

1. Aggrey, S. E., 2002. Comparison of Three Nonlinear and Spline Regression Models for Describing Chicken Growth Curves. *Poultry Science*, 81, 1782–1788.
2. Anthony, N. B., Emmerson, D. A., Nector, K. A., Bacon, W. L., Siegel, P. B. and Dunnington, E. E., 1991. Comparison of the growth curves of weight selected populations of turkeys, quail and chickens. *Poultry Science*, 70, 13–19.
3. Anthony, N. B., 2007. A review of genetic practices in poultry: Efforts to improve meat quality. *Journal of Muscle Food*, 9 (1), 25–33.
4. Barbato, G. F., Siegel, P. B. and Cherry, J. A., 1983. **Inheritance of body weight and associated traits in young chickens**. *Zeitschrift für Tierzüchtung und Züchtungsbiologie*, 100, 350-360.
5. Barbato, G. F., 1991. **Genetic architecture of growth curve parameters in chickens**. *Theoretical and Applied Genetics*, 83, 24-32.
6. Barbato, G. F., 1999. **Genetic relationships between selection for growth and reproductive effectiveness**. *Poultry Science*, 78, 444-452.
7. Chambers, J. R., 1990. **Genetics of growth and meat production in chickens**. In: R. D. Crawford (ed), **Poultry breeding and genetics**. Elsevier Science Publication, Amsterdam, 599-643.
8. Dunnington, E. A. and Siegel, P. B., 1985. **Long-term selection for eight-week body weight in chicken in chickens – direct and correlated responses**. *Theoretical and Applied Genetics*, 71, 305-313.
9. Griffin, H. D., 1996. **Understanding genetic variation in fatness in chickens**. *Annual Report, Roslin Institute, Edinburgh, UK*.
10. Grossman, M. and Bohren, B. B., 1982. **Comparison of proposed growth curve functions in chickens**. *Growth*, 46, 259-274.
11. Havenstein, G. B., Ferket, P. R. and Qureshi, M. A., 2003. **Carcass composition and yield of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets**. *Poultry Science*, 82, 1509-

1518.

12. Katanbaf, M. N., Dunnington, E. A. and Siegel, P. B., 1988. Allomorphic relationships from hatching to 56 days in parental lines and F₁ crosses of chickens selected 27 generations for high or low body weight. *Growth Development and Aging*, 52, 11–22.
13. Kaplan, S. and Gürcan, E. K., 2018. Comparison of growth curve using non-linear regression function in Japanese quail. *Journal of Applied Animal Research*, 46(1), 112–117.
14. Kestin, S. C., Gordon, S., Su, G. and Sorenson, P., 2001. Relationship in broiler chickens between lameness, liveweight, growth rate and age. *Veterinary Record*, 148, 195–197.
15. Lawrence, T. L. J. and Fowler, V. R., 2002. *Growth of farm animals*. CAB International, Wallingford, UK.
16. Mignon-Grasteau, S., Piles, M., Varona, L., de Rochambeau, H., Poivey, J. P., Blasco, A. and Beaumont, C., 2000. Genetic analysis of growth curve parameters for male and female chickens resulting from selection on shape of growth curve. *Journal of Animal Science*, 78, 2515–2524.
17. Pakdel, A., Van Arendonk, J. A. M, Vereijken, A. L. J. and Boverhuis, H., 2005. Genetic correlations among ascitis-related traits and performance traits in broilers. *British Poultry Science*, 46, 35–42.
18. Pollock, D. L., 1999. A geneticist's perspective from within a broiler primary breeder company. *Poultry Science*, 78, 414–418.
19. Praharaj, N. K., Gross, W. B., Dunnington, E. A. and Siegel, P. B., 1996. Feeding regimen by sire family interactions on growth, immunocompetence and disease resistance in chickens. *Poultry Science*, 75, 821–827.
20. Praharaj, N. K., Gross, W. B., Dunnington, E. A. and Siegel, P.B., 1996. Feeding regimen by sire family interactions on growth, immunocompetence and disease resistance in chickens. *Poultry Science*, 75, 821–827.
21. Ricklefs, R. E., 1985. Modification of growth and development of muscles of poultry. *Poultry Science*, 64, 1653–1676.
22. Siegel, P. B. and Dunnington, E. A., 1987. Selection for growth in chickens. *Critical Reviews in Poultry Biology*, 1, 1–24.
23. Siegel, P. B. and Wolford, J. H., 2003. A review of some results of selection for juvenile body weight in chickens. *Journal of Poultry Science*, 40, 81–91.
24. Walmsley, B. J., 2007. Modelling of growth and development for optimizing beef cattle production systems. PhD Thesis, University of New England, Australia.

Figures

Figure 1

Observed growth curves of four broiler genotypes

Figure 2

Observed growth curves of four broiler genotypes