

RANKL Immunisation Inhibits Prostate Cancer Metastasis by Modulating EMT Through a RANKL-dependent Pathway

Mineon Park

Chosun University Hospital

Yong Jin Cho

Chosun University Hospital

Bora Kim

Chosun University Hospital

Young Jong Ko

Chosun University Hospital

Yuria Jang

Chosun University Hospital

Hoon Hyun

Chonnam National University

Wonbong Lim (✉ wonbong@chosun.ac.kr)

Department of Premedical Sciences, College of Medicine, Chosun University, Dong-Gu, Gwangju, 501-759, Korea <https://orcid.org/0000-0001-6337-6599>

Research

Keywords: RANKL, prostate cancer, metastasis, EMT

Posted Date: October 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-98591/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Prostate cancer (PCa) morbidity in the majority of patients is due to metastatic events, which are a clinical obstacle. Therefore, a better understanding of the mechanism underlying metastasis is imperative if we are to develop novel therapeutic strategies. Receptor activator of nuclear factor kappa-B (NF- κ B) ligand (RANKL) regulates bone remodelling. RANKL was associated with epithelial-mesenchymal transition (EMT) and expression of metastasis-related genes in PC3 cells. Thus, agents that suppress RANKL signalling may be useful pharmacological treatments.

Method: In this study, we proposed a strategy to induce anti-cytokine antibodies using mutant RANKL as an immunogen. Here, we used preclinical experimental models to investigate whether an inactive form of RANKL affects bone metastasis in RANKL-induced PCa.

Results: RANKL activation was observed in human PCa tissue specimens. RANKL promoted migration and invasion of PC3 cells through EMT, and induced a significant increase in binding of β -catenin to TCF-4, an EMT-induced transcription factor in PCa cells, via mitogen-activated protein kinase and β -catenin/TCF-4 signalling. Thus, RANKL increased EMT and the metastatic properties of PC3 cells, suggesting a role as a therapeutic target to prevent PCa metastasis.

Conclusion: Treatment with mutant RANKL reduced EMT and metastasis of PC3 PCa cells in an experimental metastasis model. Thus, mutant RANKL could serve as a potential vaccine to prevent and treat metastatic PCa

Trial registration: Chosun University Hospital, CHOSUN 2020-06-001. Registered 01 June 2020-prospectively registered, https://hosp.chosun.ac.kr/medi_depart/?site=hospital&mn=151&type=view&catename=IRB

Introduction

Prostate cancer (PCa) is the second most common cancer in males and the fifth leading cause of death worldwide [1]. Metastasis of PCa cells to the skeleton occurs in a predictable manner, with lesions tending to appear first in the axial skeleton, followed by appendicular tissues [2]. Considering the effects of PCa on both haematopoiesis and bone structure, bone metastasis is a major cause of morbidity in patients with advanced disease. Replacement of haematopoietic tissue by metastatic PCa cells is associated with anaemia and increased morbidity (mortality). The 5-year survival of most patients with PCa is almost 100%; however, that of PCa patients with metastasis to distant sites is as low as 28%. Thus, PCa is one of the deadliest cancers [3]. Experimental and clinical observations reveal that treatment with anti-resorptive agents suppresses or even prevents PCa metastasis [4-7], and that accelerated bone turnover stimulates progression of skeletal secondary tumours [8-10].

During bone metastasis of PCa, cancer cell-derived cytokines stimulate expression of receptor activator of nuclear factor kappa-B ligand (RANKL), which in turn activates bone resorption. RANKL, also known as

tumour necrosis factor-related activation-induced cytokine (TRANCE) [11], osteoprotegerin ligand (OPGL) [12, 13] and osteoclastic differentiation factor (ODF) [14], interacts with RANK and is involved in all the steps related to tumour development, from initial tumour formation to migration of cancer cells and subsequent metastasis [15]. RANKL is expressed in several tissues, including brain, skin, intestine, skeletal muscle, kidney, liver, lung and mammary tissue; however, expression is very high in bone [16], lymphoid organs and the vascular system [17]. RANKL binds to RANK on the surface of pre-osteoclasts, activating them and inducing formation of osteoclasts [18].

Recent studies report expression of RANK and RANKL by various solid tumours, including breast cancer. RANKL accelerates migration and metastasis of cancer cells expressing RANK [19, 20]. Furthermore, it protects breast cancer cells from apoptosis in response to DNA damage and controls self-renewal and anchorage-independent growth of tumour-initiating cells [21]. However, it is unclear how RANKL signalling triggers metastasis of PCa. Epithelial-to-mesenchymal transition (EMT), a rapid and often reversible phenotypic change in epithelial cells, is an important phenomenon underlying cancer metastasis. Originally, EMT was described in the context of developmental processes such as heart morphogenesis and mesoderm and neural crest formation. Epithelial cells lose structures involved in cell–cell adhesion (e.g., adherens junctions and desmosomes), modulate their polarity and rearrange their cytoskeleton, which is consistent with the typical switch of intermediate filaments from cytokeratins to vimentin [22]. A recent report shows that the oncogenic c-MYC, Wnt signalling and β -catenin pathways activate the Snail/glycogen synthase kinase-3 (GSK-3) axis and induce EMT [23].

The concept of exploiting the host's immune system to treat cancer relies on the ability of immune cells to eliminate malignant cells at the early transformation stage in a process called immune surveillance [24]. Passive anti-cytokine immunotherapy with specific high-affinity antibodies has been tested in animal models and clinical trials of rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease, asthma, Crohn's disease, psoriasis and other articular autoimmune disorders. This strategy facilitates production of anti-auto-cytokine antibodies by the immune system in response to active vaccination. However, development of a desired antibody response to self-proteins necessitates suppression of immune resistance. Common anti-cytokine vaccines are prepared from autologous proteins that are converted to derivatives that lack biological activity following treatment with glutaraldehyde or formaldehyde [25]. A previous study developed a mouse RANKL mutant (mRANKL-MT) protein and confirmed its ability to inhibit osteoporosis.

Here, we used mRANKL-MT as an immunogen for RANKL-targeting immunotherapy of bone disease and investigated its potential as a cancer vaccine in a mouse model of metastatic cancer. We confirmed the activity of mRANKL-MT and clarified its effects on RANK/RANKL signalling-mediated EMT in transient RANKL-overexpressing cell lines and animal models. We found that mRANKL-MT suppressed RANKL-dependent β -catenin signalling. We also found that mRANKL-MT immunotherapy altered the characteristics of cancer cells and effectively suppressed RANKL-dependant cancer metastasis.

Methods

Patient samples

Three patient tissue samples from PCa patients not receiving chemotherapy, hormone therapy, or radiotherapy were obtained before surgery from Chosun University Hospital during 01 June 2020 and 31 July 2020 (Chosun University Hospital (CHOSUN 2020-06-001)). Tumours and normal tissues were harvested and immediately fixed in a formalin solution (neutral buffered, 10%; Sigma-Aldrich) for 24 h. Tissue specimens were successively dehydrated in ethanol and treated with xylene. Paraffin embedded tissues were sliced into 5 µm-thick sections, deparaffinised with xylene and rehydrated through graded alcohol solutions. The tissues were stained with H&E for histological analysis. Immuno-histochemical studies were performed using the Novolink™ Polymer Detection System kit (Leica Biosystems) as per the manufacturer's instructions. Antibodies specific for IL-6 (1:200; Cell Signaling Technology) and RANKL (1:200, Cell Signaling Technology) were used.

Cell lines and cultures

Human prostate adenocarcinoma luciferase-labelled PC3^{luc} cells were obtained from Professor Park and maintained at 37°C/5% CO₂ in Rowell Park Memorial Institute (RPMI) 1640 medium (Welgene, Korea) supplemented with 10% heat-inactivated foetal bovine serum (certified, GIBCO, USA) and a 10% antibiotic solution (Welgene).

Cloning of hRANKL

The RNA used to clone hRANKL cDNA was extracted from MG63 cells (ATCC® CRL-1427™) expressing RANKL. The quality of the extracted RNA was verified by agarose gel electrophoresis. The cDNA was prepared using the AccuPower RT PreMix Kit (Bioneer, Daejeon, Korea), according to the manufacturer's instructions. Amplification and cloning of the hRANKL fragment were carried out in a reaction mixture comprising KOD polymerase buffer, 10 mM dNTPs, 25 mM magnesium chloride (MgCl₂), 10 µM primers (hRANKL-*Bcl*I: 5¢-TGATCAAAGCTTGAAGCTCAGCCTTTTGC-3¢ and hRANKL-*Xho*I: 5¢-CTCGAGATCTATATCTCGAACTTTAAAAGCCCC-3¢), 2.5 U of KOD DNA polymerase (EMD Millipore, Billerica, MA, USA) and 2 µL of the RANKL gene construct (template). The thermal cycling conditions were as follows: initial denaturation at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s and extension at 70°C for 30 s. The polymerase chain reaction (PCR) product was cloned into the *Bam*H1/*Xho*I sites of a pMX vector (CELL BIOLABS, USA). Sequence analyses were carried out using programs in Vector NTI Advance 9.1.0 (Invitrogen, Carlsbad, CA, USA).

Retroviral hRANKL transduction

Plat-E cells were seeded at a density of 3×10^5 cells/well in a six-well plate for 24 h and then transiently transfected with hRANKL/pMX using 0.2 µg plasmid and 0.6 µL of the FuGENE HD transfection reagent (Promega, Madison, WI, USA), according to the manufacturer's protocol. After incubation, the DNA/FuGENE mixture was added drop-wise onto Plat-E cells. Viral supernatants were recovered from the

culture medium at 48 h after transfection. Virus-containing supernatants were filtered through 0.45 µm non-pyrogenic filters and supplemented with 10 µg/mL polybrene (Sigma-Aldrich).

Reverse-transcription quantitative PCR (RT-qPCR)

Total RNA was extracted from PCa cells using Trizol (Invitrogen) and 1 µg was used for RT-qPCR along with oligo-dT primers (10 µg) and dNTPs (10 mM). Next, qRT-PCR was performed to analyse cDNA using SYBR Green SuperMix (BIORAD, USA) on a CFX Connect Real-Time System (BIORAD, USA). All target gene primers were purchased from Bioneer Co. (Daejeon, Korea) and the cDNA was amplified using the following primer sets:

E-cadherin (h): 5'-TGGAGGAATTCTTGCTTTGC-3' (forward) and 5'-TGGAGGAATTCTTTTGC-3' (reverse); vimentin (h): 5'-GACGCCATCAACACCGAGTT-3' (forward) and 5'-GACGCCATC AACACCGAGTT-3' (reverse); β-catenin (h): 5'-ACAAACTGTTTTGAAAATCCA-3' (forward) and 5'-CGAGTCATTGCATACTGTCC-3' (reverse); MMP-9 (h): 5'-TCCAGTACCAAGACAAAG-3' (forward) and 5'-TTGCACTGCACGGTTGAA-3' (reverse); RANK (h): 5'-CAAATGCAGACCCTGGA CCA-3' (forward) and 5'-AAACGCCAAAGATGATGGCA-3' (reverse); RANKL (h), 5'-CCTGTAT GCCAACATTTGCTTTC-3' (forward) and 5'-TTCCTCTCCAGACCGTAACTTAAA-3' (reverse); IL -6 (h): 5'-AGCAAAGAGGCACTGGCAGA-3' (forward) and 5'-GTACTCATCTGCACAGCTCTGG C-3' (reverse); TCF-4 (h): 5'-GCTCAGGGTATGGAACCGGC-3' (forward) and 5'-CCCTGTAGTC CTGGTGGCATG-3' (reverse); c-MYC (h): 5'-CCTGGTGCTCCATGAGGAGAC-3' (forward) and 5'-AGACTCTGACCTTTTGCCAGG-3' (reverse); and glyceraldehyde 3-phosphate dehydrogenase (GAPDH): 5'-TCAAGAAGGTGGTGAAGCAG-3' (forward) and 5'-AGTGGGAGTTGCTGTTGAAG T-3' (reverse). Values on the vertical axis represent $2^{-\Delta Ct}$; ΔCt is the discrepancy between the target gene Ct and GAPDH Ct.

Western blot analysis

The cells were washed twice with phosphate-buffered saline (PBS; pH 7.4) and total proteins were extracted using radio immunoprecipitation assay buffer supplemented with 1% protease inhibitors, phosphatase inhibitors and phenylmethylsulfonyl fluoride (PMSF). The protein concentration was measured using the BCA Protein Assay Kit (Thermo Pierce™). The membrane was blocked with a solution containing 5% skim milk in TBS-T for 30 min and then washed in TBS-T. The membrane was incubated for overnight at 4°C with the following primary antibodies: E-cadherin (sc-7870, Santa Cruz Biotechnology), N-cadherin (ab76011, Abcam), β-catenin (#29822 94, Millipore), MMP-9 (#13667, Cell Signaling), IL-6 (#12153, Cell Signaling Technology), c-MYC (9E10, Santa Cruz Biotechnology), TCF-4 (#2565, Cell Signaling Technology), RANK (#4845, Cell Signaling Technology), P-ERK (#9101, Cell Signaling Technology), ERK (#9102, Cell Signaling Technology), GAPDH (#2118, Cell Signaling Technology), P-AKT (#9271, Cell Signaling Technology), AKT (#9272, Cell Signaling Technology), P-SRC (#2105, Cell Signaling Technology), SRC (#2108, Cell Signaling), GSK-3B (#9315, Cell Signaling Technology) and P-GSK-3B (#9336, Cell Signaling Technology). Horseradish peroxidase (HRP)-

conjugated AffiniPure goat anti-rabbit IgG (H + L) and HRP-conjugated AffiniPure goat anti-mouse IgG (H + L) were obtained from Proteintech Group, Inc (Jackson) and used as secondary antibodies.

Cell migration

Cells were seeded in 6-well plates for the cell migration assay. After each treatment, a confluent monolayer was wounded using a 200 μ L pipette tip. Images of wound closure were obtained under an inverted microscope after 48 h. The wound area was calculated using NIH ImageJ software.

Cell invasion assay

A total of 1×10^5 transfected cells were seeded into the top chamber of a 24-well polycarbonate Transwell chamber (8.0 μ m pore size; Corning Incorporated, Glendale, AZ, USA) and then treated for 24 h with hRANKL or RANKL. The number of trypan blue-stained cells in five random fields was counted using an inverted microscope.

Luciferase reporter assay to assess Wnt/ β -catenin activity

Cells were seeded into a 24-well plate 24 h prior to transient transfection with either 2 mg of TOPflash or FOPflash reporter plasmid along with 1 mg DNA using Lipofectamine 3000 (Thermo Fisher). The TOPflash luciferase reporter plasmid contains TCF-4-binding sites upstream of the luciferase gene, resulting in luciferase activity in the presence of active Wnt/ β -catenin signalling. The FOPflash reporter plasmid, on the other hand, carried mutated TCF-4-binding sites. Total cell extracts were assayed for luciferase activity according to the manufacturer's instructions (Promega).

Immunoprecipitation

Cells were lysed in lysis buffer (20 mM Tris-HCl pH 7.6–8.0, 100 mM sodium chloride NaCl, 300 mM sucrose, 3 mM MgCl₂ [buffer A]; and 20 mM Tris pH 8.0, 100 mM NaCl, 2 mM ethylenediaminetetraacetic acid [buffer B]). Whole cell lysates obtained by centrifugation were incubated with antibodies specific for active β -catenin (Millipore) and TCF-4 (Cell Signaling Technology) (dilution 1:100) and protein A Sepharose beads (Amersham Biosciences) for 2 h at room temperature. The immune complexes were washed three times using wash buffer and examined by western blotting.

Site-directed mutagenesis and production and purification of mRANKL-MT

The RNA used for the cloning of RANKL cDNA was extracted from MC3T3-E1 cells (Korean Cell Line Bank, Seoul, Korea) expressing RANKL. The quality of the extracted RNA was verified by agarose gel electrophoresis and cDNA was prepared using the AccuPower RT PreMix Kit (Bioneer, Daejeon, Korea), according to the manufacturer's instructions. Amplification and cloning of the RANKL fragment were carried out in a reaction mixture comprising KOD polymerase buffer, 10 mM dNTPs, 25 mM MgCl₂, 10 μ M primers (mRANKL-*Nde*I: 5 ϕ -CATATGAAGCCTGAGGCCAGCC ATTTGC-3 ϕ ; mRANKL-*Xho*I: 5 ϕ -CTCGAGGTCTATGTCCTGAACTTTGAAAGCC-3 ϕ ; mRANKL (K180R)-F: 5 ϕ -

CCCATCGGGTTCCCATCGAGTCACTCTGTCCTCTTG-3¢; mRANKL (K180R)-R: 5¢-CAAGAGGACAGAGTGACTIONCGATGGGAACCCGATGGG-3¢; mRANKL (D189I, R190K)-F: 5¢-CTCTTGGTACCACATCAAGGGCTGGGCCAAGAT-3¢; mRANKL (D189I, R190K)-R: 5¢-ATC TTGGCCCAGCCCTTGATGTGGTACCAAGAG-3¢; mRANKL-MT (H223F, H224Y)-F: 5¢-AA CA TTTGCTTTTCGGTTTTATGAAACATCGGGAAGCG-3¢; or mRANKL-MT (H223F, H224Y)-R: 5¢-CGCTTCCCGATGTTTCATAAAACCGAAAGCAAATGTT-3¢), 2.5 U of KOD DNA polymerase (EMD Millipore, Billerica, MA, USA) and 2 µL of RANKL gene construct as the template.

The thermal cycling conditions were as follows: initial denaturation at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s and extension at 70°C for 30 s. The PCR product obtained was cloned into the *NdeI/XhoI* site of the GST-30a vector (Novagen, Madison, WI, USA). Mutations at positions 180, 189–190 and 223–224 were introduced using megaprimers [26]. The PCR product was transformed into *Escherichia coli* BL21-CodonPlus (DE3)-RIPL (Novagen) by electroporation (5 ms, 12.5 kV/cm) and the transformed cells were cultivated in Luria-Bertani broth containing kanamycin (50 µg/mL, T&I, Daejeon, Korea). Plasmids were purified using the QIAprep Spin Miniprep Kit (Qiagen, Valencia, CA, USA). The cloned product was confirmed by sequencing. All sequence analyses were carried out using programs in Vector NTI Advance 9.1.0 (Invitrogen, Carlsbad, CA, USA). The recombinant plasmid carrying mRANKL-MT was expressed from a single *E. coli* BL21-CodonPlus (DE3)-RIPL colony using previously described methods [26].

Purification of mRANKL-MT

E. coli cells expressing mRANKL-MT were cultivated in 1 L of an auto-induction medium supplemented with kanamycin (50 µg/mL), as previously described. After centrifugation at 6000 ×g for 20 min at 4°C, the pelleted cells were resuspended in 10 mL of lysis buffer (20 mM sodium phosphate, 500 mM NaCl, 10 mM imidazole, pH 7.4) supplemented with 0.1 mg/mL lysozyme and 0.1 mM PMSF.

Glycerol (20% v/v; CARLO ERBA, France) was added to the cell suspension and the cells were sonicated and centrifuged at 15,000 ×g for 10 min at 4°C. The supernatants were passed through 0.2 µm paper filters and applied to Ni²⁺-affinity chromatography HisTrap FF columns (1 mL; GE Healthcare Life Science, Piscataway, NJ, USA) equilibrated with binding buffer (20 mM sodium phosphate, 500 mM NaCl, 10 mM imidazole, 5 mM dithiothreitol, pH 7.4). The columns were subsequently washed using binding buffer supplemented with 20 mM imidazole.

After washing, bound protein was eluted using elution buffer (Qiagen). The eluted protein was dialysed against a dialysis buffer (20% v/v glycerol in PBS) in a 10,000 MW Slide-A-Lyzer Dialysis cassette (Thermo Fisher Scientific, Waltham, MA, USA). The purified protein was vacuum concentrated (Savant Instruments, Holbrook, NY, USA) and analysed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). Protein concentrations were calculated using the Bradford assay. For endotoxin removal, an additional washing step was introduced after the initial wash for chromatography.

Animal study

The animal experimental protocol was approved by the Institutional Animal Care and Use Committee, Chosun University, Gwangju, Korea (CIACUC2019-A0015). All experiments were performed in accordance with relevant guidelines and regulations. Five-week-old male athymic nude mice (BALB-c/nu, Orient Bio Co. LTD, Seoul, Korea) were used to generate a xenograft model by intracardiac injection of PC3^{Wild}, PC3^{+RANKL} (RANKL overexpression), or PC3^{+RANKL} + IM (immunisation) cells. Following immunisation, mice were divided into an immunisation group and a non-immunisation group. The Sham group was immunised by a subcutaneous injection of PBS, while the immunisation group was injected subcutaneously with mRANKL-MT (100 mg/kg three times every 2 weeks). Mouse sera and tissue samples were collected according to indicated schedule.

***In vivo* bioluminescence measurement**

Tumour-bearing tissues were subjected to *in vivo* bioluminescence imaging using a Living Image® 4.5.4 IVIS Imaging System (Perkin Elmer). For luciferase imaging, D-luciferin (Promega) was injected intraperitoneally before imaging. Quantitative detection of luciferase was performed as follows: regions of interest (ROIs) were drawn to capture detected fluorescence, and auto-regions ROIs were used to precisely outline the target region.

Quantitative analysis of RANKL

The amount of RANKL in mouse serum was measured using a commercially available enzyme-linked immunosorbent assay (ELISA) kit (R&D Systems, USA) according to the manufacturer's protocol. Absorbance was measured in a colorimetric microplate reader (BioTek, USA) at 450 nm.

Measurement of anti-RANKL antibody titers

Serum samples obtained from immunised mice were serially diluted with PBS containing 0.02% sodium azide and 2% bovine serum albumin (BSA), and then applied to ELISA plates (Sigma-Aldrich) coated with mouse recombinant tumour necrosis factor ligand superfamily member 11 (TNFSF11; 10 µg/mL, R&D Systems). Reactivity of serum antibodies to the target protein was determined using an HRP-conjugated goat anti-mouse IgG secondary antibody (Thermo Fisher Scientific) at a dilution of 1/1000 in PBS/0.02% sodium azide/2% BSA. After development with 1,2-phenylenediamine dihydrochloride (0.4 mg/mL in 0.066 M disodium phosphate, 0.035 M citric acid and 0.01% hydrogen peroxide), absorbance was measured in an ELISA plate reader at 450 nm.

Statistical analysis

Data are expressed as the mean ± standard deviation (SD) from three independent experiments. GraphPad Prism version 6.0 software for windows was used to analyse *in vitro* and *in vivo* data. Statistical significance for pairwise comparison was evaluated using an unpaired *t*-test or one-way analysis of variance (ANOVA) with Turkey's post-hoc test. Results were considered significant at **p* < 0.05.

Results

Expression of RANKL in human PCa metastasis tissue specimens

To investigate the association between RANKL and PCa metastasis in human patients, we examined expression of RANKL and metastasis markers interleukin-6 (IL-6) in human PCa metastatic tissue specimens. Immunohistochemical analysis of tissue specimens from patients with PCa metastasis harboured irregularly shaped and dysplastic cells, whereas normal tissues did not. In addition, RANKL expression was elevated, along with that of IL-6, relative to that in normal tissue. (Fig. 1). Thus, RANKL expression was closely related to PCa metastasis, suggestive of an important role in this process.

Effects of human RANKL (hRANKL) on EMT and metastasis of PC3 cells

To explore the relationship between RANKL and EMT of PC3 cells, we performed cell migration and invasion assays with PC3 cells treated with hRANKL.

In the invasion assay, wound healing in hRANKL-treated PC3 cells was significantly better than that in control cells (Fig. 2A). Cell invasion also increased significantly following hRANKL treatment (Fig. 2B).

EMT is closely related to tumour metastasis and progression. Therefore, to determine changes at the molecular level, we measured EMT markers in PC3 cells at the mRNA (Supp. 1) and protein (Fig. 2C) levels following treatment with hRANKL. Expression of the EMT marker E-cadherin in hRANKL-treated PC3 cells fell significantly, but that of vimentin and β -catenin increased. Expression of MMP-9 and IL-6 (markers of metastasis) were significantly higher in hRANKL-treated cells than in control cells. Expression of E-cadherin protein was significantly lower in hRANKL-treated PC3 cells than in control cells. By contrast, expression of vimentin, β -catenin and MMP-9 proteins was significantly upregulated following treatment with hRANKL.

The Wnt/ β -catenin pathway regulates PCa metastasis and plays an important role in cancer progression; therefore, we examined the effects of RANKL on Wnt signalling using a TOP/FOP reporter assay (Fig. 2D). No changes in TOP/FOP reporter luciferase activity were observed in hRANKL-treated PC3 cells.

Next, we examined phosphorylation of extracellular signal-regulated kinase (ERK), protein kinase B (AKT), SRC and GSK-3B to investigate the effects of RANKL on mitogen-activated protein kinase (MAPK) and Wnt signalling in PC3 cells (Fig. 2E). PC3 cells treated with hRANKL showed a significant and time-dependent reduction in the level of phosphorylated GSK-3B. A time-dependent increase in SRC and AKT phosphorylation levels was also observed.

Next, we performed co-immunoprecipitation of β -catenin and TCF-4 to investigate the status of TCF-associated signalling in hRANKL-treated PC3 cells (Fig. 2F). The results showed a slight increase in TCF-4 levels in hRANKL-treated PC3 cells. Thus, RANKL treatment may trigger metastasis of PC3 cells by suppressing GSK-3B phosphorylation and facilitating EMT.

Overexpression of RANKL modulates EMT and metastasis of PC3 cells

To investigate whether RANKL overexpression stimulates PC3 cell growth *in vitro*, cells were transiently transfected with an overexpression plasmid containing RANKL. GFP expression by PC3^{RANKL} cells was monitored by fluorescence microscopy (Supp. 2A). We found a significant increase in expression of RANKL mRNA and protein (Supple 2B and 2C).

Next, we performed migration and invasion assays using PC3^{RANKL} cells to evaluate the effects of RANKL overexpression on EMT and metastasis. Wound healing was significantly better in PC3^{RANKL} cells than in control cells (Fig. 3A). Also, PC3^{RANKL} cells were significantly more invasive than control cells (Fig. 3B). Analysis of mRNA encoding EMT- and metastasis-related factors in PC3^{RANKL} cells revealed significant downregulation of the gene encoding E-cadherin (Supp. 2D). Furthermore, expression of genes encoding vimentin, MMP-9, IL-6 and β -catenin was upregulated significantly in PC3^{RANKL} cells. Protein expression analysis revealed that PC3^{RANKL} cells expressed significantly lower levels of E-cadherin than PC3^{Wild} cells (Fig. 3C). By contrast, expression of N-cadherin, MMP-9, IL-6, c-MYC and β -catenin was significantly higher in PC3^{RANKL} cells.

PC3^{RANKL} cells showed a significant increase in TOP/FOP luciferase reporter activity (Fig. 3D). Immunoprecipitation of β -catenin was carried out to investigate the signal transduction pathway associated with TCF in PC3^{RANKL} cells (Fig. 3E). PC3^{RANKL} cells overexpressing RANKL showed a significant increase in activation of the MAPK and β -catenin/TCF-4 signalling pathways owing to stronger binding between β -catenin and TCF-4 than in PC3^{Wild} cells. Thus, ectopic overexpression of RANKL may increase EMT and the metastatic properties of PC3 cells via the β -catenin/TCF-4 signalling pathway, suggesting the therapeutic potential of RANKL targeting for prevention of PCa metastasis.

Therapeutic effects of mRANKL-MT in PC3 cell-inoculated mice

Alignment of the mRNA sequence of mRANKL-MT with that of mRANKL-WT identified a region that could be amplified using selected primers. The recombinant mRANKL-WT sequence encoded the full-length 158 amino acid target region, which includes residues 158 to 316 (Fig. 4A). To create point mutations, Lys180, Asp189-Arg190 and His223-His224 were transformed to Arg180, Ile189-Lys190 and Phe223-Tyr224, respectively. The resulting hRANKL, mRANKL-WT and mRANKL-MT molecules had similar molecular weights (Supp. 3). Male BALB-c/nu mice were injected subcutaneously with hRANKL, mRANKL-WT, or mRANKL-MT (100 mg/kg; three times every 2 weeks). After immunisation, 1×10^6 PC3^{Wild} or PC3^{RANKL+} cells were injected into the left ventricle of Sham or immunised mice. Serum and tumour-bearing tissues were collected after 16 weeks (Fig. 4B).

To observe bone metastasis, luciferase activity in tumour-bearing tissues of Sham, PC3^{Wild}, PC3^{RANKL+} and PC3^{RANKL+} + IM mice was detected by IVIS. In PC3^{RANKL+} mice, large and strong bioluminescence spots were detected throughout the body at 16 weeks post-cancer cell injection (Fig. 4C). However, no

bioluminescence signals were detected in PC3^{RANKL+} + IM mice. The photon flux values were significantly higher in PC3^{RANKL+} mice than in PC3^{RANKL+} + IM mice (Fig. 4D).

Survival rate analysis revealed a significant decrease in the survival of animals in the PC3^{RANKL+} groups compared with that of animals from the PC3^{Wild} group. The survival rate improved higher in the PC3^{RANKL+} + IM group than that in the PC3^{RANKL+} group (Fig. 4E). The metastasis rate in the PC3^{RANKL+} group was higher than that in the PC3^{Wild} group; however, that in the PC3^{RANKL+} + IM group was significantly less than that in the PC3^{RANKL+} group (Fig. 4F).

Therapeutic effects of anti-RANKL antibodies induced by RANKL immunisation

To examine the histological characteristics of metastatic tumour-bearing tissues, metastatic lesions from each mouse were stained with haematoxylin and eosin Y (H&E). As shown in Figure 5A, gross examination of the excised tibiae from PC3^{RANKL+} mice revealed a tumour mass in the primary spongiosum (trabecular epiphysis) and bone marrow cells; this was not observed in PC3^{RANKL+} + IM mice. In particular, expressions of IL-6 and RANKL increased markedly in the trabecular epiphysis region of bones from PC3^{RANKL+} mice, but were undetectable in PC3^{RANKL+} + IM mice.

Next, we investigated whether mRANKL-MT induces production of anti-RANKL antibodies. The concentration of RANKL (Fig. 5B) was highest in PC3^{RANKL+}, and production of antibodies (Fig. 5C) was highest in PC3^{RANKL+} + IM mice. Also, we measured anti-RANKL antibody levels in the PC3^{RANKL+} + IM group with bone metastasis to investigate generation of anti-RANKL antibodies after immunisation with mRANKL-MT (Fig. 5D). The anti-RANKL titer in mice with bone metastasis was significantly higher than that in mice without bone metastasis. Also, serum RANKL levels in the PC3^{RANKL+} + IM group without metastasis were significantly lower than those in mice with metastasis (Fig. 5E). These observations suggest that anti-RANKL antibodies generated by RANKL immunisation suppress metastasis of PCa cells.

Effect of immunisation on EMT and metastasis

To investigate the effects of mRANKL-MT on PCa metastasis, sera obtained from immunised mice were used to treat RANKL-overexpressing PC3 cells. The results of cell migration assays showed that wound healing was inhibited significantly in PC3^{RANKL+} cells treated with immune serum (Fig. 6A). In addition, the invasive ability of PC3^{RANKL+} cells declined following treatment with immune sera (Fig. 6B).

Analysis of mRNA encoding EMT- and metastasis-related factors in PC3^{RANKL+} cells treated with immune sera revealed significant upregulation of E-cadherin expression (Supp. 4). By contrast, expression of vimentin, MMP-9, IL-6 and β -catenin was downregulated significantly in immune serum-treated PC3^{RANKL+} cells. Protein expression analysis showed that E-cadherin expression was higher in immune serum-treated PC3^{RANKL+} cells than in control serum-treated PC3^{RANKL+} cells (Fig. 6C). By contrast, expression of

N-cadherin, MMP-9, IL-6 and β -catenin was significantly lower in immune serum-treated PC3^{RANKL+} cells, as was expression of c-MYC.

The luciferase activity of the TOP/FOP reporter in immunised serum-treated PC3^{RANKL+} cells fell significantly (Fig. 6D). Immunoprecipitation analysis revealed a significant decrease in binding between β -catenin and TCF-4 in cells treated with immune serum (Fig. 6E). Finally, we measured expression of phosphorylated ERK, AKT, SRC and GSK-3B to investigate the effect of immune serum on MAPK and Wnt signalling in PC3^{RANKL+} cells (Fig. 6F). Cells treated with immune serum showed a significant and time-dependent increase in GSK-3B phosphorylation and a time-dependent decrease in SRC phosphorylation. These results indicate that the EMT and metastatic properties of PC3^{RANKL+} cells were inhibited by treatment with immune serum.

Discussion

PCa, which is common among men in the western world, is associated with high mortality and morbidity with respect to advanced metastasis to the bone. Evidence suggests that the RANKL signalling cascade plays a key role in proliferation, metastasis, migration and invasion of PCa [6]. The RANKL–RANK interaction plays a pivotal role in PCa metastasis; indeed, RANKL expression induces osteoclast hyperplasia and bone destruction during PCa metastasis [27]. RANKL activates RANK directly on tumour cells, as evidenced by dysregulation of several biochemical signalling pathways in PCa cells.

High expression of RANKL facilitates PCa metastasis, an idea consistent with previous studies showing that signalling through the RANK/RANKL axis is related to bone metastases of solid tumours [20, 28]. Here, we demonstrated RANKL immunoreactivity in bone metastatic lesions of PCa patients. We also found that hRANKL-treated or hRANKL-overexpressing PCa cells showed a significant increase in expression of metastasis markers such as IL-6. In particular, *in vitro* experiments show that RANKL stimulation markedly increases the migration and invasion of PC3 cells, downregulates expression of the epithelial marker E-cadherin and upregulates the mesenchymal marker vimentin. EMT correlates with tumour metastasis and progression, which is consistent with impaired cell–cell adhesion following the loss of E-cadherin expression [29, 30]. Furthermore, we show that GSK-3B phosphorylation was reduced significantly following RANKL treatment of PC3 cells due to the effect of RANKL on MAPK and Wnt signalling. Also, we observed altered expression of β -catenin and TCF-4 in PC3^{RANKL+} cells, which resulted in a highly conserved developmental signalling pathway that includes the major effector protein β -catenin. Wnt signalling is an essential pathway involved in cell development, proliferation and differentiation; indeed, regulatory abnormalities in Wnt signalling are associated with metastasis of many cancers [31]. In particular, RANKL overexpression in PC3 cells led to a significant increase in expression of Wnt3a, suggesting that RANKL is a potential target of Wnt signalling in cancer cells [32]. RANKL plays a fundamental role in osteoclastogenesis by interacting with the RANK receptor on osteoclast progenitors during bone destruction by metastatic breast cancer, thereby driving osteoclast cell lineage commitment, monocyte cell fusion and osteoclast maturation via regulation of NF- κ B-mediated gene expression;

therefore, we were intrigued to find out whether catabolic Wnt signalling mechanisms exist alongside anabolic Wnt pathways to regulate osteoclast formation in bone. β -catenin, the critical effector of the Wnt pathway, regulates a number of key processes during development, including proliferation, differentiation and cell fate determination [33]. Normally, β -catenin is localised to the cell adhesion junctions in epithelial cells and its abnormal cytoplasmic/nuclear stabilisation drives uncontrolled transcription of target genes (including c-jun, cyclin D1, c-myc, survivin and MMP-7) that regulate cell proliferation, survival and adhesion [34]. In view of cancer cell fate, it is not surprisingly that overexpression of RANKL by PC3 cells led to increased binding of β -catenin to TCF4 and to increased TOP activity. Regulation of β -catenin is linked to the pathogenesis of a number of human cancers, particularly those with an epithelial cell origin. Supporting its putative role as a Wnt signalling target, we confirmed that RANKL overexpression led to transcriptional activation of β -catenin in PC3 cells.

Over the past decades, it has become clear that the RANK/RANKL axis exerts a broad range of functions during cancer cell fate. In the cancer setting, the RANK-RANKL pathway plays a role in every stage of tumorigenesis. Therefore, inhibition of RANKL by anti-RANKL antibodies is expected to be more far-reaching than simple inhibition of cancer cell activation. Denosumab, a drug used to treat metastatic prostate bone loss, has received FDA approval; this drug inhibits the RANK-RANKL pathway [35]. Denosumab is an effective and safe drug, which is superior to zoledronic acid in terms of skeletal-related events prevention; this was borne out by a combined analysis that included three randomised phase III trials with a similar set-up [36]. These trials included patients with bone metastases due to advanced breast cancer [37], prostate cancer [38], other solid tumours or multiple myeloma [39]. However, despite medical and commercial success, passive anti-cytokine drugs such as OPG-Fc and denosumab have several limitations, including high production costs, the need for regular infusion and a limited half-life [40]. Thus, the aim is to develop a RANKL vaccine. In comparison with antibodies and other biologics, vaccines are better models for treatment of chronic disease because they are relatively cheap and small doses of protein can have a strong and long-lasting effect [41, 42]. Here, we developed a novel vaccine targeting RANKL and examined its efficacy in a murine model of prostate cancer metastasis. To circumvent the problem of the immunogen triggering cytokine activity, mutants of RANKL were generated to prevent its interaction with RANK. A previous study shows that immunisation with mutant RANKL molecules generates anti-RANKL antibodies that block the interaction between RANKL and its receptor in an animal model of osteoporosis, thereby preventing proliferation and differentiation of osteoclasts and improving bone density [43]. Therefore, to block RANKL activation during PCa metastasis, we immunised mice with mRANKL-MT followed by intracardiac injection of PC3 cells. Inhibiting RANKL in animal models of metastases exerts therapeutic effects by inhibiting cancer cell metastasis. Currently, *in vivo* tumour models that are most commonly used to study the process of cancer metastasis rely on introduction of tumour cells directly into the systemic circulation by injection into the left ventricle of laboratory rodents [44, 45]. Thus, we employed a mouse model of PCa metastasis that more accurately reflects the metastatic process of this type of cancer. Studies on the effects of mRANKL-MT in PC3^{RANKL+} mice showed that tumour growth was completely inhibited. Immunisation with mRANKL-MT effectively inhibited metastasis of tumour cells by generating anti-RANKL antibodies.

To further confirm the action of RANKL immunisation, we assessed the effects of immune serum from immunised mice on PCa cells. Anti-RANKL antibodies blocked the RANKL-mediated chemotaxis of tumour cells. Furthermore, anti-RANKL antibodies inactivated RANKL on tumour cells directly. Treatment of RANKL-overexpressing PC3 cells with immune serum almost entirely abolished cancer cell migration and invasion. Wu et al. showed that a recombinant inactive RANKL vaccine (Y234pNO2Phe) induced high antibody titers and protected mice from collagen-induced arthritis by inhibiting osteoclast function and by preventing bone erosion [46]. However, no study has reported that these types of RANKL vaccine have been used to inhibit cancer metastasis. In addition, we found that EMT and metastasis-related genes were downregulated following treatment of RANKL-overexpressing PC3 cells with immune serum. The antisera obtained from mice immunised with mRANKL-MT almost entirely inhibited the EMT process in RANKL-overexpressing PC3 cells. EMT is characterised by the loss of cell–cell adhesion and by an increase in cell motility; it is a key process in cancer progression and metastasis, making EMT inhibition an attractive therapeutic strategy [47, 48]. Deregulation of Wnt/ β -catenin signalling is a hallmark of PCa metastasis [49, 50] and β -catenin is a critical end component of the Wnt signalling pathway, which regulates cell growth, apoptosis and migratory behaviour in response to intercellular adhesion molecules [51]. Activation of β -catenin in PCa cells leads to transactivation of Wnt signalling target genes, including cyclin D1, HEF1 and matrix metalloproteinase 9 [52]. Also, previous studies show that expression of Wnt-1 and β -catenin is increased in invasive PCa cell lines and in primary prostate cancer specimens [53]. In line with these previous reports, we demonstrated that treatment of PC3 cells with immune serum led to a marked decrease in RANK/SRC and GSK-3 β signalling and β -catenin/TCF-4 transcription (Fig. 7). β -catenin forms a cell adhesion complex with E-cadherin, raising the possibility that loss of expression or a change in β -catenin distribution in the cell alters downstream signalling, decreases intercellular adhesion and promotes metastasis. These results suggest that the inhibitory effect of immune serum on PCa cell metastasis may involve suppression of the Wnt/ β -catenin signalling pathway.

In summary, this study demonstrates the protective role of mRANKL-MT against RANKL-induced PCa in mice. This effect was mediated via induction of a high-titer antibody response, inhibition of EMT and metastatic functions. Our results highlight the potential application of an anti-RANKL vaccine for treatment of metastatic RANKL-induced PCa. Moreover, the results suggest that mutant RANKL may be a potential RANKL vaccine that prevents and/or treats RANKL targeting in patients with metastatic PCa prostate cancer.

Abbreviations

PCa: Prostate cancer; NF- κ B: Nuclear factor kappa-light-chain-enhancer of activated B cells; RANKL: receptor activator of NF- κ B ligand; EMT: epithelial-mesenchymal transition; TCF-4: Transcription factor 4; TRANCE: tumour necrosis factor-related activation-induced cytokine; OPGL: osteoprotegerin ligand; ODF: osteoclastic differentiation factor; GSK-3: glycogen synthase kinase-3; mRANKL-MT: mouse RANKL mutant; IL-6: Interleukin-6; RPMI: Roswell Park Memorial Institute; FBS: Fetal bovine serum; HRP: Horseradish peroxidase; PCR: Polymerase chain reaction; IHC: Immunohistochemistry;

Declarations

Authors' contributions

M. P., Y. C. and W. L. designed and performed the research, analysed and interpreted the data, and wrote the paper; B. K., Y. K., Y. J. and Y. J. analysed and interpreted the *in vitro* data, and helped to draft some of the figures; W. L. performed and supervised RT-PCR and western blot data analysis; B. K. and Y. K. assisted with the ELISA assays; M. P., Y. C., B. K., Y. C., M. P., Y. K. and Y. J. performed *in vivo* studies under the supervision of W. L.; M. P. and Y. J. performed histologic analysis under the supervision of H. H.; M. P. and Y. C. performed pathological review of the samples; and W. L. analysed publicly available datasets and supervised the study.

Funding

This study was supported by research funding from Chosun University (awarded in 2020).

Ethics approval and consent to participate

Patient study was approved and supervised by Chosun University Hospital's institutional review board. (Chosun University Hospital, Gwangju, Korea (CHOSUN 2020-06-001)). And animal experimental protocol was approved by the Institutional Animal Care and Use Committee, Chosun University, Gwangju, Korea (CIACUC2019-A0015) and

Consent for publication

Not applicable.

Competing interests

The authors declare no financial conflicts of interest.

Author details

¹Laboratory of Orthopaedic Research, Chosun University Hospital, Dong-Gu, Gwangju, 61452, Republic of Korea

²Department of Orthopaedic Surgery, Chosun University Hospital, Dong-Gu, Gwangju, 61452, Republic of Korea

³Department of Biomedical Sciences Chonnam National University Medical School, Gwangju 61469, Republic of Korea

⁴Department of Premedical Science, College of Medicine, Chosun University, Dong-Gu, Gwangju 61452, Republic of Korea

References

1. Rawla P: Epidemiology of Prostate Cancer. *World J Oncol* 2019, 10:63-89.
2. Logothetis CJ, Lin SH. Osteoblasts in prostate cancer metastasis to bone. *Nat Rev Cancer*. 2005;5:21-28.
3. Jacobs SC: Spread of prostatic cancer to bone. *Urology*. 1983;21:337-344.
4. Neudert M, Fischer C, Krempien B, Bauss F, Seibel MJ: Site-specific human breast cancer (MDA-MB-231) metastases in nude rats: model characterisation and in vivo effects of ibandronate on tumour growth. *Int J Cancer*. 2003;107:468-477.
5. Zheng Y, Zhou H, Brennan K, Blair JM, Modzelewski JR, Seibel MJ, Dunstan CR: Inhibition of bone resorption, rather than direct cytotoxicity, mediates the anti-tumour actions of ibandronate and osteoprotegerin in a murine model of breast cancer bone metastasis. *Bone*. 2007;40:471-478.
6. Armstrong AP, Miller RE, Jones JC, Zhang J, Keller ET, Dougall WC: RANKL acts directly on RANK-expressing prostate tumor cells and mediates migration and expression of tumor metastasis genes. *Prostate*. 2008;68:92-104.
7. Ziaee S, Chu GC, Huang JM, Sieh S, Chung LW: Prostate cancer metastasis: roles of recruitment and reprogramming, cell signal network and three-dimensional growth characteristics. *Transl Androl Urol*. 2015; 4:438-454.
8. Price JT, Quinn JM, Sims NA, Vieusseux J, Waldeck K, Docherty SE, Myers D, Nakamura A, Waltham MC, Gillespie MT, Thompson EW: The heat shock protein 90 inhibitor, 17-allylamino-17-demethoxygeldanamycin, enhances osteoclast formation and potentiates bone metastasis of a human breast cancer cell line. *Cancer Res*. 2005;65:4929-4938.
9. Zheng Y, Zhou H, Ooi LL, Snir AD, Dunstan CR, Seibel MJ: Vitamin D deficiency promotes prostate cancer growth in bone. *Prostate*. 2011;71:1012-1021.
10. Ooi LL, Zheng Y, Zhou H, Trivedi T, Conigrave AD, Seibel MJ, Dunstan CR: Vitamin D deficiency promotes growth of MCF-7 human breast cancer in a rodent model of osteosclerotic bone metastasis. *Bone*. 2010;47:795-803.
11. Wong BR, Josien R, Lee SY, Sauter B, Li HL, Steinman RM, Choi Y: TRANCE (tumor necrosis factor [TNF]-related activation-induced cytokine), a new TNF family member predominantly expressed in T cells, is a dendritic cell-specific survival factor. *J Exp Med*. 1997;186:2075-2080.
12. Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, et al: Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell*. 1998;93:165-176.
13. Kong YY, Boyle WJ, Penninger JM: Osteoprotegerin ligand: a common link between osteoclastogenesis, lymph node formation and lymphocyte development. *Immunol Cell Biol*. 1999;77:188-193.
14. Kodaira K, Kodaira K, Mizuno A, Yasuda H, Shima N, Murakami A, Ueda M, Higashio K: Cloning and characterization of the gene encoding mouse osteoclast differentiation factor. *Gene*. 1999;230:121-

127.

15. Renema N, Navet B, Heymann MF, Lezot F, Heymann D: RANK-RANKL signalling in cancer. *Biosci Rep.* 2016;36.
16. Kartsogiannis V, Zhou H, Horwood NJ, Thomas RJ, Hards DK, Quinn JM, Niforas P, Ng KW, Martin TJ, Gillespie MT: Localization of RANKL (receptor activator of NF kappa B ligand) mRNA and protein in skeletal and extraskeletal tissues. *Bone.* 1999;25:525-534.
17. Collin-Osdoby P, Rothe L, Anderson F, Nelson M, Maloney W, Osdoby P: Receptor activator of NF-kappa B and osteoprotegerin expression by human microvascular endothelial cells, regulation by inflammatory cytokines, and role in human osteoclastogenesis. *J Biol Chem.* 2001;276:20659-20672.
18. Wang XF, Zhang YK, Yu ZS, Zhou JL: The role of the serum RANKL/OPG ratio in the healing of intertrochanteric fractures in elderly patients. *Mol Med Rep.* 2013;7:1169-1172.
19. Tan W, Zhang W, Strasner A, Grivennikov S, Cheng JQ, Hoffman RM, Karin M: Tumour-infiltrating regulatory T cells stimulate mammary cancer metastasis through RANKL-RANK signalling. *Nature.* 2011;470:548-553.
20. Jones DH, Nakashima T, Sanchez OH, Kozieradzki I, Komarova SV, Sarosi I, Morony S, Rubin E, Sarao R, Hojilla CV, et al: Regulation of cancer cell migration and bone metastasis by RANKL. *Nature.* 2006;440:692-696.
21. Schramek D, Leibbrandt A, Sigl V, Kenner L, Pospisilik JA, Lee HJ, Hanada R, Joshi PA, Aliprantis A, Glimcher L, et al: Osteoclast differentiation factor RANKL controls development of progestin-driven mammary cancer. *Nature.* 2010;468:98-102.
22. Savagner P: The epithelial-mesenchymal transition (EMT) phenomenon. *Ann Oncol.* 2010;21 Suppl 7:vii89-92.
23. Cho KB, Cho MK, Lee WY, Kang KW: Overexpression of c-myc induces epithelial mesenchymal transition in mammary epithelial cells. *Cancer Lett.* 2010;293:230-239.
24. Sharma P, Wagner K, Wolchok JD, Allison JP: Novel cancer immunotherapy agents with survival benefit: recent successes and next steps. *Nat Rev Cancer.* 2011;11:805-812.
25. Liu C, Zhao Y, He W, Wang W, Chen Y, Zhang S, Ma Y, Gohda J, Ishida T, Walter TS, et al: A RANKL mutant used as an inter-species vaccine for efficient immunotherapy of osteoporosis. *Sci Rep.* 2015;5:14150.
26. Skeletal Complications of Cancer. Special Issue dedicated to Gregory Robert Mundy. Proceedings of the 10th International Conference on Cancer-Induced Bone Disease. Sheffield, United Kingdom. September 22-25, 2010. *Bone.* 2011;48:5-166, S162-155.
27. Roodman GD: Mechanisms of bone metastasis. *N Engl J Med* 2004;350:1655-1664.
28. Santini D, Perrone G, Roato I, Godio L, Pantano F, Grasso D, Russo A, Vincenzi B, Fratto ME, Sabbatini R, et al: Expression pattern of receptor activator of NFkappaB (RANK) in a series of primary solid tumors and related bone metastases. *J Cell Physiol.* 2011;226:780-784.

29. Nieto MA: Epithelial plasticity: a common theme in embryonic and cancer cells. *Science*. 2013;342:1234850.
30. Tan EJ, Kahata K, Idas O, Thuault S, Heldin CH, Moustakas A: The high mobility group A2 protein epigenetically silences the *Cdh1* gene during epithelial-to-mesenchymal transition. *Nucleic Acids Res*. 2015;43:162-178.
31. Lamouille S, Xu J, Derynck R: Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol*. 2014;15:178-196.
32. Spencer GJ, Utting JC, Etheridge SL, Arnett TR, Genever PG: Wnt signalling in osteoblasts regulates expression of the receptor activator of NFkappaB ligand and inhibits osteoclastogenesis in vitro. *J Cell Sci*. 2006;119:1283-1296.
33. Pedone E, Marucci L: Role of beta-Catenin Activation Levels and Fluctuations in Controlling Cell Fate. *Genes (Basel)*. 2019;10.
34. Anand M, Lai R, Gelebart P: beta-catenin is constitutively active and increases STAT3 expression/activation in anaplastic lymphoma kinase-positive anaplastic large cell lymphoma. *Haematologica*. 2011;96:253-261.
35. Goldstein DA: Denosumab for bone lesions in multiple myeloma - what is its value? *Haematologica*. 2018;103:753-754.
36. Lipton A, Fizazi K, Stopeck AT, Henry DH, Brown JE, Yardley DA, Richardson GE, Siena S, Maroto P, Clemens M, et al: Superiority of denosumab to zoledronic acid for prevention of skeletal-related events: a combined analysis of 3 pivotal, randomised, phase 3 trials. *Eur J Cancer*. 2012;48:3082-3092.
37. Stopeck AT, Lipton A, Body JJ, Steger GG, Tonkin K, de Boer RH, Lichinitser M, Fujiwara Y, Yardley DA, Viniegra M, et al: Denosumab compared with zoledronic acid for the treatment of bone metastases in patients with advanced breast cancer: a randomized, double-blind study. *J Clin Oncol*. 2010;28:5132-5139.
38. Fizazi K, Carducci M, Smith M, Damiao R, Brown J, Karsh L, Milecki P, Shore N, Rader M, Wang H, et al: Denosumab versus zoledronic acid for treatment of bone metastases in men with castration-resistant prostate cancer: a randomised, double-blind study. *Lancet*. 2011;377:813-822.
39. Henry DH, Costa L, Goldwasser F, Hirsh V, Hungria V, Prausova J, Scagliotti GV, Sleeboom H, Spencer A, Vadhan-Raj S, et al: Randomized, double-blind study of denosumab versus zoledronic acid in the treatment of bone metastases in patients with advanced cancer (excluding breast and prostate cancer) or multiple myeloma. *J Clin Oncol*. 2011;29:1125-1132.
40. Krishna M, Nadler SG: Immunogenicity to Biotherapeutics - The Role of Anti-drug Immune Complexes. *Front Immunol*. 2016;7:21.
41. Semerano L, Assier E, Boissier MC: Anti-cytokine vaccination: a new biotherapy of autoimmunity? *Autoimmun Rev*. 2012;11:785-786.
42. Bachmann MF, Dyer MR: Therapeutic vaccination for chronic diseases: a new class of drugs in sight. *Nat Rev Drug Discov*. 2004;3:81-88.

43. Ko Y, Lee G, Kim B, Park M, Jang Y, Lim W: Modification of the RANKL-RANK-binding site for the immunotherapeutic treatment of osteoporosis. *Osteoporos Int.* 2020; 31:983-993.
44. Rosol TJ, Tannehill-Gregg SH, Corn S, Schneider A, McCauley LK: Animal models of bone metastasis. *Cancer Treat Res.* 2004;118:47-81.
45. Simmons JK, Hildreth BE, 3rd, Supsavhad W, Elshafae SM, Hassan BB, Dirksen WP, Toribio RE, Rosol TJ: Animal Models of Bone Metastasis. *Vet Pathol.* 2015;52:827-841.
46. Wu T, Li F, Sha X, Li F, Zhang B, Ma W, Liu M, Yang W, Li H, Tao H: A novel recombinant RANKL vaccine prepared by incorporation of an unnatural amino acid into RANKL and its preventive effect in a murine model of collagen-induced arthritis. *Int Immunopharmacol.* 2018;64:326-332.
47. Wang Y, Zhou BP: Epithelial-mesenchymal Transition—A Hallmark of Breast Cancer Metastasis. *Cancer Hallm.* 2013;1:38-49.
48. Montanari M, Rossetti S, Cavaliere C, D'Aniello C, Malzone MG, Vanacore D, Di Franco R, La Mantia E, Iovane G, Piscitelli R, et al: Epithelial-mesenchymal transition in prostate cancer: an overview. *Oncotarget.* 2017;8:35376-35389.
49. Vatansever HS, Gumus B, Aydogdu O, Sivrikoz ON, Turkoz-Uluer E, Kivanc M, Atesci YZ, Bugdayci H: The role of stem/progenitor cells and Wnt/beta-catenin signaling pathway in the patients with prostate cancer. *Minerva Urol Nefrol.* 2014;66:249-255.
50. Jiang YG, Luo Y, He DL, Li X, Zhang LL, Peng T, Li MC, Lin YH: Role of Wnt/beta-catenin signaling pathway in epithelial-mesenchymal transition of human prostate cancer induced by hypoxia-inducible factor-1alpha. *Int J Urol.* 2007;14:1034-1039.
51. Harjunpaa H, Llort Asens M, Guenther C, Fagerholm SC: Cell Adhesion Molecules and Their Roles and Regulation in the Immune and Tumor Microenvironment. *Front Immunol.* 2019;10:1078.
52. Liu Z, Rebowe RE, Wang Z, Li Y, Wang Z, DePaolo JS, Guo J, Qian C, Liu W: KIF3a promotes proliferation and invasion via Wnt signaling in advanced prostate cancer. *Mol Cancer Res.* 2014;12:491-503.
53. Francis JC, Thomsen MK, Taketo MM, Swain A: beta-catenin is required for prostate development and cooperates with Pten loss to drive invasive carcinoma. *PLoS Genet.* 2013;9:e1003180.

Figures

Fig.1




Figure 1

Histopathology of metastatic prostate cancer tissues. Representative H&E stained and immunostained images showing expression of RANKL and IL-6 in normal (no histopathological change) or metastasis tissues. Magnification: 200 \times . Scale = 100 μ m.

Fig.2




Figure 2

EMT and metastatic properties of PC3 cells following *hRANKL* treatment. (A) Treatment with *hRANKL* led to a significant increase in the migratory capacity of PC3 cells, as evident from the decrease in the wound gap distance at 48 h. Magnification, 100 \times ; scale bar, 100 μm . The data in the associated graphs are expressed as the mean \pm SD. (B) Transwell invasion assays were performed to compare the invasiveness of *hRANKL*-treated PC3 cells and untreated cells. Treatment with *hRANKL* led to a significant reduction in the invasiveness of PC3 cells. Magnification, 100 \times ; scale bar, 100 μm . Data in the associated graphs are expressed as the mean \pm SD. (C) Expression of EMT and metastasis-related proteins E-cadherin, N-cadherin, β -catenin, MMP-9, IL-6, c-MYC, TCF-4 and β -actin in *hRANKL*-treated PC3 cells was measured by western blotting. β -actin was used as a loading control. (D) TOP/FOP luciferase reporter assays in PC3Wild and PC3RANKL+ cells. (E) Western blot analysis of MAPK phosphorylation in PC3 cells incubated with 1 $\mu\text{g/mL}$ *hRANKL* for 0, 5, 15 and 30 min. (F) Co-immunoprecipitation for β -catenin and

TCF-4 in hRANKL-treated PC3 cells. Results are representative of three separate experiments with comparable results.

Fig.3




Figure 3

Modulation of the EMT and metastatic properties of PC3 cells following overexpression of RANKL. (A) A cell migration assay was performed to compare wound healing in PC3^{Wild} and PC3^{RANKL+} cells. PC3^{RANKL+} cells showed a significant increase in migratory capacity, as evident from the decreased wound gap at 48 h. Magnification, 100 \times ; scale bar, 100 μ m. The data in the associated graphs are expressed as the mean \pm SD. (B) Transwell invasion assays were performed to compare the invasiveness of PC3^{Wild} and PC3^{RANKL+} cells. PC3^{RANKL+} cells showed a significant increase in invasiveness. Magnification, 100 \times ; scale bar, 100 μ m. Data in the associated graphs are expressed as the mean \pm SD. (C) Expression of E-cadherin, N-cadherin, β -catenin, MMP-9, IL-6 and β -actin in PC3^{Wild} and PC3^{RANKL+} cells, as measured by western blotting. β -actin was used as a loading control. (D) TOP/FOP luciferase reporter assays in PC3^{Wild} and PC3^{RANKL+} cells. Significant differences were observed at * $p < 0.05$, compared with the control. (E) Immunoprecipitation for β -catenin and TCF-4 in PC3^{Wild} and PC3^{RANKL+}

cells. Each blot was obtained under the same experimental conditions and the data were representative of three separate experiments with comparable results.

Fig.4




Figure 4

Effects of immunisation with mutant RANKL. (A) Comparison of the sequences of hRANKL, mRANKL-WT and mRANKL-MT. (B) In vivo experimental flow. Male BALB-c/nu mice (6-weeks-old) were immunised with a subcutaneous injection of test agent (20 µg/kg). (C) Tumour growth and metastases were monitored via in vivo bioluminescence imaging of the reporter activity induced by PC3^{Wild}, PC3^{RANKL+} and PC3^{RANKL+} + IM treatment. Data per representative mouse are shown. A large hot-spot of bioluminescence was observed in vivo following inoculation of mice with PC3^{RANKL+} cells. Multiple

localised and distant metastases were observed in vivo after the injection of PC3RANKL+ cells into the hearts of nude mice. Coloured bars indicate the bioluminescence signal intensity (photon/s/cm²/steradian). (D) The bioluminescence photon flux that appeared after inoculation of cells into mice is shown graphically. Data from related graphs are displayed as the mean \pm SD. (E) Survival and (F) metastasis rate in mice from the SHAM, PC3Wild, PC3RANKL+ and PC3RANKL+ + IM groups.

Fig.5




Figure 5

Effects of immunisation with mutant RANKL on PC3-innoculated metastatic model. (A) representative H&E stained and immunostained images of RANKL and IL-6 in the PC3Wild, PC3RANKL+ and PC3RANKL+ + IM groups. Magnification: 200 \times . Scale bar = 100 μ m. (B) Concentration of RANKL in mouse serum. The mean \pm SD values were obtained by densitometry, as shown in the analysis. Significant differences were observed at * p < 0.05 and ** p < 0.01 vs. control. (C) Serum samples from mice were obtained after immunisation. Anti-RANKL values in PC3Wild, PC3RANKL+ and PC3RANKL+ + IM groups. The mean \pm SD values were obtained by densitometry, as shown in the analysis. Significant differences were observed at * p < 0.05 and ** p < 0.01 vs. the control. (D) Anti-RANKL values and (E) RANKL concentration in the serum of RANKL-immunized mice with metastasis (+) or metastasis (-). Bar graphs show the mean \pm standard deviation (SD). Significant differences were observed at * p < 0.05, metastasis (+) vs. no metastasis (-).

Fig.6




Figure 6

EMT and metastatic properties of PC3 cells treated with immune sera. (A) Cell migration assays were performed to compare the wound healing capacity of immune serum-treated PC3^{RANKL+} cells and untreated cells. Immune serum treatment decreased the migration capacity of PC3 cells, as evident from the increase in the wound healing gap at 48 h. Magnification, 100 \times ; scale bar, 100 μ m. Data from related graphs are displayed as the mean \pm SD. (B) Transwell invasion assays were performed to compare the invasiveness of immunised serum-treated PC3^{RANKL+} and untreated cells. Serum-treated PC3^{RANKL+}

cells showed a significant decrease in invasiveness. Magnification, 100 \times ; scale bar, 100 μ m. Data in the associated graphs are expressed as the mean \pm SD. (C) Expression of E-cadherin, N-cadherin, β -catenin, MMP-9, IL-6, c-MYC, TCF-4 and β -actin in immune serum-treated PC3RANKL+ cells, as measured by western blotting. β -actin was used as a loading control. Similar results were obtained in three independent experiments. (D) TOP/FOP luciferase reporter assays in immune serum-treated PC3RANKL+ cells. Significant differences were observed at * $p < 0.05$ vs. the control. (E) Co-immunoprecipitation of β -catenin and TCF-4 from PC3RANKL+ and immune serum-treated PC3RANKL+ cells. Each blot was obtained under the same experimental conditions. (F) Western blot analysis of MAPK phosphorylation levels in serum-treated PC3RANKL+ cells. Similar results were obtained in three independent experiments.

Fig.7




Figure 7

Schematic diagram illustrating anti-RANKL-mediated inhibition of RANK signaling. Activation of RANK by RANKL activates AKT and ERK, thereby liberating the active form of β -catenin. Generation of anti-RANKL

antibodies by mRANKL-MT inactivates RANKL, thereby downregulating AKT/ERK activity and preventing downstream of β -catenin.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryLegend.docx](#)
- [supple4.jpg](#)
- [supple3.jpg](#)
- [supple2.jpg](#)
- [supple1.jpg](#)