

Identification of the Prognostic LncRNA Biomarkers and Comprehensive Analysis of LncRNA-Mediated ceRNA Network for Uterine Corpus Endometrial Carcinoma

Yiran Cai

Medical School of Nantong University

Jin Cui

Medical School of Nantong University

Huiqun Wu (✉ wuhuiqun@ntu.edu.cn)

Medical School of Nantong University

Research Article

Keywords: Uterine corpus endometrial cancer, prognostic biomarker, long non-coding RNA, competing endogenous RNA network, The Cancer Genome Atlas(TCGA), Clinical Proteomic Tumor Analysis Consortium(CPTAC)

Posted Date: November 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-989585/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Given that long non-coding RNAs (lncRNAs) involved in the tumor initiation or progression of the endometrium and that competing endogenous RNA (ceRNA) plays an important role in increasingly more biological processes, lncRNA-mediated ceRNA is likely to function in the pathogenesis of uterine corpus endometrial carcinoma (UCEC). Our present study aimed to explore the potential molecular mechanisms for the prognosis of UCEC through an lncRNA-mediated ceRNA network.

Methods

The transcriptome profiles and corresponding clinical profiles of UCEC dataset were retrieved from CPTAC and TCGA databases respectively. Differentially expressed genes (DEGs) in UCEC samples were identified via “Edge R” package. Then, an integrated bioinformatics analysis including functional enrichment analysis, tumor infiltrating immune cell (TIIC) analysis, Kaplan-Meier curve, Cox regression analysis were conducted to analyze the prognostic biomarkers.

Results

In the CPTAC dataset of UCEC, a ceRNA network comprised of 36 miRNAs, 123 lncRNAs and 124 targeted mRNAs was established, and 8 of 123 prognostic-related DElncRNAs (Differentially Expressed long noncoding RNA) were identified. While in the TCGA dataset, a ceRNA network comprised of 38 miRNAs, 83 lncRNAs and 110 targeted mRNAs was established, and 2 of 83 prognostic-related DElncRNAs were identified. After filtered by risk grouping and Cox regression analysis, 10 prognostic-related lncRNAs including LINC00443, LINC00483, C2orf48, TRBV11-2, MEG-8 were identified. In addition, 33 survival-related DE mRNAs (Differentially Expressed messenger RNA) in two ceRNA networks were further validated in the HPA database. Finally, six lncRNA/miRNA/mRNA axes were established to elucidate prognostic regulatory roles in UCEC.

Conclusion

Several prognostic lncRNAs are identified and prognostic model of lncRNA-mediated ceRNA network is constructed, which promotes the understanding of UCEC development mechanisms and potential therapeutic targets.

Introduction

Uterine corpus endometrial cancer (UCEC), one of globally common gynecological malignancies, presents a possibly upward trend of with the increase of obese women. [1–3] The choice for treatment of UCEC has

considerable exploration and development prospects for the perspective of molecular biology. In previous studies, risk factors including p53 expression[4], and estrogen receptor (ER) and progesterone receptor (PR) expression[5], as well as clinical treatment manners have been identified[6]. The discovery of these factors provides access to take advantage of underlying therapeutic biomarkers for personalized treatment strategies. miRNA is a family of small non-coding RNA molecules of about 21 to 25 nucleotides long. miRNA inhibits translation of targeted mRNA or affects its stability by specific identifications, and down regulates its expression by combining at its 3'UTR site[7]. The abnormal miRNA expression in development of tumors has been confirmed by many studies [8]. LncRNA is known to play a role as key signal transduction mediators in the occurrence, progression and treatment of numerous malignancies[9–11]. According to the ceRNA hypothesis[12], lncRNA is a molecular sponge of miRNA, which suppresses the activity of miRNA by binding with microRNA response element (MRE) and down regulates the expression of the target genes indirectly. Based on this argument, the ceRNA network has been extensively researched and verified in lung cancer[13], breast cancer[14] and so on, while there have been little discussion on lncRNA-mediated ceRNA networks of UCEC.

Therefore, in this study, we retrieved and analyzed lncRNA expression in UCEC from TCGA and CPTAC database separately and performed an integrated bioinformatics analysis including functional enrichment analysis, Tumor Infiltrating Immune Cell (TIIC) analysis, and constructed the UCEC-specific ceRNA network and figured out the underlying association between those ceRNAs and the progression of UCEC.

Methods

Data Collection

Transcriptional and clinical data of UCEC in both The Cancer Genome Atlas data portal(TCGA; <https://portal.gdc.cancer.gov/>) and Clinical Proteomic Tumor Analysis Consortium(CPTAC; <https://cptac-data-portal.georgetown.edu/data-use-agreement>) were retrieved. There were 587 samples downloaded from TCGA, containing 555 UCEC patients and 32 normal specimens, and 116 samples downloaded from CPTAC, containing 101 UCEC patients and 15 normal specimens, contributing to the UCEC and normal control group as a cohort. The clinical features of UCEC patients from 2 databases were respectively shown (Table 1). 12 other samples of originally 555 UCEC samples in TCGA database were omitted in this whole study due to missing associated information. Therefore, actually a total of 575 samples from TCGA were conducted to the further analyses. The clinical features of UCEC patients including age, gender, race, pathology stage, histological type and vital status were extracted. Transcriptome data were annotated with the Genecode website (<https://www.gencodegenes.org/>). No samples were excluded when to screen for differentially expressed RNAs (DERNAs, including three ones: differentially expressed long noncoding RNA(DELncRNA), differentially expressed microRNA (DEmiRNA), differentially expressed messenger RNA(DEmRNA)). Both databases were publically available and were released in compliance with ethical approvals; therefore no further application from University Ethics Committee was obtained.

Table 1
 Characteristics of 543 UCEC patients from TCGA and 101 UCEC patients from
 CPTAC database

TCGA features	Variables	Patients (n, %)
Age	>60	334(61.51)
	≤60	206(37.94)
	NA	3(0.55)
Gender	Female	543(100.00)
Race	White	372(68.51)
	NA	32(5.89)
	Black or African American	106(19.52)
	Native Hawaiian or other pacific islander	9(1.66)
	American indian or Alaska native	3(0.55)
	Asian	20(3.68)
	Pathology stage	Stage I
	Stage II	51(9.39)
	Stage III	124(22.84)
	Stage IV	29(5.34)
Primary diagnosis	Endometrioid carcinoma	399(73.48)
	Serous cystadenocarcinoma	133(24.49)
	Other types	11(2.03)
Survival	Alive	452(83.24)
	Dead	91(16.76)
CPTAC features	Variables	Patients (n, %)
Age	>60	64(63.37)
	≤60	37(36.63)
Gender	Female	101(100.00)
Race	White	59(58.42)
	NA	38(37.62)
	Black or African American	3(2.97)

TCGA features	Variables	Patients (n, %)
	Asian	1(0.99)
Pathology stage	Stage I	75(74.26)
	Stage II	8(7.92)
	Stage III	15(14.85)
	Stage IV	3(2.97)
Histological type	Endometrioid carcinoma	85(84.16)
	Other types	16(15.84)
Tumor grade	G1 + G2	71(70.30)
	G3	27(26.73)
	Other	3(2.97)
Survival	Alive	84(83.17)
	Dead	10(9.90)
	Not reported	7(6.93)

Identification of DEGs

The "edge R" was utilized to identify differentially expressed genes (DEGs) between the normal samples and UCEC patients by the criteria (false discovery rate (FDR) or adjusted $P < 0.01$, and $|\log_2FC| > 2$ (FC is fold change)). All of the statistical tests were conducted and the heatmap and volcano plot were displayed by ggplot2 package in R software package (version:4.0.3; <https://www.r-project.org/>) and statistical significance was defined as a P-value < 0.05 unless otherwise stated.

Construction of Protein-Protein Interaction Network

In this study, a total of 1064 and 917 DEGs filtered by $|\log_2FC| > 3$ from the two databases were subjected to perform protein-protein interaction (PPI) network analysis using Search Tool for the Retrieval of Interacting Genes (STRING; <https://string-db.org/>) [15]. An interaction with a combined score by default > 0.4 was considered statistically significant. Cytoscape (version 3.8.2) is a bioinformatics software platform publically used for visualizing molecular interaction networks [16]. To find hub genes actively participated in UCEC progression, we employed the maximal clique centrality (MCC) algorithm to represent 20 key mRNAs with important biological functions via CytoHubba in Cytoscape [17].

Tumor Infiltrating Immune Cells Profiling

To characterize proportions of tumor infiltrating immune cells (TIICs) in UCEC, CIBERSORT (<http://cibersort.stanford.edu/>) algorithm in combination with a LM22 gene signature matrix was used to assess the relative fractions of 22 invasive immune cell subtypes in each UCEC sample. Regarding the results of the algorithm, we also accumulated the percentage of each immune cell theoretically calculated from each patient sample and presented those top ranked immune cell types in bar graphs. Worthy to point out, if these UCEC samples lack of associated transcriptome information, they were then omitted. Additionally, to investigate the immune infiltration landscape of UCEC, gene set enrichment analysis (ssGSEA) was performed by GSVA package in R (<https://bioconductor.org/packages/release/bioc/html/GSVA.html>) to calculate the score of immune infiltration in each sample on the basis of immune cell-specific gene expression levels. Standardized profiles of gene expression data in both groups were extracted and immune scores were evaluated, scoring types including different cell clusters and respective expression values corresponding to different colors.

Construction of ceRNA Network and Extraction of Survival-related LncRNA-miRNA-mRNA subnetwork

In this study, the lncRNA-mediated ceRNA (competing endogenous RNA) network of UCEC was constructed as lncRNA-miRNA and miRNA-mRNA pairs. To build a ceRNA network, following steps were conducted. Firstly, based on miRNAs provided by the miRcode website (<http://www.mircode.org/>), lncRNA-targeted miRNAs simultaneously in our DE miRNAs were filtered. Secondly, to identify the miRNA-targeted mRNAs, we searched miRTarBase[18] (https://mirtarbase.cuhk.edu.cn/~miRTarBase/miRTarBase_2019/php/index.php), miRDB[19] (<http://www.mirdb.org/>), TargetScan[20] (<http://www.targetscan.org/>) in combination to obtain targeted mRNAs. Finally, Cytoscape software v3.8.2 (<https://cytoscape.org/>) was adopted to visualize the UCEC-related ceRNA network. The paired interactions in above network were analyzed in two UCEC datasets from TCGA and CPTAC, respectively. Subsequently, we took the intersection of the two ceRNA networks.

Furthermore, on the foundation of overall ceRNA network, we firstly conducted survival analyses and Cox regression analyses of these genes, both results of which were considered as hub genes in our ceRNA network. Thereafter, we input these hub genes to visualize their lncRNA-miRNA-mRNA regulatory relationships via Cytohubba (<http://hub.iis.sinica.edu.tw/cytohubba>) in Cytoscape. Eventually, we constructed a novel ceRNA subnetwork composed of lncRNA-miRNA-mRNA pairs, which may provide prognostic molecular values.

Survival Analysis and LncRNA-mediated Prognostic Model Construction

On one hand, survival R was operated for survival analysis of all DE RNAs in the ceRNA network. Kaplan-Meier (K-M) curves were plotted with DE RNAs by log-rank test. On the other hand, we identified the lncRNAs linked with total survival ($p < 0.05$) to act as prognostic lncRNA signature candidates and then imported them for Cox regression analyses. According to the median risk score, the UCEC patients were divided into the high risk and low risk groups. To evaluate the accuracy of models survival-related DE lncRNAs, we carried out the receiver operating characteristic (ROC) curve analysis (UCEC data in CPTAC using 3 years as the predicted time, UCEC data in TCGA using 5 years as the predicted time),

along with the area under the receiver operating characteristic curve (AUC) analysis at a criterion of $AUC > 0.7$. Besides, we further retrieved analyses of survival-related lncRNAs in the GEPIA database (GEPIA; <http://gepia.cancer-pku.cn/index.html>) by p values ≤ 0.1 (although 0.11 was also considered as significant in this study). Subsequently, we compared these results of survival-related lncRNAs with our analyses thus to validate and prove its reliability.

Validation of Dysregulated mRNAs via HPA and GEPIA databases

The Human Protein Atlas Portal (HPA) (www.proteinatlas.org) [21] which contains different genes in specific cancer types and publicly available information of Immunohistochemistry (IHC) staining was used for survival analyses. Gene Expression Profiling Interactive Analysis, an interactive website, composed of 9736 patients and 8587 normal samples from TCGA and GTEx projects (The Genotype-Tissue Expression project) were utilized for the analysis of RNA sequencing expression. According to prognostic lncRNA-mRNA or mRNA-miRNA signatures in the lncRNA-mediated network, we took 33 mRNAs resulted from UCEC data of TCGA into two external databases mentioned above for further retrieval and analysis.

Functional and Pathway Enrichment Analysis

To analyze functions represented in two profiles of identified DEmRNAs in the ceRNA network, Gene Ontology (GO), Kyoto Encyclopedia of genes and Genomes (KEGG) were performed by cluster Profiler R package and plotted by GO plot R package and KEGG plot R package. Three methods were utilized to enrich meaningful biological pathways by the standard of p value less than 0.05.

Results

Differentially expressed RNAs in UCEC

In a study of 575 UCEC samples collected from TCGA (http://cancergenome.nih.gov/publications/publication_guide_lines), "edge R" (Adjust $P < 0.01$ and $|\log_2 FC| > 2$) was adjusted to identify the DERNAs, including DElncRNAs, DEmiRNAs, DEmRNAs. A total of 1513 up-regulated DEmRNAs, 914 down-regulated DEmRNAs, 686 up-regulated DElncRNAs, 330 down-regulated DElncRNAs, 124 up-regulated DEmiRNAs, and 50 down-regulated DEmiRNAs were identified in TCGA database. Similarly, a total of 116 samples were identified with the same criteria in the CPTAC database, including 2463 differential DEmRNAs, 1741 DElncRNAs and 204 DEmiRNAs. The heatmap and volcano plot of those highest dysregulated DElncRNAs, DEmiRNAs and DEmRNAs were shown in Fig.1 and the top 10 up and down regulated DElncRNAs, DEmiRNAs and DEmRNAs were shown in supplement Tables 1.

Construction of PPI Network

The 1981 DEmRNAs ($|\log_2 FC| > 3$) were further selected to construct PPI network to select hub genes that play crucial roles in UCEC genesis. Given the large number of DEmRNAs in this module, we used

MCC algorithm in “Cytohubba” of Cytoscape software to visualize and select hub genes in the PPI network. The top 20 high score genes in in CPTAC were shown (Fig.2A). Similarly, the top 20 high score genes in TCGA belong to Histone cluster 1 H family which was shown (Fig.2B).

Two graphs separately describe the top 20 most dynamic hub genes and their intersection relationships evaluated by MCC algorithm in “Cytohubba”. These sub-graphs of these selected mRNA-coding protein nodes are shown from highly essential (red) to essential (yellow).

TILCs Enrichment Analysis

Using CIBERSORT algorithm, we evaluated 101 tumor transcriptome profiles from CPTAC database and 543 tumor transcriptome profiles from TCGA database (Fig.3A,3B). In addition, we also performed ssGSEA analysis by GSVA package to score the corresponding TILCs in each simple sample, and finally we found that TILCs in both CPTAC and TCGA database sources expressed well (Fig.3C,3D). On account of the top five immune cell components in UCEC patients by CIBERSORT algorithm, 2 bar graphs were then visualized. In CPTAC, there were abundant CD8 T cells (28.5%) and plasma cells (20.5%) and other TILCs. In TCGA samples, naive CD4 cells (18.9%) and CD8 cells (15.2%) were well infiltrated, accompanied by slightly increased activated NK cells (10.2%), plasma cells (9.1%) and macrophages M0(7.5%) (Fig.3E,3F).

Construction of ceRNA Network and hub LncRNA-miRNA-mRNA subnetwork

In order to better comprehend the interactions of mRNAs, lncRNAs, and miRNAs in UCEC, we constructed an lncRNA-mediated ceRNA regulatory network. To begin with, 1741 DElncRNAs in CPTAC database succeeded to match with 123 lncRNAs in the miRCODE database. Considering 123 of 1741 DElncRNAs could interact with DEmiRNAs, 36 miRNAs in both miRCODE and CPTAC database were selected to construct lncRNA-miRNA pairs. Meanwhile, to interplay with 204 DEmiRNAs acquired from CPTAC database, we retrieved 1420 mRNAs in three databases (miRTarBase, miRDB and TargetScan). The 1420 miRNA-targeted mRNAs predicted in these databases were intersected with 2463 DEmRNAs thus to obtain 124 miRNA-targeted mRNAs belonging to CPTAC database. Finally, an lncRNA-mediated ceRNA network consisting of 36 miRNAs, 123 lncRNAs and 124 mRNAs were achieved (Fig.4A). Meanwhile, the same workflow of UCEC-specific ceRNA network construction was repeated in data from TCGA. We obtained an lncRNA-mediated ceRNA network consisting of 38 miRNAs, 83 lncRNAs and 110 mRNAs (Fig.4B).

Furthermore, cytohubba was applied to visualize our extracted hub genes composing of lncRNA, miRNA and mRNAs and derived regulatory ceRNA network thus to identify potentially prognostic molecular pathways of UCEC in CPTAC and TCGA (Fig.4C,4D). A total of 6 hub lncRNA-miRNA-mRNA regulatory relationships from 2 databases were shown (Table 2). Moreover, coincident ceRNA results in the overall ceRNA network from both CPTAC and TCGA databases were shown in the Venn diagram (Fig.4E).

Table 2 CeRNA subnetwork of prognostic regulatory DEGs in UCEC from CPTAC (up) and TCGA (down) database

LncRNA	miRNA	mRNA
TRBV11-2	has-mir-363	SOX11
MEG8	has-mir-424	CCNE1, CBX2
	has-mir-363	SOX11
	has-mir-183	DLX4, NR3C1
lncRNA	miRNA	mRNA
LINC00443, C2orf48, LINC00483	miR-183	DLX4, NR3C1
DGCR5	has-mir-195	CCNE1
	has-mir-383	
	has-mir-424	

Identification and Validation of Prognostic lncRNAs in ceRNA networks

In order to figure out the effects of interactions for survival between lncRNAs, miRNAs and mRNAs, we imported survival-related data of UCEC and genes in ceRNA to analyze its prognosis. Survival R were operated for DERNA significantly correlated with overall survival in the ceRNA network ($p < 0.05$), the results of which were plotted by Kaplan–Meier (K-M) curves (Fig.5A-5W).

As shown in CPTAC, 4 survival-related DElncRNAs on the level of 3-year survival were identified in DElncRNA-mediated ceRNA networks, including FREM2-AS1, HPYR1, LINC00028, MIR205HG (Fig.5A-5D). Similarly, 19 survival-related DElncRNAs, 9 DEmiRNAs and 33 DEmRNAs on the level of 5-year survival were revealed in the ceRNA network for TCGA. Fig.5E-5W illustrated Kaplan–Meier curve analyses about 10 lncRNAs of 19 survival-related DElncRNAs, 5 mRNAs of 33 DEmRNAs and 2 of 9 DEmiRNAs derived from TCGA database. Besides, we successfully validated 5 survival-related lncRNAs in GEPIA database by p values ≤ 0.1 (Fig.6A-6E).

To further identify DElncRNAs with prognostic features in a more accurate way, multi-Cox regression analyses and corresponded ROC curves were carried out. After eliminating some samples lacking in survival time, 94 complete samples in CPTAC were divided into the high-risk ($n=47$) and low-risk ($n=47$) groups (cutoff value = -0.78) and 543 samples with complete survival information in TCGA into the high-risk ($n=272$) and low-risk ($n=272$) cohort by median value (cutoff value = -0.18; one sample of survival data was just in the median and counted in both groups). We performed a multi-factor COX regression analysis and a global survival analysis of the model thus separately identified two lncRNAs of 3-year

survival data in CPTAC and eight lncRNA prognosis candidates of 5-year survival UCEC data in TCGA by $p < 0.05$ (Fig.7A,7B,7C,7F)). Receiver operating characteristic (ROC) curves tested the influence on their lncRNA signatures associated with overall survival in UCEC. Area under ROC curve of 3-year survival rate (AUC) and 5-year survival rate (AUC) were respectively 0.967 and 0.751. (Fig.7B,7E) Besides, multivariate cox regression analysis of totally 10 prognostic lncRNAs associated with overall survival in UCEC patients generated from 2 databases were shown in Table 3.

Table 3 Multivariate Cox regression analysis of totally 10 prognostic lncRNAs associated with overall survival (OS)

LncRNA(in TCGA)	Coef	HR	95%CI_LL	95%CI_HL	p-value
FAM41C	0.017169325	1.017317565	1.007489748	1.027241249	5.27E-04
MIR7_3HG	0.003483874	1.00348995	1.001209451	1.005775643	0.002688886
LINC00483	0.010635045	1.010691798	1.003527131	1.017907617	0.003389722
ABHD11_AS1	8.26E-04	1.000826495	1.00038596	1.001267224	2.35E-04
LINC00443	0.007147984	1.007173592	1.001433816	1.012946267	0.014233205
OXCT1_AS1	0.026906491	1.027271739	1.006299256	1.048681314	0.010568912
PRICKLE2_AS2	0.273785608	1.314932861	1.133483915	1.525428288	3.02E-04
GLIS3_AS1	9.53E-04	1.000953853	1.000210868	1.001697389	0.011853011
LncRNA(in CPTAC)	Coef	HR	95%CI_LL	95%CI_HL	p-value
MEG8	0.007280976	1.007307546	1.003839185	1.010787892	3.51E-05
TRBV11_2	0.027816351	1.028206838	1.012679086	1.043972683	3.40E-04

In the multivariate Cox regression analysis derived from TCGA database, 8 lncRNAs including FAM41C, MIR7_3HG, LINC00483, ABHD11_AS1, LINC00443, OXCT1_AS1, PRICKLE2_AS2 and GLIS3_AS1 were identified to construct the OS prediction model. OS-related prediction model=(0.017169325* expression value of FAM41C)+(0.003483874* expression value of MIR7_3HG)+(0.010635045* expression value of LINC00483)+(8.26E-04* expression value of ABHD11_AS1)+(0.007147984* expression value of TRBV11_2)+(0.027816351* expression value of TRBV11_2)+(0.027816351* expression value of LINC00443)+(0.026906491* expression value of OXCT1_AS1)+(0.273785608* expression value of PRICKLE2_AS2)+(9.53E-04* expression value of GLIS3_AS1). We divided the 543 UCEC cases into the high and low-risk groups according to the median values of the OS-related prediction model.

In the multivariate Cox regression analysis derived from CPTAC database, 2 lncRNAs including MEG8 and TRBV11_2 were identified to construct the OS prediction model. OS-related prediction model=(0.007280976* expression value of MEG8)+(0.027816351* expression value of TRBV11_2). We divided the 94 UCEC cases into the high and low-risk groups according to the median values of the OS-related prediction model.

Validations of survival analysis and mRNA Expression at the Transcriptional Level

To further demonstrate the prognostic significance of 33 mRNAs screened from the ceRNA network, we selected external databases for survival analysis and validation with IHC images. Firstly, we input 33 screened mRNAs into HPA database (version 20.1; https://www.ProteinAtlas.org/about/assays+annotation#tcga_survival) to validate whether they were associated with the prognosis of UCEC. Consequences revealed that 8 mRNAs (CBX2, CCL22, CCNE1, DLX4, IGFBP5, NR3C1, SOX11, POLQ) highly expressed in UCEC were closely related with its prognosis (log rank P values ≤ 0.001). Subsequently, we retrieved overall survival analyses of 8 mRNAs generated from GEPIA by filtered criteria of P values ≤ 0.1 (although 0.11 was also considered as significant in this study) and verified 5 mRNAs (CCNE1, CCL22, NR3C1, IGFBP5 and POLQ).

Based on two previous steps for external verifications, two IHC images of the last screened mRNAs (CCNE1, NR3C1) in the HPA database approved the same results (Fig.8A). Survival validations of 5 mRNAs including CCNE1, CCL22, NR3C1, IGFBP5 and POLQ from GEPIA were shown in the Fig.8B-8F. In this study, we identified 5 survival-related mRNAs, there were no related IHC samples of CCL22, IGFBP5 and POLQ but CCNE1 and NR3C1 to further validate in the HPA database. The translational expression level of CCNE1 and NR3C1 was positively linked with disease status, as they were up-regulated in UCEC samples.

Enrichment Analyses of Functional Pathways

To elucidate the biological functions represented in two profiles of identified DEmRNAs, we performed enrichment analyses mainly by "cluster profiler", with the standard of $p < 0.05$. In this study, GO analyses disclosed that top significant GO terms ($p < 0.05$) commonly obtained from UCEC data in CPTAC (Fig.9A,9B) and TCGA database (Fig.9E,9F). The KEGG analyses revealed that what closely related to DEmRNAs originated from CPTAC were mainly enriched in pathways such as "cellular senescence", "proteoglycans in cancer" and "microRNAs in cancer" (Fig.9C,9D)). The KEGG results derived from DEmRNAs in TCGA database were shown (Fig.9G,9H). The top 20 GO and KEGG results for TCGA and CPTAC database were provided in supplement Table 2.

Discussion

Recently, with the increase of obese women, UCEC has become one of the leading gynecologic tumors[22]. Although some diagnostic markers like CA125, CA199, and CEA are clinically used, survival results are not optimistic after routine diagnosis and therapy. Therefore, it is worthy to discover and

analyze biomarkers for prognosis prediction of UCEC. LncRNAs have increasingly seized the attention of cancer research fields because of serving as regulating biomarkers[23]. But due to experimental complexity, functional studies related to lncRNAs have limitations to carry out in comparison with those of protein-encoding RNAs. As is illustrated in accumulating researches, molecular mechanisms underlying ceRNA network provide an explanation for carcinogenesis and its associated development. LncRNAs act as key components of ceRNA family, through miRNA response elements (MREs), compete with molecules binding to the same miRNAs to achieve regulation of expression levels between each other.

As our lncRNA-mediated ceRNA network of UCEC respectively constructed from the CPTAC and TCGA databases indicated, there were a total of 23 lncRNAs, 9 miRNAs, and 33 mRNAs correlated with the overall survival results and served as promising biomarkers for predicting prognosis of UCEC. Conventional prognostic model constructions often make inadequate risk groupings and estimates of clinical outcomes[24, 25]. Nonetheless, this ceRNA hypothesis provides us a novel predictive insight from the angle of heterogeneity between UCEC patients to analyze overall survival (OS) results. For bioinformatics analysis conducted in multiple databases, it is common practice to combine sample profiles from multiple databases for further analysis after standardized quality control treatment. However, the follow-up time length of UCEC profiles from CPTAC is shorter than that from TCGA database. So we calculated on the level of three-year survival results thus didn't combine it with that from TCGA database.

In our present study, MIR205HG and ADARB2-AS1 were significantly correlated with survival. High expression of MIR205HG was found at the first time to predict a good prognosis, while high expression of ADARB2-AS1 had an opposite effect on the survival outcome of patients. Dong et al. showed that lncRNA MIR205HG depleted SRSF1 to increase KRT17 expression[26], while KRT17 silencing impaired cervical cancer cell proliferation and migration and activated apoptosis. lncRNA MIR205HG also acts as a ceRNA to accelerate tumor growth and progression in cervical cancer through spongyform Mir-122-5p[27]. ADARB2-AS1 has been reported as a prognostic related lncRNA in UCEC[28], which again echoed the reliability of our results.

In addition, in order to make our results convincing, we put lncRNA biomarkers into GEPIA for external verification. The results showed that 5 highly expressed lncRNAs of DGCR5, GLIS3-AS1, UPK1A-AS1, MEG8, TPTEP1 indicated poor prognosis, which is in agreement with our results. To clarify our findings, we comprehensively analyzed survival results of lncRNA-regulated mRNAs in GEPIA and HPA. Through these external databases, we also concluded that high expression of CCL22, CCNE1, IGFBP5, NR3C1 and POLQ genes were associated with poor prognosis of UCEC. Furthermore, we mirrored methods in foregoing study[29] to identify our mRNA results and obtained poor survival results of NR3C1 and CCNE1 by validations of IHC images, which suggested their tumor promoter roles. Some researches has verified that CCNE1 amplification is associated with aggressive potential in UCEC tumorigenesis[30–32]. CCNE1, known as Cyclin E1, is a member of Cyclins to function as regulators of CDK kinases. The protein encoded by Cyclin belongs to the highly conserved Cyclin family, characterized in its dramatic periodicity

in protein abundance through cell cycle. With respect to other carcinoma progression, patients with over-expressed CCNE1 were reportedly at increased threat for poor endings of cervical cancer[33] and triple-negative breast cancer[34]. Functioned as regulatory genes in the downstream, mRNA CCNE1 and NR3C1 brought potential reference values for our presently identified lncRNA-mediated ceRNA pathways.

Furthermore, on the basis of overall ceRNA network, we constructed a novel prognostic ceRNA subnetwork, which were composed of lncRNA-miRNA-mRNA axes. We firstly conducted survival analysis and multivariate analysis on lncRNAs in the ceRNA network derived from TCGA database, and identified them as hub genes in the following subnetwork. We firstly identified these key lncRNAs in the ceRNA network of TCGA, then paired their corresponding key miRNAs. Then we matched the key miRNAs with survival-related mRNAs thus to construct the survival-related subnetwork. For survival analysis in CPTAC database, although we failed to identify survival-related mRNAs and miRNAs, we surprisingly discovered that survival-related mRNAs and miRNAs in TCGA database also existed in the ceRNA network constructed by CPTAC database. Therefore, we chose to use these mRNAs, miRNAs and key lncRNAs derived from CPTAC to jointly construct survival-related subnetworks of CPTAC. In our constructed ceRNA subnetwork, there were 6 lncRNA-mediated lncRNA-miRNA-mRNA axes to role in survival outcomes of UCEC patients. A novel survival-related lncRNA DGCR5 could up-regulate CCNE1 expression by binding to miR195, miR383 and miR424, although DGCR5 had not been directly recorded in UCEC tumorigenesis procedures. DiGeorge syndrome critical region gene 5 (DGCR5), a molecular sponge to regulate cancerous signaling pathways, has been previously discovered to be extremely dysregulated in various tumors and induce the malignant phenotypes of organs such as liver, pancreas and lungs.etc. Except for DGCR5, lncRNA LINC00443, LINC00483, C2orf48, TRBV11-2 and MEG8 were identified by multivariate Cox regression analysis, which reveals more accurate ability to predict prognosis.

From multivariate Cox regression analysis in our constructed ceRNA network, we totally identified 10 lncRNA prognostic signature candidates to predict the survival events of UCEC patients. As shown in our ceRNA subnetwork diagram for TCGA, lncRNAs such as LINC00443, C2orf48, LINC00483 could regulate DLX4 and NR3C1 expression by binding to miR-183. LINC00443, LINC00483 and C2orf48 were previously proved to promote carcinoma progression[35], the same consequence of which were validated by our experiment. DLX4 has been shown to cause tumor migration, invasion, and metastasis[36]. Previous *in vivo* studies on UCEC reported that DLX4 promoted cell proliferation, migration, and suggested poor prognosis, which is consistent with our findings[37]. NR3C1 encodes glucocorticoid receptors to affect glucocorticoid response and participates in other transcription regulatory procedures. Former study on miRNA-mRNA regulatory network in UCEC found that over expression of mRNA NR3C1 led to poor prognosis[38, 39]. Previously *in vivo* experiments proved that over-expressed LINC00483 promoted UCEC tumorigenesis, the mechanism of which was mainly to sponge with miR-508-3p to regulate RGS17 expression levels[40]. In other cancer researches, LINC00483 also acted as a strong ceRNA molecule and exhibited its regulatory ability to mediate tumor progression and prognosis, such as lung adenocarcinoma[41] and gastric cancer[42]. Our clinical survival and transcriptome results revealed that patients with over-expression of LINC00443, C2orf48 and LINC00483 had poor prognostic outcomes. The two lncRNAs filtered in multi-Cox regression analysis in CPTAC, TRBV11-2 and MEG8 provided potential

pathways to explain ceRNA regulatory network despite of none associated reports for UCEC. Firstly, they could up-regulate SOX11 expression by binding to hsa-mir-363 to bring about poor prognosis, and survival related mRNA SOX11 hypermethylation was reported as a tumor biomarker in UCEC[43]. Secondly, MEG8 competed with has-mir-424 to CBX2 and CCNE1 expression as well as competing with has-mir-183 to up-regulate DLX4 and NR3C1 expression. Therefore, we predicted that TRBV11-2 and MEG8 worked as prognostic ceRNAs to up-regulate SOX11, CCNE1 and NR3C1 expression thus resulting in poor prognosis, suggesting that these lncRNAs could promote UCEC development. Furthermore, LINC00028 existed in our survival analysis results rooted from both databases, and the over-expression of LINC00028 indicated a poor prognosis. Besides, LINC00028 has been reported to be involved in ceRNA regulatory network of osteosarcoma recurrence[44]. However, its mechanisms related to UCEC pathogenesis remains unclear.

As illustrated in former studies[45, 46], UCEC carcinogenesis is promoted by cell cycle acceleration. Similarly, our KEGG analyses in both databases indicated that DE mRNAs were mostly enriched in pathways such as microRNAs in cancer, cellular senescence and proteoglycans in cancer. Deoxyribonucle, and deoxyRNA were identified in tumor and adjacent normal tissue samples in a large cohort of UCEC patients.

Even though we identified the molecular mechanism of ceRNA from 587 samples in TCGA and 116 samples in CPTAC database, there was still a limited sample size for more reliable biomarkers that hindered us from incorporating data profiles originated from CPTAC and TCGA database into a comprehensive study ideally. Concerning the pilot study limited by failure to closely link the analysis of the two databases, multicentric studies are supposed to carry out to support our new researches inevitably.

Conclusion

Using data obtained from CPTAC and TCGA, we screened out lncRNA prognostic signatures on the basis of ceRNA completely composed of hub genes. Besides, an lncRNA-mediated ceRNA network reveals the molecular mechanism that facilitates UCEC pathological progress. lncRNAs including DGCR5, LINC00443, C2orf48, LINC00483, TRBV11-2 and MEG8 involved in lncRNA-miRNA-mRNA regulatory network were identified as promising diagnostic, therapeutic or prognostic biomarkers. Further studies are warranted to explore meaningful biological functional pathways underlying these lncRNA roles for UCEC.

Abbreviations

ceRNA:competing endogenous RNA; DEGs: Differentially expressed genes;lncRNA:long noncoding RNA;mRNA:messenger RNA; miRNA :microRNA;UCEC:uterine corpus endometrial carcinoma;TIIc:tumor infiltrating immune cell;TCGA:The Cancer Genome Atlas; CPTAC:Clinical Proteomic Tumor Analysis Consortium;STRING:Search Tool for the Retrieval of Interacting Genes; HPA:Human Protein Atlas Portal;ssGSEA:single sample gene set enrichment analysis;IHC:Immunohistochemistry; GEPIA: Gene

Expression Profiling Interactive Analysis;GO: Gene ontology; KEGG: Kyoto encyclopedia of genes and genomes; PPI: Protein–protein interact

Declarations

Ethics approval and consent to participate

Not applicable.

TCGA and CPTAC belong to public databases. The patients involved in the database have obtained ethical approval. Users can download relevant data for free for research and publish relevant articles. Our study is actually an integrated bioinformatics analysis based on open source data, so there are no ethical issues and other conflicts of interest. It is specially noted that the Immunohistochemistry images mentioned in the article are retrieved and downloaded from the CPTAC website requiring no additional ethical approval.

Consent for publication

Not applicable.

Availability of data and materials

In our study, different web-based datasets were used for our further data analysis. The web links to all the original data sources were listed as below: The datasets including the RNA-seq transcriptome data and clinical data of UCEC cohort were respectively obtained from The Cancer Genome Atlas Program (TCGA) (<https://portal.gdc.cancer.gov/>) and National Cancer Institute Clinical Proteomic Tumor Analysis Consortium (<https://cptac-data-portal.georgetown.edu/>) data portal. All data generated from the analysis process of this study are available from authors listed above on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by the grant from Excellent Key Teachers in the “Qing Lan Project” of Jiangsu Colleges and Universities and “226 Project” of Nantong[(2018)III-436]. The funding body listed above played no role in the design of the study, collection, analysis, or interpretation of data, nor was it involved in the preparation of this manuscript.

Author s' contributions

YRC prepared the dataset and analyzed the data. JC wrote and revised the manuscript. HQW reviewed, modified and provided experimental conditional support. All authors have read, revised, and approved the

final manuscript.

Acknowledgements

Not applicable.

Authors' information

¹Medical School of Nantong University, Nantong, 226001, China.

²Department of Medical Informatics, Medical School of Nantong University, Nantong, 226001, China.

References

1. Calle EE, Kaaks R: Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 2004, 4(8):579–591.
2. Janda M, McGrath S, Obermair A: Challenges and controversies in the conservative management of uterine and ovarian cancer. *Best Pract Res Clin Obstet Gynaecol* 2019, 55:93–108.
3. Lu KH, Broaddus RR: Endometrial Cancer. *N Engl J Med* 2020, 383(21):2053–2064.
4. Engelsen IB, Stefansson I, Akslen LA, Salvesen HB: Pathologic expression of p53 or p16 in preoperative curettage specimens identifies high-risk endometrial carcinomas. *Am J Obstet Gynecol* 2006, 195(4):979–986.
5. Smith D, Stewart CJR, Clarke EM, Lose F, Davies C, Armes J, Obermair A, Brennan D, Webb PM, Nagle CM et al: ER and PR expression and survival after endometrial cancer. *Gynecol Oncol* 2018, 148(2):258–266.
6. Wright JD, Burke WM, Wilde ET, Lewin SN, Charles AS, Kim JH, Goldman N, Neugut AI, Herzog TJ, Hershman DL: Comparative effectiveness of robotic versus laparoscopic hysterectomy for endometrial cancer. *J Clin Oncol* 2012, 30(8):783–791.
7. Bartel DP: MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 2004, 116(2):281–297.
8. Peng Y, Croce CM: The role of MicroRNAs in human cancer. *Signal Transduct Target Ther* 2016, 1:15004.
9. Zhang Z, Gu M, Gu Z, Lou YR: Role of Long Non-Coding RNA Polymorphisms in Cancer Chemotherapeutic Response. *J Pers Med* 2021, 11(6).
10. Schmitt AM, Chang HY: Long Noncoding RNAs in Cancer Pathways. *Cancer Cell* 2016, 29(4):452–463.
11. Ahadi A: Functional roles of lncRNAs in the pathogenesis and progression of cancer. *Genes Dis* 2021, 8(4):424–437.
12. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP: A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell* 2011, 146(3):353–358.

13. Yang J, Qiu Q, Qian X, Yi J, Jiao Y, Yu M, Li X, Li J, Mi C, Zhang J et al: Long noncoding RNA LCAT1 functions as a ceRNA to regulate RAC1 function by sponging miR-4715-5p in lung cancer. *Mol Cancer* 2019, 18(1):171.
14. Sun M, Gomes S, Chen P, Frankenberger CA, Sankarasharma D, Chung CH, Chada KK, Rosner MR: RKIP and HMGA2 regulate breast tumor survival and metastasis through lysyl oxidase and syndecan-2. *Oncogene* 2014, 33(27):3528–3537.
15. Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P, von Mering C et al: STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 2013, 41(Database issue):D808-815.
16. Smoot ME, Ono K, Ruscheinski J, Wang PL, Ideker T: Cytoscape 2.8: new features for data integration and network visualization. *Bioinformatics* 2011, 27(3):431–432.
17. Chin CH, Chen SH, Wu HH, Ho CW, Ko MT, Lin CY: cytoHubba: identifying hub objects and sub-networks from complex interactome. *BMC Syst Biol* 2014, 8 Suppl 4(Suppl 4):S11.
18. Huang HY, Lin YC, Li J, Huang KY, Shrestha S, Hong HC, Tang Y, Chen YG, Jin CN, Yu Y et al: miRTarBase 2020: updates to the experimentally validated microRNA-target interaction database. *Nucleic Acids Res* 2020, 48(D1):D148-d154.
19. Wong N, Wang X: miRDB: an online resource for microRNA target prediction and functional annotations. *Nucleic Acids Res* 2015, 43(Database issue):D146-152.
20. Fromm B, Billipp T, Peck LE, Johansen M, Tarver JE, King BL, Newcomb JM, Sempere LF, Flatmark K, Hovig E et al: A Uniform System for the Annotation of Vertebrate microRNA Genes and the Evolution of the Human microRNAome. *Annu Rev Genet* 2015, 49:213–242.
21. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A et al: Proteomics. Tissue-based map of the human proteome. *Science* 2015, 347(6220):1260419.
22. Siegel RL, Miller KD, Jemal A: Cancer statistics, 2019. *CA Cancer J Clin* 2019, 69(1):7–34.
23. Shi X, Sun M, Liu H, Yao Y, Song Y: Long non-coding RNAs: a new frontier in the study of human diseases. *Cancer Lett* 2013, 339(2):159–166.
24. Huo X, Sun H, Cao D, Yang J, Peng P, Yu M, Shen K: Identification of prognosis markers for endometrial cancer by integrated analysis of DNA methylation and RNA-Seq data. *Sci Rep* 2019, 9(1):9924.
25. Ouyang D, Li R, Li Y, Zhu X: A 7-lncRNA signature predict prognosis of Uterine corpus endometrial carcinoma. *J Cell Biochem* 2019, 120(10):18465–18477.
26. Dong M, Dong Z, Zhu X, Zhang Y, Song L: Long non-coding RNA MIR205HG regulates KRT17 and tumor processes in cervical cancer via interaction with SRSF1. *Exp Mol Pathol* 2019, 111:104322.
27. Li Y, Wang H, Huang H: Long non-coding RNA MIR205HG function as a ceRNA to accelerate tumor growth and progression via sponging miR-122-5p in cervical cancer. *Biochem Biophys Res Commun* 2019, 514(1):78–85.

28. Xia L, Wang Y, Meng Q, Su X, Shen J, Wang J, He H, Wen B, Zhang C, Xu M: Integrated Bioinformatic Analysis of a Competing Endogenous RNA Network Reveals a Prognostic Signature in Endometrial Cancer. *Front Oncol* 2019, 9:448.
29. Zhang X, Feng H, Li Z, Li D, Liu S, Huang H, Li M: Application of weighted gene co-expression network analysis to identify key modules and hub genes in oral squamous cell carcinoma tumorigenesis. *Onco Targets Ther* 2018, 11:6001–6021.
30. Kuhn E, Bahadirli-Talbott A, Shih le M: Frequent CCNE1 amplification in endometrial intraepithelial carcinoma and uterine serous carcinoma. *Mod Pathol* 2014, 27(7):1014–1019.
31. Leskela S, Pérez-Mies B, Rosa-Rosa JM, Cristobal E, Biscuola M, Palacios-Berraquero ML, Ong S, Matias-Guiu Guia X, Palacios J: Molecular Basis of Tumor Heterogeneity in Endometrial Carcinosarcoma. *Cancers (Basel)* 2019, 11(7).
32. Nakayama K, Rahman MT, Rahman M, Nakamura K, Ishikawa M, Katagiri H, Sato E, Ishibashi T, Iida K, Ishikawa N et al: CCNE1 amplification is associated with aggressive potential in endometrioid endometrial carcinomas. *Int J Oncol* 2016, 48(2):506–516.
33. Zhang Y, Li X, Zhang J, Mao L: E6 hijacks KDM5C/Inc_000231/miR-497-5p/CCNE1 axis to promote cervical cancer progression. *J Cell Mol Med* 2020, 24(19):11422–11433.
34. Yang R, Xing L, Zheng X, Sun Y, Wang X, Chen J: The circRNA circAGFG1 acts as a sponge of miR-195-5p to promote triple-negative breast cancer progression through regulating CCNE1 expression. *Mol Cancer* 2019, 18(1):4.
35. Liu J, Nie S, Liang J, Jiang Y, Wan Y, Zhou S, Cheng W: Competing endogenous RNA network of endometrial carcinoma: A comprehensive analysis. *J Cell Biochem* 2019, 120(9):15648–15660.
36. Zhang L, Yang M, Gan L, He T, Xiao X, Stewart MD, Liu X, Yang L, Zhang T, Zhao Y et al: DLX4 upregulates TWIST and enhances tumor migration, invasion and metastasis. *Int J Biol Sci* 2012, 8(8):1178–1187.
37. Zhang L, Wan Y, Jiang Y, Zhang Z, Shu S, Cheng W, Lang J: Overexpression of BP1, an isoform of Homeobox Gene DLX4, promotes cell proliferation, migration and predicts poor prognosis in endometrial cancer. *Gene* 2019, 707:216–223.
38. Ding H, Fan GL, Yi YX, Zhang W, Xiong XX, Mahgoub OK: Prognostic Implications of Immune-Related Genes' (IRGs) Signature Models in Cervical Cancer and Endometrial Cancer. *Front Genet* 2020, 11:725.
39. Sun R, Liu J, Nie S, Li S, Yang J, Jiang Y, Cheng W: Construction of miRNA-mRNA Regulatory Network and Prognostic Signature in Endometrial Cancer. *Onco Targets Ther* 2021, 14:2363–2378.
40. Hu P, Zhou G, Zhang X, Song G, Zhan L, Cao Y: Long non-coding RNA Linc00483 accelerated tumorigenesis of cervical cancer by regulating miR-508-3p/RGS17 axis. *Life Sci* 2019, 234:116789.
41. Yang S, Liu T, Sun Y, Liang X: The long noncoding RNA LINC00483 promotes lung adenocarcinoma progression by sponging miR-204-3p. *Cell Mol Biol Lett* 2019, 24:70.
42. Li D, Yang M, Liao A, Zeng B, Liu D, Yao Y, Hu G, Chen X, Feng Z, Du Y et al: Linc00483 as ceRNA regulates proliferation and apoptosis through activating MAPKs in gastric cancer. *J Cell Mol Med*

2018, 22(8):3875–3886.

43. Shan T, Uyar DS, Wang LS, Mutch DG, Huang TH, Rader JS, Sheng X, Huang YW: SOX11 hypermethylation as a tumor biomarker in endometrial cancer. *Biochimie* 2019, 162:8–14.
44. Zhang S, Ding L, Li X, Fan H: Identification of biomarkers associated with the recurrence of osteosarcoma using ceRNA regulatory network analysis. *Int J Mol Med* 2019, 43(4):1723–1733.
45. Wang Y, Qiu H, Hu W, Li S, Yu J: RPRD1B promotes tumor growth by accelerating the cell cycle in endometrial cancer. *Oncol Rep* 2014, 31(3):1389–1395.
46. Xiong H, Li Q, Chen R, Liu S, Lin Q, Xiong Z, Jiang Q, Guo L: A Multi-Step miRNA-mRNA Regulatory Network Construction Approach Identifies Gene Signatures Associated with Endometrioid Endometrial Carcinoma. *Genes (Basel)* 2016, 7(6).

Figures




Figure 1

DERNAs between UCEC and normal tissues in the CPTAC and TCGA database. Volcano plot of UCEC-specific lncRNAs (A), mRNAs (B) and miRNAs (C) from CTPAC and lncRNAs (D), mRNAs(E), miRNAs(F) from TCGA. Red dot indicated high expression, and blue dot indicated low expression. Grey showed that those genes showed no difference between cancer and non-cancer tissues. The hierarchical clustering heatmaps of DERNAs between the UCEC and normal tissues from the CPTAC(G-I) and TCGA database(J-L)




Figure 3

The immune infiltration landscape in UCEC. A, B: Estimation of 22 immune leucocytes fractions with different colors in UCEC by CIBERSORT algorithm. The vertical axis represented the cell proportions. C, D: The score of 31 immune signatures in each sample by heatmap by ssGSEA. E, F: The top five immune cell components were derived from CIBERSORT algorithm and then visualized in the bar chart.




Figure 4

Overall construction of lncRNA-mediated ceRNA network in UCEC from CPTAC (A) and TCGA database (B). Diamonds, rounded rectangles and ellipses respectively represented lncRNAs, miRNAs and mRNAs. The red indicated high expression while the blue indicated low expression. Potentially prognostic ceRNA network of lncRNA-miRNA-mRNA pairs and derived regulatory genes arranged by Cytohubba from CPTAC (A) and TCGA database (B). E: Venn diagram of genes completely involved in the overall ceRNA network

in CPTAC and TCGA database. The overlapped area in the middle indicated intersected genes of two databases.




Figure 5

Kaplan–Meier curves of DERNAs and overall survival rate in UCEC samples.




Figure 6

Kaplan–Meier curves of survival-related lncRNAs validated in GEPIA database. Graphs of lncRNAs including MEG8 (A), DGCR5 (B), GLIS3-AS1(C), TPTEP1(D), and UPK1A-AS1 (E)




Figure 7

LncRNA signature prognostic risk models for UCEC. For CPTAC dataset, Kaplan–Meier curves showed that low-risk group had a lower mortality rate than high-risk group ($P=0.0046<0.05$) (A). The ROC curves of OS-related predictive signatures ($AUC:0.967$)(B). The high expression level of MEG8 and TRBV11_2 were significantly associated with worse OS in Kaplan–Meier curves (both $P<0.05$) (C). While for TCGA dataset, Kaplan–Meier curves showed that low-risk group had a lower mortality rate than high-risk group ($P<0.0001$)(D). The ROC curves of OS-related predictive signatures ($AUC: 0.751$)(E). The high expression level of the eight lncRNAs was significantly associated with worse OS in Kaplan–Meier curves. (F)




Figure 8

GESA of functional pathways for DEmRNAs in UCEC. The GO chord (A: CPTAC, E: TCGA) for top 30 DEmRNAs and their GO terms. The GO circle (B: CPTAC, F: TCGA) and the outer circle represented the expression ($|\log_2FC|$) of DEmRNAs in each enriched GO terms. To be specific, red dots on respective GO term indicated the up-regulated DEmRNAs. Blue dots indicated the down-regulated DEmRNAs. The inner circle represented the significance of GO terms(log10-adjusted p values). The bar plot (C: CPTAC, G: TCGA) and dot plot (D: CPTAC, H: TCGA) of 10 pathways lncRNA-related DEmRNAs from KEGG pathway analyses. For the bar plot, color intensity represented the significance of ranked functional pathways top to bottom while the horizontal axis represents the enriched mRNAs. For the dot plot, the bubble size was ordered by the number of enriched mRNAs and the color from red to blue also represented its significance.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementtable1.docx](#)
- [supplementtable2.docx](#)