

Angioplasty for the Treatment of Left Brachiocephalic Vein Tumor: A Case Report

Long Gui

Lu'an People's Hospital affiliated to Anhui Medical University <https://orcid.org/0000-0002-2728-2939>

Hongbo Xu (✉ 416042457@qq.com)

Lu'an People's Hospital affiliated to Anhui Medical University <https://orcid.org/0000-0001-6819-8672>

Chengdong Ning

Lu'an people's Hospital affiliated to Anhui Medical University

Niuliu Huang

Lu'an People's Hospital affiliated to Anhui Medical University

Yu Pan

Lu'an People's Hospital affiliated to Anhui Medical University

Case report

Keywords: brachiocephalic vein tumor, innominate vein aneurysm, angioplasty

Posted Date: October 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-98968/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Cardiothoracic Surgery on March 31st, 2021. See the published version at <https://doi.org/10.1186/s13019-021-01440-y>.

Abstract

Background: Brachiocephalic vein tumor is a rare vascular malformation, which is often reported in case reports. At present it has attracted much attention due to the serious complications, such as vein tumor rupture, pulmonary embolism, venous thrombosis, etc. We report a case of left brachiocephalic vein tumor with compression symptoms.

Case presentation: a 52-year-old male who was admitted to our hospital with irritating cough for more than one month. Chest contrast-enhanced CT showed a localized expansion of 5.2cm in diameter of the left brachiocephalic vein in the anterior mediastinum. The patient received venous angioplasty with brachiocephalic vein tumor resection, and the postoperative recovery was well.

Conclusion: Surgical operation is an effective treatment method for brachiocephalic vein tumor, but it is still necessary to choose the appropriate way according to the type, size, location, lesion scope and complications of the vein tumor.

Background

Brachiocephalic vein tumor (also known as innominate vein aneurysm) is a relatively rare vascular malformation, which mostly appears in the literature as case reports [1]. The exact etiology of brachiocephalic vein tumor remains undetermined. Possible etiologies include congenital developmental defects, malformations, trauma, inflammation, and degenerative changes of the vascular wall [2,3]. It can be divided into primary and secondary. Primary venous tumors are due to the uneven development of vascular structure and defects in the adventitia muscle layer, which lead to the localized weakness and bulging of the vascular wall, in addition to cystic hygroma-related cases. The secondary is mainly due to excessive pressure in the brachiocephalic vein, excessive filling, or heart disease accompanied by blood return disorder. Clinically, it is divided into two subtypes, fusiform and cystic, according to its morphology. The former shows fusiform expansion in the brachiocephalic vein, and the latter shows cystic expansion, connected with the brachiocephalic vein with the tumor neck [4].

Brachiocephalic venous tumors often have no obvious clinical symptoms and signs, and are usually found due to abnormal imaging findings during physical examination. Some patients gradually develop compression symptoms due to the increase in tumor volume, such as post-activity chest tightness, wheezing, coughing after and pain behind the breastbone, etc. There are also a very small number of patients who seek medical treatment due to related complications (such as tumor rupture, pulmonary embolism, venous thrombosis, etc.). The incidence of complications and mortality increases with the increase of tumor cavity [5].

Here we report a case of surgical treatment due to brachiocephalic vein tumor.

Case Presentation

A 52-year-old male patient was admitted to the hospital due to an irritating cough for more than one month. He had no recent history of chest trauma surgery or respiratory diseases. A chest CT scan showed anterior mediastinal space, and showed differential diagnosis of teratoma, thymoma and lymphomas, in order to clarify the relationship between the space-occupying lesions and blood vessels and surrounding tissues, a chest-enhanced CT examination was performed. After intravenous injection of contrast medium, a dual-phase scan was performed. It was found that there was a localized tumor-like expansion in the left brachiocephalic vein and its diameter was 5.2cm with the smooth edge (Fig.1), which confirmed the diagnosis of venous anoma. After careful evaluation of the patient, surgery was performed.

A median sternum incision was selected in the surgery. During the operation, a huge tumor is found to be located in front of the left brachiocephalic vein. The tumor and surrounding tissues were carefully separated to prevent uncontrollable bleeding during the process, and to mechanical damage caused by extracorporeal circulation. We first performed heparinization of the whole body blood, and blocked both ends of the brachiocephalic vein, separated the fibrous membrane of the venous blood vessel, and found that the tumor wall was thin. The tumor was cystic expansion, and there as a larger tumor neck connected to it. After the tumor was removed, a double-layer continuous suture with 5-0 prolene thread was used to suture the brachiocephalic vein. We paid attention to the shape of the normal brachiocephalic vein during the suture process. To prevent vascular distortion and lumen stenosis, we carefully loosened the blocking bands at both ends before knotting, determined the shape of the formed blood vessel, and expelled the possible air (Fig. 2). After confirming that there is no obvious active bleeding, heparin was neutralized with protamine, and the operation ended after the sternum is fixed by the wire.

The mediastinal drainage tube was removed on the third day after the operation, and the patient was discharged from the hospital two days later. He was followed up for 6 months, and there was no obvious abnormality in the chest enhanced CT, with good recovery.

Discussion

Brachiocephalic venous tumors often display no specific clinical symptoms. They are often found by further examinations after physical examination of abnormal mediastinum [5]. In this case, the patient had an irritating cough due to the large tumor and compression. The chest CT scan only detected abnormal mediastinal masses, which requires enhanced CT or MRI to identify. When thrombosis occurs, it is manifested as a localized filling defect in the lumen. If blind biopsy is performed, serious or even fatal complications may occur [6]. MRI of the chest shows abnormal blood flow at the base of the venous tumor, which shows better imaging than CT.

The reports of brachiocephalic vein tumors are often based on individual cases, lacking of unified treatment guidelines. Conservative treatment includes: for fusiform and small cystic brachiocephalic vein tumors, which rarely cause tumor enlargement, compression, rupture, etc., therefore, long-term antiplatelet conservative treatment and regular review are sufficient if there is no significant change in the size and

shape of the tumor [7]. But if the symptoms are aggravated, the tumor is enlarged, or imaging suggests that related complications occur, surgical intervention should be applied.

Scholars are constantly exploring that endovascular treatment is less traumatic for patients, and recovery is quicker after surgery. Gaopo Cai [8] reported that the use of an endovascular stent to treat a case of left brachiocephalic vein tumor have achieved satisfactory early results, but during the 12-18 months follow-up stent thrombosis gradually appeared, and its long-term effect needs further checked. In 2014, Jargiello [9] reported percutaneous transcatheter thrombin injection for the treatment of cystic venous tumors. After intraoperative injection of thrombin, balloon expansion is used to seal the entrance of the tumor to achieve the therapeutic effect. It is suitable for patients with narrow and long tumor necks, or small tumors, but their allergic reactions and intraoperative and postoperative embolization complications need to be vigilant. For this case, because of our insufficient experience in endovascular treatment and the patient was unwilling to try this new technique, surgery was therefore used.

We believe that although the surgical path of median thoracotomy is more traumatic, it gives the surgeon a full view during the operation and facilitates the observation of the anatomical relationship between the venous tumor and the surrounding tissues, and can provide help for the extracorporeal circulation surgery when necessary. This method is safe and effective. After blocking the blood flow on both sides of the tumor, we clamped the neck of the tumor, excised the tumor, and did double-layer continuous suture with 5-0 prolene thread for angioplasty. During the process, we paid attention to the diameter of the venous lumen, to prevent excessive sutures from causing stenosis. The results of the postoperative follow-up visit were satisfactory. The patient's symptoms were significantly improved after the operation. Cheng Fang [10] also reported a case of left brachiocephalic vein tumor with accumulated superior vena cava. They used autologous pericardium tissue from the patient to reconstruct venous blood vessels, and also achieved a good therapeutic effect. In addition, autologous pericardium has the advantages of strong plasticity, easy access, and better immunocompatibility [11], so that patients do not need anticoagulant treatment after surgery. This method is beneficial when the tumor is large and other blood vessels are dilated.

Conclusion

In summary, once the mediastinal space is found and the diagnosis of venous tumors is not ruled out, it is necessary to further examine the chest with enhanced CT or MRI. The treatment methods are diverse, and the clinical needs should be based on the type, size, location, scope of the venous tumor, and complications. Appropriate method should be chosen. For larger cystic brachiocephalic venous tumors, it is feasible to perform surgical resection and angioplasty to reconstruct the venous lumen.

Abbreviations

CT: computed topography; MRI: magnetic resonance imaging.

Declarations

Acknowledgements

Not applicable.

Authors' contributions

Long Gui wrote the manuscript and participate in patient management. Hongbo Xu ,Niuliu, Huang, Yu Pan performed the operation. Chengdong Ning performed manuscript revision. The authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

Not applicable.

Ethics approval and consent to participate

Ethics approval not required.

Consent for publication Written

Consent for publication of information and images was given by the patient.

Competing interests

The authors declare that they have no competing interests.

References

1. Lohrenz C , Daniel Rückner, Wintzer O, et al. Large left innominate vein aneurysm presenting as an anterior mediastinal tumour in a young female[J]. VASA, 2018, 47(6):515-7.
2. Hosaka A, Kato M,Kato I, et al. Brachiocephalic venous aneurysm with unusual clinical observations[J].Journal of Vascular Surgery, 2011;54(6):77S-79S.
3. Yilmaz N , Pelouze A , Murith N , et al. Incidental diagnosis of a silent brachiocephalic vein aneurysm and its surgical treatment[J]. Journal of Vascular Surgery Cases and Innovative Techniques, 2020, 6(1):21-3.
4. Nguyen Q , Olive J K , Vervoort D , et al. Brachiocephalic vein aneurysm: a systematic review of the literature[J]. Journal of Thoracic Disease, 2020, 12(5):2747-58.

5. Schaffer J C , Raudenbush B L , Molinari C , et al. Left innominate vein aneurysm: diagnostic imaging and pitfalls.[J]. Int J Angiol, 2013, 22(02):127-30.
6. Huang Wei, Jiang Ge Ning. A rare case of left innominate vein aneurysm mimicking thymoma[J]. Interact Cardiovasc Thorac Surg 2017;25:669-70.
7. Mikroulis D , Vretzakis G , Eleftheriadis S , et al. Long-term antiplatelet treatment for innominate vein aneurysm[J]. Vasa Ztschrift Für Gefsskrankhten, 2010, 39(3):262-4.
8. Gaopo Cai,Zhaohui Hua,Peng Xu,et al.Endovascular treatment for left innominate vein aneurysm: Case report and literature review[J].Journal of Interventional Medicine,2019,2(01):35-7.
9. Jargiello T , Durakiewicz M, Sojka M,et al. Saccular aneurysm of superior vena cava treated with percutaneous,transcatheter thrombin injection[J]. Cardiovascular & Interventional Radiology, 2014, 37(2):529-32.
10. Cheng Fang, Hui Pan, Zhoubin Li, et al. Autologous pericardium used for reconstruction of left innominate vein in patient with mediastinal venous aneurysm:a case report[J]. 2020, 20(1):656-9.
11. Lam BK, Pettersson GB, Vogt DP. Urgent inferior vena cava replacement with an autologous pericardium tube graft. J Thorac Cardiovasc Surg. 2003; 126:2101–3.

Figures




Figure 1

Chest-enhanced CT scan showed the location of the vein tumor, marked with a pentacle




Figure 2

Intra-operative images of left brachiocephalic vein tumor. A: The tumor and surrounding tissues were carefully separated. B: Block both ends of the brachiocephalic vein. C: The left brachiocephalic vein tumor after angioplasty

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CAREChecklist.pdf](#)