

Baicalein Inhibits Heparin-induced Tau Aggregation by Initializing Tau Oligomer Formation

Shweta K Sonawane

CSIR-National Chemical Laboratory: National Chemical Laboratory CSIR

Vladimir N Uversky

University of South Florida

Subashchandra Bose Chinnathambi (✉ s.chinnathambi@ncl.res.in)

Neurobiology Group, Division of Biochemical Sciences, CSIR-National Chemical Laboratory, Dr. Homi Bhabha Road, 411008 Pune, India <https://orcid.org/0000-0002-5468-2129>

Research

Keywords: Baicalein, Tau inhibition, Tau oligomers, Tau dissolution, Alzheimer's disease

Posted Date: November 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-99003/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 12th, 2021. See the published version at <https://doi.org/10.1186/s12964-021-00704-3>.

Abstract

Background

Amyloid aggregate deposition is the key feature of Alzheimer's disease. The proteinaceous aggregates found in the afflicted brain are the intra-neuronal neurofibrillary tangles formed by the microtubule-associated protein Tau and extracellular deposits, senile plaques, of amyloid beta (A β) peptide proteolytically derived from the amyloid precursor protein. Accumulation of these aggregates has manifestations in the later stages of the disease, such as memory loss and cognitive inabilities originating from the neuronal dysfunction, neurodegeneration, and brain atrophy. Treatment of this disease at the late stages is difficult, and many clinical trials have failed. Hence, the goal is to find means capable of preventing the aggregation of these intrinsically disordered proteins by inhibiting the early stages of their pathological transformations. Polyphenols are known to be neuroprotective agents with the noticeable potential against many neurodegenerative diseases, such as Alzheimer's disease, Parkinson's disease, and Prion diseases.

Methods

We analyzed the capability of Baicalein to inhibit aggregation of human Tau protein by a multifactorial analysis that included several biophysical and biochemical techniques.

Results

The potency of baicalein, a polyphenol from the *Scutellaria baicalensis Georgi*, against many neurodegenerative diseases has been screened and validated. ThS fluorescence assay revealed the potent inhibitory activity of Baicalein, whereas ANS revealed its mechanism of Tau inhibition *viz.* by oligomer capture and dissociation. In addition, Baicalein dissolved the preformed mature fibrils of Tau thereby possessing a dual target action. Tau oligomers formed by Baicalein were non-toxic to neuronal cells, highlighting its role as a potent molecule to be screened against AD.

Conclusion

In conclusion, Baicalein inhibits aggregation of hTau40 by enhancing the formation of SDS-stable oligomers and preventing fibril formation. Baicalein-induced oligomers do not affect the viability of the neuroblastoma cells. Therefore, Baicalein can be considered as a lead molecule against Tau pathology in AD.

Background

Alzheimer's disease (AD) is the leading cause of dementia and cognitive inabilities in aged population.^{1,2} The disease results in a progressive neuronal dysfunction and neuronal atrophy followed by death.³ Protein misfolding and aggregation within the neurons and on the neuronal membranes hamper cellular transport and synaptic transmission and cause neuronal dysfunction. Two functionally distinct and

discrete proteins are involved in the neuronal dysfunction associated with AD. The intra-neuronal functional loss is caused by the neurofibrillary tangles of microtubule-associated protein Tau⁴. The proteinaceous deposits of A β peptide proteolytically derived from the amyloid precursor protein (APP) on the neuronal membrane affects its synaptic transmission.⁵ Both pathological processes together cause neuronal death. The amyloid cascade hypothesis was widely accepted as the key pathological mechanism triggering AD until recent finding that the aberrant APP metabolism triggers the disease and not the A β plaques deposition. This reconsideration of A β hypothesis has also pointed out to the crucial role of Tau pathology as one of the key determinants of AD.⁶ The reconsideration of A β as a trigger in AD was necessitated by the fact that the therapies against A β accumulation,⁷ though were effective in reducing the plaque load, were not able to improve the symptoms of AD.⁸ Hence, the focus of AD therapeutics has shifted towards Tau as a target. Progression of AD can be described based on the Braak stages, which in turn mark the spread of Tau lesions and proportionate the symptoms of the disease.^{9,10} Being intrinsically disordered, Tau is a highly flexible protein, which compacts on temperature variations and adopts transient structures on interaction with its partners.^{11,12} Tau is modified post-translationally, which can either aid in its function or leads to its aggregation.¹³ As a result of aberrant post-translational modifications (PTMs), Tau detaches from microtubules and self-assemble, resulting in the increased intracellular accumulation of aggregated forms.¹⁴ The misfolded and aggregated Tau is also postulated to spread between neurons and aggravate the pathology.¹⁵ The most extensive and important PTM involved in the pathological Tau aggregation is phosphorylation, and hence it is a greatly targeted modification to alleviate the disease.¹⁶ Tau therapeutics can be categorized into wide array of molecules belonging to diverse classes that have specific interaction sites on Tau (Fig. 1A).

There is a repertoire of molecules that were found to be effective against Tau pathology. These compounds include cinnamaldehyde,¹⁷ olecanthal,¹⁸ azaphilones, resveratrol,¹⁹ curcumin,²⁰ quercetin,²¹ polyphenols,²² liminoids,²³ etc.^{24,25} Natural polyphenolic compounds are emerging as neuroprotective molecules because of their ability to abrogate oxidative stress, which is one of the major causative agents of the age-related neurodegeneration.^{26,27} Polyphenols also have multitude of intracellular targets, which makes them efficient candidates for improving cognitive abilities and blocking neurodegeneration.²⁸ Though polyphenols are found to be effective against various molecular targets in neurodegenerative diseases, they have drawbacks that need to be addressed in depth.²⁹ Baicalein is the active flavone purified from the dried roots of *Scutellaria baicalensis* Georgi. This molecule has widespread functions in myriad of pathologies, such as cancer,³⁰ oxidative stress,³¹ inflammation,³² bacterial³³ and viral infections,³⁴ etc. Baicalein have been widely studied in neurodegenerative diseases and found to be potent against protein misfolding and aggregation.³⁵ Baicalein protects against the cerebral ischemia-mediated neuronal death, which is one of the major causes of death and disability world-wide.³⁶ Baicalein attenuates disease pathology in *in vitro* and *in vivo* models of PD. It inhibits aggregation of α -synuclein and abates inflammasome activation in the nigro-striatal dopaminergic system *in vivo*.^{37,38} The Prion protein-induced neurotoxicity, mitochondrial dysfunction and ROS

production are suppressed by Baicalein in the SH-SY5Y and SK-N-SH cells.³⁹ Baicalein also plays a major role in overcoming AD pathologies. It improved A β -induced LTP impairments in mouse model *via* the phosphorylation of Akt, which is involved in cell survival and growth.⁴⁰ Baicalein prevented Tau hyperphosphorylation by inhibiting GSK-3 β activity. This flavone also rescued the behavioral deficits in AD model induced by AlCl₃.⁴¹ All these studies depicted the role of Baicalein in animal models of the neurodegenerative diseases at the physiological level. Our aim is to study the direct effect of Baicalein in inhibiting Tau aggregation and a putative mechanism for its action. Here, we report that Baicalein inhibits heparin-induced Tau aggregation by initializing Tau oligomer formation, dissolving these oligomers and preventing their further fibrillization.

Experimental Procedures

Chemicals and reagents

Baicalein, ThS, ANS, BES, and MTT were purchased from Sigma. Other reagents such as NaCl, Sodium azide, heparin, MgCl₂, EGTA and PMSF were obtained from MP Biomedicals. DTT was obtained from Calbiochem and protease inhibitor cocktail was purchased from Roche. pan-Tau K9JA was purchased from Dako (A-0024) and secondary goat anti-rabbit antibody conjugated to HRP was purchased from Thermo Fisher Scientific (A16110). Hybond-PVDF 10600087 membrane 0.45 μ m was obtained from Amersham Biosciences. The neuroblastoma N2a (ATCC: CCL-131 Neuro-2a Neuroblastoma mouse) cells were purchased from ATCC. The cell culture reagents were purchased from Invitrogen. 400 carbon coated copper grids were purchased from Ted Pella (01822-F). 10 mM and 1 mM stock solutions of Baicalein were prepared in ethanol. 12.5 mM mother stock of ThS was prepared in 1:1 ethanol: miliQ water, which was further diluted to 200 μ M stock in filtered miliQ water. ANS was prepared as 10 mM stock in filtered miliQ.

Tau purification

The recombinant wild type human full-length Tau (hTau40WT) was expressed in *E. coli* BL21* cells by 0.5 mM IPTG induction at 37° C for 4 hours^{11,54-56}. The cells were harvested after induction and mechanically disrupted in a homogenizer (Constant Cell Disruption Systems) at 15000-psi pressure. The cell lysate was supplemented with 0.5M NaCl and 5 mM DTT and heated at 90 °C for 15 minutes. After cooling, the precipitated proteins were separated by centrifugation at 40,000 rpm for 50 minutes. The supernatant containing Tau was dialyzed overnight in Sepharose A buffer (20 mM MES, 1 mM EGTA, 1 mM MgCl₂, 2 mM DTT, 0.1 mM PMSF pH 6.8). The dialysate was further cleared by centrifugation at 40,000 rpm for 50 minutes. Tau was further purified by cation-exchange column chromatography (Sepharose fast flow GE healthcare). The eluate obtained after cation-exchange chromatography was further purified by size-exclusion chromatography (16/600 Superdex 75 pg GE healthcare). The protein concentration was estimated by BCA assay. The protein was concentrated and stored in -80 °C.

Tau aggregation inhibition assay

Aggregation inhibition of full-length Tau in presence of Baicalein was performed as follows.^{25,54} Fresh aliquot of full-length Tau was taken from - 80 °C and centrifuged at 60,000 rpm for 1 hour (Optima Max XP Beckman Coulter). The supernatant was separated and used for assay. The aggregation assay was set up with 20 µM of Tau in 20 mM BES buffer pH 7.4, protease inhibitor cocktail, 0.01% Sodium azide. 5 µM of Heparin (17,500 Da) was used as an inducer of aggregation. 25 mM NaCl and 1 mM DTT were added to the reaction mixture to provide ionic strength and reducing conditions respectively. Baicalein was added in the concentrations of 0-500 µM and the respective reaction mixtures were incubated at 37 °C.

Soluble protein assay

In order to check the effect of Baicalein in absence of the aggregation inducer heparin, a soluble protein assay was set up. The assay composition was same as mentioned above. Baicalein 25 and 100 µM respectively were incubated with 20 µM Tau in presence and absence of inducer heparin in separate reaction mixtures. Two control reaction mixtures were set up for without Baicalein but presence and absence of heparin with 20 µM Tau. All four reaction mixtures were incubated at 37 °C.

PHFs formation in vitro and Tau Disaggregation assay

The mature fibrils for full-length Tau were prepared as follows. 100 µM soluble Tau was incubated with 25 µM Heparin 17,500 Da in the aforementioned reaction buffer. The mature fibril formation was allowed for 8 days at 37 °C. The disaggregation assay was set up by incubating 20 µM of mature fibrils with varied concentrations of Baicalein (0 to 500 µM) and the dissolution of fibrils was recorded by fluorescence, SDS and TEM analysis.

ThS fluorescence assay

The progression of Tau aggregation was monitored by Thioflavin S fluorescence assay^{23,57}. 2 µM of Tau from the reaction mixture was incubated with 8 µM ThS in 50 mM ammonium acetate pH 7.0 for 10 minutes. The readings were recorded in triplicates by exciting the fluorophore at 440 nm and obtaining the emission readings at 521 nm in Tecan Infinite 200 Pro series plate reader. The buffer background fluorescence was subtracted from each reading. Initial measurements were taken within an interval of 5 hours till 25 hours and later readings were continued at an interval of 12 hours each.

ANS fluorescence assay

The hydrophobicity changes in Tau protein during the process of aggregation were monitored by ANS fluorescence measurement. 2 µM Tau was incubated with 40 µM ANS in 50 mM ammonium acetate pH 7.0 for 20 minutes. The fluorescence measurements were carried out at excitation/emission wavelengths of 390 and 475 nm respectively in Tecan Infinite 200 Pro series plate reader.

SDS-PAGE analysis

SDS-PAGE analysis was carried out for the detection confirming Tau aggregation inhibition on 10% polyacrylamide gels at 3 time intervals (0, 48, and 120 hours). The SDS-PAGE gels were quantified by Image Lab software (Bio-Rad). The quantification for complete individual lanes was carried out using the software and the obtained intensities were plotted in the form of the bar graphs.

Immunoblotting

The samples at regular time intervals were confirmed for aggregation inhibition (full-length) by immunoblotting using the antibody against total pan-Tau K9JA (Dako A-0024). The samples from 20 μM reaction mixtures were diluted 10 folds and loaded onto the SDS-PAGE for electrophoresis. The subsequent proteins were transferred onto Hybond-PVDF membrane 0.45 μm (Amersham Biosciences) at 300 mA for 90 minutes. After the transfer the blot was blocked in 5% milk in PBST (0.1% Tween 20) for 1 hour at room temperature. Subsequently, the blot was incubated with primary antibody rabbit polyclonal anti-human Tau K9JA diluted 1:8000 for 1 hour at room temperature. The unbound antibody was removed by washing the blot in PBST (0.1% Tween 20) thrice. The blot was further probed with secondary goat anti-rabbit antibody conjugated to HRP for one hour at room temperature in 1:10000 dilutions. The unbound antibody was washed thrice with PBST for 10 min each and the blot was developed using ECL-Plus reagent (Thermo Fisher Scientific). The chemiluminescence was detected on Amersham Imager 600 and quantified using Image Quant TL software.

Circular Dichroism Spectroscopy

The changes in Tau conformation during the process of aggregation in presence and absence of Baicalein were monitored by CD spectroscopy. The measurements were carried out in quartz cuvette (1 mm) in Jasco J-815 CD spectrometer. The scan was carried out at a speed of 100 nm/min in the range of 190–250 nm at 1 nm bandwidth. A total of 5 acquisitions were obtained for each sample. The samples were diluted to 3 μM in sodium phosphate buffer pH 6.8 for CD measurements.

Electron microscopy measurements

Tau filaments in absence and presence of Baicalein were observed by transmission electron microscope Tecnai G2 20 S-Twin. 400 carbon coated copper grids (Ted Pella 01822-F) were spotted with 2 μM Tau. The excess was blotted on a Whatman paper. Two quick washes of filtered MilliQ water were given for 45 seconds each and the grids were stained in 2% uranyl acetate for 1 minute. The grids were properly dried before measurement.

Cell culture and toxicity assays

The neuroblastoma N2a (ATCC: CCL-131 Neuro-2a Neuroblastoma mouse) cells were used for the toxicity assays. 10,000 cells/well were seeded in 96 well culture plates and grown in complete DMEM F12 media supplemented with antibiotic Penstrep⁵⁸ for 24 hours. The cells were treated with Baicalein (0 to 20 μM) with 5 μM full-length soluble Tau or 5 μl of Tau aggregates (50 μM total concentration) in serum-starved media (0.5% FBS) for 24 hours. 0.5 mg/ml MTT (Methylthiazolyldiphenyl-tetrazolium bromide) was added to each well and incubated for four hours at 37 °C. The viable cells enzymes reduce MTT into

formazan crystals which were dissolved in 100 μ l of 100% DMSO. The reading was taken at 570 nm in Tecan Infinite 200 Pro series plate reader. The viability of untreated cells was considered as 100% and the viability of treatment groups was calculated in accordance with this untreated group.

Statistical analysis

The error bars represent mean \pm SD values. 95% confidence intervals. The statistical analyses were carried out by Sigma Plot 10.0. Unpaired t-test was used to calculate the p-values. *: p value \leq 0.05, **: p value \leq 0.01, ***: p value \leq 0.001. Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| >$ Tukey's criterion.

Results

Baicalein inhibits hTau40 assembly

Screening of Tau aggregation inhibitors (TAI) in cell-free assays represents a well-established approach. These primary screens include ThS fluorescence to study the progress of aggregation in solution. We adopted a similar preliminary screening approach to check the inhibitory effects of Baicalein (0-500 μ M) on full-length Tau aggregation (Fig. 1B). We monitored ThS fluorescence initially at every 5 hours till 24 hours followed by the interval of 12 hours. The initial 5 hours interval measurements helped capturing inhibition by higher concentrations of Baicalein (100, 500 μ M). The fluorescence increased till 5 hours for the higher concentrations followed by gradual decrease (Fig. 1C). This pattern of changes in fluorescence intensity was observed at lower Baicalein concentrations as well, but at increased incubation time. The fluorescence intensity did not decrease for the untreated Tau showing steady aggregation. This decrease in ThS fluorescence intensity in Baicalein-treated samples suggested concentration-dependent inhibition of Tau aggregation. The kinetics was monitored till 120 hours and the percent inhibition at the end of 120 hours was maximum 85% at 500 μ M of Baicalein and 75% at 100 μ M Baicalein (Fig. 1D). The IC_{50} value for inhibition of aggregation for full-length Tau was found to be 27.6 μ M of Baicalein (Fig. 1E). Therefore, Baicalein was found to inhibit the assembly of full-length Tau aggregates at extended incubation time of 120 hours.

Baicalein is involved in the formation of Tau oligomers

The ThS fluorescence studies revealed a distinct pattern of initial surge followed by gradual decrease, which led us to study the transition species formed in presence of Baicalein by ANS fluorescence. ANS binds to intermediate partially folded states with transient secondary structure with less compactness. ANS fluorescence showed a similar pattern like ThS with an initial increase and a further decrease in fluorescence (Fig. 2A, B). The hydrophobicity changes mapped by ANS showed that 500 μ M of Biacalein increased Tau hydrophobicity as compared to lower concentrations (Fig. 2C). But at the higher concentration of Baicalein, ANS showed rapid and substantial increase in fluorescence at the initial time

points and then the decrease before becoming stagnant (Fig. 2A). The kinetics of ANS suggests that Baicalein might be involved in initial Tau oligomer formation restricting further fibrillization.

SDS-PAGE and Immunoblot analysis confirms Tau oligomerization in the presence of Baicalein

The fluorescence assays suggested role of Baicalein in inducing Tau oligomers. In order to reaffirm this hypothesis, reaction mixtures were analyzed at regular time intervals using the 10% SDS-PAGE. After 48 hours of incubation, we observed a clear increase in higher molecular weight oligomers in sample treated with 100 μ M Baicalein as compared to the control (Fig. 3A,B; 48 hours red asterisk). At the same time point, ThS fluorescence decreased and ANS fluorescence began to increase suggesting the arrest of Tau oligomers by Baicalein. But the Tau sample treated with 500 μ M Baicalein showed complete absence of higher order aggregates at 48 hours suggesting complete inhibition of Tau assembly. As the time advanced, at the end of 120 hours of incubation, we observed a clear decrease in the higher order aggregates in the 100 μ M treated sample as compared to control (Fig. 3A, B; 120 hours green asterisk). To confirm the aggregation inhibition of full-length Tau by Baicalein via promotion of oligomerization, we further carried out immunoblotting for these higher order aggregates by probing them with the total pan-Tau antibody (K9JA) at various time intervals. At zero time point, only soluble Tau protein could be visualized (Fig. 3C). The 24 hours incubation showed complete oligomerization in the Tau samples treated with 200 μ M Baicalein (all the material was present in the stacking gel part of the blot) as opposed to control, which showed mixed population of oligomers in the form of a smear in the resolving as well as stacking gel part of blot. The ANS fluorescence revealed that the treated samples showed increased oligomerization as the time of incubation progressed.

Baicalein arrests Tau oligomers with partial secondary structure

For many aggregating intrinsically disordered proteins, oligomers are partially folded intermediate species formed during the process of aggregation. In order to check for the conformational states of Baicalein-induced oligomers, far-UV circular dichroism (CD) spectroscopic analysis was performed. Tau is a natively unfolded (intrinsically disordered) proteins with a characteristic far-UV CD spectrum possessing prominent minimum at 198 nm and not showing significant signal at 220 nm (Fig. 4A). On the other hand, the far-UV CD spectrum of Tau protein in the presence of Baicalein is consistent with the presence of partial β -sheet structure, suggesting that Baicalein was able to induce some partial folding of tau protein, likely leading to the formation of oligomeric species (Fig. 4B). Next, these Baicalein-induced (or Baicalein-stabilized) oligomers were visualized by transmission electron microscopy (see Fig. 4C). The untreated Tau showed long filamentous aggregates at 120 hours. Baicalein-treated Tau showed small filaments at 48 and 120 hours with lack of intact fibrillar aggregates (Fig. 4C). Therefore, these qualitative analyses along with the conformational studies suggested that Baicalein arrests Tau aggregation at the stage of oligomer formation and prevents complete fibrillation of this protein.

Baicalein does not alter Tau conformation

All the biochemical and biophysical experiments suggested induction (or stabilization) of Tau oligomers by Baicalein in the presence of heparin. In order to check whether Baicalein in the absence of heparin can cause conformational changes in Tau, this protein was incubated with Baicalein in absence of heparin. The ThS and ANS fluorescence did not increase for Baicalein-treated samples (in absence of heparin) (Fig. 5A, B). The SDS-PAGE analysis showed the presence of higher order structures in the heparin treated Tau. The sample treated with higher concentration of Baicalein (100 μM) showed presence of very faint higher order bands at a later time point 120 hours (Fig. 5C, D). The fluorescence kinetics and the SDS-PAGE analysis clearly suggest that Baicalein might be able to induce Tau oligomerization but at a much slower pace. The structural analysis of these oligomers revealed that in the absence of heparin, Baicalein does not induce transition of Tau to β -sheet structure and maintains the native random coil conformation (Fig. 5E). The qualitative electron microscopy analysis showed that Baicalein-treated Tau without heparin does not form fibrils (Fig. 5F).

Baicalein dissolves the mature fibrils of the aggregated Tau protein

Although, Tau oligomers act as a toxic species, the filamentous Tau increase the aggregate load in the neuron, which hampers the cellular functioning, as well as increases the load on the clearance machinery.⁴² Hence, it is important to have some efficient means that could dissolve these filamentous Tau aggregates to non-harmful oligomers. Since Baicalein had the inhibitory effect on Tau aggregation, we were interested to see whether this flavone can dissolve the pre-formed mature fibrils. To this end, the mature fibrils of full-length Tau were treated with series of concentration of Baicalein (0-500 μM). ThS fluorescence was used to monitor the disaggregation. The fluorescence intensity was found to decrease in the Baicalein-treated samples (Fig. 6A), with disaggregation approaching 90% in the presence of 500 μM Baicalein whereas the fluorescence of untreated control remained almost stagnant (Fig. 6B). The DC_{50} value for the disaggregation of fibrils formed by the full-length Tau was found to be 24.6 μM (Fig. 6C). The conformational analysis of disaggregated samples suggested the presence of mixed population of Tau and its aggregates (Fig. 6D). The SDS-PAGE analysis revealed a time- and concentration-dependent decrease in the aggregate load in the Baicalein-treated samples (Fig. 7A,B). The electron micrographs showed the presence of fibrillar aggregates in the control, but only amorphous aggregates and oligomers in the Baicalein-treated samples (Fig. 7C). These observations suggest that Baicalein has a dual role, being able to inhibit Tau fibrillation and to disaggregate the preformed Tau fibrils.

Baicalein is biocompatible and rescues Tau aggregate-mediated toxicity in neuronal cells

To assess the toxicity of Baicalein to neuronal cells, we screened a broad range of Baicalein concentrations from 5 nM to 20 μM for toxicity to neuroblastoma (N2a) cells. Baicalein was found to be

non-toxic even at 25 μM concentration (Fig. 8A). To demonstrate the toxicity of Baicalein-treated Tau, cells were exposed to the Tau aggregation reaction mixtures containing various concentrations of Baicalein. Tau:Baicalein ratios were maintained as (1:0.25, 1:0.5, 1:1, 1:2 and 1:4) for 120 hours. Baicalein-treated full-length Tau was found to be non-toxic to the neuronal cells (Fig. 8B), and the morphology of treated cells remained unchanged. As Baicalein disaggregated preformed Tau fibrils, we tested the effect of such disaggregated Tau on the N2a cells. Tau aggregates were treated with various concentrations of Baicalein and given to cells at 0 and 120 hours. The highest concentration of Baicalein was shown to rescue the toxicity of Tau aggregates at 120 hours and the cell morphology remained intact (Fig. 8C). Therefore, Baicalein-induced oligomers as well as Baicalein-treated mature fibrils are non-toxic to N2a cells.

Discussion

Tau as a target in AD has been extensively studied, and a great number of molecules were found to be potent against Tau-mediated pathologies. Baicalein has been implicated in overcoming AD pathologies with the major targets being ROS generation and rescuing of the behavioral and cognitive deficits.⁴⁰ The behavioral and cognitive deficits are late manifestations of Tau pathology, which starts with abrupt aggregation followed by gradual accumulation of aggregated material. In the current scenario, the pathological transitions of Tau at the initial stages needs to be inhibited. Few molecules, including methylene blue, azaphilones, anthraquinones, *etc.* have been found to inhibit Tau aggregation at the initial stages. Baicalein, a natural small molecule was shown to be inhibit the aggregation of α -synuclein, a key protein involved in the PD pathogenesis.⁴³ We report analogous results for human Tau protein, wherein Baicalein efficiently inhibits the proteins aggregation *in vitro* (Fig. 9). Though, the IC_{50} of Baicalein for Tau inhibition is reported to be 2.7 μM ,⁴⁴ no sufficient evidence is available for the same. We report an IC_{50} of 27.8 μM for inhibition of Tau and DC_{50} of 24.6 μM . Similar for α -synuclein, Baicalein enhanced Tau oligomerization. In the presence of Baicalein, the aggregation process of Tau is halted at the stage of the oligomer formation, without allowing protein to form fibrils. This is evident from ANS fluorescence, which increases initially and then decreases at the extended incubation time. ANS fluorescence increases due to the binding of fluorophore to the exposed hydrophobic regions of Tau during aggregation.⁴⁵ Thus, Baicalein enhances Tau oligomerization and captures these intermediate species. The structural studies revealed that the Baicalein-stabilized α -synuclein oligomers have a β -sheet-rich structure,⁴⁶ which is in agreement with our data, wherein the conformational analyses of Baicalein-induced Tau oligomers revealed the presence of the partial β -sheet structure. Baicalein-induced Tau oligomers were found to be SDS-resistant. The SDS-stable $\text{A}\beta$ oligomers are formed by green tea polyphenol EGCG *via* polar and non-polar interactions.^{47,48} Several other polyphenols, such as nordihydroguaiaretic acid (NDGA), resveratrol, and myricetin divert the $\text{A}\beta$ monomers and fibrils towards SDS-stable oligomers.⁴⁹ The catecholamine dopamine inhibits α -synuclein fibrillization, as well as forms covalent adducts with this protein resulting in the formation of the SDS- and heat-stable oligomers.⁵⁰ The oligomers induced by these compounds are off the fibrillation pathway and non-toxic, which is similar to

Baicalein-induced hTau40 oligomers characterized in our study. Baicalein treatment inhibited the mature fibril formation for Tau, which was visualized by TEM. The treatment with bioflavonoid cinnamaldehyde show presence of broken Tau filaments and absence of intact fibrils.¹⁷ Thus, bioflavonoids might have similar mechanisms for inhibiting the amyloidogenic aggregation of proteins. Baicalein disaggregated the pre-formed fibrils of Tau similar to α -synuclein fibril dissolution.⁴³ This shows that Baicalein might have a common pathway of inhibiting misfolded protein aggregation and dissolution of mature fibrils. Baicalein interacts with proteins in a residue-specific manner. For example, in human serum albumin it interacts with Leu, Arg, and Ala residues.⁵¹ On the other hand, Baicalein binds close to the tyrosine residue of A β and forms Schiff's base with the lysine residue.⁵² Furthermore, the Baicalein-induced Tau oligomers were found to be non-toxic to the neuronal cells. Similarly, Baicalein rescued the toxicity of α -Synuclein oligomers.⁵³ Further, it also protected SH SY-5Y and HeLa cells from toxicity of α -Synuclein and A β oligomers, thus proving to be a potential molecule against amyloid aggregation in general.

Conclusion

In conclusion, Baicalein inhibits aggregation of hTau40 by enhancing the formation of SDS-stable oligomers and preventing fibril formation. Baicalein-induced oligomers do not affect the viability of the neuroblastoma cells. Therefore, Baicalein can be considered as a lead molecule against Tau pathology in AD (Fig. 9).

Abbreviations

IDPs

Intrinsically disorder proteins

PHFs

Paired helical filaments

ThS

Thioflavin S

ANS

8-Anilinonaphthalene-1-sulfonic acid

CD

Circular dichroism

TEM

Transmission election microscopy

MTT

Methylthiazolyldiphenyl-tetrazolium bromide

Declarations

Ethics declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported in part by grants from the Department of Science and Technology-Science and Engineering Research Board SERB/HS-EMR000306/2018 and in-house CSIR-National Chemical Laboratory grant MLP029526. SS greatly acknowledges the Department of Biotechnology, Government of India for her fellowship.

Author contributions

S.C. designed the work, conceived and supervised this project, and wrote the paper; S.S. and S.C carried out the experiments and analyzed the data. S.C. and S.S. wrote the paper. V.N.U. helped in the discussion and edited the manuscript. All authors read and approved the final version of the manuscript.

Corresponding author

Correspondence to **Subashchandrabo**se Chinnathambi, email s.chinnathambi@ncl.res.in

Acknowledgements

The authors greatly acknowledge the Chinnathambi lab member for critical reading and suggestions.

Author information

Affiliations

Shweta Kishor Sonawane, Subashchandrabose Chinnathambi

Neurobiology Group, Division of Biochemical Sciences, CSIR-National Chemical Laboratory,

Dr. Homi Bhabha Road, 411008 Pune, India

Shweta Kishor Sonawane, Subashchandrabose Chinnathambi

Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, 201002, India

Vladimir N. Uversky

Department of Molecular Medicine and USF Health Byrd Alzheimer's Research Institute, Morsani College of Medicine, University of South Florida, Tampa, Florida, 33612, USA

References

- 1 Association, A. s. 2016 Alzheimer's disease facts and figures. *Alzheimer's & Dementia***12**, 459-509 (2016).
- 2 Scheltens, P. *et al.* Alzheimer's disease. *The Lancet***388**, 505-517, doi:10.1016/S0140-6736(15)01124-1 (2016).
- 3 Bondi, M. W., Edmonds, E. C. & Salmon, D. P. Alzheimer's Disease: Past, Present, and Future. *Journal of the International Neuropsychological Society***23**, 818-831, doi:10.1017/S135561771700100X (2017).
- 4 Wang, Y. & Mandelkow, E. Tau in physiology and pathology. *Nature Reviews Neuroscience***17**, 22 (2016).
- 5 Sheng, M., Sabatini, B. L. & Südhof, T. C. Synapses and Alzheimer's disease. *Cold Spring Harbor perspectives in biology***4**, a005777 (2012).
- 6 Wang, J., Gu, B. J., Masters, C. L. & Wang, Y.-J. A systemic view of Alzheimer disease—insights from amyloid- β metabolism beyond the brain. *Nature Reviews Neurology***13**, 612 (2017).
- 7 Kametani, F. & Hasegawa, M. Reconsideration of amyloid hypothesis and tau hypothesis in Alzheimer's disease. *Frontiers in neuroscience***12**, 25 (2018).
- 8 Ricciarelli, R. & Fedele, E. The amyloid cascade hypothesis in Alzheimer's disease: it's time to change our mind. *Current neuropharmacology***15**, 926-935 (2017).
- 9 Braak, H. & Braak, E. Staging of Alzheimer's disease-related neurofibrillary changes. *Neurobiology of aging***16**, 271-278 (1995).
- 10 Braak, H. & Del Tredici, K. Spreading of tau pathology in sporadic Alzheimer's Disease along Cortico-cortical top-down connections. *Cerebral Cortex***28**, 3372-3384 (2018).
- 11 Gorantla, N. V., Shkumatov, A. V. & Chinnathambi, S. in *Tau Protein* 3-20 (Springer, 2017).
- 12 Avila, J. *et al.* Tau structures. *Frontiers in aging neuroscience***8**, 262 (2016).

- 13 Martin, L., Latypova, X. & Terro, F. Post-translational modifications of tau protein: implications for Alzheimer's disease. *Neurochemistry international***58**, 458-471 (2011).
- 14 Mietelska-Porowska, A., Wasik, U., Goras, M., Filipek, A. & Niewiadomska, G. Tau protein modifications and interactions: their role in function and dysfunction. *International journal of molecular sciences***15**, 4671-4713 (2014).
- 15 Sonawane, S. K. & Chinnathambi, S. Prion-Like Propagation of Post-Translationally Modified Tau in Alzheimer's Disease: A Hypothesis. *Journal of Molecular Neuroscience***65**, 480-490 (2018).
- 16 Hanger, D. P., Anderton, B. H. & Noble, W. Tau phosphorylation: the therapeutic challenge for neurodegenerative disease. *Trends in molecular medicine***15**, 112-119 (2009).
- 17 Peterson, D. W. *et al.* Cinnamon extract inhibits tau aggregation associated with Alzheimer's disease in vitro. *Journal of Alzheimer's Disease***17**, 585-597 (2009).
- 18 Li, W. *et al.* Inhibition of tau fibrillization by oleocanthal via reaction with the amino groups of tau. *Journal of neurochemistry***110**, 1339-1351 (2009).
- 19 He, X. *et al.* Resveratrol attenuates formaldehyde induced hyperphosphorylation of tau protein and cytotoxicity in N2a cells. *Frontiers in neuroscience***10**, 598 (2017).
- 20 Rane, J. S., Bhaumik, P. & Panda, D. Curcumin inhibits tau aggregation and disintegrates preformed tau filaments in vitro. *Journal of Alzheimer's Disease***60**, 999-1014 (2017).
- 21 Shen, X. Y. *et al.* Quercetin inhibits okadaic acid-induced tau protein hyperphosphorylation through the Ca²⁺-calpain-p25-CDK5 pathway in HT22 cells. *International journal of molecular medicine***41**, 1138-1146 (2018).
- 22 Syarifah-Noratiqah, S.-B., Naina-Mohamed, I., Zulfarina, M. S. & Qodriyah, H. Natural polyphenols in the treatment of Alzheimer's disease. *Current drug targets***19**, 927-937 (2018).
- 23 Gorantla, N. V., Das, R., Mulani, F. A., Thulasiram, H. V. & Chinnathambi, S. Neem derivatives inhibits tau aggregation. *Journal of Alzheimer's disease reports***3**, 169-178 (2019).
- 24 Spagnuolo, C. *et al.* Neuroprotective role of natural polyphenols. *Current topics in medicinal chemistry***16**, 1943-1950 (2016).
- 25 Dubey, T. & Chinnathambi, S. Brahmi (*Bacopa monnieri*): An ayurvedic herb against the Alzheimer's disease. *Archives of Biochemistry and Biophysics*, 108153 (2019).
- 26 Darvesh, A. S., Carroll, R. T., Bishayee, A., Geldenhuys, W. J. & Van der Schyf, C. J. Oxidative stress and Alzheimer's disease: dietary polyphenols as potential therapeutic agents. *Expert review of neurotherapeutics***10**, 729-745 (2010).

- 27 Freysson, A., Page, G., Fauconneau, B. & Bilan, A. R. Natural polyphenols effects on protein aggregates in Alzheimer's and Parkinson's prion-like diseases. *Neural regeneration research***13**, 955 (2018).
- 28 Ramassamy, C. Emerging role of polyphenolic compounds in the treatment of neurodegenerative diseases: a review of their intracellular targets. *European journal of pharmacology***545**, 51-64 (2006).
- 29 Ebrahimi, A. & Schluesener, H. Natural polyphenols against neurodegenerative disorders: potentials and pitfalls. *Ageing research reviews***11**, 329-345 (2012).
- 30 Liu, H. *et al.* The fascinating effects of baicalein on cancer: a review. *International journal of molecular sciences***17**, 1681 (2016).
- 31 Kang, K. A. *et al.* Baicalein inhibits oxidative stress-induced cellular damage via antioxidant effects. *Toxicology and industrial health***28**, 412-421 (2012).
- 32 Li, F.-Q., Wang, T., Pei, Z., Liu, B. & Hong, J.-S. Inhibition of microglial activation by the herbal flavonoid baicalein attenuates inflammation-mediated degeneration of dopaminergic neurons. *Journal of Neural Transmission***112**, 331-347 (2005).
- 33 Górnjak, I., Bartoszewski, R. & Króliczewski, J. Comprehensive review of antimicrobial activities of plant flavonoids. *Phytochemistry Reviews***18**, 241-272 (2019).
- 34 Moghaddam, E. *et al.* Baicalin, a metabolite of baicalein with antiviral activity against dengue virus. *Scientific reports***4**, 5452 (2014).
- 35 Li, Y., Zhao, J. & Hölscher, C. Therapeutic potential of baicalein in Alzheimer's disease and Parkinson's disease. *CNS drugs***31**, 639-652 (2017).
- 36 Liang, W., Huang, X. & Chen, W. The effects of Baicalin and Baicalein on cerebral ischemia: a review. *Aging and disease***8**, 850 (2017).
- 37 Hung, K.-C., Huang, H.-J., Wang, Y.-T. & Lin, A. M.-Y. Baicalein attenuates α -synuclein aggregation, inflammasome activation and autophagy in the MPP⁺-treated nigrostriatal dopaminergic system in vivo. *Journal of ethnopharmacology***194**, 522-529 (2016).
- 38 Zhao, W.-Z., Wang, H.-T., Huang, H.-J., Lo, Y.-L. & Lin, A. M.-Y. Neuroprotective effects of baicalein on acrolein-induced neurotoxicity in the nigrostriatal dopaminergic system of rat brain. *Molecular neurobiology***55**, 130-137 (2018).
- 39 Moon, J.-H. & Park, S.-Y. Baicalein prevents human prion protein-induced neuronal cell death by regulating JNK activation. *International journal of molecular medicine***35**, 439-445 (2015).
- 40 Gu, X.-H. *et al.* The flavonoid baicalein rescues synaptic plasticity and memory deficits in a mouse model of Alzheimer's disease. *Behavioural brain research***311**, 309-321 (2016).

- 41 Zhou, L. *et al.* Baicalein improves behavioral dysfunction induced by Alzheimer's disease in rats. *Neuropsychiatric disease and treatment***12**, 3145 (2016).
- 42 Tai, H.-C. *et al.* The synaptic accumulation of hyperphosphorylated tau oligomers in Alzheimer disease is associated with dysfunction of the ubiquitin-proteasome system. *The American journal of pathology***181**, 1426-1435 (2012).
- 43 Zhu, M. *et al.* The flavonoid baicalein inhibits fibrillation of α -synuclein and disaggregates existing fibrils. *Journal of Biological Chemistry***279**, 26846-26857 (2004).
- 44 Masuda, M. *et al.* Small molecule inhibitors of α -synuclein filament assembly. *Biochemistry***45**, 6085-6094 (2006).
- 45 Kumar, S. *et al.* Stages and conformations of the Tau repeat domain during aggregation and its effect on neuronal toxicity. *Journal of Biological Chemistry***289**, 20318-20332 (2014).
- 46 Hong, D.-P., Fink, A. L. & Uversky, V. N. Structural characteristics of α -synuclein oligomers stabilized by the flavonoid baicalein. *Journal of molecular biology***383**, 214-223 (2008).
- 47 del Amo, J. M. L. *et al.* Structural properties of EGCG-induced, nontoxic Alzheimer's disease A β oligomers. *Journal of molecular biology***421**, 517-524 (2012).
- 48 Hyung, S.-J. *et al.* Insights into antiamyloidogenic properties of the green tea extract (-)-epigallocatechin-3-gallate toward metal-associated amyloid- β species. *Proceedings of the National Academy of Sciences***110**, 3743-3748 (2013).
- 49 Taniguchi, S. *et al.* Inhibition of Heparin-induced Tau Filament Formation by Phenothiazines, Polyphenols, and Porphyrins. *Journal of Biological Chemistry***280**, 7614-7623, doi:10.1074/jbc.M408714200 (2005).
- 50 Conway, K. A., Rochet, J.-C., Bieganski, R. M. & Lansbury, P. T. Kinetic stabilization of the α -synuclein protofibril by a dopamine- α -synuclein adduct. *Science***294**, 1346-1349 (2001).
- 51 Liu, H., Bao, W., Ding, H., Jang, J. & Zou, G. Binding Modes of Flavones to Human Serum Albumin: Insights from Experimental and Computational Studies. *The Journal of Physical Chemistry B***114**, 12938-12947, doi:10.1021/jp102053x (2010).
- 52 Song, S.-m., Wang, Y.-x., Xiong, L.-m., Qu, L.-b. & Xu, M.-t. Interaction between baicalein and amyloid- β fibrils studied by fluorescence spectroscopy. *Chemical Research in Chinese Universities***29**, 20-25, doi:10.1007/s40242-013-2180-1 (2013).
- 53 Lu, J.-H. *et al.* Baicalein Inhibits Formation of α -Synuclein Oligomers within Living Cells and Prevents A β Peptide Fibrillation and Oligomerisation. *ChemBioChem***12**, 615-624, doi:10.1002/cbic.201000604 (2011).

54 Sonawane, S. K., Balmik, A. A., Boral, D., Ramasamy, S. & Chinnathambi, S. Baicalein suppresses Repeat Tau fibrillization by sequestering oligomers. *Archives of Biochemistry and Biophysics* **675**, 108119 (2019).

55 Sonawane, S. K., Ahmad, A. & Chinnathambi, S. Protein-Capped Metal Nanoparticles Inhibit Tau Aggregation in Alzheimer's Disease. *ACS omega* **4**, 12833-12840 (2019).

56 Gorantla, N. V., Khandelwal, P., Poddar, P. & Chinnathambi, S. in *Tau Protein* 21-31 (Springer, 2017).

57 Gorantla, N. V. *et al.* Molecular Cobalt (II) Complexes for Tau Polymerization in Alzheimer's Disease. *ACS omega* (2019).

58 Balmik, A. A., Sonawane, S. K. & Chinnathambi, S. Modulation of Actin network and Tau phosphorylation by HDAC6 ZnF UBP domain. *bioRxiv*, 702571 (2019).

Figures




Figure 1

Tau aggregation inhibition by Baicalein. A) Tau protein domain organization with the known binding sites for the small molecule inhibitors of Tau aggregation. B) ThS fluorescence for aggregation kinetics of full-length Tau in presence of varying concentrations of Baicalein. The ThS fluorescence decreased with increase in dose of Baicalein showing concentration dependent inhibition of Tau aggregation. C) ThS fluorescence for Baicalein treated samples at the initial time point revealed that the higher concentrations had rapid effect on inhibiting Tau aggregation within first 24 hours. The ThS fluorescence intensity

decreased rapidly in the 24 hours time window and remained at basal level throughout the kinetics. D) The ThS percent inhibition of Tau aggregation at different concentrations of Baicalein. The highest concentration of Baicalein 500 μM showed around 85% inhibition followed by 75% by 100 μM of Baicalein. E) The IC₅₀ value for full-length Tau aggregation inhibition, which was found to be 27.69 μM of Baicalein. (The values are mean \pm std. deviation of two independent experiments. The statistical analysis was carried out by Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| >$ Tukey's criterion.






Figure 2

Tau oligomerization-induced by Baicalein. A) ANS fluorescence for aggregation kinetics of full-length Tau in presence of varying concentrations of Baicalein. B) The highest concentration (500 μM) of Baicalein

induced Tau oligomerization within 15 hours of incubation as evidenced by sharp rise in ANS fluorescence. C) The hydrophobicity changes at the end of 120 hours revealed that 500 μM of Baicalein showed increased hydrophobicity which is characteristic property of oligomers. The statistical analysis was carried out by Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| > \text{Tukey's criterion}$.

A



C




Figure 3

SDS-resistant Tau oligomers-induced by Baicalein. A) SDS-PAGE analysis of Tau oligomers. The zero time point did not show presence of any aggregates in control as well as treated samples. At the end of 48 hours higher order aggregates (red asterisk) were observed in the 100 μ M Baicalein treated sample as compared to control. The 120 hours incubated sample showed a decrease in aggregates (green asterisk) in treated samples as compared to control. B) The SDS-PAGE quantification for different time points indicate the initial increase followed by the decrease in the aggregates at higher concentrations of Baicalein shows significant decrease in 500 μ M Baicalein at 48 and 120 hours. C) The formation of Tau oligomers induced by Baicalein in presence of heparin was also confirmed by immunoblot analysis using the pan-Tau antibody K9JA for total Tau. At 24 hours of incubation higher order oligomers were observed in Baicalein treated samples (lane 6, 7) as opposed to control (lane 2). After 48 hours of incubation the oligomers population was seen to increase (lane 3, 4) but at 120 hours the decrease in the oligomer load was observed (lane 7). The statistical analysis was carried out by Student's unpaired T- test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| >$ Tukey's criterion.




Figure 4

Tau aggregation inhibition studied by CD spectroscopy. A) The conformational analysis of Baicalein treated and control samples showed the typical spectra of a random coil structure. The Sol. represents soluble protein without heparin. B) At the end of 120 hours the pattern of spectrum remained same with respect to soluble (black arrow) and treated samples (pink arrow) showed a partial β -sheet conformation. C) The control untreated sample shows the presence of long filaments (white arrows) at day 5. The

treatment with 100 μM Baicalein at the initial time points showed presence of intermediately long filaments (white arrow) and small broken filaments. As the time advanced, Baicalein was shown to inhibit Tau aggregation as only broken pieces (highlighted by black border and black arrow) of Tau filaments were visualized in the electron micrographs.




Figure 5

Interaction of Baicalein with soluble Tau. A) The aggregation of Tau was monitored by ThS fluorescence. The positive control (heparin) showed an increase in intensity with respect to time as opposed to heparin negative control. The Baicalein treated soluble protein did not show any increase in fluorescence as compared to positive control suggesting that Baicalein does not lead to Tau fibrillization in absence of heparin. B) The hydrophobicity imparted by Baicalein alone without heparin was monitored by ANS fluorescence. The intensity did not increase significantly as compared to the positive control. C) The SDS-PAGE analysis revealed a time dependent increase in higher molecular aggregates in the positive control from 24 hours of incubation (lane 2). No higher molecular weight aggregates were observed in negative control as well as sample treated with 25 μ M of Baicalein (lanes 3, 4 respectively) but faint bands of higher order oligomers were observed in 100 μ M treated sample from 72 hours onwards (lane 5), suggesting formation of oligomers. D) Quantification for SDS-PAGE for soluble Tau with Baicalein. E) Conformational analysis of soluble Tau with Baicalein. The CD analysis showed normal spectra for negative control and Baicalein treated soluble Tau but a slight shift in the positive control (aggregated) Tau. F) The electron micrographs at different time points reveal presence of amorphous aggregates at the initial time points and clumps of oligomers at the end point. The statistical analysis was carried out by Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| > \text{Tukey's criterion}$.




Figure 6

Dissolution of Tau aggregates by Baicalein. A) The effect of Baicalein on the dissolution of mature Tau aggregates was studied using ThS fluorescence, which showed a decrease after 5 hours of incubation in a concentration-dependent manner. B) The plot shows the DC50 value for Tau disaggregation as 24.6 μM. C) The percent disaggregation plot shows a maximum of 85% disaggregation by 500 μM of Baicalein treatment. D) The CD analysis of disaggregated Tau by Baicalein shows the presence of mixed structures. The Sol. represents for soluble protein. The statistical analysis was carried out by Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| >$ Tukey's criterion.




Figure 7

Baicalein-mediated Tau dissolution by SDS-PAGE A) Figure shows the time-dependent SDS-PAGE analysis of Tau aggregates dissolution. After 72 hours, the higher concentrations of Baicalein showed a maximum decrease in the aggregates (lane 7, 8, 9 green asterisk) as compared to control (lane 2 red asterisk). B) The SDS-PAGE quantification for different time points indicate the initial increase followed by the decrease in the aggregates at the higher concentrations of Baicalein. C) Transmission electron micrographs showed the untreated mature fibrils (white arrow) versus disaggregated amorphous (encircled black), Baicalein treated fibrils of full-length Tau. The statistical analysis was carried out by

Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| >$ Tukey's criterion.




Figure 8

Cell toxicity of Baicalein. A) N2a cells treated with a broad range of Baicalein shows no toxicity till 25 μM . B) Baicalein treated soluble Tau shows no significant effect on cell viability implying Baicalein-induced oligomers are non-toxic. C) Baicalein treated Tau aggregates shows a mild rescue of toxicity at 120 hours with the highest concentration of Baicalein suggesting that disaggregated Tau mature filaments are non-toxic to neuro2a cells. The viability of untreated cells was considered as 100% and the viability of treatment groups was calculated in accordance with this untreated group. The statistical analysis was carried out by Student's unpaired T-test with respect to Baicalein untreated control. *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$. ns: non-significant p value). Further, post hoc analysis was carried by one-way ANOVA and Tukey's criterion was determined for honestly significant difference (HSD). The data was considered significant if $|X-X'| > \text{Tukey's criterion}$.




Figure 9

Dual effect of Baicalein on Tau. Tau in presence of heparin forms mature fibrils via intermediate oligomers formation. On the other hand, Tau in presence of Baicalein and heparin is restricted to form Tau oligomers without further mature fibril formation. These PHFs when exposed with Baicalein are

dissolved into smaller fragments. The exposure of heparin-induced Tau oligomers and PHFs to healthy neuronal cells imparts toxicity in neurons and leads to neuronal death. The Baicalein-induced Tau oligomers as well as the dissolved fragments of PHFs do not affect the neuronal health and maintains the morphology. These suggest nontoxic nature of Baicalein-induced Tau oligomers.