

Impact of dietary Mannan-oligosaccharide and β -Glucan supplementation on growth, histopathology, E-coli colonization and hepatic transcripts of TNF- α and NF- κ B of broiler challenged with E. coli O78

Sabreen Ezzat Fadl (✉ nourmallak@yahoo.com)

Animal Health Research Institute <https://orcid.org/0000-0001-5541-6159>

Ghada Ahmed El-Gammal

Animal Health Research Institute, Giza, Egypt

Osama Atia Sakr

Animal Health Research Institute, Giza, Egypt

Abdelbary Mohammed Prince

Cairo University

Aly A.B.S. Salah

Animal Health Research Institute, Giza, Egypt

Ayman Ali Atia

Animal Health Research Institute, Giza, Egypt

Abdelhaleem Mohamed Hegazy

Animal Health Research Institute, Giza, Egypt

Research article

Keywords: Antioxidant, Gene Expression, Growth Performance, Histopathology, Immunostimulant, Agrimos

Posted Date: December 19th, 2019

DOI: <https://doi.org/10.21203/rs.2.19307/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on June 19th, 2020. See the published version at <https://doi.org/10.1186/s12917-020-02423-2>.

Abstract

Background: Using probiotics have become popular. They are considered an alternative to Antibiotic Growth Promoters (AGP). Probiotics are supplemented into animal feed for improving growth performance along with preventing and controlling enteric pathogens. The aim of this work was to study the impact of dietary supplementation of Mannan-oligosaccharide and β -Glucan (Agrimos®) on broiler challenged with *Escherichia coli* O₇₈ (*E. coli* O₇₈ - marked with an antibiotic (320 μ g ciprofloxacin/ml broth) on growth performance, serum biochemistry, immune organs-histopathology, *E. coli* colonization, and hepatic transcripts of Tumor necrosis factor-alpha (TNF- α) and Nuclear factor-kappa B (NF- κ B).

Methods: A total of 125 one-day-old chicks were used for conducting the experiment. Five chicks were slaughtered for measuring the initial weight of lymphoid tissue; the remaining chicks (120) were allotted into four groups according to Mannan-oligosaccharide and β -Glucan supplementation and *E. coli* infection. The data were analyzed using SPSS version 16.

Results: Results indicated significant alteration of growth performance, serum biochemistry, selected liver gene expression with pathological lesions, especially in lymphoid organs due to *E. coli* infection, which were mitigated by Mannan-oligosaccharide and β -Glucan supplementation.

Conclusion: It could be concluded Mannan-oligosaccharide and β -Glucan supplementation in broiler's diet improved the immune response of broilers and mitigated pathological lesion resulted from *E. coli* infection.

Background

Prebiotics has been shown to alter the immune system, modify gastrointestinal microflora, and reduce invasion of the pathogen such as Salmonella spp. and *E. coli* [1]. The major function of prebiotics is to activate the metabolism of some groups of beneficial bacteria in the intestinal tract and/or stimulate their growth. Pelicano et al. [2], Spring et al. [3], and Xu et al. [4] have shown that prebiotics in addition to broilers diet results into the improvement of the gut microflora and so the improvement of growth. The gut microflora composition plays an important role in digestion; which can be performed in a positive, negative or neutral manner. Gastrointestinal microflora modifications reduce attachment of the pathogen and may have a beneficial effect on the nutrients digestibility [5]. Administration of Agrimos (Mannan-oligosaccharides (MOS) and β -Glucans), which obtained from a specific strain of *Saccharomyces cerevisiae* cell wall was found to improve the productive performance and immune functions in broiler chickens [6].

Hence, this study was conducted to examine agrimos (MOS and β -Glucans) effect on growth, immunity, serum biochemistry, histopathology, selected liver gene expressions and colonization of *E. coli* in broilers.

Results

Clinical Signs and Postmortem Lesions of *E. coli* Infection: Experimental infection with *E. coli* revealed suggestive clinical signs and PM lesions after 48 hr. post infection in the form of depression with diarrhea whitish in color.

PM lesions revealed liver enlarged and congested and distended gallbladder and cecum. Such changes were less prominent in the agrimos-infected group (see Table 3).

Growth Performance: The feed supplemented with agrimos in the third and fourth groups significantly ($P \leq 0.05$) increased the BW and total gain along the whole experiment and improved FCR compared to the other groups (see Table 4). On the other hand, the above mentioned parameters were decreased significantly ($P \leq 0.05$) in the control-infected group as compared to the other groups.

Serum Parameters Related to Liver Function: In Table (5) there were a significant ($P \leq 0.05$) increase in the activities of AST and ALT enzymes in the group of broiler chickens infected with *E. coli* compared to the control non-infected group. However, serum total protein and albumin were significantly ($P \leq 0.05$) decreased. Agrimos supplementation did not affect serum ALT, AST, and albumin but the serum total protein and globulin affected significantly ($P \leq 0.05$) when compared with the control non-infected group.

Serum Antioxidant Enzymes Activity: Serum MDA level was significantly ($P \leq 0.05$) increased in broiler chickens infected with *E. coli* (see Table 6). However, serum CAT and SOD activities were significantly ($P \leq 0.05$) reduced compared to the other groups. Agrimos supplementation significantly ($P \leq 0.05$) decreased serum MDA but significantly, ($P \leq 0.05$) increased serum CAT and SOD when compared to the control-infected group.

Immune Response against Newcastle Disease (ND): Immune response to vaccination of ND, which evaluated by HI titer revealed a difference in log₂ of GM. There was a significant difference in HI titer in the second week after infection between each of the *E. coli* infected group and the control negative group, and the agrimos infected group. In addition, the agrimos group significantly differs about the other three groups.

After vaccination in the second and third week, this pattern was observed where no difference was recorded between each of the *E. coli* infected group, the control non-infected, and the agrimos infected group. Besides, the agrimos group significantly differs about the other three groups (see Table 7).

Colonization of *E. coli*: The colonization of *E. coli* in the control infected group in different organs showed several rates of 67%, 44%, 22%, 44%, and 53% in the respiratory tract, liver, gallbladder, spleen, and fecal swab respectively (see Table 8). However, in the agrimos infected group, the colonization of *E. coli* in different organs showed reduced rates of 22%, 33%, 22%, 11%, and 33% in the respiratory tract, liver, gallbladder, spleen, and fecal swab respectively.

The Weight of Immune Organs: At zero, 15, and 35 days of age, the weight of the immune organ versus body weight was evaluated (see Table 9 and Fig. 1). At days 15 and 35, the results of thymus weight showed a significant increase ($P \leq 0.05$) in the agrimos non-infected and infected groups in comparison with the

other groups. However, there was a significant decrease ($P \leq 0.05$) in the control infected group compared to the control non-infected group.

The results of spleen at day 15 showed that there was a significant increase ($P \leq 0.05$) in the agrimos infected group compared to the other groups. Furthermore, there was a significant increase ($P \leq 0.05$) in the agrimos non-infected group compared to the control non-infected and infected groups. However, at day 35, the agrimos non-infected group witnessed a significant increase in the spleen's weight ($P \leq 0.05$) compared to the other groups. As well, the control infected group had a significant increase ($P \leq 0.05$) compared to the control non-infected group at day 15 but a significant decrease ($P \leq 0.05$) at Day 35.

The bursa's weight was significantly increased ($P \leq 0.05$) in the agrimos non-infected group compared to the other groups but it was significantly decreased ($P \leq 0.05$) in the control infected group compared to the other groups at days 15 and 35.

The thymus' and bursa's weight in the control non-infected group were significantly ($P \leq 0.05$) decreased but the spleen's weight was significantly ($P \leq 0.05$) increased. The control-infected group showed a significant ($P \leq 0.05$) increase in the weights of thymus, spleen, and bursa at day 15, and then showed a significant ($P \leq 0.05$) decrease at day 35. The agrimos non-infected and infected groups showed a significant ($P \leq 0.05$) increase in the weights of spleen and bursa at days 15 and 35 but showed a significant ($P \leq 0.05$) increase in the weight of thymus at day 15 only which returned to its normal size at day 35.

The Pathological Findings of Broiler Chickens: The histopathological examination of the spleen, bursa, thymus, liver, and duodenum on the chicks of the control non-infected group (at days 15 and 35 after infection) showed no obvious histopathological alterations (see Fig. 2a and 2b). Concerning the control infected group (at day 15), the bursa showed variable degrees of epithelial hyperplasia, degeneration, and ulceration in some cases. These changes were associated with subcortical fibrous tissues proliferation in some cases (see Fig. 2c, 2d and 2e). The thymus of this group showed a narrowing in the cortical width and an increase in the medulla in most cases. Some other cases had the appearance of clear areas or holes that contained small dark nuclei (a defining characteristic of the apoptosis in the lymphoid organ) (see Fig. 2f). No microscopical changes were detected on the spleen of this group. Regarding the control infected group (at day 35), similar changes were seen at day 15 but more holes were detected in the thymus. Also, the focal area of round cells aggregation was seen in the liver of this group (see Fig. 3a). The duodenum of this group showed hyperplasia on the epithelial cells lining the intestinal villi accompanied with vaculation (see Fig. 3b)

The histopathological examination of the bursa of Fabricius in most cases in the agrimos non-infected group at day 15 (4 out of 5) showed normal atypical fold (plica) (see Fig. 3c). Some of these cases showed highly dilated germinal center and narrow cortex, associated with a mild degree of epithelial lobulation. One case showed follicular lymphocytic depletion. Similar histopathological changes were seen on the bursa of Fabricius in the agrimos non-infected group at day 35. One case showed pores within the cortex of many plicae (see Fig. 3d), and this was associated with interfollicular edema. Regarding the thymus gland in both times (days 15 and 35), no obvious histopathological alterations were seen on the thymic lobe except in few cases (2 out of 10) which showed thrombus formation within the thymic vasculature (see Fig. 3e). Concerning the caecal tonsil of the agrimos non-infected group at days 15 and 35, just epithelial sloughing was seen in some cases (see Fig. 3f), and no obvious histopathological alterations were viewed on the lymphoid nodules (lymphoid aggregation) (see Fig. 4a). Two cases (out of 10) showed necrotic cyst (see Fig. 4b). No obvious microscopical changes were seen on the spleen of this group (see Fig. 4c). The microscopical examination of the liver of this group showed mild focal areas of round cells aggregation (see Fig. 4d) in two cases (out of 10).

The histopathological examination of the bursa of Fabricius in most cases in the agrimos infected group at days 15 and 35 after infection (2 out of 4) showed normal morphological appearances but 2 cases showed interfollicular edema (see Fig. 4e) and epithelial hyperplasia and folding associated with mucous (goblet) cell activation. Regarding the thymus gland in both times (days 15 and 35), no obvious histopathological alterations were seen on the thymic lobe except in few cases (1 out of 4) which showed focal areas of necrosis and hyalinization (see Fig. 4f). No obvious microscopical changes were seen on the spleen of this group (see Table 10 and 11).

Gene Expression Related to *E. coli* Infection and Agrimos Supplementation: The results of Q-PCR revealed that the expression levels of NF and TNF genes of liver tissues in the group infected by *E. coli* were significantly ($P \geq 0.05$) increased (Fig 5) compared to the negative control group.

In the agrimos non-infected group, the gene expression of NF and TNF in liver tissues did not show significant differences in comparison with the control non-infected group. The levels of such genes were similar to those of the control non-infected group.

It was found that the NF and TNF expression levels were significantly ($P \geq 0.05$) reduced due to agrimos supplementation when compared to those of the infected group without supplementation. This shows the protective effect of agrimos supplementation on the expression of the above-mentioned genes of the infected broiler chicks.

Discussion

In the present study, the birds infected with *E. coli* showed varied clinical signs according to their age, an organ involved, and concurrent disease conditions. Clinical signs in the *E. coli* infected groups (even in the treated one) similar to results of Remus et al. [7] who recorded severe liver damage in cases of *E. coli* infection. However, dietary supplementation of agrimos® decreased the clinical signs and PM lesions. This result is consistent with Sohail et al. [8]. On the other hand, results of the growth performance in the C+ group showed significant retardation, which may be attributable to the *E. coli* infection and are compatible with that of Manafi et al. [9]. Meanwhile, agrimos® dietary supplementation significantly improved growth performance which may be ascribed to the increased feed intake due to the increased digestibility. Jahanian and Ashnagar [10] reported that digestibility coefficients of dry matter and crude protein were improved because of using dietary MOS. These results in harmony with results of El-Far et al. [11] and Mousa et al. [12] in broilers and Japanese quail respectively.

Results of biochemical parameters showed significant increase in serum AST and ALT activities and globulin concentration meanwhile serum total protein and albumin were significantly decreased in the C+ group. These results agree with Manafi et al. [9] who reported the same results when birds challenged with *E. coli*. However dietary supplementation of agrimos® not affect serum AST and ALT activities and albumin concentration but significantly increased total protein and globulin of the serum. The above results are consonant with those of Mousa et al. [12] meanwhile Yalçinkaya et al. [13] found a significant decreased AST and ALT activities but Rokade et al. [14] found significant increased AST and ALT activities and protein concentration of the serum in broilers dietary supplemented with MOS. On the other hand, results of antioxidant enzymes and MDA in harmony with results of Zhou et al. [15] who found significant decrease of antioxidant enzymes in birds infected with necrotic enteritis. On the other hand, Abudabos et al. [16] reported that *Salmonella typhimurium* had no impact on total antioxidant capacity or oxidative stress in challenged broilers. However, agrimos® supplementation significantly decreased serum MDA and increased CAT. This result is compatible with that of El-Kader et al. [17] in broilers.

The reduction of HI titer in the *E. coli* infected groups might be attributed to the stress of infection and diarrhea, which might change the acid-base balance [18]. In addition, the increase in HI titer due to the nutrition of agrimos® appears to be insignificant. This might be attributed to the fact that agrimos® tend to initiate non-specific immunity rather than humoral immunity [19]. The dietary MOS stimulates humoral immune responses against infectious bursal disease (IBDV) and Newcastle disease (NDV) vaccine viruses [20]. Regarding results of *E. coli* colonization, the higher colonization rates of *E. coli* in different organs of C+ group were in the respiratory tract, fecal swab, liver, and spleen where avian pathogenic *E. coli* cause severe respiratory and systemic diseases in poultry [21]. This result is in harmony with that of Manafi et al. [9]. However, this colonization was decreased by agrimos® dietary supplementation which was supported by results of Sohail et al. [8].

Cloacae bursa plays a crucial role in the poultry immune system. In addition, the bursal weight reflects the anatomical response to immune status in broilers [9]. In the present investigation, there were significant decreases in thymus and bursa's weights versus body weight in the C+ group, which was supported by Gottardo et al. [22]. However, these results are incompatible with those obtained by Manafi et al. [9]. Also, the spleen's weight versus body weight in the C+ group significantly increased at day 15 then decreased at day 35. These results might be attributable to the infection, which was supported by results of Huff et al. [23] in turkeys.

On the other hand, there were significant increases in thymus', spleen, and bursa's weights versus body weight in the agrimos® groups. These results are in harmony with that of Sohail et al. [8].

Malfunctions of lymphoid organs reduce resistance to bacterial, viral, parasitic and fungal infections. As a result, birds become more susceptible to other infections, leading to high economic losses due to morbidity and mortality [24]. Therefore, the rational assessment of the immunosuppression in the poultry warrants rapid and accurate diagnostic approaches. Thus, the present study focused on the pathology of lymphoid organs.

The pathological changes in the C+ group were apoptosis in the lymphoid organ and focal area of round cells aggregation in the liver. These results are in harmony with that of Kumar et al. [25] meanwhile spleen revealed depletion of lymphocytes, areas of congestion and necrosis of lymphocytes from the white pulp, bursa of Fabricius showed lymphoid depletion in follicles, and thymus showed some lesions like medullary congestion and mild cortical depletion [24]. Remus et al. [7] reported *E. coli* caused severe liver and epithelial damage. However, agrimos® supplementation to broilers revealed no pathological changes. This result is compatible with that of Awaad et al. [6]. As well, in the infected group, the supplementation of agrimos® mitigated the pathological changes caused by infection. This result might be due to the composition of agrimos® which contains β -Glucans and MOS.

Pro-inflammatory cytokines such as nuclear factor (NF) and tumor necrosis factor (TNF) produced during the acute phase response can induce the production of several acute-phase proteins (APPs) by hepatocytes [26]. In the present study, it was found that agrimos® had no clear effect on liver NF or TNF expression levels of the uninfected broiler chicks. However, the reduction in liver enzyme activities and the down-regulation of NF and TNF gene expressions indicate the protective effects and anti-inflammatory properties of agrimos® in broiler chicks with *E. coli* infection. These results are in consistent with that obtained by Markazi [27] in *Salmonella* challenge broilers.

Most of the previous research has been performed on healthy chicks without infection conditions or has lacked to clear some gene expressions at the site of infection. *E. coli* infection is considered a common disease in poultry production, which can induce different oxidative and inflammatory stress. Thus, in the present study, *E. coli*-inoculated chickens were used to investigate the effects of agrimos® on healthy chickens and under such conditions.

Together with TNF or NF data, agrimos® can apparently regulate the cytokine responses of broiler chicks with *E. coli* infection and its immunomodulatory effects seem to be protective by repressing the ongoing inflammation. In brief, the decreased TNF and NF expressions of liver tissues in the agrimos® fed broiler chicks with *E. coli* infection might be an indication of slowdown inflammation and gradual restoration of the infected broiler chicks' health status.

Socio-Economic Importance

Veterinary medicine plays a crucial role in public health. The scope of veterinary public health includes food hygiene practices development and supervision in addition to zoonotic disease control and eradication [28]. In addition, antibiotic usage in animals plays a major role in the emerging public health crisis of antibiotic resistance [29]. Thus, nowadays, using probiotics or prebiotics have become popular. They are considered an alternative to Antibiotic Growth Promoters (AGP). Probiotics are supplemented into animal feed for improving growth performance along with preventing and controlling enteric pathogens [30].

Attention should be given to the enteric pathogens in the poultry field due to their economic effects (death, loss of weight, and poor conversion ratio). Besides, the spread of *E. coli* and the high usage of antibiotics in the poultry industry can result in bacterial resistance to such antibiotics. Instead, it is recommended to use prebiotics.

Conclusion

In the present study, agrimos supplementation increased growth performance. In case of infection with *E. coli*, agrimos improved the non-specific immunity which in turn led to an improvement in the specific immunity against ND which was reported in the discussion above. Consequently, there is no need to use antibiotics.

Methods

Agrimos: The combination of Manno-oligosaccharides (MOS) and Beta-Glucans, which extracted from *Saccharomyces cerevisiae* with a specific dedicated manufacturing process produced agrimos, which modulates the non-specific immune response and favors the development of the specific response. It was made in Lallemand Company (a Canadian company) and imported by Egavet (Egyptian Group for Trade and veterinary Services) Company, Egypt.

Experimental Design, Feeding Program, and Management: 125 mixed-sex chicks (age one-day) were used (Cobb-505 broiler strain). The experiment was done at the Animal Health Research Institute, Kafrelsheikh branch. The chicks obtained from a private farm at Kafrelsheikh governorate. At the beginning of the experiment, five chicks were weighed and slaughtered. Their bursa, spleen, and thymus gland were also weighed. The remaining broiler chicks (120) were kept in a clean room furnished with sawdust with good ventilation and supplemented with ad libitum feed and water under good hygienic management and sanitation with light/dark 23/1 hrs./day. Chicks (average body weight = 40-50 g/chick) after 3 days acclimatization period were equally allotted into (ranking method) 4 groups each group contains 3 replicates (10 chicks/replicate). The negative control group (first group), was fed on a diet [31] without any additives (basal) or infection (Table 1). The positive control (second group), was fed on a diet without any additives but infected with *E. coli*. The third group was fed on agrimos (2kg/ton) supplemented diet but without infection. The fourth group was fed on agrimos (2kg/ton) supplemented diet and infected. All chicks groups were vaccinated against ND and IBD at 7 & 12 days respectively. The chemical analysis of feed was determined according to **AOAC** [32]. Individual bird body weight was recorded at the beginning of the experiment. All performance parameters (body weight (BW), weight gain (WG), and feed conversion ratio (FCR)) were calculated according to **Vohra and Roudybush** [33], **Castell and Tiews** [34], and **Tacon** [35] respectively. The feed intake for each pen was recorded weekly with the observation of chicks for symptoms, postmortem (PM), and mortality along the period of the experiment.

The sacrificed birds from the experimental groups were used for sample collection. The remaining live birds were euthanized in strong bags by CO₂ suffocation. all dead birds and slaughtered ones, as well as remnants of samples and bedding material, were buried in the strict hygienically controlled properly constructed burial pit.

Experimental Infection: At the beginning of the experiment, all chicks in the two experimentally infected groups were infected with *E. coli* O₇₈ via the oral route (1 ml containing 3x10⁸). This strain was supplied from Animal Health Research Institute, Kafrelsheikh Regional Lab, Poultry and Rabbit Disease Department). *E. coli* O₇₈ was isolated from poultry and contained the virulence genes, which are a defining characteristic of Avian Pathogenic *E. coli* O₇₈ (APEC). *E. coli* O₇₈ was marked with an antibiotic (320 µg ciprofloxacin/ml broth) and prepared according to **McFarland** [36]. All groups (30 birds/group) were vaccinated against ND and IBD at Days 7 and 12 respectively.

Sample Collection and Measurements of the Serum Parameters: At day 35, samples of the blood were taken from 5 birds/group after slaughtering (the birds were euthanized by intraperitoneal injection of sodium pentobarbital (50 mg/kg) to minimize suffering during slaughtering for blood collection). After coagulation of the blood, serum was separated (centrifugation at 3000 rpm for 15 min.) and kept in a freezer (-20° C) until analysis. Activities of the serum catalase (CAT), superoxide dismutase (SOD), alanine transaminase (ALT), and aspartate aminotransferase (AST) and concentration of lipid peroxidation (MDA), total protein, and albumin were measured by using commercial kits obtained from BIODIAGNOSTIC Company.

Immune Response against Newcastle Disease (ND): Hemagglutination Inhibition (HI) titer was performed against ND [37]. The Newcastle disease virus and positive control serum were kindly provided by the Animal Health Research Institute, Kafrelsheikh.

Colonization of *E. coli*: Shedding of the *E. coli* in the fecal matter and its colonization in organs was recorded and confirmed by the re-isolation of *E. coli* from infected birds.

The Weight of Immune Organs: Five chicks at zero day of age and five chicks from each group were weighted then slaughtered at 15 and 35 days of age. Each organ including thymus, spleen, and bursa of Fabricius was isolated and weighed. The weight of the targeted immune organs was calculated as described by **Keil et al.**[38] following the equation below:

$$\text{Immune Organ Weight/Body Weight} = \text{Organ Weight (gm)}/\text{Body Weight (gm)} \times 100$$

Necropsy and Histopathology: Sections of liver, intestine, bursa, thymus gland and spleen were taken after necropsy and fixed immediately in 10% buffered formalin and processed for histopathological evaluation, using routine paraffin embedding section. Sections of 3µm thickness were cut and stained using H&E according to **Bancroft and Gamble** [39].

Q-PCR (Real-Time Polymerase Chain Reaction): Collection of the Sample: At Day 35, samples of the liver were collected on liquid nitrogen from 4 birds/group into clean Eppendorf tubes and stored at -80 °C until use.

Q-PCR (Real-Time Polymerase Chain Reaction): The liver samples were collected (35 days) on liquid nitrogen from 4 birds/group into clean Eppendorf tubes and stored at -80 °C until use.

Total RNA from the liver samples was extracted using easy RED total RNA extraction kits (iNtRON Biotechnology, Inc.) according to the manufacturer's instructions. The RNA integrity was verified by using agarose gel electrophoresis. The first-strand cDNA was synthesized with total RNA using Intron-Power cDNA synthesis kit (Cat. no. 25011).

The specific primers were used to amplify the selected genes of the chicken (*Gallus gallus*) with GAPDH as housekeeping (internal standard) gene-primer sequence (Table 2). The qRT-PCR assay was carried out using a Stratagene MX300P Q-PCR system (Agilent Technologies), using Real MODTM Green FAST qPCR master mix (S) following the manufacturer's recommendations. MxPro QPCR Software was used for data collection.

The relative gene expression levels were evaluated using the $2^{-\Delta\Delta Ct}$ method [40].

Statistical Analysis: The obtained results were statistically analyzed by one-way ANOVA using **SPSS** version 16.

Declarations

Ethics approval and consent to participate

The current study was approved by the Ethical Committee for live birds sampling at the Animal Health Research Institute, Egypt (License No. AHRI 42102017), according to local Egyptian laws.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This research did not receive any specific financial support from funding agencies in public, commercial, or non-profit sectors.

Authors' Contributions

All authors contributed equally to this work whereas they designed, conducted the experiment and wrote the manuscript.

SF and AS measured serum biochemistry

GE followed up isolation of *E. coli*, colonization and reisolation of *E. coli*

OS designed diet formulation and measured growth performance parameters.

AA examined pathological changes

AP measured Q-PCR for the collected samples

HAM followed up immune response against Newcastle disease, clinical signs and mortalities

SF made interpretation of the results

Acknowledgments:

We would like to thanks Prof. Rawhya Emran, Dep. of Pathology, Molecular Pathology Unit, Animal Health Research Institute for her valuable support.

Abbreviations

AGP: Antibiotic Growth Promoters; ALT: alanine transaminase; APPs: acute-phase proteins; AST: aspartate aminotransferase; BW: body weight; CAT: catalase; DNA: deoxyribonucleic acid; *E. coli*: *Escherichia coli*; FCR: feed conversion ratio; HI: Hemagglutination Inhibition; IBDV: infectious bursal disease; Q-PCR: Real-Time Polymerase Chain Reaction; MDA: lipid peroxidation; MOS: mannan oligosaccharide; ND: Newcastle Disease; NDV: Newcastle disease virus; NF: nuclear factor; PM: postmortem; RNA: ribonucleic acid; SOD: superoxide dismutase; TNF: tumor necrosis factor

References

1. Cummings JH, Macfarlane GT. Gastrointestinal effects of prebiotics. *Br J Nutr.* 87(Suppl.2), 2002;145-151 (Abstr.). DOI: <https://doi.org/10.1079/BJN/2002530>
2. Pelicano ER, Souza PA, Souza HB, Figueiredo DF, Boiago MM, Carvalho SR, Bordon VF. Intestinal mucosa development in broiler chickens fed natural growth promoters. *Braz J Poultry Sci.* 2005 Dec;7(4):221-9. <http://dx.doi.org/10.1590/S1516-635X2005000400005>
3. Spring P, Wenk C, Dawson KA, Newman KE. The effects of dietary mannaoligosaccharides on cecal parameters and the concentrations of enteric bacteria in the ceca of salmonella-challenged broiler chicks. *Poult sci.* 2000 Feb 1;79(2):205-11. <https://doi.org/10.1093/ps/79.2.205>

4. Xu ZR, Hu CH, Xia MS, Zhan XA, Wang MQ. Effects of dietary fructooligosaccharide on digestive enzyme activities, intestinal microflora and morphology of male broilers. *Poult sci.* 2003 Jun 1;82(6):1030-6. <https://doi.org/10.1093/ps/82.6.1030>
5. Hajati H, Rezaei M. The application of prebiotics in poultry production. *Int J Poult Sci.* 2010;9(3):298-304. DOI: 10.3923/ijps.2010.298.304 · Source: DOAJ
6. Awaad MH, Atta AM, El-Ghany WA, Elmenawey M, Ahmed K, Hassan AA, Nada AA, Abdelaleem GA. Effect of a specific combination of mannan-oligosaccharides and β -glucans extracted from yeast cell wall on the health status and growth performance of ochratoxicated broiler chicken. *J Am Sci.* 2011;7(3):82-96. Available in research gate net.
7. Remus A, Hauschild L, Andretta I, Kipper M, Lehnen CR, Sakomura NK. A meta-analysis of the feed intake and growth performance of broiler chickens challenged by bacteria. *Poult sci.* 2014 Apr 28;93(5):1149-58. <https://doi.org/10.3382/ps.2013-03540>
8. Sohail MU, Ijaz A, Younus M, Shabbir MZ, Kamran Z, Ahmad S, Anwar H, Yousaf MS, Ashraf K, Shahzad AH, Rehman H. Effect of supplementation of mannan oligosaccharide and probiotic on growth performance, relative weights of viscera, and population of selected intestinal bacteria in cyclic heat-stressed broilers. *J Appl Poult Res.* 2013 Oct 1;22(3):485-91. <https://doi.org/10.3382/japr.2012-00682>
9. Manafi M, Hedayati M, Khalaji S, Kamely M. Assessment of a natural, non-antibiotic blend on performance, blood biochemistry, intestinal microflora, and morphology of broilers challenged with *Escherichia coli*. *Rev Bras Zootec.* 2016 Dec;45(12):745-54. <http://dx.doi.org/10.1590/s1806-92902016001200003>
10. Jahanian R, Ashnagar M. Effect of dietary supplementation of mannan-oligosaccharides on performance, blood metabolites, ileal nutrient digestibility, and gut microflora in *Escherichia coli*-challenged laying hens. *Poult sci.* 2015 Aug 22;94(9):2165-72. <https://doi.org/10.3382/ps/pev180>
11. El-Far AH, Ahmed HA, Shaheen HM. Dietary supplementation of phoenix dactylifera seeds enhances performance, immune response, and antioxidant status in broilers. *Oxid Med Cell Longev.* 2016;2016. <http://dx.doi.org/10.1155/2016/5454963>
12. Mousa SM, Soliman MM, Bahakaim AS. Effect of mannan oligosaccharides and β -glucans on productive performance and some physiological and immunological parameters of growing Japanese quail chicks. *Egypt Poult Sci J.* 2014 Apr 1;34(2). © 2019 EBSCO Industries, Inc. All rights reserved
13. Yalçinkaya İ, Güngör T, Başalan M, Erdem E. Mannan oligosaccharides (MOS) from *Saccharomyces cerevisiae* in broilers: Effects on performance and blood biochemistry. *Turk J Vet Anim Sci.* 2008 Feb 25;32(1):43-8. This research was supported by the Scientific Research Fund of the University of Kırıkkale, Project No: 03/09-03-02
14. Rokade JJ, Kagate M, Bhanja SK, Mehra M, Goel A, Vispute M, Mandal AB. Effect of mannan-oligosaccharides (MOS) supplementation on performance, immunity and HSP70 gene expression in broiler chicken during hot-dry summer. *Indian J Anim Res.* 2018 Jun 1;52(6):868-74. DOI:10.18805/ijar.B-3291
15. Zhou M, Zeng D, Ni X, Tu T, Yin Z, Pan K, Jing B. Effects of *Bacillus licheniformis* on the growth performance and expression of lipid metabolism-related genes in broiler chickens challenged with *Clostridium perfringens*-induced necrotic enteritis. *Lipids Health Dis.* 2016 Dec;15(1):48. <https://doi.org/10.1186/s12944-016-0219-2>
16. Abudabos AM, Alyemni AH, Zakaria HA. Effect of two strains of probiotics on the antioxidant capacity, oxidative stress, and immune responses of *Salmonella*-Challenged broilers. *Braz J Poult Sci.* 2016 Mar;18(1):175-80. <http://dx.doi.org/10.1590/18069061-2015-0052>
17. El-Kader HA, Alam SS, Nafeaa AA, Mahrous KF. Influence of dietary supplementation of mannanoligosaccharide (bio-mos®) prebiotic on the genotoxic and antioxidant status in Japanese quail. *Int. J. pharm. Sci. Rev. Res.* 2014;27:289-95. Available online at www.globalresearchonline.net
18. Sil GC, Das PM, Islam MR, Rahman MM. Management and disease problems of cockrels in some farms of Mymensingh, Bangladesh. *Int J Poult Sci.* 2002;1(4):102-5. © Asian Network for Scientific Information 2002
19. Eddie Ip WK, Takahashi K, Alan Ezekowitz R, Stuart LM. Mannose-binding lectin and innate immunity. *Immunol Rev.* 2009 Jul;230(1):9-21. <https://doi.org/10.1111/j.1600-065X.2009.00789.x>
20. Oliveira MC, Figueiredo-Lima DF, Faria Filho DE, Marques RH, Moraes VM. Effect of mannanoligosaccharides and/or enzymes on antibody titers against infectious bursal and Newcastle disease viruses. *Arq Bras Med Vet Zoo.* 2009 Feb;61(1):6-11. <http://dx.doi.org/10.1590/S0102-09352009000100002>
21. Sadeyen JR, Kaiser P, Stevens MP, Dziva F. Analysis of immune responses induced by avian pathogenic *Escherichia coli* infection in turkeys and their association with resistance to homologous re-challenge. *Vet Res.* 2014 Dec;45(1):19. doi:10.1186/1297-9716-45-19
22. Gottardo ET, Burin Junior ÁM, Lemke BV, Silva AM, Pasa B, Cassiano L, Muller Fernandes JI. Immune response in *Eimeria* sp. and *E. coli* challenged broilers supplemented with amino acids. *Austral J Vet Sci.* 2017 Sep;49(3):175-84. <http://dx.doi.org/10.4067/S0719-81322017000300175>
23. Huff GR, Huff WE, Balog JM, Rath NC. The effects of dexamethasone immunosuppression on turkey osteomyelitis complex in an experimental *Escherichia coli* respiratory infection. *Poult Sci.* 1998 May 1;77(5):654-61. <https://doi.org/10.1093/ps/77.5.654>
24. Wani BM, Darzi MM, Masood Saleem Mir SA, Shakeel I. Pathological and Pharmacochemical Evaluation of Broiler Chicken Affected Naturally with Colibacillosis in Kashmir Valley. *Int J Pharmacol.* 2017 Jan 1;13(4):388-95. Doi: 10.3923/ijp.2017.388.395
25. Kumar A, Jindal N, Shukla CL, Asrani RK, Ledoux DR, Rottinghaus GE. Pathological changes in broiler chickens fed ochratoxin A and inoculated with *Escherichia coli*. *Avian Pathol.* 2004 Aug 1;33(4):413-7. <https://doi.org/10.1080/03079450410001724021>
26. Petersen HH, Nielsen JP, Heegaard PM. Application of acute phase protein measurements in veterinary clinical chemistry. *Vet Res.* 2004 Mar 1;35(2):163-87. <https://doi.org/10.1051/vetres:2004002>
27. Markazi A. *Effects of Whole Yeast Cell Product Supplementation in Chickens Post-coccidial and Post-Salmonella Challenge* (Doctoral dissertation, The Ohio State University). http://rave.ohiolink.edu/etdc/view?acc_num=osu1437497725
28. Lathers CM. Role of veterinary medicine in public health: antibiotic use in food animals and humans and the effect on evolution of antibacterial resistance. *J Clin Pharmacol.* 2001 Jun;41(6):595-9. <https://doi.org/10.1177/00912700122010474>
29. Landers TF, Cohen B, Wittum TE, Larson EL. A review of antibiotic use in food animals: perspective, policy, and potential. *Public Health Rep.* 2012 Jan;127(1):4-22. <https://doi.org/10.1177/003335491212700103>

30. Bajagai YS, Klieve AV, Dart PJ, Bryden WL. Probiotics in animal nutrition: production, impact and regulation. FAO; 2016. Available in Google scholar.
31. Nutrient requirements of poul., 9th ed. National Academy of Sci., Washington, DC. 1994. Available in Google scholar.
32. Official methods of analysis. 13th Ed., w. Horwitz. w, (Editor), Academic Press, Washington, D.C., USA, no. 925.09. 2000. Available in Google scholar.
33. Vohra P, Roudybush T. The effect of various levels of dietary protein on the growth and egg production of *Coturnix coturnix japonica*. *Poult Sci.* 1971 Jul 1;50(4):1081-4. <https://doi.org/10.3382/ps.0501081>
34. Castell JD, Tiewes K. Report of the EIFAC. IUNS and ICES working group on the standardization of methodology in fish nutrition research. Hamburg, Federal Republic of Germany, EIFAC. Technology. 1980;36, 24. Available in Google scholar.
35. Tacon A. The nutrition and feeding of farmed fish and shrimp a training manual. V61. 1. The essential nutrients FAO. 1987;PP. 117-130. Available in Google scholar.
36. McFarland J. Nephelometer: an instrument for media used for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. *The Journal of the American Medical Association*, 1907;14, 1176-8. 1907;XLIX(14):1176-1178. doi:10.1001/jama.1907.25320140022001f
37. King DJ, Seal BS. (1998). Biological and molecular characterization of ND. virus field isolates with comparison to reference NDV strain and pathogenicity after chicken or embryo passage of selected isolates. *Avian Dis.* 1998;42, 507-516. DOI: 10.2307/1592677. <https://www.jstor.org/stable/1592677>
38. Keil DE, Mehlmann T, Butterworth L, Peden-Adams MM. Gestational exposure to perfluorooctane sulfonate suppresses immune function in B6C3F1 mice. *Toxicol Sci.* 2008 Feb 5;103(1):77-85. <https://doi.org/10.1093/toxsci/kfn015>
39. Bancroft JD, Gamble M, editors. Theory and practice of histological techniques. Elsevier health sciences; 2008. Available in Google scholar
40. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻ ΔΔCT method. *methods.* 2001 Dec 1;25(4):402-8. Available in Google scholar

Tables

Table 1: Experimental feeding program.

Physical composition	Basal diet (0 - 3) weeks	Basal diet (3 - 6) weeks
Yellow corn	55	61.35
Soybean meal 48%	32	30.6
Corn gluten 62%	4.42	0
Sunflower oil	4.4	4.4
Methionine	0.16	0.08
Dicalcium phosphate	1.85	1.4
Lime stone	1.25	1.3
Choline 60%	0.22	0.17
Common salt	0.4	0.4
Premix*	0.3	0.3
Chemical composition %		
ME Kcal/kg	3218.5	3229.7
Crude protein	23.02	20.03
Calcium	1	.9
Available phosphorus	0.45	0.35
Lysine	1.1	1
Methionine + cysteine	0.9	.72
Choline	1300mg/1kg	1000mg/1kg

* The used premix (Multivita Co.) composed of retinol 12000000 IU, cholecalciferol 2200000 IU, tocopherol 10000 mg, menadione 2000 mg, thiamin 1000 mg, riboflavin 5000 mg, pyridoxal 1500 mg, cobalamin 10 mg, Niacin 30000 mg, Biotin 50 mg, Folic acid 1000 mg, Pantothenic acid 10000 mg, Iron 30000 mg, Manganese 60000 mg, Copper 4000 mg, Zinc 50000 mg, Iodine 1000 mg, Cobalt 100 mg, Selenium 100 mg, calcium carbonate (CaCO₃) carrier to 3000g.

Table 2: Primers used for qPCR analysis

Primer	Sequence	Reference
TNF- α	for: 5'-GAGCTGTGGGGAGAACAAAAGGA-3' Rev: 5'-TTGGCCCTTGAAGAGGACCTG-3'	Tabatabaei et al. (2015)
NF- κ B	for: 5'-CAAGGCAGCAATAGACGAG-3' Rev: 5'-GTTGAGAGTTAGCAGTGAGGCA-3'	
GAPDH	For- 5'CCTCTCTGGCAAAGTCCAAG3' Rev- 5'CAACATCAAATGGGCAGATG3'	

TNF- α : Tumor necrosis factor-alpha, NF- κ B: Nuclear factor-kappa B, GAPDH: Glyceraldehyde-3 phosphate dehydrogenase.

Table 3 The mortality rate of broiler chicken infected with *E. coli* and fed on agrimos[®] at 35 days.

Groups	Control noninfected	Control infected	Agrimos [®] noninfected	Agrimos [®] infected
Parameter				
Total No.	30	30	30	30
Dead No.	1	2	0	1
Survival %	96.67	93.33	100	96.67
Mortality %	3.33	6.67	0	3.33

Table 4 Growth performance of broiler chicken infected with *E. coli* and fed on agrimos[®] at 35 days.

Groups	Control noninfected	Control infected	Agrimos [®] noninfected	Agrimos [®] infected
Parameters				
Initial body weight (g/chick)	45.33 \pm 0.88 ^a	45 \pm 1.00 ^a	45.67 \pm 0.88 ^a	45.67 \pm 1.20 ^a
Final body weight (g/bird)	1544.33 \pm 1.76 ^b	1358.67 \pm 2.03 ^c	1736.33 \pm 2.33 ^a	1547.00 \pm 1.73 ^b
Total Weight gain (g/bird)	1499 \pm 1.53 ^b	1313.67 \pm 2.19 ^c	1690.67 \pm 1.45 ^a	1501.33 \pm 0.67 ^b
FCR value	1.67 \pm 0.005 ^b	1.87 \pm 0.006 ^a	1.54 \pm 0.003 ^c	1.66 \pm 0.003 ^b

Values are expressed as mean \pm standard errors. Means in the same row (a-c) with different letters significantly differ at ($p \leq 0.05$).

Table 5

Serum liver function of broiler chicken infected with *E. coli* and fed on agrimos[®] at 35 days.

Groups	Control noninfected	Control infected	Agrimos [®] noninfected	Agrimos [®] infected
Parameters				
ALT (u/l)	19.33 \pm 0.88 ^b	26.33 \pm 0.88 ^a	18.67 \pm 0.88 ^b	20.67 \pm 0.67 ^b
AST (u/l)	53.33 \pm 1.76 ^b	92.67 \pm 0.88 ^a	51.33 \pm 0.88 ^b	52.67 \pm 1.45 ^b
Total protein (g/dl)	2.27 \pm 0.08 ^b	1.9 \pm 0.06 ^c	2.9 \pm 0.12 ^a	2.7 \pm 0.11 ^a
Albumin (g/dl)	1.20 \pm 0.02 ^a	0.47 \pm 0.05 ^b	1.24 \pm 0.03 ^a	1.21 \pm 0.01 ^a
Globulin (g/dl)	1.06 \pm 0.09 ^b	1.43 \pm 0.01 ^a	1.66 \pm 0.12 ^a	1.49 \pm 0.11 ^a

Values are expressed as mean \pm standard errors. Means in the same row (a-c) with different letters significantly differ at ($p \leq 0.05$).

Table 6

Serum antioxidant of broiler chicken infected with *E. coli* and fed on agrimos® at 35 days.

Parameters	Control noninfected	Control infected	Agrimos® noninfected	Agrimos® infected
MDA (n.mol/l)	0.65 ± 0.02 ^c	1.17 ± 0.08 ^a	0.65 ± 0.01 ^c	0.8 ± 0.03 ^b
CAT (u/ml)	12.7 ± 0.01 ^b	9.73 ± 0.15 ^c	15.36 ± 0.13 ^a	13.03 ± 0.09 ^b
SOD (u/ml)	12.18 ± 0.0 ^c	9.37 ± 0.0 ^d	17.2 ± 0.89 ^a	15.0 ± 0.0 ^b

Values are expressed as mean ± standard errors. Means in the same row (a-c) with different letters significantly differ at (p≤0.05).

Table 7 HI titer for ND at different periods

Groups	Control infected	Control noninfected	Agrimos® infected	Agrimos® noninfected
Age				
0 day	4.6±0.07 ^b	4.8±0.06 ^b	4.8±0.06 ^b	5.2±0.06 ^a
	C.V=11.6%	C.V=9.03%	CV=9.3%	CV=8.3%
1st weeks p. v	5.2±0.06 ^c	5.6±0.7 ^b	5.6±0.07 ^b	6.2±0.06 ^a
	C.V=8.6%	CV=9.5%	CV=9.5%	CV=7.2%
2nd weeks P. v	6±0.09 ^b	6.2±0.11 ^b	6.2±0.06 ^b	7±0.09 ^a
	C.V=11.5	CV=13.4	CV=6.9%	CV=10%
3rd weeks P. v	4.95±0.09 ^b	4.6±0.24 ^b	4.8±0.06 ^b	6±0.09 ^a
	C.V=14.5%	CV=11.6%	CV=9%	CV=11.7%
Average of log₂GM±SEM	5.19±0.3 ^b	5.3±0.37 ^b	5.35 ±0.34 ^b	6.1 ±0.37 ^a

GM: geometric mean. CV: coefficient of variation. p. v: post vaccination

No difference between item carries the same letter in the same row.

Table 8: Colonization of *E. coli* and rate of shedding as judged by intestinal colonization.

	R. T. swab			Liver			G. bladder			Spleen			Fecal swab		
	+	T	%	+	T	%	+	T	%	+	T	%	+	T	%
<i>E. coli</i>	6	9	67	4	9	44	2	9	22	4	9	44	8	15	53
<i>E. coli</i> + A*	2	9	22	3	9	33	2	9	22	1	9	11	5	15	33

R. T. swab= respiratory tract swab

* *E. coli* + A = *E. coli* + Agrimos®

Table 9: The weight of Immune organs of broilers at 35 days.

Organs	Treatment and age											
	C-			C+			A-			A+		
	Zero-day	At 15 days	At 35 days	Zero-day	At 15 days	At 35 days	Zero-day	At 15 days	At 35 days	Zero-day	At 15 days	At 35 days
T/B.W.	0.287	0.59	0.134	0.287	0.377	0.069	0.287	0.65	0.282	0.287	0.627	0.276
	±	±	±	±	±	±	±	±	±	±	±	±
S/B.W.	0.033 ^Y	0.012 ^{bX}	0.008 ^{BZ}	0.033 ^Y	0.013 ^{cX}	0.006 ^{CZ}	0.033 ^Y	0.006 ^{aX}	0.012 ^{AY}	0.033 ^Y	0.009 ^{aX}	0.011 ^{AY}
	0.036	0.046	0.132	0.036	0.094	0.056	0.036	0.11	0.229	0.036	0.127	0.159
	±	±	±	±	±	±	±	±	±	±	±	±
	0.007 ^Y	0.003 ^{dY}	0.011 ^{BX}	0.007 ^Z	0.004 ^{cX}	0.006 ^{CY}	0.007 ^Z	0.006 ^{bY}	0.009 ^{AX}	0.007 ^Z	0.003 ^{aY}	0.005 ^{BX}
B/B.W.	0.157	0.329	0.121	0.157	0.259	0.064	0.157	0.394	0.355	0.157	0.338	0.317
	±	±	±	±	±	±	±	±	±	±	±	±
	0.024 ^Y	0.003 ^{bX}	0.004 ^{CY}	0.024 ^Y	0.01 ^{cX}	0.006 ^{DZ}	0.024 ^Y	0.003 ^{aX}	0.014 ^{AX}	0.024 ^Y	0.017 ^{bX}	0.013 ^{BX}

T = thymus B. W. = body weight S = spleen B = bursa

Values are expressed as mean ± standard errors. Means in the same row (a-d), (A-C) and (X-Z) with different letters significantly differ at (p≤0.05).

le (10): Histopathological assessment of dietary agrimos[®] supplementation on *E. coli* challenged broiler chicken.

	Infected group						Agrimos non-infected group						Agrimos infected group					
	At 15 days			At 35 days			At 15 days			At 35 days			At 15 days			At 35 days		
	B	T	S	B	T	S	B	T	S	B	T	S	B	T	S	B	T	S
ular	-		Normal	-		Normal	Normal	Normal	Normal	Normal	Normal	Normal	-	Normal	Normal	-		Norr
ilic																		
ion																		
and	-			-									+			+		
al																		
al	-			-									-			+		
ages																		
al	-			-									-			-		
ve																		
tion																		
l	+			-									-			-		
and																		
n																		
l	+			+									+			+		
sia																		
l	+			+									-			+		
rtion																		
n in	-			-									-			-		
and																		
of																		
r	-			-	+								-			+	+	
or																		
r	-			-									-			+		
and																		
ytic	-			-									-			+		
l																		
nce	-	+		-	++								-			-		
s in																		
	-			-									-			-		
tion																		
s	-			-									-			-		
	-			-									-			-		
c	-	+		-									-			-		
xis																		
ation	-			-									-			-		
sis																		
lood																		

Sever ++ Moderate + Mild B= Bursa T=Thymus S=Spleen

Table (11): Incidence of pathological changes in organs at different periods

Parameter	Infected group				Agrimos non-infected group				Agrimos infected group															
	At 15 days		At 35 days		At 15 days		At 35 days		At 15 days		At 35 days													
	N	A	T	%	N	A	T	%	N	A	T	%												
Bursa	2	2	4	50	1	3	4	75	4	1	5	20	4	1	5	20	2	2	4	50	2	2	4	50
Thymus	4	0	4	0	2	2	4	50	3	2	5	40	4	1	5	20	3	1	4	25	3	1	4	25
Spleen	3	1	4	25	4	0	4	0	5	0	5	0	5	0	5	0	4	0	4	0	4	0	4	0

B = Bursa T=Thymus S=Spleen N=Normal A=Abnormal T=Total %= Ratio

Figures




Figure 1
 Thymus of one-day-old chicken (A1). Thymus of control non-infected group at 35 days (A2). Thymus of control infected group at 35 days (A3). Thymus of agrimos non-infected group at 35 days (A4). Thymus of agrimos infected group at 35 days (A5). Bursa of one-day-old chicken (B1). Bursa of control non-infected group at 35 days (B2). Bursa of control infected group at 35 days (B3). Bursa of agrimos non-infected group at 35 days (B4). Bursa of agrimos infected group at 35 days (B5). Spleen of one-day-old chicken (C 1). Spleen of control non-infected group at 35 days (C 2). Spleen of control infected group at 35 days (C 3). Spleen of agrimos non-infected group at 35 days (C 4). Spleen of agrimos infected group at 35 days (C 5).




Figure 2

a. showing normal bursa of Fabricius. Control negative one-day-old chick. b. showing normal thymus. Control negative one-day-old chick. c. Bursa of Fabricius showing variable degree of epithelial hyperplasia (arrow a) and ulceration (arrow b). Control infected group at 15 days post infection. d. Bursa of Fabricius showing variable degree of epithelial hyperplasia (arrow a) associated with degeneration and ulceration (arrow b). Control infected group at 15 days post infection. e. Bursa of Fabricius showing variable degree of epithelial hyperplasia associated with degeneration and sub cortical fibrous tissues proliferation. Control infected group at 15 days post infection. f. Thymus showing numerous holes on the cortex and medulla. Control infected group at 15 days post infection. H&E. a and b X 100 other figures X 200.




Figure 3

a. Liver showing focal area of round cells aggregation. Control infected group at 35 days post infection. b. Intestine (duodenum) showing hyperplasia of the epithelial lining the intestinal villi associated with vacuolation. Control infected group at 35 days. c. Bursa of Fabricius showing normal atypical fold. Agrimos® non-infected group at 15 days. d. Bursa of Fabricius showing pores within the cortex. Agrimos® non-infected group at 35 days. e. Thymus showing thrombus formation (arrow). Agrimos® non-infected group at 15-35 days. f. Caecal tonsil showing epithelial sloughing (arrow). Agrimos® non-infected group at 15-35 days. H&E. X 200 for all figures except c X 100.




Figure 4
 a. Caecal tonsil showing normal lymphoid aggregation (arrow). Agrimos® non-infected group at 15-35 days. b. Caecal tonsil showing necrotic cyst (arrow). Agrimos® non-infected group at 15-35 days. c. Spleen showing normal appearance. Agrimos® non-infected group at 15-35 days. d. Liver showing mild focal areas of round cells aggregation (arrow). Agrimos® non-infected group at 15-35 days. e. Bursa of Fabricius showing inter follicular edema. Agrimos® infected group at 15 days post infection. f. Thymus showing numerous cortical holes in cortex and medulla (these holes containing apoptotic bodies). Agrimos® infected group at 15 days post infection. H&E. X 200 for all figures except e X 100.




Figure 5
 Changes in the gene expression of NF (right) and TNF-α (left) among experimental groups. C (negative control), C+ (positive control), A (Agrimos® treated without infection), A+ (Agrimos® treated group with E. coli infection)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [NC3RsARRIVEGuidelines20131.docx](#)