

Genomics Analysis and Degradation Characteristics of Lignin by *Streptomyces Thermocarboxydus* Strain DF3-3

Fangyun Tan

Beijing University of Agriculture

Jun Cheng

Beijing University of Agriculture

Yu Zhang

Beijing University of Agriculture

Xingfu Jiang

Chinese Academy of Agricultural Sciences Institute of Plant Protection

Yueqiu Liu (✉ liuyueqiu@bua.edu.cn)

Beijing University of Agriculture <https://orcid.org/0000-0002-9152-4752>

Research

Keywords: *Streptomyces thermocarboxydus* strain DF3-3, alkali lignin, enzyme activity, genomics, metabolic pathways

Posted Date: November 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-996090/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Lignocellulose is an important raw material for biomass-to-energy conversion, and it exhibits a complex but inefficient degradation mechanism. Microbial degradation is promising due to its environmental adaptability and biochemical versatility, but the pathways used by microbes for lignin degradation have not been fully studied. Degradation intermediates and complex metabolic pathways require more study.

Results: A novel actinomycete DF3-3, with the potential for lignin degradation, was screened and isolated. After morphological and molecular identification, DF3-3 was determined to be *Streptomyces thermocarboxydus*. The degradation of alkali lignin reached 31% within 15 days. Manganese peroxidase and laccase demonstrated their greatest activity levels, 1821.66 UL⁻¹ and 1265.58 UL⁻¹, respectively, on the sixth day. The highest lignin peroxidase activity was 480.33 UL⁻¹ on the fourth day. A total of 19 lignin degradation intermediates were identified by gas chromatography-mass spectrometry (GC-MS), including 10 aromatic compounds. Genome sequencing and annotation identified 107 lignin-degrading enzyme-coding genes containing three core enzymatic systems for lignin depolymerization: laccases, peroxidases and manganese peroxidase. In total, 7 lignin metabolic pathways were predicted.

Conclusions: *Streptomyces thermocarboxydus* strain DF3-3 has good lignin degradation ability. Degradation products and genomics analyses of DF3-3 show that it has a relatively complete lignin degradation pathway, including the β -keto adipate pathway and peripheral reactions; gentisate pathway; anthranilate pathway; homogentisic pathway; and catabolic pathway for resorcinol. Two other pathways, the phenylacetate-CoA pathway and the 2,3-dihydroxyphenylpropionic acid pathway, are predicted based on genome data alone. This study provides the basis for future characterization of potential biotransformation enzyme systems for biomass energy conversion.

Background

Lignocellulosic biomass is an easily available, low cost and renewable alternative to fossil fuels. Currently, the process for pretreatment of lignocellulosic biofuel requires removal or delocalization of lignin, which might generate aromatic compounds that inhibit enzymatic hydrolysis and fermentation [1]. Due to its rich aromatic content, lignin is a valuable waste from the biomass industry [2]. Researchers worldwide are focusing on lignin and its components for conversion into value-added products. Lignin is a complex aromatic heteropolymer derived from the condensation of hydroxyphenylpropane monomers and comprises various ether bonds and carbon-carbon bonds [3]. It is mainly composed of three basic monomers: guaiacyl (G) units, syringyl (S) units and p-hydroxyphenyl (H) units [4, 5]. Microorganisms that degrade lignocellulose are widely distributed in nature, and known examples include bacteria, fungi and actinomycetes. Among them, fungi and some bacteria have mainly been studied, and white-rot fungi, and brown rot fungi in particular have obvious degradation effects on lignin [6, 7], but the development of industrial applications has been difficult [8]. Some actinomycetes and bacteria also participate in lignin degradation. Several typical lignin-degrading bacteria, such as *Rhodococcus*, *Pseudomonas*, *Sphingobium* and *Sphingomonas*, have been identified [9–12]. *Streptomyces* are among the identified bacteria capable of degrading lignin [13]. Pasti et al. [14] isolated 11 strains of actinomycetes from the intestines of termites and analysed their ability to degrade lignocellulose, lignin and carbohydrates. Other researchers have also screened streptomyces from the soil that degrade lignin [15–17].

There are many kinds of enzymes involved in the degradation of lignin, including laccase (Lac), lignin peroxidase (LiP), manganese peroxidase (MnP), multifunctional peroxidase (VP), and dye decolouring peroxidase (DyPs) [18]. The specific mechanism for biological degradation of lignin needs to be studied further. Lignin is rich in high-value

degradation intermediates, such as vanillin, guaiacol, catechin and protocatechin [19]. At the same time, the study of the metabolic mechanism of lignin and its transformation and utilization is an important part of the research needed for production of "second-generation biofuels" [20].

Hundreds of lignin derivatives have been identified in studies of bacterial degradation of lignin [21]. Because of the complex degradation mechanism, the interpretation of metabolic pathways and intermediate products is an important part of understanding the degradation of lignin. With the development of sequencing technology and bioinformatics, genomics research has become an important method for studying degradation mechanisms. Researchers are paying increasing attention to analysing metabolism of intermediate products through annotations of related degradation genes. *Pseudomonas putida* is a lignin-degrading bacterium that has been studied earlier. Lin et al. [22] identified several lignin-degrading enzymes, including haem peroxidase, from its genome and constructed five lignin metabolic pathways. Masai et al. [23] conducted a series of studies on lignin degradation and related genes in *Sphingomonas paucimobilis* SYK-6 and established a relatively complete lignin degradation metabolic pathway. Niewerth et al. [24] determined and analysed the whole genome sequence of *Arthrobacter* sp. Rue61a in soil, and the results showed that it has an aromatic degradation pathway that utilizes the characteristic products of lignin degradation, reflecting the saprophytic capabilities and nutritional diversity of organisms.

Several *Streptomyces* strains, such as *Streptomyces viridosporus* T7A [25] and *Streptomyces setonii* 75Vi2 [26], have been reported to degrade lignin. However, the actual catabolic pathways of lignin derivatives and the responsible enzymes and genes have not been investigated by using molecular methods with *Streptomyces*. In our present study, a novel isolate, DF3-3, identified as *Streptomyces thermocarboxydus*, was found to degrade lignin. Alkali lignin is a model compound with a structure similar to that of lignocellulose and is often used as a raw material for lignin degradation studies [27]. This study used alkali lignin to investigate the characteristics of alkali lignin degradation by *Streptomyces thermocarboxydus* strain DF3-3. GC–MS combined with genomics was used to identify the genes responsible for lignin degradation and explore the metabolic pathway for lignin degradation by *Streptomyces thermocarboxydus* strain DF3-3.

Results

Morphological and physiological characteristics of DF3-3

A strain of actinomycete named DF3-3 was isolated from the greening litter of Beijing University of Agriculture. The scribing form of the plate shows a rough white surface, the upper and lower surfaces are inconsistent in colour, and the hyphae in the base are obvious on Gause's medium. (Figure 1a). When DF3-3 was inoculated on Gause's guaiacol medium, a clear colour reaction appeared. When inoculated on Gause's-Azure B medium, a transparent fading circle appeared (Figure 1b and c). This shows that DF3-3 has the ability to degrade lignin [28]. Microscopy was used to observe the morphology of DF3-3 (Figure 1d). DF3-3 grew luxuriantly on the plate, hyphae were developed, and aerial hyphae were slender. The spore filaments were spiral-shaped with obvious characteristics of *Streptomyces*. Figure 1d shows a scanning electron micrograph of DF3-3 grown on Gause's medium. Mature spore chains were moderately long, with 40 to 80 spores per chain. The single spores were oval or cylindrical with diameters of 0.5 to 0.7 μm and lengths of 1.1 to 1.3 μm and have rough surfaces.

While keeping the conditions of the basal medium otherwise unchanged, different nitrogen sources and carbon sources were added to observe the growth of DF3-3 with different nitrogen sources. The results are shown in Table 1.

Table 1
Utilization characteristics of the nitrogen source and carbon source of DF3-3.

Trait	Result	Trait	Result	Trait	Result
Nitrogen source					
NH ₄ Cl	+	(NH ₄) ₂ SO ₄	+	Acrylamide	+
Potassium nitrate	+	Ammonium tartrate	+	Peptone	+
Carbon source					
Glucose	+	Alpha-D-methylglucoside	+	Sodium lactate	-
Mannose	+	trehalose	+	Sodium acetate	-
Melibiose	+	Cellobiose	+	Sodium formate	-
L-arabinose	-	Xylose	+	Sodium malate	+
Starch	+	Ribose	-	Sodium succinate	+
Melezitose	-	Inulin	-	Sodium malonate	+
Erythritol	-	Salicin	+	Sodium tartrate	-
Maltose	+	Glycerin	+	Sodium tyrosine	+
Sucrose	-	Sodium butyrate	-	Amylase	+

Molecular identification

Genome de novo sequencing was performed on DF3-3 cells using second- and third-generation sequencing methods, namely, Illumina HiSeq+PacBio, and the gene location and sequence information of the samples were obtained through de novo assembly and gene prediction. According to the whole genome sequencing results (Table 2), strain DF3-3 has a chromosome with a total genome length of 7311713 bp. GeneMarkS predicted and annotated a total of 6929 coding sequences (CDSs), with a G+C content of 72.24%.

Table 2
Genome-wide characteristics of DF3-3.

	value
Genome Size (bp)	7311713
Chrom No.	1
GC Content (%)	72.24
Gene No.	6929
GC Content in Gene Region (%)	72.52
Gene Average Len(bp)	927.29
Gene/Genome (%)	87.88
GC Content in Intergenetic Region (%)	70.17
tRNA No.	66
rRNA No.	18

At present, 95% of the average nucleotide identity (ANI) is often used as the standard for species classification and species clustering [29]. The whole genomes of eight strains with high homology to strain DF3-3 were selected and compared and analysed with DF3-3 using ANI. The results are shown in Figure 2. The similarity between DF3-3 and *Streptomyces thermocarboxydus* reached 98.96%, and it can now be identified as *Streptomyces thermocarboxydus*.

Biodegradation of alkali lignin by DF3-3

To investigate lignin degradation by strain DF3-3, cells were incubated at 30°C in medium with alkali lignin as the carbon source. The growth curve and degradation rate of alkali lignin are shown in Figure 3. The degree of degradation of alkaline lignin by DF3-3 shows that the degradation rate increased significantly in days 1–4, and the increase was the largest on the fourth day. After 4–7 days, degradation continued, but the rate slowed down, and the efficiency for degradation of alkali lignin reached 31% on day 15. The degradation of lignin by microorganisms requires a relatively slow process to reach a significant level. The white-rot fungus *Phanerochaete chrysosporium* was used for degradation of lignin, and the efficiency reached approximately 20% on day 15 [30], so DF3-3 exhibited a better performance. This result is also similar to the degradation results seen with some *Streptomyces* strains, such as *S. viridosporus* T7A (lignin loss 30.9%) and *S. setonii* 75Vi2 (lignin loss 34.1%) [26, 31].

Analysis of lignin-degrading enzymes

Lignin molecules are not easily taken passively into the cell; therefore, *Streptomyces thermocarboxydus* DF3-3 might produce extracellular enzymes for synergistic degradation. Three major types of enzymes responsible for the degradation of lignin are lignin peroxidase (LiP), manganese peroxidase (MnP) and laccase (Lac) [32]. MnPs oxidize Mn(II) to Mn(III), and Mn(III) oxidizes phenolic compounds and generates phenoxy radicals that in turn undergo a variety of reactions, resulting in depolymerization. In the presence of Mn(II), MnP oxidizes nonphenolic lignin model compounds via peroxidation of unsaturated lipids. LiP is the most effective peroxidase and can oxidize phenolic and nonphenolic compounds, amines, aromatic ethers, and polycyclic aromatics [33]. Lac is a copper oxidoreductase [34] that can degrade refractory polyphenols and nonphenolics in lignin, and the expression of its coding genes in bacteria has also been reported [35]. The activities of these three enzymes from DF3-3 are shown in Figure 4. MnP and Lac activity increased constantly during the initial 6 days, with maxima of 1821.66 U/L and

1265.58 U/L seen at day 6, followed by slight decreases from day 7. Lip activity was maintained at a low level, with a maximum of 480.33 U/L on day 4. These results indicated that MnP and Lac play crucial roles during the entire process of alkaline lignin degradation by DF3-3.

Aromatic intermediates identified by GC-MS analysis

Using ethyl acetate as the solvent for GC-MS determination [36], the degradation products of alkali lignin produced in incubation with *Streptomyces thermocarboxydus* strain DF3-3 were analysed from the first day to the 15th day. In total, 19 degradation intermediate products were identified by GC-MS, including seven different types of compounds, including eight organic acids (butanoic acid (**4**), lactic acid (**8**), 2-hydroxypropanoic acid (**9**), pyrrole-2-carboxylic acid (**10**), 3-phenylpyruvic acid (**12**), 4-hydroxybenzoic acid (**13**), 4-hydroxyphenylpyruvate (14), palmitic acid (**16**), stearic acid (**17**)), three esters (acetic acid butyl ester (**1**), dibutyl phthalate (**15**), bis(2-ethylhexyl) phthalate (**19**)), two ethers (1-(1-ethoxy)propane (**2**) and 1-(1-propoxyethoxy)propane (**6**)), two alcohols (2-ethoxyethanol (**5**) and ethylene glycol (**7**)), one alkane (m-xylene (**2**)), and 1-(1-propylene (6-di-butylphenol phenol (2,2-**ethylethylenes(11-methyl)**). These 19 low molecular weight products are related to lignin metabolism, and their serial numbers, name, and retention times are shown in Table 3.

Table 3
Compounds identified from degradation of alkali lignin by DF3-3.

	Retention time	Compounds	Molecular formula	Control group
1	4.530	Acetic acid, butyl ester	C ₆ H ₁₂ O ₂	+
2	4.651	1-(1-Ethoxyethoxy) propane	C ₇ H ₁₆ O ₂	+
3	5.690	M-Xylene	C ₈ H ₁₀	+
4	6.055	Butyric acid	C ₄ H ₈ O ₂ *	+
5	6.251	2-Ethoxyethanol	C ₄ H ₁₀ O ₂ *	+
6	6.642	1-(1-Propoxyethoxy) propane	C ₈ H ₁₈ O ₂	+
7	8.793	Ethylene glycol	C ₂ H ₆ O ₂ *	+
8	11.058	Lactic acid	C ₃ H ₆ O ₃ *	+
9	13.038	2-Hydroxybutyric acid	C ₄ H ₈ O ₃ *	-
10	19.375	Pyrrole-2-carboxylic acid	C ₅ H ₅ NO ₂	-
11	23.225	2,4-Di-tert-butylphenol	C ₁₄ H ₂₂ O	-
12	25.121	3-Phenylpyruvic acid	C ₉ H ₈ O ₃ *	-
13	25.996	4-Hydroxybenzoic acid	C ₇ H ₆ O ₃ *	-
14	32.028	4-Hydroxyphenylpyruvate	C ₉ H ₈ O ₄ *	-
15	32.970	Dibutyl phthalate	C ₁₆ H ₂₂ O ₄	+
16	34.543	Palmitic acid	C ₁₆ H ₃₂ O ₂ *	+
17	38.084	Stearic acid	C ₁₈ H ₃₆ O ₂ *	+
18	41.008	2,2'-Methylenebis(6-tert-butyl-4-methyl-phenol)	C ₂₃ H ₃₂ O ₂	-
19	43.033	Bis(2-ethylhexyl) phthalate	C ₂₄ H ₃₈ O ₄	-

“+” indicates that the product was detected in the uninoculated control group, “-” indicates that the product was not detected in the uninoculated control group.

The variations in the peak areas for products **3**, **8**, **10**, **11**, **12**, **13**, **14**, **15**, **18**, and **19** were plotted using the peak areas and culture times (days) as the ordinate and abscissa, respectively, and the result is shown in Figure 5. The peak areas for products **12**, **15**, and **18** varied greatly; in comparison, the peak areas for products **3**, **8**, **10**, **11**, **13**, **14**, and **19** exhibited small changes. The content of product **12** gradually increased within 1–4 days after culture, reached a maximum on the 4th day, and then decreased to a minimum on the 6th day. The content of product **15** quickly increased to a high value on the first day of culture and then gradually decreased to the lowest value until the 7th

day. The content of product **18** increased to the first high value on the 1st day after culture, dropped to the lowest point on the 3rd day, then rose to another highest point on the 4th day, and finally decreased to the lowest value. These data indicate that the contents of **12**, **15**, and **18** underwent two cycles in the first 8 days of culture.

The process of lignin degradation can be divided into two main parts. First, through the rupture of aromatic ether bonds and carbon-carbon bonds, lignin polymers are depolymerized to form various oligomers. Then, these depolymerized aromatic organic compounds form other small molecular benzene compounds [37]. Here, m-xylene (**3**), 2,4-di-tert-butylphenol (**11**), 3-phenylpyruvic acid (**12**), 4-hydroxybenzoic acid (**13**), 4-hydroxyphenylpyruvate (**14**), dibutyl phthalate (**15**), 2,2'-methylenebis(4-methyl-6-tert-butylphenol), and bis(2-ethylhexyl) phthalate (**19**) were detected among the products produced by DF3-3 in degradation of alkaline lignin. Dibutyl phthalate (**15**) is a common lignin degradation intermediate [38–40] that can be metabolized to produce phthalic acid, which is further converted into syringyl, protocatechin and other phenolic compounds and then degraded [41]. According to the metabolic pathway for formation of dibutyl phthalate (**15**) and diisooctyl phthalate (**19**), product **19** is thought to have a similar degradation process.

Both 4-hydroxyphenylpyruvate (**14**) and 3-phenylpyruvic acid (**12**) were detected among the products of the experimental group, and their aromatic structures were related to alkali lignin. 4-Hydroxybenzoic acid is an important intermediate in many lignin metabolic pathways, including the coumaric acid pathway, gentisic acid pathway, and cinnamic acid pathway [42]. This indicated that DF3-3 uses the β -keto adipate metabolic pathway or other similar degradation pathways.

In addition, small molecular compounds such as ethylene glycol (**7**) and 2-ethoxyethanol (**5**) were detected in the experiment, indicating that DF3-3 degradation of lignin has great potential for energy conversion.

Analyses of the lignin metabolism pathways based on whole genome sequencing and annotation

Initial degradation of lignin into low molecular weight compounds by extracellular phenoloxidases

To identify the genes related to lignin degradation by DF3-3, the genes encoding known lignin enzymes of other bacterial species were selected as query sequences, and a BLASTp search of the DF3-3 genome was carried out. The results showed that DF3-3 has abundant genes encoding lignin-degrading enzymes, and a total of 107 different genes involved in the degradation process have been annotated (Tables 4-11).

Table 4
Genes responsible for lignin depolymerization.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 2915	2288	copA	carbonate dehydratase	<i>Actinospica acidiphila</i>	WP_058917537.1	96.19
gene 5998	2267	copA	carbonate dehydratase	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028959238.1	99.74
gene 3898	2225	katG	catalase-peroxidase	<i>Streptomyces</i> sp. Akac8	WP_136237873.1	99.87
gene 3899	380	fur	transcriptional repressor	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028958816.1	99.21
gene 0904	1400	YhjG	pentachlorophenol monooxygenase	<i>Streptomyces</i> sp. 4F	ALV51870.1	95.49
gene 1050	533	ahpD	alkyl hydroperoxide reductase	<i>Actinospica acidiphila</i>	WP_163089316.1	99.44
gene 6181	1415	glcD	FAD-linked oxidase	<i>Actinospica acidiphila</i>	NEC50278.1	98.73
gene 5348	716	yfiH	laccase	<i>Streptomyces griseorubens</i>	GGQ64023.1	96.64
gene 4491	197	-	multicopper oxidase domain-containing protein	unclassified <i>Streptomyces</i>	WP_106959434.1	98.46
gene 2609	989	-	multicopper oxidase domain-containing protein	<i>Streptomyces</i> sp. XHT-2	WP_161108161.1	99.39
gene 2857	983	-	Phenoxybenzoate dioxygenase	<i>Streptomyces tuius</i>	WP_190903372.1	88.07
gene 5116	1529	aldH	aldehyde dehydrogenase	<i>Streptomyces</i> sp. XHT-2	MXQ60577.1	99.61
gene 1234	1451	katE	catalase	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028959332.1	99.79
gene 3983	1655	katE	catalase	<i>Streptomyces</i> sp. GESEQ-13	WP_210638384.1	99.09
gene 3220	2276	katE	catalase HP11	<i>Streptomyces</i> sp. XHT-2	WP_161108660.1	99.74
gene 3744	959	-	Pimeloyl-ACP methyl ester carboxylesterase	<i>Streptomyces</i> sp. di188	SCD58877.1	88.43
gene 6937	1277	-	Dye-decolorizing peroxidase	<i>Streptomyces pharetrae</i> CZA14	OSZ61955.1	83.61

Table 5
Genes responsible for the β -ketoadipate pathway and peripheral reactions.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 3935	1001	-	Vanillate O-demethylase oxygenase subunit	<i>Streptomyces</i> sp. SID4956	WP_161232727.1	99.40
gene 2826	773	hpaI	4-hydroxy-2-oxovalerate aldolase	<i>Streptomyces</i> sp. XHT-2	MXQ61669.1	99.61
gene 0228	335	-	phenylpropionate dioxygenase-like ring-hydroxylating dioxygenase	<i>Streptomyces paradoxus</i>	MBB6077390.1	94.59
gene 0557	599	ubiX	aromatic acid decarboxylase	<i>Streptomyces</i> sp. 4F	ALV54514.1	97.99
gene 0021	377	pcaC	4-carboxymuconolactone decarboxylase	<i>Streptomyces</i> sp. ok210	WP_093930409.1	98.40
gene 0859	506	-	carboxymuconolactone decarboxylase family protein	<i>Streptomyces</i> sp. GESEQ-13	WP_210634556.1	99.40
gene 2323	1199	pcaL	4-carboxymuconolactone decarboxylase	<i>Streptomyces</i> sp. Akac8	WP_136239018.1	99.75
gene 2838	1121	pcaL	3-oxoadipate enol-lactonase	<i>Streptomyces</i> sp. SID4956	WP_161232455.1	99.73
gene 2839	1331	pcaB	3-carboxy-cis, cis-muconate cycloisomerase	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028959090.1	99.55
gene 2840	605	pcaG	protocatechuate 3,4-dioxygenase subunit alpha	<i>Streptomyces</i> sp. GESEQ-13	WP_210638048.1	99.50
gene 2841	773	pcaH	protocatechuate 3,4-dioxygenase subunit beta	<i>Streptomyces</i> sp. GESEQ-13	WP_210638047.1	99.22
gene 2842	1208	-	β -ketoadipyl CoA thiolase	<i>Streptomyces</i> sp. B9173	OQR62711.1	88.25
gene 2831	1364	pcaK	4-hydroxybenzoate transporter PcaK	<i>Streptomyces</i> sp. GESEQ-13	WP_210638055.1	98.02
gene 2847	1439	pcaK	4-hydroxybenzoate transporter PcaK	<i>Streptomyces</i> sp. Akac8	WP_136238348.1	99.79
gene 2852	734	pcaR	Pca regulon regulatory protein	<i>Streptomyces cellulosae</i>	GHE61509.1	97.13
gene 2848	1013	hcaD	Terephthalate 1,2-dioxygenase	<i>Streptomyces cellulosae</i>	GHE61489.1	97.63
gene 4139	1148	hcaD	hypothetical protein	<i>Streptomyces</i> sp. GESEQ-13	WP_210637914.1	98.43
gene 5175	317	hcaC	3-phenylpropionate/trans-cinnamate dioxygenase ferredoxin subunit	<i>Streptomyces</i> sp. di50b	SCD40618.1	97.14

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 5716	1265	hcaD	pyridine nucleotide-disulfide oxidoreductase	<i>Streptomyces</i> sp. 4F	ALV50055.1	96.67
gene 5542	1025	-	Trans-1,2-dihydrobenzene-1,2-diol dehydrogenase	<i>Streptomyces afghaniensis</i> 772	EPJ42281.1	82.27
gene 4140	923	-	4,5-dihydroxyphthalate decarboxylase	<i>Streptomyces</i> sp. 3212.3	REE57871.1	84.69
gene 4458	1220	pobA	4-hydroxybenzoate 3-monooxygenase	<i>Streptomyces</i> sp. XHT-2	WP_161108189.1	99.01
gene 6344	1175	pobA	4-hydroxybenzoate 3-monooxygenase	<i>Streptomyces</i> sp. XHT-2	WP_161107724.1	99.49
gene 6436	1517	-	cyclohexanone monooxygenase	<i>Streptomyces</i> sp. 4F	ALV51359.1	97.03
gene 6471	518	pat	phosphinothricin N-acetyltransferase	<i>Streptomyces werraensis</i>	GHE99025.1	94.19
gene 4600	1322	-	4-methylmuconolactone transporter	<i>Streptomyces afghaniensis</i> 772	EPJ35848.1	87.02

Table 6
Genes responsible for gentisate pathways.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 1961	2366	-	bifunctional salicylyl-CoA 5-hydroxylase/oxidoreductase	<i>Streptomyces</i> sp. 4F	ALV52714.1	97.07
gene 0811	1613	gabD	succinate-semialdehyde dehydrogenase	<i>Streptomyces</i> sp. UNC401CLCol	WP_028960221.1	99.81
gene 4361	1388	gabD	NADP-dependent succinic semialdehyde dehydrogenase	<i>Streptomyces cellulosa</i>	GHE69900.1	98.70
gene 3797	1187	-	fumarylacetoacetate hydrolase	<i>Streptomyces</i> sp. SID8376	WP_205526335.1	99.75
gene 2855	1286	-	aromatic ring-hydroxylating dioxygenase subunit alpha	<i>Streptomyces</i> sp. GESEQ-13	WP_137208924.1	99.53
gene 4131	1280	-	Phthalate 4,5-dioxygenase oxygenase subunit	<i>Streptomyces</i> sp. MBT84	MBW8706675.1	70.79
gene 4137	620	-	gentisate 1,2-dioxygenase	<i>Robiginitomaculum</i> sp.	PHR56577.1	81.03
gene 4928	1085	frmA	alcohol dehydrogenase	<i>Streptomyces cellulosa</i>	GHE58756.1	97.23
gene 5219	689	-	maleylpyruvate isomerase	<i>Streptomyces</i> sp. XHT-2	WP_161107348.1	99.13

Table 7
Genes responsible for pathways for anthranilate.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 1963	1610	abmG	2-aminobenzoate-CoA ligase	<i>Streptomyces</i> sp. 4F	ALV52716.1	94.03
gene 1964	1128	-	acyl-CoA dehydrogenase	<i>Streptomyces</i> sp. XHT-2	WP_161107119.1	99.47
gene 0037	638	pabA	aminodeoxychorismate/anthranilate synthase component II	<i>Streptomyces</i> sp. GESEQ-13	WP_210635465.1	99.53
gene 5387	1880	phzE	anthranilate synthase	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028960786.1	99.52
gene 0262	1202	kynU	L-kynurenine hydrolase	<i>Streptomyces</i> sp. UNC401CLCoI	WP_028959521.1	99.25
gene 0261	830	TDO2	tryptophan 2,3-dioxygenase	<i>Streptomyces</i> sp. GESEQ-13	WP_210635572.1	99.28
gene 4644	1697	-	Tryptophan 2-monooxygenase	<i>Streptomyces</i> sp. di50b	SCD87347.1	97.35
gene 0499	1217	-	Aromatic-amino-acid aminotransferase 1	<i>Streptomyces</i> sp. UNC401CLCoI	WP_064743320.1	99.49
gene 2090	873	-	indole-3-glycerol phosphate synthase	<i>Streptomyces</i> sp. I4 (2020)	WP_199204736.1	96.82
gene 5302	806	trpC	indole-3-glycerol phosphate synthase	<i>Streptomyces</i> sp. I4 (2020)	WP_199207289.1	99.63
gene 5306	1493	trpE	anthranilate synthase component I	<i>Streptomyces</i> sp. XHT-2	WP_161107330.1	99.8

Table 8
Genes responsible for pathways for resorcinol.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 0128	971	-	LysR family transcriptional regulator	<i>Streptomyces</i> sp. UNC401CLCol	WP_078611393.1	99.69
gene 2842	1208	-	β -ketoacyl-CoA thiolase	<i>Streptomyces</i> sp. B9173	OQR62711.1	88.25
gene 1964	1128	-	acyl-CoA dehydrogenase	<i>Streptomyces</i> sp. XHT-2	WP_161107119.1	99.47
gene 2768	1091	-	aryl-alcohol dehydrogenase	<i>Streptomyces</i> sp. di50b	SCD35921.1	93.66
gene 2849	962	pdxA	Terephthalate dihydrodiol dehydrogenase	<i>Streptomyces</i> sp. McG7	WP_215047915.1	98.75
gene 2850	470	andAd	terephthalate 1,2-dioxygenase	<i>Streptomyces</i> sp. XHT-2	MXQ61646.1	100.00
gene 2851	1283	andAc	Terephthalate 1,2-dioxygenase	<i>Rhodococcus pyridinivorans</i> AK37	EHK83592.1	83.05
gene 3600	1502	-	2-polyprenyl-6-methoxyphenol hydroxylase	<i>Streptomyces</i> sp. di50b	SCD47389.1	85.06
gene 5989	1268	-	dimethylaniline monooxygenase	<i>Streptomyces</i> sp. 4F	ALV54407.1	96.91
gene 3045	383	-	extradiol dioxygenase	<i>Streptomyces</i> sp. 4F	ALV53653.1	93.70
gene 4526	476	-	putative dehydrochlorinase	<i>Streptomyces</i> sp.	AUR34009.1	81.65
gene 5542	1025	-	Trans-1,2-dihydrobenzene-1,2-diol dehydrogenase	<i>Streptomyces afghaniensis</i> 772	EPJ42281.1	82.27
gene 6472	785	-	extradiol ring-cleavage dioxygenase	<i>Streptomyces griseorubens</i>	KEG41654.1	98.85

Table 9
Genes responsible for catabolic pathways for homogentisic.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 4937	1361	hmgA	homogentisate 1,2-dioxygenase	<i>Streptomyces</i> sp. XHT-2	WP_161108369.1	99.78
gene 1961	2366	-	bifunctional salicylyl-CoA 5-hydroxylase/oxidoreductase	<i>Streptomyces</i> sp. 4F	ALV52714.1	97.07
gene 0499	1217	-	Aromatic-amino-acid aminotransferase 1	<i>Streptomyces</i> sp. UNC401CLCol	WP_064743320.1	99.49
gene 6956	1079	hisC	Putative phenylalanine aminotransferase	<i>Streptomyces werraensis</i>	GHF12148.1	96.93
gene 0811	1613	gabD	succinate-semialdehyde dehydrogenase	<i>Streptomyces</i> sp. UNC401CLCol	WP_028960221.1	99.81
gene 4361	1388	gabD	NADP-dependent succinic semialdehyde dehydrogenase	<i>Streptomyces cellulosa</i>	GHE69900.1	98.70
gene 0636	1214	fahA	Fumarylacetoacetate hydrolase	<i>Streptomyces</i> sp. SID4956	WP_161232839.1	99.01
gene 3797	1187	-	fumarylacetoacetate hydrolase	<i>Streptomyces</i> sp. SID8376	WP_205526335.1	99.75
gene 2443	305	phhB	4a-hydroxytetrahydrobiopterin dehydratase	<i>Streptomyces griseorubens</i>	WP_033273716.1	98.02
gene 4811	2576	pheT	phenylalanine-tRNA ligase subunit beta	<i>Streptomyces</i> sp. GESEQ-13	WP_210636334.1	99.77
gene 4812	1121	pheS	phenylalanine-tRNA ligase subunit alpha	<i>Streptomyces cellulosa</i>	GHE73990.1	100
gene 3632	1814	hppD	4-hydroxyphenylpyruvate dioxygenase	<i>Streptomyces cellulosa</i>	GHE74791.1	97.35
gene 3633	876	aroE	shikimate dehydrogenase	<i>Streptomyces</i> sp. FxanaD5	WP_019522987.1	99.66
gene 6183	1145	hppD	4-hydroxyphenylpyruvate dioxygenase	<i>Actinospica acidiphila</i>	WP_163089359.1	98.69
gene 4928	1085	frmA	alcohol dehydrogenase	<i>Streptomyces cellulosa</i>	GHE58756.1	97.23
gene 5219	689	-	maleylpyruvate isomerase	<i>Streptomyces</i> sp. XHT-2	WP_161107348.1	99.13

Table 10
Genes responsible for pathways for phenylacetate-CoA.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 0050	1061	paaE	phenylacetic acid degradation protein	<i>Streptomyces cellulosa</i>	GHE31167.1	99.15
gene 0051	539	paaD	phenylacetate-CoA oxygenase subunit	<i>Streptomyces cellulosa</i>	GHE31162.1	97.77
gene 0052	737	paaC	phenylacetate-CoA oxygenase subunit	<i>Streptomyces</i> sp. GESEQ-13	WP_210635474.1	98.78
gene 0053	287	paaB	phenylacetate-CoA oxygenase subunit PaaB	<i>Streptomyces griseoruber</i>	WP_055636201.1	98.95
gene 0054	1016	paaA	phenylacetate-CoA oxygenase subunit PaaA	<i>Streptomyces</i> sp. SID4956	WP_161232185.1	99.41
gene 0058	1691	PaaN	phenylacetic acid degradation protein	<i>Streptomyces</i> sp. Akac8	WP_136238706.1	99.82
gene 0059	1514	paaH	3-hydroxyacyl-CoA dehydrogenase	<i>Streptomyces</i> sp. SID4956	WP_161233084.1	100.00
gene 1159	803	paaG	1,2-epoxyphenylacetyl-CoA isomerase	<i>Streptomyces cellulosa</i>	GHE51256.1	98.13
gene 1381	848	paaH	3-hydroxybutyryl-CoA dehydrogenase	<i>Streptomyces</i> sp. F-7	WP_093768152.1	99.65
gene 2386	1805	paaH	3-hydroxybutyryl-CoA dehydrogenase	<i>Streptomyces</i> sp. SMS_SU21	WP_102640749.1	97.17
gene 2736	764	paaF	3-hydroxypropionyl-CoA dehydratase	<i>Streptomyces</i> sp. GESEQ-13	WP_210638121.1	99.61
gene 4099	1523	paaK	phenylacetate-CoA ligase	<i>Streptomyces</i> sp. GESEQ-13	WP_210637941.1	99.77
gene 4804	860	paaH	3-hydroxybutyryl-CoA dehydrogenase	<i>Streptomyces</i> sp. l4 (2020)	WP_199207038.1	99.30
gene 6544	734	paaF	enoyl-CoA hydratase	<i>Streptomyces cellulosa</i>	GHE27114.1	98.77
gene 4828	1376	Pad	Phenylacetaldehyde dehydrogenase	<i>Streptomyces</i> sp. DI166	SBT93309.1	87.53
gene 1454	767	-	enoyl-CoA hydratase	<i>Streptomyces</i> sp. Akac8	WP_136239245.1	99.61
gene 1960	827	-	enoyl-CoA hydratase	<i>Streptomyces griseorubens</i>	WP_033274100.1	97.82
gene 1962	846	PaaX	phenylacetic acid degradation operon negative regulatory protein	<i>Streptomyces aureorectus</i>	MBA8975440.1	90.71

Table 11
Genes responsible for catabolic pathways for 2,3-dihydroxyphenylpropionate.

Gene ID	Size (aa)	Gene Name	Encode protein	Species of reference gene	Accession no. (NCBI)	BLAST identity (%)
gene 3746	1028	-	3-hydroxycinnamic acid hydroxylase	<i>Streptomyces albogriseolus</i>	GHB97425.1	91.81
gene 1753	1640	mhpA	3-(3-hydroxy-phenyl) propionate acid hydroxylase	<i>Streptomyces</i> sp. di50b	SCE12787.1	96.15
gene 2825	1613	mhpA	3-(3-hydroxy-phenyl) propionate hydroxylase	<i>Streptomyces griseorubens</i>	GGQ79721.1	97.77
gene 2828	794	mhpD	fumarylacetoacetate hydrolase	<i>Streptomyces</i> sp. GESEQ-13	WP_210638174.1	99.62
gene 2830	935	mhpB	3-carboxyethylcatechol 2,3-dioxygenase	<i>Streptomyces</i> sp. GESEQ-13	WP_210638056.1	99.04
gene 2826	773	hpaI	4-hydroxy-2-oxovalerate aldolase	<i>Streptomyces</i> sp. XHT-2	MXQ61669.1	99.61
gene 5487	359	hpaF	isomerase	<i>Streptomyces cellulosa</i>	GHE29375.1	97.48
gene 0171	431	-	aldehyde dehydrogenase	<i>Streptomyces werraensis</i>	GHF03416.1	88.46
gene 0388	1460	-	aldehyde dehydrogenase	<i>Streptomyces</i> sp. GESEQ-13	WP_210635659.1	99.18

In the initial stages of lignin degradation, extracellular enzymes are responsible for depolymerization of lignin. Degradation enzyme systems involved in lignin degradation include the lignin oxidase system, hydrogen peroxide production enzyme system and other enzyme systems. In this study, multiple lignin depolymerization-related genes were found in the genome of DF3-3 (Table 4), including genes encoding multicopper oxidase (gene 5348, gene 4491, gene 2609), alcohol dehydrogenase (gene 2768), alcohol dioxygenase (gene 3045), aldehyde dehydrogenase (aldH), pentachlorophenol monooxygenase (gene 0904) and terephthalate dihydrodiol dehydrogenase (pdxA).

Catabolism pathways for lignin components

Many low molecular weight compounds are produced by initial depolymerization of lignin. Based on GC-MS and genome analysis, 5 metabolic pathways of lignin-based derivatives were predicted: β -keto adipate pathway and peripheral reactions; gentisate pathway; anthranilate pathway; homogentisic pathway; and catabolic pathway for resorcinol (shown in Figure 7). Based on genome data alone, we predicted two other pathways: the phenylacetate-CoA pathway and the 2,3-dihydroxyphenylpropionic acid pathway (Figure 8).

β -keto adipate pathway and peripheral reactions

The β -keto adipate pathway is an important aromatic metabolic pathway for many microorganisms. There are 26 genes involved in the degradation process of lignin in DF3-3 (Table 5), and they are scattered throughout the genome of DF3-3 (Figure 9). The key genes of the protocatechuic acid 3,4-dioxygenase (pcaG, pcaH) gene appear in the whole genome of DF3-3 and can be imagined to metabolize lignin intermediates through the β -keto adipate

pathway branched by protocatechin. There are two 4-hydroxycinnamic acid dioxygenases (hcaC, hcaD) in DF3-3, and 4-hydroxycinnamic acid can initially be degraded to generate 4-hydroxycinnamic acid coenzyme A and further converted into p-hydroxybenzoic acid, which was found among the degradation products of alkaline lignin produced by DF3-3. 3-Phenylpyruvic acid and 4-hydroxyphenylpyruvate were also detected as metabolites of DF3-3. We speculate that they can also be degraded by the hcaC gene. Then, 4-hydroxyphenylpyruvate undergoes a series of reactions to produce p-hydroxybenzoic acid. The pcaG and pcaH genes cause benzene epoxidation to enable a ring-opening reaction generating cis-hexadienedioic acid; this further generates β -keto adipate, which is degraded and finally enters the tricarboxylic acid cycle. Unlike 4-hydroxyphenylpyruvate, 3-phenylpyruvic acid may not be converted only into 4-hydroxyphenylpyruvate by hcaC but could also be converted into benzoic acid and catechol, and then a ring-opening reaction takes place to enter the β -keto adipate pathway or generate 4-hydroxy-2-oxovalerate and further degradation.

The gentisate pathway

The gentisate pathway usually starts from the degradation of salicylic acid. Ethyl salicylate is a precursor in the production of salicylic acid. The pathway for degradation of phthalic acid by some fungi has been explained, in which dibutyl phthalate or diisooctyl phthalate is converted to ethyl salicylate [43]. Some aromatic oxygenases may play important roles in this process. In the DF3-3 genome, the genes encoding salicylic acid-5-hydroxylase (gene 1961) and gentisic acid 1,2-dioxygenase (gene 4137) were identified (Table 6). Gentianic acid 1,2-dioxygenase oxygenates the benzene ring of gentisic acid to produce maleylpyruvate, which is then further metabolized by maleate pyruvate isomerase (gene 5219). Dibutyl phthalate and diisooctyl phthalate were detected among the metabolites of DF3-3. Phthalate dioxygenase oxygenase (gene 4131) detected in the DF3-3 genome gene also indicated that it may appear as an intermediate product of the salicylic acid metabolic pathway.

The anthranilate pathway

The specific mechanism of the anthranilic acid pathway in bacteria has not been fully elucidated [42, 44]. A 2-aminobenzoate-CoA ligase (abmG) and an L-kynurenine hydrolase (kynU) gene were identified in DF3-3, and a tryptophan 2,3-dioxygenase (TDO2) gene was also observed (Table 7). It is speculated that the anthranilic acid pathway of DF3-3 generates formyl kynurenine through the degradation of tryptophan, which is further converted to kynurenine and then generates anthranilic acid under the action of kynU. In addition, tryptophan may undergo a series of redox reactions at its side chain to produce indole derivatives. The pyrrole-2-carboxylic acid detected among the metabolites of DF3-3 can also be degraded by the anthranilate synthase (phzE) gene and abmG gene through a similar pathway.

Catabolic pathway for resorcinol

Due to the existence of the metabolite m-xylene and related genes (Table 8), it is speculated that DF3-3 has related metabolic pathways. The resorcinol pathways for other strains involve the participation of oxidoreductases. The functions of some of these genes, such as aromatic compound monooxygenase (gene 3600)-coding genes, have not been fully clarified. Aryl-alcohol dehydrogenase and extradiol dioxygenase may effect the degradation of aromatic phenols. It is speculated that the metabolism of m-xylene by DF3-3 may take place through oxidation of

methyl groups to produce resorcinol. Then, resorcinol undergoes further oxygenation on the branch chain and ring-opening oxidation with the action of oxygenases (Figure 7).

Homogentisic pathway

The homogentisic pathway is the central pathway for catabolism of phenylalanine and tyrosine. Genomic analysis of DF3-3 shows that phenylalanine is converted to tyrosine by phenylalanine hydroxylase and auxiliary methanolamine dehydratase (PhhB), and tyrosine aminotransferase (hisC) is converted to tyrosine 4-hydroxyphenylpyruvate, which is further converted into homogentisate by 4-hydroxyphenylpyruvate dioxygenase (hppD) and degraded by the action of homogentisate 1,2-dioxygenase (hmgA) to produce maleacetoacetate (Figure 9). A fumarylacetoacetate hydrolase (fahA) gene was identified in the DF3-3 genome (Table 9), indicating that hydrolysis of maleacetoacetate occurs through the degradation of fumarate acetoacetate. The genes involved in the homogentisate pathway are scattered throughout the genome, and the processes are carried out with the joint action of hmgA and other related genes. This non-linkage also appears in other bacteria [44]. 4-Hydroxyphenylpyruvate, as the precursor of homogentisic acid, was observed among the products detected by GC-MS, and two hppD genes observed in the genome are related to the 4-hydroxy phenylpyruvate dioxygenase in *Streptomyces cellulosae* and *Actinospica acidiphila* with high similarities (97.35% and 98.69%).

In addition to the above metabolic pathways, the following two possible lignin degradation pathways were found for DF3-3 through a gene search. A relatively complete set of coding genes is present in the genome to form the pathway, but the related metabolites were not observed in this study.

Phenylacetate-CoA pathway

Phenylacetate can be derived from lignin-related phenylpropane units [45], and the general pathway for aerobic metabolism has just been discovered and studied with some bacteria [46, 47]. There were at least 15 genes in DF3-3 involved in this process (Table 10). The phenylacetate coenzyme A oxygenase gene is organized into clusters (paaABCDE) and exists in the DF3-3 genome (Figure 9), downstream from which there are a phenylacetic acid degradation protein (paaN) and a 3-hydroxyacyl-CoA dehydrogenase (paaH) encoding gene; a 1,2-epoxyphenylacetyl-coenzyme A isomerase (paaG) gene exists a little further downstream, and the isomerase it encodes can convert 3-hydroxypropionyl-CoA into 1,2-epoxyphenylacetyl-CoA. At the same time, 3-hydroxypropionyl-CoA can also be further degraded by 3-hydroxypropionyl-CoA dehydrogenase (paaH) and 3-hydroxypropionyl-CoA dehydrogenase (paaF) and finally enter the tricarboxylic acid cycle.

2,3-Dihydroxyphenylpropionic acid pathway

The 2,3-dihydroxyphenylpropionic acid pathway-related mhpABD gene cluster was observed in the DF3-3 genome (Table 11), including encoding for 3-(3-hydroxy-phenyl) propionate hydroxylase (mhpA), 2,3-dihydroxyphenylpropionate 1,2-dioxygenase (mhpB) and fumarate acetoacetate hydrolase (mhpD) genes, which are located upstream of the 4-hydroxy-2-oxovalerate aldolase (hpaI) gene (Figure 9). DF3-3 lacks the mhpE and mhpF genes, and the generated 4-hydroxy-2-oxovalerate is decomposed by the aldolase encoded by the hpaI gene. It was also determined that other aldehyde dehydrogenases may effect the degradation of acetaldehyde and generate pyruvate and acetyl-CoA to enter the tricarboxylic acid cycle for metabolism.

Discussion

Biofuel production by biodegradable lignocellulose has far-reaching prospects [13]. Research on the degradation system of lignocellulose-degrading microorganisms has great significance to its practical application and development. However, few microorganisms, only white rot fungi, have been reported that can degrade lignin and cellulose at the same time [18]. In the actual degradation of biomass materials, the degradation of lignin, cellulose and hemicellulose often occurs simultaneously and is interrelated. In our research, we found that DF3-3 has good degradation performance on both lignin and cellulose. More experiments and new approaches are needed in future work to better develop and apply DF3-3.

The enzymology for bacterial lignin degradation has been well-studied in recent years, and some bacterial specific enzymes for lignin degradation have been reported. Like Co-dehydrogenase (LigD) [48], glutathione-dependent β -etherase enzymes (LigE, F, G) have been identified from *Sphingomonas paucimobilis* SYK-6 [49], a demethylase enzyme (LigX) from *Pseudomonas paucimobilis* [50], and DyP-type peroxidases from *Rhodococcus jostii* RHA1 [51]. Recent reports have shown multicopper oxidases that demonstrate laccase activities [52, 53]. In our study, we found that DF3-3 had Lac, Mnp and Lip activities. Genome research identified three multicopper oxidase coding genes (gene 5348, gene 4491, gene 2609), and gene 5348 has high similarity (96.64%) to the laccase structural protein gene of *Streptomyces griseorubens* (GGQ64023.1) [54], showing that DF3-3 has the ability to encode laccase at the genetic level. No gene encoding manganese peroxidase has been detected, but a catalase/peroxidase gene (gene3898) was found in DF3-3, which was 99.87% similar to KatG from *Streptomyces* sp. Akac8 [8]. The fur gene (gene3899) encoding a transcription regulator appears downstream, and these were reported as possible manganese peroxidase-encoding genes in *Streptomyces reticuli* [43, 44], suggesting that DF3-3 can exhibit manganese peroxidase activity. In addition, a gene (gene 6937) encoding a dye decolouring peroxidase was observed. The peroxidase encoded by it has a broad spectrum of substrates, which is also commonly reported in the depolymerization of lignin in some bacteria [22, 45, 46].

The hydrogen peroxide-producing enzyme system mainly participates in lignin degradation in the capacity of auxiliary enzymes [47]. These enzymes include glyoxal oxidase, aryl alcohol oxidase, quinone reductase and related dehydrogenases [48]. They produce hydrogen peroxide to support degradation by other peroxidases. The alcohol dehydrogenase (gene 2768), alcohol dioxygenase (gene 3045), and aldehyde dehydrogenase (aldH) annotated in the DF3-3 genome are thought to be involved in the process of lignin degradation. In addition, catalase removes the hydrogen peroxide produced by these reactions quickly enough to prevent oxidative damage to [4Fe-4S]-clusters in proteins and protect the body from toxification [24, 49]. The catalase-encoding gene (katE) identified in DF3-3 is thought to be involved in lignin degradation.

According to the GC-MS results, 2,4-di-tert-butylphenol and 2,2'-methyl bis(4-methyl-6-tert-butyl phenol) were involved in the metabolic processes of DF3-3. Recent studies have also shown that there is a 2,4-di-tert-butylphenol metabolic pathway for microbial degradation of lignin [55]. Considering the resorcinol pathway and its correlative gene, it is speculated that there may be a similar pathway in DF3-3. It may retain the structure of the tert-butyl side face and be metabolized by the meta-cleavage pathway. The formation of 2,2'-methyl bis(4-methyl-6-tert-butyl phenol) may come from the same metabolic intermediate as 2,4-di-tert-butylphenol. However, there are few studies on microbial degradation of this kind of structure at present. As an environmental pollutant, 4-tert-butylphenol can be degraded by several reported bacteria [56, 57]. However, to better understand this purification process, its specific metabolic process and some of the enzymes involved require further study.

The lignin degradation pathway of actinomycetes was first studied through research on the culture medium and metabolites [31, 58]. As the understanding of molecular biology increased, people began to seek more direct evidence. Masai et al. [23] first established a relatively complete pathway of lignin degradation and metabolism by means of enzymology and genomics. In *Sphingomonas paucimobilis* SYK-6, β -aryl ether cleavage catalysis [59], the biphenyl ring cleavage pathway [60], the ferulate catabolic pathway [61], the O-demethylation systems of vanillate and syringate [62], the protocatechuate 4,5-cleavage pathway [63], and multiple 3-O-methylgallate catabolic pathways [64] were described. Eleven lignin metabolic pathways were found in the genome of *Cupriavidus necator*, among which the β -keto adipate pathway also included four branches: catechol, chlorocatechol, methylcatechol and protocatechuate ortho ring-cleavage [42]. In our study, evidence of a lignin degradation pathway was found in the metabolites and genetics. The products detected by GC-MS are also related to five lignin metabolic pathways. However, some unusual products detected may point to new branches. We speculate that 4-hydroxyphenylpyruvate (14) and 3-phenylpyruvic acid (12) may be transformed from phenylalanine and enter the homologous pathway and β -keto adipate pathway, respectively, for further metabolism. A large number of genes related to the β -keto adipate pathway have been detected in DF3-3, but there are still some genes involved in the reaction process that have not been compared, and for some genes, the reaction process in which they specifically participate has not been further elucidated. Perhaps there are related reactions in DF3-3 which are different from those in other bacteria. The heterologous expression of these gene fragments is helpful to the establishment and production of efficient engineered bacteria with biological enzymes.

Recent studies have shown that bacterial degradation of lignin has complex growth condition-specific regulation [65]. Because of the diversity of structure of lignan compounds, different reactions may occur in the degradation process of different substrate. Based on genome data alone, we predicted two other pathways the phenylacetate-CoA pathway and the 2,3-dihydroxyphenylpropionic acid pathway, but did not find the related metabolites by GC-MS. This absence might be related to the substrates. To further verify the degradation process, transcriptome analysis, proteomic analysis and other biological methods are needed to study the enzymes and the genes involved in their degradation pathways to understand the biological function of DF3-3 in the degradation of lignin.

Conclusions

Based on the above data and analyses, we isolated a bacterial strain identified as *Streptomyces thermocarboxydus* strain DF3-3 from greening litter and concluded that it degraded alkaline lignin, and the degradation efficiency reached 31% within 15 days. In total, 19 alkaline lignin degradation intermediates were identified by GC-MS, and 107 possible lignin-degrading enzyme encoding genes in the DF3-3 genome were annotated; 7 pathways for metabolism of lignin and its intermediates were predicted, including the β -keto adipate pathway and peripheral reactions; gentisate pathway; anthranilate pathway; catabolic pathway for resorcinol; homogentisic pathway; phenylacetate-CoA pathway; and the 2,3-dihydroxyphenylpropionic acid pathway. Intermediates in the first five metabolic pathway were detected by GC-MS. The degradation products and genomics analyses show that DF3-3 has a relatively complete lignin degradation pathway.

Methods

Sampling, isolation and screening of bacterial strain

Samples were collected from greening litter of Beijing University of Agriculture (40.0947° N, 116.3151° E). One gram of sample was placed in a 50-mL sterile centrifuge tube containing 10 mL of sterile distilled water and shaken at

120 rpm overnight. Next, 10^{-1} and 10^{-2} serial dilutions of each sample suspension were spread as 0.1-mL aliquots on Gause's synthetic medium with the formula (g/L): 0.5 NaCl; 1 KNO₃; 0.5 K₂HPO₄·3H₂O; 0.5 MgSO₄·7H₂O; 0.01 FeSO₄·7H₂O; 20 soluble starch [8]. The plates were incubated at 30°C for one week, and distinct colonies were picked and subcultured for further analysis. Gause's guaiacol medium and Gause's Azure B medium used for lignin degradation screening contained 0.1% guaiacol and 0.1% aniline blue, respectively, added to Gause's medium. Different external nitrogen sources (20 g/L), such as acrylamide and potassium nitrate, and additional carbon sources (1 g/L), such as glucose and mannose, were used to replace soluble starch culture strains in studies of their utilization of nitrogen sources and carbon sources. The culture medium for detecting lignin degradation and enzyme activity was kraft lignin-MSM medium (3 g of kraft lignin L, 2 g of [NH₄]₂SO₄, 1 g of K₂HPO₄, 1 g of KH₂PO₄, 0.2 g of MgSO₄, 0.1 g of CaCl₂, 0.05 g of FeSO₄, and 0.02 g of MnSO₄ in 1 L distilled water, pH 7.0).

Scanning electron microscope observations

The shapes of the bacteria were observed by scanning electron microscopy. A cover glass was inserted into the solid medium to cultivate the strain, and the insert was removed after the bacterial body climbed onto the glass slide. The precipitate was washed by adding a phosphate buffer solution (pH 7.2), added to 2.5% glutaraldehyde, fixed at room temperature for 2~4 hours, and then placed in a refrigerator at 4°C overnight. After elution with a 30–95% ethanol gradient, the material was rinsed with tert-butanol, then 20 µL of tert-butanol was added and the mixture was put into a refrigerator at -20°C until it froze and solidified. Using critical point drying (HITACHI HCP-2 Critical Point Dryer) and gold sputter coating (Eiko IB-3 ion plating machine), the sample was observed by scanning electron microscopy (SEM, JSM-6360LV, JEOL, Japan) [66].

Strain growth curve determination

To assess the growth of bacteria, an equal quantity of bacteria was inserted into Gause's liquid medium and cultured on a shaker. The culture solution was removed and centrifuged every 24 hours. The supernatant was discarded, and the filter paper was placed into an oven. The mixture was dried to a constant weight, and the filter paper and the bacteria were weighed. The weight of the filter paper was compared with the weight of the bacteria. All assays were performed with three replicates.

Biodegradation of alkali lignin

To determine the lignin loss from alkaline lignin cause by various strains, samples (1.5 mL) were centrifuged at 12,000 ×g for 10 min. One millilitre of supernatant was diluted by adding 2 mL of phosphate buffer (pH 7.6). The lignin concentration was determined by measuring the absorbance at 280 nm with a UV-Vis spectrophotometer (Shimadzu UN-1900i) [67]. The calculated standard curve for lignin was $y = 0.0786x - 0.0245$, $R^2 = 0.9988$.

Enzyme assay

Samples were centrifuged at 12000 rpm for 5 min, and the supernatant was used for lignin peroxidase (Lip), laccase and manganese peroxidase (MnP) enzyme assays. Laccase activity was determined by monitoring the oxidation of ABTS at 420 nm ($\epsilon_{420} = 36000 \text{ M}^{-1} \text{ cm}^{-1}$) [68]. A lignin peroxidase assay was carried out by using peroxidase oxidation of Azure B. LiP activity was determined by measuring the absorbance at 651 nm ($\epsilon_{651} = 48.8 \text{ M}^{-1} \text{ cm}^{-1}$) [69]. Manganese peroxidase activity was determined from the change in absorbance occurring when Mn²⁺ is oxidized to Mn³⁺ and forms a complex with malonate, which produced absorbance at 270 nm ($\epsilon_{270} = 11590 \text{ M}^{-1} \text{ cm}^{-1}$) [70].

Genome sequencing and functional annotation

The genome of DF3-3 was sequenced at Major Biomedical Technology Co., Ltd. (Shanghai, China). Genomic DNA was extracted using a Wizard® Genomic DNA Purification Kit (Promega). Purified genomic DNA was quantified by a TBS-380 fluorometer (Turner BioSystems Inc., Sunnyvale, CA). The genome was sequenced by adopting the second-generation + third-generation sequencing method of Illumina HiSeq+PacBio, with a shotgun library of 400 bp insertion size. Assembly software canu, SPAdes, etc. was used for three-generation sequence assembly [71], and GeneMarkS software was used to predict the coding sequence (CDS) in the genome [72]. The prediction and annotation of genes were carried out using Prodigal Son (prokaryotic dynamic programming gene discovery algorithm). GeneMarkS was used to predict the plasmid genome. tRNAscan-SE v2.0 software was used to predict the tRNA contained in the genome, and Barrnap software was used to predict the rRNA contained in the genome. Functional annotation of the predicted coding gene was carried out by comparison with 6 major databases (NR, Swiss-Prot, Pfam, EggNOG, GO and KEGG) [73–76].

Alkali lignin degradation products determined by GC-MS

DF3-3 was inoculated in 100 ml of medium with AL as the carbon source and cultured on a shaker for seven days. Samples were collected every 24 hours, and a number of control groups was set up. The sample was centrifuged (10,000 rpm, 15 min) to remove the bacteria, the supernatant was acidified with HCl to pH 2–3, and it was thoroughly extracted with a three-fold volume of ethyl acetate. The extract was rotary evaporated to 10 ml at 37°C and dried with anhydrous Na₂SO₄. After evaporating the solvent in a nitrogen stream, 100 µl of the organic layer was derivatized. Then, 100 µl of dioxane and 10 µl of pyridine were added to the sample and vortexed, and 50 µl of bis(trimethylsilyl)trifluoroacetamide (BSTFA) was added. The mixed solution was placed in a water bath at 80°C for 45 min and shaken regularly. The silanized sample was tested by GC-MS [77].

The analytical column was a DB-5 capillary column (30 m length, 0.25 mm inner diameter, 0.25 mm film thickness). The carrier gas was helium. The column temperature was initially 50°C (5 min), then it was raised to 280°C (10°C/min, holding time of 5 min). The transmission line and ion source temperatures were 200 and 250°C, respectively. The solvent delay time was 4.0 min. The injection volume was 1 µl. Electron ionization mass spectra were recorded in the range 30–550 (m/z) in full scan mode.

Abbreviations

AL: Alkali lignin; GC-MS: Gas chromatography-mass spectrometry; MnP: Manganese peroxidase; Lac: Laccase; LiP: Lignin peroxidase.

Declarations

Fangyun Tan and Jun Cheng are co-first authors.

Ethics approval and consent to participate

Ethical approval and consent to participate are not required.

Consent for publication

All authors agree to the submission and publication of the manuscript in the journal *Biotechnology for Biofuel*.

Availability of data and materials

The additional data generated during this study are available in the Additional file.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the National Key Research and Development Program of China (code: 2017YFF0207800) and the Scientific Research Program of Beijing Municipal Education Commission (code: KM201810020012).

Authors' contributions

YQL designed the experimental strategy. FYT, JC, YZ and XFJ carried out the experiments and performed the bioinformatics data analysis. FYT and JC wrote the manuscript. YQL supervised the overall research. This manuscript was proofread by all the authors. All authors read and approved the final manuscript.

Acknowledgements

The authors thank Beijing Key Laboratory of New Technology in Agricultural Application for their support of the part of the work involving scanning electron microscopy. And genome analysis was performed using the free online platform of Majorbio Cloud Platform (www.majorbio.com).

Author information

¹School of Landscape Architecture, Beijing University of Agriculture, Beijing 102206 PR China; ²School of Bioscience and Resource Environment, Beijing University of Agriculture, Beijing 102206 PR China; ³Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193 PR China.

References

1. Zeng Y, Zhao S, Yang S, Ding SY. Lignin plays a negative role in the biochemical process for producing lignocellulosic biofuels. *Curr Opin Biotechnol*. 2014;27:38-45.
2. Kumar M, Singhal A, Thakur IS. Comparison of submerged and solid state pretreatment of sugarcane bagasse by *Pandora* sp. ISTKB: enzymatic and structural analysis. *Bioresour Technol*. 2016;203:18-25.
3. Kamimura N, Sakamoto S, Mitsuda N, Masai E, Kajita S. Advances in microbial lignin degradation and its applications. *Curr Opin Biotechnol*. 2019;56:179-86.
4. Ralph J, Lundquist K, Brunow G, Lu F, Kim H, Schatz PF, et al. Lignins: natural polymers from oxidative coupling of 4-hydroxyphenyl- propanoids. *Phytochem Rev*. 2004;3:29-60.
5. Abdelaziz OY, Brink DP, Prothmann J, Ravi K, Sun M, García-Hidalgo J, et al. Biological valorization of low molecular weight lignin. *Biotechnol Adv*. 2016;34:1318-46.
6. Tuor U, Winterhalter K, Fiechter A. Enzymes of white-rot fungi involved in lignin degradation and ecological determinants for wood decay. *J Biotechnol*. 1995;41:1-17.
7. Sánchez C. Lignocellulosic residues: biodegradation and bioconversion by fungi. *Biotechnol Adv*. 2009;27:185-94.

8. Yang YS, Zhou JT, Lu H, Yuan YL, Zhao LH. Isolation and characterization of *Streptomyces* spp. strains F-6 and F-7 capable of decomposing alkali lignin. *Environ Technol.* 2012;33:2603-9.
9. Mcleod MP, Warren RL, Hsiao W, Araki N, Myhre M, Fernandes C, et al. The complete genome of *Rhodococcus* sp. RHA1 provides insights into a catabolic powerhouse. *Proc Natl Acad Sci U S A.* 2006;103:15582-7.
10. Masai E, Kamimura N, Kasai D, Oguchi A, Ankai A, Fukui S, et al. Complete genome sequence of *Sphingobium* sp. strain SYK-6, a degrader of lignin-derived biaryls and monoaryls. *J Bacteriol.* 2012;194:534-5.
11. Lin L, Cheng Y, Pu Y, Sun S, Li X, Jin M, et al. Systems biology-guided biodesign of consolidated lignin conversion. *Green Chem.* 2016;18:5536-47.
12. Janshekar H, Fiechter A. On the bacterial degradation of lignin. *Eur J Appl Microbiol Biotechnol.* 1982;14:47-50.
13. Bugg TD, Ahmad M, Hardiman EM, Singh R. The emerging role for bacteria in lignin degradation and bio-product formation. *Curr Opin Biotechnol.* 2011;22:394-400.
14. Pasti MB, Pometto AL, Nuti MP, Crawford DL. Lignin-solubilizing ability of actinomycetes isolated from termite (*Termitidae*) gut. *Appl Environ Microbiol.* 1990;56:2213-8.
15. Vishwanatha BT, Babu G, Malagi SB, Dandin CJ, Nayaka S. Isolation, identification and characterization of streptomyces sp. SN-2. *Biosci Biotechnol Res Asia.* 2017;14:1401-7.
16. Feng H, Sun Y, Zhi Y, Mao L, Luo Y, Wei X, et al. Lignocellulose degradation by the isolate of *Streptomyces griseorubens* JSD-1. *Funct Integr Genom.* 2015;15:163-73.
17. Feng H, Zhi Y, Shi W, Mao L, Zhou P. Isolation, identification and characterization of a straw degrading *Streptomyces griseorubens* JSD-1. *Afr J Microbiol Res.* 2013;7:2730-5.
18. Lambertz C, Ece S, Fischer R, Commandeur U. Progress and obstacles in the production and application of recombinant lignin-degrading peroxidases. *Bioengineered.* 2016;7:145-54.
19. Cragg SM, Beckham GT, Bruce NC, Bugg TDH, Distel DL, Dupree P, et al. Lignocellulose degradation mechanisms across the Tree of Life. *Curr Opin Chem Biol.* 2015;29:108-19.
20. Bugg TDH, Ahmad M, Hardiman EM, Rahmanpour R. Pathways for degradation of lignin in bacteria and fungi. *Nat Prod Rep.* 2011;28:1883-96.
21. Zakzeski J, Bruijninx PC, Jongerius AL, Weckhuysen BM. The catalytic valorization of lignin for the production of renewable chemicals. *Chem Rev.* 2010;110:3552-99.
22. Lin L, Wang X, Cao L, Xu M. Lignin catabolic pathways reveal unique characteristics of dye-decolorizing peroxidases in *Pseudomonas putida*. *Environ Microbiol.* 2019;21:1847-63.
23. Masai E, Katayama Y, Fukuda M. Genetic and biochemical investigations on bacterial catabolic pathways for lignin-derived aromatic compounds. *J Agric Chem Soc Jpn.* 2007;71:1-15.
24. Niewerth H, Schuldes JR, Parschat K, Kiefer P, Vorholt JA, Daniel R, et al. Complete genome sequence and metabolic potential of the quinaldine-degrading bacterium *Arthrobacter* sp. Rue61a. *BMC Genom.* 2012;13:534.
25. Ramachandra M, Crawford DL, Hertel G. Characterization of an extracellular lignin peroxidase of the lignocellulolytic actinomycete *Streptomyces viridosporus*. *Appl Environ Microbiol.* 1988;54:3057-63.
26. Antai SP, Crawford DL. Degradation of softwood, hardwood, and grass lignocelluloses by two streptomyces strains. *Appl Environ Microbiol.* 1981;42:378-80.
27. Chakar FS, Ragauskas AJ. Review of current and future softwood kraft lignin process chemistry. *Ind Crops Prod.* 2004;20:131-41.
28. Lei C, Junfeng Y, Liping Y, Keqin Z. Several qualitative methods for the screening of fungi to decompose lignin. *Wei Sheng wu xue Tong bao.* 2002;29:67-9.

29. Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun.* 2018;9:5114.
30. Zhao M, Zhang C, Zeng G, Huang D, Xu P, Cheng M. Growth, metabolism of *Phanerochaete chrysosporium* and route of lignin degradation in response to cadmium stress in solid-state fermentation. *Chemosphere.* 2015;138:560-7.
31. Crawford DL, Pometto AL, Crawford RL. Lignin degradation by *Streptomyces viridosporus*: isolation and characterization of a new polymeric lignin degradation intermediate. *Appl Environ Microbiol.* 1983;45:898-904.
32. Kumar M, Mishra A, Singh SS, Srivastava S, Thakur IS. Expression and characterization of novel laccase gene from *Pandoraea* sp. ISTKB and its application. *Int J Biol Macromol.* 2018;115:308-16.
33. Pérez J, Muñoz-Dorado J, De la Rubia T, Martínez J. Biodegradation and biological treatments of cellulose, hemicellulose and lignin: an overview. *Int Microbiol.* 2002;5:53-63.
34. de Gonzalo G, Colpa DI, Habib MHM, Fraaije MW. Bacterial enzymes involved in lignin degradation. *J Biotechnol.* 2016;236:110-9.
35. Wu J, Kim KS, Lee JH, Lee YC. Cloning, expression in *Escherichia coli*, and enzymatic properties of laccase from *Aeromonas hydrophila* WL-11. *J Environ Sci.* 2010;22:635-40.
36. Zhu D, Zhang P, Xie C, Zhang W, Sun J, Qian WJ, et al. Biodegradation of alkaline lignin by *Bacillus ligniniphilus* L1. *Biotechnol Biofuels.* 2017;10:44.
37. Chen YH, Chai LY, Zhu YH, Yang ZH, Zheng Y, Zhang H. Biodegradation of kraft lignin by a bacterial strain *Comamonas* sp. B-9 isolated from eroded bamboo slips. *J Appl Microbiol.* 2012;112:900-6.
38. Raj A, Reddy MMK, Chandra R, Purohit HJ, Kapley A. Biodegradation of kraft-lignin by *Bacillus* sp. isolated from sludge of pulp and paper mill. *Biodegradation.* 2007;18:783-92.
39. Ksibi M, Amor SB, Cherif S, Elaloui E, Houas A, Elaloui M. Photodegradation of lignin from black liquor using a UV/TiO₂ system. *J Photochem Photobiol A Chem.* 2003;154:211-8.
40. Shin KS, Lee YJ. Depolymerisation of lignosulfonate by peroxidase of the white-rot basidiomycete, *Pleurotus ostreatus*. *Biotechnol Lett.* 1999;21:585-8.
41. Xiong YI, Zhao Y, Ni K, Shi Y, Xu Q. Characterization of ligninolytic bacteria and analysis of alkali-lignin biodegradation products. *Pol J Microbiol.* 2020;69:339-47.
42. Pérez-Pantoja D, De la Iglesia R, Pieper DH, González B. Metabolic reconstruction of aromatic compounds degradation from the genome of the amazing pollutant-degrading bacterium *Cupriavidus necator* JMP134. *FEMS Microbiol Rev.* 2008;32:736-94.
43. Xu G, Li F, Wang Q. Occurrence and degradation characteristics of dibutyl phthalate (DBP) and di-(2-ethylhexyl) phthalate (DEHP) in typical agricultural soils of China. *Sci Total Environ.* 2008;393:333-40.
44. Schühle K, Jahn M, Ghisla S, Fuchs G. Two similar gene clusters coding for enzymes of a new type of aerobic 2-aminobenzoate (anthranilate) metabolism in the bacterium *Azoarcus evansii*. *J Bacteriol.* 2001;183:5268-78.
45. Liu D, Yan X, Si M, Deng X, Min X, Shi Y, et al. Bioconversion of lignin into bioplastics by *Pandoraea* sp. B-6: molecular mechanism. *Environ Sci Pollut Res Int.* 2019;26:2761-70.
46. Mohamed Mel S, Ismail W, Heider J, Fuchs G. Aerobic metabolism of phenylacetic acids in *Azoarcus evansii*. *Arch Microbiol.* 2002;178:180-92.
47. Ferrández A, Miñambres B, García B, Olivera ER, Luengo JM, García JL, et al. Catabolism of phenylacetic acid in *Escherichia coli*. Characterization of a new aerobic hybrid pathway. *J Biol Chem.* 1998;273:25974-86.

48. Brown ME, Chang MCY. Exploring bacterial lignin degradation. *Current Opinion in Chemical Biology*. 2014;19:1-7.
49. Masai E, Katayama Y, Kubota S, Kawai S, Yamasaki M, Morohoshi N. A bacterial enzyme degrading the model lignin compound beta-etherase is a member of the glutathione-S-transferase superfamily. *FEBS Lett*. 1993;323:135-40.
50. Masai E, Kubota S, Katayama Y, Kawai S, Yamasaki M, Morohoshi N. Characterization of the C alpha-dehydrogenase gene involved in the cleavage of beta-aryl ether by *Pseudomonas paucimobilis*. *Biosci Biotechnol Biochem*. 1993;57:1655-9.
51. Colpa DI, Fraaije MW, van Bloois E. DyP-type peroxidases: a promising and versatile class of enzymes. *J Ind Microbiol Biotechnol*. 2014;41:1-7.
52. Majumdar S, Lukk T, Solbiati JO, Bauer S, Nair SK, Cronan JE, et al. Roles of small laccases from *Streptomyces* in lignin degradation. *Biochemistry*. 2014;53:4047-58.
53. Ihssen J, Reiss R, Luchsinger R, Thöny-Meyer L, Richter M. Biochemical properties and yields of diverse bacterial laccase-like multicopper oxidases expressed in *Escherichia coli*. *Sci Rep*. 2015;5:10465.
54. Medema MH, Trefzer A, Kovalchuk A, van den Berg M, Müller U, Heijne W, et al. The sequence of a 1.8-mb bacterial linear plasmid reveals a rich evolutionary reservoir of secondary metabolic pathways. *Genome Biol Evol*. 2010;2:212-24.
55. Toyama T, Momotani N, Ogata Y, Miyamori Y, Inoue D, Sei K, et al. Isolation and characterization of 4-tert-butylphenol-utilizing *Sphingobium fuliginis* strains from *Phragmites australis* rhizosphere sediment. *Appl Environ Microbiol*. 2010;76:6733-40.
56. Gabriel FL, Giger W, Guenther K, Kohler HPE. Differential degradation of nonylphenol isomers by *Sphingomonas xenophaga* Bayram. *Appl Environ Microbiol*. 2005;71:1123-9.
57. Gabriel FL, Cyris M, Jonkers N, Giger W, Guenther K, Kohler HPE. Elucidation of the ipso-substitution mechanism for side-chain cleavage of α -quaternary 4-nonylphenols and 4-t-butoxyphenol in *Sphingobium xenophagum* Bayram. *Appl Environ Microbiol*. 2007;73:3320-6.
58. Crawford DL, Pometto AL, 3rd, Crawford RL. Production of useful modified lignin polymers by bioconversion of lignocellulose with *Streptomyces*. *Biotechnol Adv*. 1984;2:217-32.
59. Masai E, Katayama Y, Nishikawa S, Yamasaki M, Morohoshi N, Haraguchi T. Detection and localization of a new enzyme catalyzing the beta-aryl ether cleavage in the soil bacterium (*Pseudomonas paucimobilis* SYK-6). *FEBS Lett*. 1989;249:348-52.
60. Sonoki T, Obi T, Kubota S, Higashi M, Masai E, Katayama Y. Coexistence of two different O demethylation systems in lignin metabolism by *Sphingomonas paucimobilis* SYK-6: cloning and sequencing of the lignin biphenyl-specific O-demethylase (LigX) gene. *Appl Environ Microbiol*. 2000;66:2125-32.
61. Masai E, Harada K, Peng X, Kitayama H, Katayama Y, Fukuda M. Cloning and characterization of the ferulic acid catabolic genes of *Sphingomonas paucimobilis* SYK-6. *Appl Environ Microbiol*. 2002;68:4416-24.
62. Masai E, Sasaki M, Minakawa Y, Abe T, Sonoki T, Miyauchi K, et al. A novel tetrahydrofolate-dependent O-demethylase gene is essential for growth of *Sphingomonas paucimobilis* SYK-6 with syringate. *J Bacteriol*. 2004;186:2757-65.
63. Kersten PJ, Dagley S, Whittaker JW, Arciero DM, Lipscomb JD. 2-pyrone-4,6-dicarboxylic acid, a catabolite of gallic acids in *Pseudomonas* species. *J Bacteriol*. 1982;152:1154-62.

64. Kasai D, Masai E, Miyauchi K, Katayama Y, Fukuda M. Characterization of the 3-O-methylgallate dioxygenase gene and evidence of multiple 3-O-methylgallate catabolic pathways in *Sphingomonas paucimobilis* SYK-6. *J Bacteriol.* 2004;186:4951-9.
65. Wöhlbrand L, Kallerhoff B, Lange D, Hufnagel P, Thiermann J, Reinhardt R, et al. Functional proteomic view of metabolic regulation in "Aromatoleum aromaticum" strain EbN1. *Proteomics.* 2007;7:2222-39.
66. Castillo U, Myers S, Browne L, Strobel G, Hess WM, Hanks J, et al. Scanning electron microscopy of some endophytic streptomyces in snakevine- *Kennedia nigricans*. *Scanning.* 2005;27:305-11.
67. Lara MA, Rodríguez-Malaver AJ, Rojas OJ, Holmquist O, González AM, Bullón J, et al. Black liquor lignin biodegradation by *Trametes elegans*. *Int Biodeterior Biodegrad.* 2003;52:167-73.
68. Kinnunen A, Majjala P, JArvinen P, Hatakka A. Improved efficiency in screening for lignin-modifying peroxidases and laccases of basidiomycetes. *Curr Biotechnol.* 2017;6:105-15.
69. Morya R, Kumar M, Thakur IS. Utilization of glycerol by *Bacillus* sp. ISTVK1 for production and characterization of polyhydroxyvalerate. *Bioresour Technol Rep.* 2018;2:1-6.
70. Kumar M, Singh J, Singh MK, Singhal A, Thakur IS. Investigating the degradation process of kraft lignin by β -proteobacterium, *Pandoraea* sp. ISTKB. *Environ Sci Pollut Res.* 2015;22:15690-702.
71. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 2012;19:455-77.
72. Besemer J, Borodovsky M. GeneMark: web software for gene finding in prokaryotes, eukaryotes and viruses. *Nucleic Acids Res.* 2005;33:W451-4.
73. Bairoch A, Apweiler R. The SWISS-PROT protein sequence database and its supplement TrEMBL in 2000. *Nucleic Acids Res.* 2000;28:45-8.
74. Jensen LJ, Julien P, Kuhn M, von Mering C, Muller J, Doerks T, et al. eggNOG: automated construction and annotation of orthologous groups of genes. *Nucleic Acids Res.* 2008;36:D250-4.
75. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 2000;28:27-30.
76. Finn RD, Alex B, Jody C, Penelope C, Eberhardt RY, Eddy SR, et al. Pfam: the protein families database. *Nucleic Acids Research.* 2014;42:D222-30.
77. Raj A, Chandra R, Reddy MMK, Purohit HJ, Kapley A. Biodegradation of kraft lignin by a newly isolated bacterial strain, *Aneurinibacillus aneurinilyticus* from the sludge of a pulp paper mill. *World J Microbiol Biotechnol.* 2007;23:793-9.

Figures




Figure 1

DF3-3 colour reaction and morphological characteristics. a Colony morphology streaked on a plate of DF3-3. b Transparent fading circle on Gause's- Azure B medium. c Colour reaction on Gause's-guaiacol medium. d Scanning electron micrograph of DF3-3.




Figure 2

ANI analysis results for DF3-3 and other bacteria.




Figure 3

Growth curve and alkaline lignin degradation curve for DF3-3. Average values of three replicates are shown with the standard error of the mean as error bars (data are contained in S1).




Figure 4

Enzyme activities of LiP, Lac, and MnP of DF3-3 during 11 days of incubation. Average values of three replicates are shown with the standard errors of the mean shown as error bars.




Figure 5

Variations in peak areas for products identified by GC-MS before TAC




Figure 6

Variations in the peak areas for products identified by GC-MS after TAC.




Figure 7

Predicted lignin degradation pathway for *Streptomyces thermocarboxydus* strain DF3-3. Numbers used to represent the compounds are the same as those in Table 3.




Figure 8

Lignin metabolic pathway of DF3-3 to be further verified.




Figure 9

Locations of genes encoding aromatic catabolic pathways are indicated in the genome of DF3-3; orange, genes for the β -ketoadipate central pathway; green, genes for the phenylacetyl-CoA ring-cleavage pathway; yellow, genes for the 2,3-dihydroxyphenylpropionate pathway; purple, genes for the anthranilate pathway; dark green, genes for the homogentisate pathway; dark blue, genes for the gentisate pathway; blue, genes for the resorcinol pathway.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementarymaterial9.23.docx](#)